

2021 KSCCM clinical practice guidelines for pain, agitation, delirium, immobility, and sleep disturbance in the intensive care unit

Yijun Seo¹, Hak-Jae Lee², Eun Jin Ha³, Tae Sun Ha⁴

¹Department of Anesthesiology and Pain Medicine, and Anesthesia and Pain Research Institute, Yonsei University College of Medicine, Seoul, Korea

²Division of Acute Care Surgery, Department of Surgery, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

³Department of Critical Care Medicine, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, Korea

⁴Department of Surgery, Soonchunhyang University Bucheon Hospital, Soonchunhyang University College of Medicine, Bucheon, Korea

Acute and Critical Care 2022;37:1-25
<https://doi.org/10.4266/acc.2022.00094>

In the article entitled “2021 KSCCM clinical practice guidelines for pain, agitation, delirium, immobility, and sleep disturbance in the intensive care unit,” Figure 4 was incorrectly presented. The correct figure is as follows.




Figure 4. Delirium assessment tool: the Confusion Assessment Method for the intensive care unit (CAM-ICU) flowchart. RASS: Richmond Agitation-Sedation Scale.

Corrigendum

Received: February 22, 2023

Revised: February 22, 2023

Accepted: February 22, 2023

Corresponding author

Tae Sun Ha
Department of Surgery,
Soonchunhyang University Bucheon
Hospital, 170 Jomaru-ro, Wonmi-gu,
Bucheon 14584, Korea
Tel: +82-32-621-6265
Fax: +82-32-621-5016
Email: applips@schmc.ac.kr

Copyright © 2023 The Korean Society of Critical Care Medicine

This is an Open Access article distributed under the terms of Creative Attributions Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.