



Published in final edited form as:

*Obesity (Silver Spring)*. 2023 April ; 31(4): 912–922. doi:10.1002/oby.23680.

## Sleep duration and timing are prospectively linked with insulin resistance during late adolescence

Erica C. Jansen, PhD, MPH<sup>a,b,\*</sup> [Research Assistant Professor of Nutritional Sciences], Helen J. Burgess, PhD<sup>c</sup> [Professor of Psychiatry], Ronald D. Chervin, MD, MS<sup>b</sup> [Professor of Neurology and Michael S. Aldrich Collegiate Professor of Sleep Medicine], Dana Dolinoy, PhD<sup>a,d</sup> [Professor and Chair of Environmental Health Sciences and Professor of Nutritional Sciences], Martha María Téllez-Rojo, ScD<sup>e</sup> [Professor], Alejandra Cantoral, ScD<sup>f</sup> [Assistant Professor], Libni Olascoaga-Torres, MS<sup>e</sup> [Research Coordinator], Joyce Lee, MD, MPH<sup>g</sup> [Research Professor of Pediatrics], Galit Levi Dunietz, PhD, MPH<sup>a,b</sup> [Associate Professor of Neurology], Louise M. O'Brien, PhD, MS<sup>b</sup> [Associate Professor of Neurology], Karen E. Peterson, ScD<sup>a,d</sup> [Professor and Chair of Nutritional Sciences, Professor of Global Public Health]

<sup>a</sup>Sleep Disorders Center and Department of Neurology, University of Michigan, Ann Arbor, MI

<sup>b</sup>Department of Nutritional Sciences, University of Michigan School of Public Health, Ann Arbor, MI

<sup>c</sup>Department of Psychiatry, University of Michigan, Ann Arbor, MI

<sup>d</sup>Department of Environmental Health Sciences, University of Michigan School of Public Health, Ann Arbor, MI

<sup>e</sup>Center for Research on Nutrition and Health, National Institute of Public Health, Cuernavaca, Mexico

<sup>f</sup>Department of Health, Iberoamerican University, Mexico City, Mexico

<sup>g</sup>Department of Pediatrics, University of Michigan, Ann Arbor, MI

### Abstract

**Objective:** To evaluate whether short sleep duration or later sleep timing are risk factors for insulin resistance (IR) in late adolescence.

**Methods:** Mexico City adolescents enrolled in a longitudinal birth cohort (ELEMENT) took part in 2 study visits during peri-puberty that occurred approximately 2 years apart. IR was assessed with serum glucose and insulin. Four groups were defined using puberty-specific cutpoints: no

\*Corresponding author: Erica C. Jansen; 3863 SPH I, University of Michigan School of Public Health, 1415 Washington Heights, Ann Arbor, MI 48109- janerica@umich.edu.

**Disclosures:** Dr. Chervin reports patents and copyrighted material for the assessment and treatment of patients with obstructive sleep apnea, has received consulting fees from Eli Lilly and Company, and acts as treasurer and board member for the International Pediatric Sleep Association and as an Advisory Board member for the non-profit Pajama Program. Dr. Lee reports support from the NIH (P30GRANT12959224) and the Elizabeth Weiser Caswell Diabetes Institute, has received consulting fees from Tandem Diabetes Care, and is on the Medical Advisory Board for GoodRx. Dr. Burgess serves on the scientific advisory board for Natrol, LLC, and is a consultant for F. Hoffmann-La Roche Ltd. None of these disclosures played a role in the analysis, interpretation, decision to publish, or preparation of this manuscript. None of the other authors has any disclosures to make.

IR over the follow-up period, transition from normal to IR, transition from IR to normal, and IR at both timepoints. Baseline sleep assessments were measured with 7-day wrist actigraphy. Multinomial logistic regression models were used to evaluate associations between sleep duration and timing with HOMA-IR categories, adjusting for age, sex, and baseline pubertal status.

**Results:** Adolescents 1 hour below the sleep duration recommendations-for-age were 2.74 times more likely to develop IR (95% CI [1.0, 7.4]). Similarly, adolescents who were in the latest category of sleep midpoint (>4:33 AM) were more likely than those with earliest midpoints (1 to 3 AM) to develop IR (OR=2.63, [1.0, 6.7]). Changes in adiposity over follow-up did not mediate sleep and IR.

**Conclusions:** Insufficient sleep duration and late sleep timing were associated with development of IR over a 2-year period in late adolescence.

### Keywords

actigraphy; midpoint; insufficient sleep; diabetes; puberty

---

### Introduction

Type 2 diabetes (T2D), characterized by insulin resistance (IR), is one of the leading causes of death and morbidity worldwide (1, 2), and Mexico is no exception. The prevalence of diabetes in the Mexican adult population is 14%, and in 2016 the disease was declared a public health emergency (3, 4). While the disease typically is detected during adulthood, the origins may be traced back to earlier life. Indeed, literature on the Developmental Origins of Health and Disease (DOHaD) has revealed that gestational diabetes and excessive weight gain during pregnancy can predispose offspring to the development of T2D in adulthood (5)(6). During childhood, unhealthy dietary patterns(7) and higher body size(8) have been related to later T2D.

The late pubertal and emerging adulthood stage may be another sensitive period for development of T2D later in life. During puberty, adolescents experience transient IR that typically normalizes upon full maturation (9). Specifically, insulin resistance begins to increase early in puberty (Tanner stage 2, on the pubertal Tanner scale that delineates the stages of puberty from 1 (pre-pubertal) to 5 (fully mature)), reaches a peak in mid-puberty (Tanner stage 3 and 4) and declines again by full maturation (Tanner stage 5)(10). Mechanisms remain unclear but could involve altered fuel metabolism meant to preserve lean muscle mass and maximize fat as a fuel source(11, 12). Of relevance for adolescent health, IR may continue to remain high or even rise for some adolescents (9). Although persistent post-pubertal IR has not been studied extensively in large-scale longitudinal studies, prior work suggests that weight status is one important indicator (13). Specifically, adolescents who enter puberty with obesity were more likely to remain insulin resistant and to develop other metabolic complications(9). Other lifestyle factors including diet (e.g., high glycemic index diets(14)) and low physical activity during puberty have also been related to diabetes and cardiovascular disease risk in adulthood (vegetable consumption and physical activity (15)). In addition to diet and physical activity, poor sleep is emerging as an important risk factor for IR during adolescence. Short sleep duration, delayed sleep timing,

and poor sleep quality have each been related to higher IR among adolescents, primarily in cross-sectional investigations (16–20). In Mexican adolescents who were 14 years of age on average, we recently showed that both delayed and short sleep duration were associated cross-sectionally with higher IR, an association that was stronger in girls (21). Variability in sleep duration and timing has also been related to other markers of metabolic health among adolescents (e.g. triglycerides(22) and adiposity(23)), although associations with IR are unclear. Furthermore, the influence of sleep on IR has been rarely examined in longitudinal studies, i.e., whether sleep predicts persistent or emerging IR in the late adolescence period.

Thus, the aim of our study was to assess whether sleep characteristics, including short sleep duration, sleep duration variability, later sleep timing, and social jetlag (>2 hour difference between weekend and weekday sleep timing) were associated with persistent or increasing IR in late adolescence in a Mexican cohort. Further, given that we found some evidence of mediation by higher adiposity (BMI z scores) in our cross-sectional study, we sought to evaluate the extent to which changes in adiposity over the follow-up period mediated relationships between sleep characteristics and IR.

## Methods

### Study population

The study sample included adolescent participants from 2 of 3 sequentially-enrolled cohorts of the Early Life Exposure in Mexico to ENvironmental Toxicants (ELEMENT) study (24). Between 1997 and 2004, 1012 mother/child dyads were recruited from prenatal clinics of the Mexican Social Security Institute in Mexico City, which serves low- to middle-income populations formally employed in the private sector. At baseline, mothers reported sociodemographic and health information. In 2015, a subset of 554 adolescents from the original birth cohorts 2 and 3 between the ages of 9 and 17 years were selected to participate in a follow-up visit (called time 1 (T1) in the present study). Participants from cohort 1 were not included in this follow-up visit since they were older and most were post-pubertal. Starting in 2017, 519 (94%) of the same adolescents participated in an additional follow-up visit that was identical to the 2015 visit when they were between 11 and 19 years (time 2 (T2)). Among these, there were 362 adolescents who had wrist actigraphy data at T1 as well as insulin and glucose measurements at both T1 and T2 (See Figure S1 for flowchart). Compared to the full sample of 519 adolescents, the analytic sample was younger (–2.4 years with 95% CI –2.7, –2.0;  $P<0.0001$ ) and reported less screen time (–1.4 hours/day with 95% CI –2.1, –0.8;  $P<0.0001$ ). The institutional review boards at the Mexico National Institute of Public Health and the University of Michigan approved the research protocols (CI 599 and HUM00034344). Informed consent was obtained from participants aged 18 years. For those aged <18 years, both participant assent and parental informed consent were obtained.

### Sleep Measures

During the in-person follow-up visits, adolescents were given an ActiGraph GT3X-BT; (ActiGraph LLC, Pensacola, FL) to wear on their non-dominant wrist continuously for 7 days and provided with a sleep diary to record bedtimes and wake times. For this

analysis, we used sleep data from T1 participants who provided 4 days of valid wear days (a valid wear day was defined as <4 hours of non-wear time); of those who agreed to wear the actigraph at this time, 96% wore it for 7 days. Nightly sleep measures were estimated from the actigraphy data with 60-second epoch lengths using a pruned dynamic programming (PDP) algorithm developed in R (R Foundation for Statistical Computing, Vienna, Austria)(25). This method has been evaluated against polysomnography, the gold standard for sleep assessment, and compared to manual detection of sleep/wake times in the present cohort with >95% correlation (25). The algorithm incorporated the self-reported bedtimes and wake times to improve accuracy and was used to calculate weekday (Sunday through Thursday night) and weekend (Friday and Saturday) sleep duration (minutes), and weekday and weekend sleep midpoint (the midpoint of sleep onset and wake time; reported in decimal hours). Naps were not included as one of the sleep parameters since the algorithm could not differentiate between naps and daytime sedentary behavior. The primary sleep exposures included weekday sleep duration and weekday sleep midpoint at T1 (i.e., averaged across the weekdays), variability in sleep duration (represented by the standard deviation of the 7 days) and the difference between weekend sleep midpoint and weekday sleep midpoint, a marker of circadian misalignment known as social jetlag (26). Sleep duration was categorized as: meeting the age-specific duration recommendations defined by the American Academy of Sleep Medicine(27), being within 1 hour below the recommendation, or 1 hour below the recommendation. Sleep duration variability was categorized as <1-hour SD, 1 to 2-hour SD, or >2-hour SD. Weekday sleep midpoint was divided evenly into tertiles (as there are currently no specific guidelines for optimal sleep midpoints for adolescents), and social jetlag was divided into: a difference of 1 hour from weekend to weekdays, >1 hour later to 2 hours later on the weekend, or >2 hours later on the weekend.

## HOMA-IR

At both T1 and T2, venous whole blood was taken during the clinic visit after an overnight fast (at least 8 hours), and immediately separated and stored at  $-80^{\circ}\text{C}$ . Serum glucose and insulin were measured at the Michigan Diabetes Research Center (P30DK020572). The insulin assay is a double-antibody radioimmunoassay (RIA) that uses  $^{125}\text{I}$ -Human insulin tracer (Millipore Sigma), a guinea pig anti-porcine insulin first antibody (MDRTC, 68.5% cross-reaction to human proinsulin), and a goat anti-guinea pig gamma globulin (Antibodies Inc.)-PEG second antibody. It was standardized against the Human Insulin International Reference Preparation (NIBSC) and run in duplicate. The glucose assay was run on our automated chemistry analyzer (Randox) in singlicate. HOMA-IR (Homeostatic Model Assessment of Insulin Resistance) was calculated from the product of the fasting serum insulin and fasting serum glucose divided by 405 (28). A cutpoint developed in Chilean adolescents that accounted for pubertal stage was used to define IR (29). A HOMA-IR of >2.1 is considered insulin resistant among adolescents with Tanner stages 1 and 2 (Tanner breast development for girls and Tanner genital development for boys), while HOMA-IR>3.3 is considered insulin resistant among adolescents in Tanner stage 3 or higher. Tanner staging was assessed by a trained clinician during a physical examination (30). Four groups were defined using the cutpoints: normal HOMA-IR over the follow-up

period, transition from normal to IR, transition from IR to normal, and IR at both time points.

### Covariates

Possible baseline confounders were selected *a priori* and included sex, age, pubertal status, maternal education, physical activity, total energy intake, screen time, and smoking status (ever/never). For descriptive statistics, age at T1 was divided into 9–12 years and 13 years in accordance with the AASM sleep recommendation age cut points (27). In addition to the Tanner staging (described above), pubertal status was assessed through questions about menstruation (girls), and estimation of testicular volume in boys (30). Since the IR categories already incorporated adjustment for Tanner stages, we examined other indicators of puberty as potential residual confounders, given that not all pubertal indicators are always aligned within individual and that there may be measurement error. Specifically, pubertal status was considered dichotomously as either being in late or post- puberty versus earlier puberty based on menarche and testicular volume. For girls, late or post-puberty was considered as already having experienced menarche. For boys, late or post-puberty was considered a physician-assessed testicular volume  $\geq 15$  mL. Intensity and duration of physical activity was obtained from the actigraphs and classified as moderate or vigorous, and as minutes/day (31). Total energy intake was estimated by a food frequency questionnaire regarding the previous week's intake as described in detail elsewhere (32). Total screen time was assessed with a questionnaire adapted for and validated in Mexican adolescents (33), and was divided into quartiles. Smoking behavior was self-report with a single question "Have you ever tried smoking?" and was categorized dichotomously. Maternal education was reported by mothers at the original cohort enrollment visit and was classified into 4 categories: <9 years (attended primary or secondary school only), 9–11 years (completed secondary or completed secondary and at least some high school), 12 years (completed high school), and >12 years (post-high school).

Potential mediators of fat mass or distribution included changes in BMI, body fat percentage, waist circumference and triceps skinfolds, measured at T1 and T2 by trained research assistants using standard protocols described previously (34). Duplicate height was measured to the nearest 0.1 cm (Tonelli E120 A), weight to the nearest 0.1 kg (InBody270), waist circumference to the nearest 0.1 cm (QM2000; QuickMedical), triceps skinfold in mm (Lange calipers, Beta Technology), and body fat percentage with bioelectrical impedance (InBody270). All height measures were averaged and BMI was computed as  $\text{kg/m}^2$ . Changes in BMI and other adiposity measures were computed by subtracting the T2 measure from T1.

### Statistical Analysis

To assess potential confounding, we examined associations between covariates and HOMA-IR outcomes by computing the percentages of adolescents in each category of change in HOMA-IR across covariates. Next, bivariate associations between sleep variables and HOMA-IR outcomes were evaluated by estimating the percentages of adolescents in each HOMA-IR category according to sleep variables. P-values in all categorical analyses were obtained with Chi Square Tests. In multivariable complete-case analysis, multinomial

logistic regression models were run with HOMA-IR change categories as the outcome (“normal HOMA-IR at T1 and T2” was the reference group) and categorical T1 sleep characteristics as the exposure, adjusting for age, sex, and pubertal status. Additional adjustment for physical activity and total energy intake did not alter the estimates and thus these variables were not retained. Due to a positive correlation ( $r=0.46$ ), sleep duration and sleep midpoint were run in the same model whereas sleep duration variability and social jetlag were run in separate models (the latter were not strongly correlated). P-values for trend (P, trend) were obtained by running the exact same models but with a variable representing ordinal categories of each sleep category. Age and sex were evaluated as possible effect modifiers by including interaction terms for these variables along with each sleep measure. To investigate mediation of sleep and IR by adiposity changes, we fitted linear regression models with change in waist circumference, change in BMI, change in body fat percentage, and change in triceps skinfolds as the outcomes and sleep characteristics as predictors. A formal mediation model based on the methods of Vanderweele (35, 36) was run on a subsample, where IR was considered as a dichotomous outcome (1 for became insulin resistant at T2 among those who did not have IR at T1, and 0 for those who were never insulin resistant). All analyses were conducted in Stata 14.0, and a P-value<0.05 was considered statistically significant.

## Results

At the T1 visit, the average age was  $14.4 \pm 2.1$  years; 79% of the sample had reached the latter stages of puberty (menarche for girls and 15 mL testicular volume for boys). Within the younger participants (9–12 years old), 38% met the sleep duration recommendations for this age group and 53% had <1 hour of difference between weekdays and weekends (i.e. no social jetlag) (Table 1). Among the adolescents who were 13 years or older at T1, 6% met the sleep duration recommendations and 54% had no social jetlag. None of the younger adolescents exceeded the sleep duration recommendations, while 3% of the older adolescents slept longer than recommended.

The median (Q25, Q75) HOMA-IR at T1 was 3.0 (2.2, 4.5), and 51% were considered insulin resistant. At the T2 follow-up visit approximately 2 years later ( $\pm 0.4$  years), 98% of the sample were in late or post-puberty based on menarche or testicular volume, and the median (Q25, Q75) HOMA-IR was 3.4 (2.6, 4.8) with 57% considered as insulin resistant. Forty percent of participants were insulin resistant at both time points, 17% became insulin resistant from T1 to T2, 11% changed from insulin resistant to normal, and 32% were non-insulin resistant at both time points. Males were more likely to be have normal IR at both time points, while females were more likely to transition into and out of IR (Table 2). Participants who were younger at T1 were more likely to transition from IR to normal over the follow-up period. Those who reported to have ever tried smoking and those who had higher increases in adiposity were more likely to become insulin resistant.

In unadjusted analyses, there were no statistically significant associations between sleep characteristics and categories of IR (Table 3). However, in adjusted multinomial models, there were associations between sleep duration and timing with transitions to IR (Table 4). Specifically, adolescents who were 1 hour below the sleep duration recommendation in

comparison to those who met it had a 2.74 higher odds of becoming insulin resistant over follow-up (95% CI 1.01 to 7.43; P, trend=0.05). Considered as a continuous variable, each additional hour of sleep was related to 0.71 (0.52, 0.96; P=0.02) lower odds of developing IR, as well as 0.76 (0.59, 0.97; P=0.04) lower odds of being insulin resistant at both T1 and T2 and 0.70 (0.49, 0.99; P=0.03) lower odds of transitioning from insulin resistant to normal compared to normal HOMA-IR at both T1 and T2 (Table S1). Those with a sleep midpoint later than 4:33 AM had 2.63 higher odds of becoming insulin resistant (95% CI 1.03 to 6.72; P, trend=0.04). As a continuous variable, each hour later sleep midpoint was related to a 1.31 (0.97, 1.75) times higher odds of becoming insulin resistant, although this was not a statistically significant association (P=0.08). Interaction analysis showed no evidence of interaction by sex or age. There were no associations between sleep duration variability or social jetlag with changes in IR.

We next evaluated whether changes in adiposity mediated the relationships between sleep duration and timing with IR categories. As a first step, we examined associations between sleep duration and timing with continuous adiposity changes, but all were null. In formal mediation models, there was no evidence of mediation by adiposity changes for either sleep duration or timing with development of IR (e.g., P=0.9 for the natural indirect effect of sleep duration and odds of developing IR through changes in waist circumference and P=0.8 for sleep midpoint; data not shown). Finally, sensitivity analyses evaluating whether changes in sleep characteristics from T1 to T2 were related to IR revealed no evidence of associations (not shown).

## Discussion

Within this cohort of Mexican adolescents, shorter sleep duration and later sleep timing at baseline (T1) were each associated with developing IR during late adolescence. Specifically, adolescents who were 1 hour below the sleep duration recommendation or had later sleep midpoints than their peers at T1 had a 2.74 and 2.63 higher odds, respectively, of becoming insulin resistant over the next two years. In contrast, neither social jetlag nor sleep duration variability were associated with IR categories. Although higher changes in adiposity were associated with higher likelihood of developing IR and persistent IR across 2 years of follow-up, adiposity change was not a mediator in the sleep timing or duration and IR pathways.

Longitudinal investigations of actigraphy-based sleep and IR during adolescence are rare. Previous cross-sectional work, including ours, has linked insufficient sleep duration with IR among adolescents (16–20). Delayed sleep timing has also been related IR independently of sleep duration in few cross-sectional studies (18, 21). Existing longitudinal studies were conducted in children and focused on sleep duration. Among 8–11 year old Danish children, declines in sleep duration over a period of approximately 200 days were associated with increases in HOMA-IR after accounting for physical activity and sedentary time (37). Another study of 3900 European children who were between 2 and 11 years old at baseline found that shorter self-reported sleep duration was associated with higher HOMA-IR measured 4 years later, and this was mediated by waist circumference (20). Our study is an extension to the literature with regards to the focus on late adolescence.

Puberty is a unique period for sleep since it marks the beginning of a circadian phase delay that continues into the young adult period(38). Increased independence and new social or academic activities during this time period could also contribute to shorter sleep duration. Moreover, our evidence suggests that sleep duration and timing during puberty may contribute to IR that persists beyond the typical transient increase in IR. During typical development, as adolescents progress through the Tanner stages, IR rises (it may or may not reach the threshold for “insulin resistant”), but then falls again by Tanner stage 5 (9). Within our sample, 11% of adolescents who started out as insulin resistant at the first visit became non-insulin-resistant by the T2 visit. Nonetheless, there was a substantial number of adolescents in our sample who either remained insulin resistant or became insulin resistant over follow-up. Since the majority of these participants were classified as late puberty or fully mature by the follow-up visit, the fact that they continued as insulin resistant likely signals increased risk for cardiometabolic complications as they enter young adulthood. From a clinical viewpoint, our results may suggest that more extreme challenges to healthy sleep during late puberty could serve as an early indicator for later cardiometabolic risk, suggesting the need for sleep intervention.

Although we were underpowered to examine sex differences in sleep and IR relationships, we did not find statistical evidence that associations were stronger in females. This is in contrast to our earlier findings whereby sleep timing and duration were associated cross-sectionally with HOMA-IR in females but not males (21). While the literature is somewhat mixed (17), one other pediatric study suggested similar associations between sleep duration or timing with IR in girls but not boys (16). These dissimilar findings within the same cohort may suggest that relationships vary across the pubertal transition.

The fact that changes in measures of fat mass and distribution over the follow-up period were not statistical mediators also suggests that the potential mechanisms to explain a relationship between baseline sleep and development of IR could be independent of body size and composition, at least over this relatively short-term period of 2 years. Diet and physical activity are two potential behavioral mediators. A large body of evidence now shows that insufficient sleep duration and late sleep timing among children and adolescents are associated with higher caloric intake(39, 40), especially calories from carbohydrates(41, 42) and sugar-sweetened beverages (39–42). These sleep characteristics have also been associated with lower fruit and vegetable intake(41). Moreover, each of these dietary factors have been associated with IR (43–46). Insufficient sleep is also related to declines in moderate and vigorous activity, and higher sedentary behavior(47, 48), factors that are also independent predictors of IR (49). It is worth pointing out that in the present analysis, moderate/vigorous physical activity and total energy intake were related to development of IR (although not statistically significantly), but the addition of these variables to the sleep and IR model did not alter estimates. This suggests that the quantity of food and physical activity may not play a strong mediating role in the sleep and IR relationship, although further examination of potential modifying roles for these factors is warranted. However, it is important to highlight the likely role of circadian disruption for the observed relationship between sleep timing and IR. Circadian disruption can refer to differences between the circadian rhythms of the central brain clock (the suprachiasmatic nucleus) and the peripheral clocks in tissues such as liver, muscle, adipose tissue, and the pancreas, or between the



central brain clock and behavioral sleep/wake patterns (50). These differences can arise from delayed sleep patterns as well as delayed eating(51) and mistimed physical activity (52), and may contribute directly to IR by altering daily rhythms of glucose metabolism(50).

A primary strength of this study is the repeated measures of IR over late adolescence. Additional studies that follow these participants into young adulthood will provide further insights into potential cardiometabolic ramifications associated with persistent IR in late adolescence. The fact that sleep duration and timing were measured with wrist actigraphy rather than self-report is another strength. There are also limitations to consider. HOMA-IR is not the gold standard to measure IR, although it is a feasible measure in the context of a large cohort study and has been shown to be highly correlated with euglycemic-hyperinsulinemic clamp measurements(53). Further, clear consensus does not exist on which HOMA-IR value should demarcate IR among adolescents. We used cutpoints from a Chilean study(29) that accounted for pubertal status, but we acknowledge there may be limitations in this approach. In particular, the cutpoints were not validated against euglycemic-hyperinsulinemic clamps, although they had a sensitivity of 0.65 and specificity of 0.69 for metabolic syndrome (54). Of note, these cutpoints were more conservative than others reported in the literature among healthy populations of adolescents (55), suggesting that we may be underestimating IR in the present study. In addition, even though sex-stratified analyses of sleep and IR were warranted based on prior literature, the study was not adequately powered to detect these differences. Furthermore, because sleep was not a primary exposure of interest for the original grant, we did not have self-reported information on sleep quality, which could be an independent predictor of IR and an unmeasured confounder in this analysis. Finally, study findings may not be generalizable to adolescents in other populations, especially in settings where cultural norms and customs regarding sleep may differ. On a related note, findings may not be generalizable to adolescents who exceed sleep duration recommendations; there were too few adolescents in our cohort to examine relationships between excessive sleep duration and IR.

In summary, within a cohort of Mexican adolescents, we found that insufficient sleep duration and later sleep timing were each associated with over 2.5 times higher odds of developing IR over a 2-year follow-up period. These associations were of higher magnitude among females and were independent of changes in adiposity over the follow-up period. This prospective, longitudinal study raises the possibility of causal relationships, but cannot prove them. Whether intervening to improve sleep specifically around the time of pubertal transition can reduce risk for IR in late adolescence deserves further investigation.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgements:

We gratefully acknowledge the American British Cowdray Hospital for use of their research facilities, and we are extremely grateful to the research staff and participants for their commitment to the project.

**Funding:**

This work was supported by the US Environmental Protection Agency grant RD 83543601; National Institute of Environmental Health Sciences grants: P01 ES02284401, P30 ES017885, and R24 ES028502; National Heart, Lung, Blood Institute grant K01 HL151673. This study was also supported and partially funded by the National Institute of Public Health/Ministry of Health of Mexico. The funding sources had no involvement in the study design, collection, analysis, or interpretation of the data, the writing of the report, and the decision to submit the article for publication.

**REFERENCES**

1. Anon. Global Burden of Disease Study 2019. [WWW document]. URL <https://vizhub.healthdata.org/gbd-results/>
2. Sarwar N, Gao P, Kondapally Seshasai SR, et al. Diabetes mellitus, fasting blood glucose concentration, and risk of vascular disease: A collaborative meta-analysis of 102 prospective studies. *Lancet* 2010;375:2215–2222. [PubMed: 20609967]
3. Anon. (2022). Vista de Prevalencia de diabetes y descontrol glucémico en México: resultados de la Ensanut 2016 [WWW document]. URL <https://saludpublica.mx/index.php/spm/article/view/10752/11830>
4. Anon. (2022). Emite la Secretaría de Salud emergencia epidemiológica por diabetes mellitus y obesidad | Secretaría de Salud | Gobierno | gob.mx [WWW document]. URL <https://www.gob.mx/salud/prensa/emite-la-secretaria-de-salud-emergencia-epidemiologica-por-diabetes-mellitus-y-obesidad>
5. Fabricius-Bjerre S, Jensen RB, Færch K, et al. Impact of Birth Weight and Early Infant Weight Gain on Insulin Resistance and Associated Cardiovascular Risk Factors in Adolescence. *PLoS One* 2011;6:e20595. [PubMed: 21655104]
6. Fall CHD, Kumaran K. Metabolic programming in early life in humans. *Philos Trans R Soc Lond B Biol Sci* 2019;374.
7. Kaikkonen JE, Mikkilä V, Magnussen CG, Juonala M, Viikari JSA, Raitakari OT. Does childhood nutrition influence adult cardiovascular disease risk?—Insights from the Young Finns Study. *Ann Med* 2013;45:120–128. [PubMed: 22494087]
8. Al Mamun A, Cramb SM, O’Callaghan MJ, Williams GM, Najman JM. Childhood Overweight Status Predicts Diabetes at Age 21 Years: A Follow-up Study. *Obesity* 2009;17:1255–1261. [PubMed: 19214172]
9. Kelsey MM, Zeitler PS. Insulin Resistance of Puberty. *Curr Diabetes Reports* 2016 167 2016;16:1–6.
10. Moran A, Jacobs DR, Steinberger J, et al. Insulin resistance during puberty: Results from clamp studies in 357 children. *Diabetes* 1999;48:2039–2044. [PubMed: 10512371]
11. Arslanian SA, Kalhan SC. Correlations between fatty acid and glucose metabolism. Potential explanation of insulin resistance of puberty. *Diabetes* 1994;43:908–914. [PubMed: 8013756]
12. Arslanian SA, Kalhan SC. Protein turnover during puberty in normal children. *Am J Physiol* 1996;270.
13. Qi Q, Hua S, Perreira KM, et al. Sex Differences in Associations of Adiposity Measures and Insulin Resistance in US Hispanic/Latino Youth: The Hispanic Community Children’s Health Study/Study of Latino Youth (SOL Youth). *J Clin Endocrinol Metab* 2017;102:185–194. [PubMed: 27802095]
14. Goletzke J, Herder C, Joslowski G, et al. Habitually Higher Dietary Glycemic Index During Puberty Is Prospectively Related to Increased Risk Markers of Type 2 Diabetes in Younger Adulthood. *Diabetes Care* 2013;36:1870–1876. [PubMed: 23349549]
15. Kähönen E, Aatola H, Lehtimäki T, et al. Influence of early life risk factors and lifestyle on systemic vascular resistance in later adulthood: the cardiovascular risk in young Finns study. *Blood Press* 2021;30:367–375. [PubMed: 34605743]
16. Hitze B, Bosy-Westphal A, Bielfeldt F, et al. Determinants and impact of sleep duration in children and adolescents: Data of the Kiel Obesity Prevention Study. *Eur J Clin Nutr* 2009;63:739–746. [PubMed: 18665183]

17. Matthews KA, Dahl RE, Owens JF, Lee L, Hall M. Sleep Duration and Insulin Resistance in Healthy Black and White Adolescents. *Sleep* 2012;35:1353–1358. [PubMed: 23024433]
18. Simon SL, Behn CD, Cree-Green M, et al. Too Late and Not Enough: School Year Sleep Duration, Timing, and Circadian Misalignment Are Associated with Reduced Insulin Sensitivity in Adolescents with Overweight/Obesity. *J Pediatr* 2019;205:257–264.e1. [PubMed: 30497764]
19. Javaheri S, Storfer-Isser A, Rosen CL, Redline S. Association of short and long sleep durations with insulin sensitivity in adolescents. *J Pediatr* 2011;158:617–623. [PubMed: 21146189]
20. Thumann BF, Michels N, Felso R, et al. Associations between sleep duration and insulin resistance in European children and adolescents considering the mediating role of abdominal obesity. *PLoS One* 2020;15.
21. Chen P, Baylin A, Lee J, et al. The Association Between Sleep Duration and Sleep Timing and Insulin Resistance Among Adolescents in Mexico City. *J Adolesc Health* 2021;69:57–63. [PubMed: 33221190]
22. Spruyt K, Molfese DL, Gozal D. Sleep Duration, Sleep Regularity, Body Weight, and Metabolic Homeostasis in School-aged Children. *Pediatrics* 2011;127:e345–e352. [PubMed: 21262888]
23. He F, Bixler EO, Liao J, et al. Habitual sleep variability, mediated by nutrition intake, is associated with abdominal obesity in adolescents. *Sleep Med* 2015;16:1489–94. [PubMed: 26611945]
24. Perng W, Tamayo-Ortiz M, Tang L, et al. Early Life Exposure in Mexico to ENvironmental Toxicants (ELEMENT) Project. *BMJ Open* 2019;9.
25. Baek J, Banker M, Jansen EC, et al. An Efficient Segmentation Algorithm to Estimate Sleep Duration from Actigraphy Data. *Stat Biosci* 2021 2021:1–21.
26. Wittmann M, Dinich J, Mellow M, Roenneberg T. Social jetlag: misalignment of biological and social time. *Chronobiol Int* 2006;23:497–509. [PubMed: 16687322]
27. Paruthi S, Brooks LJ, D'Ambrosio C, et al. Recommended amount of sleep for pediatric populations: A consensus statement of the American Academy of Sleep Medicine. *J Clin Sleep Med* 2016;12:785–786. [PubMed: 27250809]
28. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and  $\beta$ -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985;28:412–419. [PubMed: 3899825]
29. Burrows AR, Leiva BL, Burgueño AM, et al. Insulin sensitivity in children aged 6 to 16 years: Association with nutritional status and pubertal development. *Rev Med Chil* 2006;134:1417–1426. [PubMed: 17277855]
30. Chavarro JE, Watkins DJ, Afeiche MC, et al. Validity of Self-Assessed Sexual Maturation Against Physician Assessments and Hormone Levels. *J Pediatr* 2017;186:172–178.e3. [PubMed: 28438374]
31. Chandler JL, Brazendale K, Beets MW, Mealing BA. Classification of physical activity intensities using a wrist-worn accelerometer in 8–12-year-old children. *Pediatr Obes* 2016;11:120–127. [PubMed: 25893950]
32. Jansen EC, Marcovitch H, Wolfson JA, et al. Exploring dietary patterns in a Mexican adolescent population: A mixed methods approach. *Appetite* 2020;147.
33. Hernández B, Gortmaker SL, Laird NM, Colditz GA, Parra-Cabrera S, Peterson KE. Validez y reproducibilidad de un cuestionario de actividad e inactividad física para escolares de la ciudad de Mexico. *Salud Publica Mex* 2000;42:315–323. [PubMed: 11026073]
34. Rodríguez-Carmona Y, Meijer JL, Zhou Y, et al. Metabolomics reveals sex-specific pathways associated with changes in adiposity and muscle mass in a cohort of Mexican adolescents. *Pediatr Obes* 2022;17:e12887. [PubMed: 35023314]
35. VanderWeele TJ. Mediation Analysis: A Practitioner's Guide. *Annu Rev Public Health* 2016;37:17–32. [PubMed: 26653405]
36. Anon. (2022). EconPapers: PARAMED: Stata module to perform causal mediation analysis using parametric regression models [WWW document]. URL <https://econpapers.repec.org/software/bocbocode/s457581.htm>
37. Hjorth MF, Chaput JP, Damsgaard CT, et al. Low physical activity level and short sleep duration are associated with an increased cardio-metabolic risk profile: A longitudinal study in 8–11 year old Danish children. *PLoS One* 2014;9.

38. Carpenter JS, Robillard R, Hickie IB. Variations in the sleep-wake cycle from childhood to adulthood: chronobiological perspectives. *ChronoPhysiology Ther* 2015;5:37–49.
39. Hart CN, Spaeth AM, Egleston BL, et al. Effect of changes in children’s bedtime and sleep period on targeted eating behaviors and timing of caloric intake. *Eat Behav* 2022;45:101629. [PubMed: 35390756]
40. Thellman KE, Dmitrieva J, Miller A, Harsh JR, LeBourgeois MK. Sleep timing is associated with self-reported dietary patterns in 9- to 15-year-olds. *Sleep Heal* 2017;3:269–275.
41. Duraccio KMR, Whitacre C, Krietsch KN, et al. Losing sleep by staying up late leads adolescents to consume more carbohydrates and a higher glycemic load. *Sleep* 2022;45.
42. Beebe DW, Simon S, Summer S, Hemmer S, Strotman D, Dolan LM. Dietary intake following experimentally restricted sleep in adolescents. *Sleep* 2013;36:827–834. [PubMed: 23729925]
43. Romero-Polvo A, Denova-Gutiérrez E, Rivera-Paredes B, et al. Association between dietary patterns and insulin resistance in mexican children and adolescents. *Ann Nutr Metab* 2012;61:142–150. [PubMed: 23037180]
44. Perng W, Fernandez C, Peterson KE, et al. Dietary Patterns Exhibit Sex-Specific Associations with Adiposity and Metabolic Risk in a Cross-Sectional Study in Urban Mexican Adolescents. *J Nutr* 2017;147:1977–1985. [PubMed: 28855420]
45. Steffen LM, Jacobs DR, Murtaugh MA, et al. Whole Grain Intake Is Associated with Lower Body Mass and Greater Insulin Sensitivity among Adolescents. *Am J Epidemiol* 2003;158:243–250. [PubMed: 12882946]
46. Karatzi K, Moschonis G, Barouti AA, Lionis C, Chrousos GP, Manios Y. Dietary patterns and breakfast consumption in relation to insulin resistance in children. *The Healthy Growth Study. Public Health Nutr* 2014;17:2790–2797. [PubMed: 24477051]
47. Mead MP, Baron K, Sorby M, Irish LA. Daily Associations Between Sleep and Physical Activity. *Int J Behav Med* 2019.
48. Master L, Nye RT, Lee S, et al. Bidirectional, Daily Temporal Associations between Sleep and Physical Activity in Adolescents. *Sci Reports* 2019 91 2019;9:1–14.
49. Berman LJ, Weigensberg MJ, Spruijt-Metz D. Physical activity is related to insulin sensitivity in children and adolescents, independent of adiposity: a review of the literature. *Diabetes Metab Res Rev* 2012;28:395–408. [PubMed: 22389103]
50. Stenvers DJ, Scheer FAJL, Schrauwen P, la Fleur SE, Kalsbeek A. Circadian clocks and insulin resistance. *Nat Rev Endocrinol* 2019.
51. Rangaraj VR, Siddula A, Burgess HJ, Pannain S, Knutson KL. Association between Timing of Energy Intake and Insulin Sensitivity: A Cross-Sectional Study. *Nutr* 2020, Vol 12, Page 503 2020;12:503.
52. Gabriel BM, Zierath JR. Circadian rhythms and exercise — re-setting the clock in metabolic disease. *Nat Rev Endocrinol* 2018 154 2019;15:197–206. [PubMed: 30655625]
53. Lorenzo C, Haffner SM, Stanáková A, Laakso M. Relation of Direct and Surrogate Measures of Insulin Resistance to Cardiovascular Risk Factors in Nondiabetic Finnish Offspring of Type 2 Diabetic Individuals. *J Clin Endocrinol Metab* 2010;95:5082–5090. [PubMed: 20702522]
54. Burrows RA, Leiva LB, Weisstaub G, et al. High HOMA-IR, adjusted for puberty, relates to the metabolic syndrome in overweight and obese Chilean youths. *Pediatr Diabetes* 2011;12:212–218. [PubMed: 21426454]
55. Andrade MIS De Oliveira JS, Leal VS, et al. Identification of cutoff points for Homeostatic Model Assessment for Insulin Resistance index in adolescents: systematic review. *Rev Paul Pediatr* 2016;34:234. [PubMed: 26559605]

### Study Importance

- Sleep has emerged as a possible risk factor for insulin resistance (IR) among adolescents, but longitudinal investigations that consider the multidimensionality of sleep are lacking.
- Both short sleep duration and later sleep timing were related to higher odds of developing IR over a 2-year period during late adolescence.
- Whether intervening to improve both sleep timing and duration can reduce risk for IR in late adolescence deserves further investigation.

**Table 1.**

Sleep characteristics at baseline, stratified by sex and age

Baseline sleep characteristics	n	Boys, %	Girls, %	Younger age group 9–12, %	Older age group, 13+, %
Weekday sleep duration <sup>1</sup>					
Recommended sleep duration	196	54.5	53.8	38.0	65.6
Within 1 hour below recommendation	103	29.8	27.2	35.3	23.6
1 hour below recommendation	63	15.7	19.0	26.7	10.8
Sleep duration variability over 7 days					
<1 hour SD	97	28.65	25.00	36.67	19.81
1–2 hour SD	190	51.69	53.26	49.33	54.72
>2 hour SD	75	19.66	21.74	14.00	25.47
Weekday sleep midpoint					
Tertile 1– 12:55 AM to 2:58 AM	127	38.20	32.07	44.00	28.77
Tertile 2– 2:59 AM to 4:33 AM	128	30.34	40.22	36.67	34.43
Tertile 3– 4:34 AM to 7:59 AM	107	31.46	27.72	19.33	36.79
Social jetlag					
Within 1 hour, weekend to weekday	193	56.82	50.54	53.33	53.81
>1 hour later on weekend	96	26.14	27.17	28.00	25.71
>2 hours later on weekend	71	17.05	22.28	18.67	20.48

<sup>1</sup>Recommended sleep duration for ages 6–12 is 9 to 12 hours per 24 hours, while recommended sleep duration for 13–18 is 8 to 10 hours per 24 hours

**Table 2.**

Associations between baseline sociodemographic or lifestyle characteristics and categories for changes in insulin resistance (IR) during the 2 years between baseline (T1) and follow-up (T2) assessments

	N (Row Total)	Normal HOMA-IR at T1 and T2 N=115	Transition from normal to IR N=61	Transition from IR to normal N=40	IR at T1 and T2 N=146
Sex					
Male	178	41.4 <sup>†</sup>	12.9	11.8	34.8
Female	184	23.4	20.7	10.3	45.7
P-value				0.002	
Age group, years (y)					
9 to <12 years	82	23.2	15.9	9.8	51.2
12 to <14 years	136	31.6	11.8	11.0	45.6
14 to 18 years	144	36.8	22.2	11.8	29.2
P-value				0.02	
Testicular volume (boys only)					
<15 mL	35	31.4	14.3	11.4	42.9
15 mL (latter stages of puberty)	132	43.2	13.6	10.6	32.6
P-value				0.61	
Menarche status (girls only)					
Had not experienced	38	15.8	13.2	7.9	63.2
Had experienced	147	25.2	21.1	11.6	42.2
P-value				0.15	
Maternal education, years (y)					
8 y or less (secondary or primary)	39	28.2	25.6	15.4	30.8
9 to 11 y (some high school)	145	26.9	20.7	9.0	43.5
12 y (completed high school)	121	36.4	11.6	12.4	39.7
>12 y	54	37.0	11.1	9.3	42.6
P-value				0.24	
Moderate/vigorous physical activity, min/day, quartiles					
Q (Quartile)1, 15 to 60.22	89	25.8	23.6	11.2	39.3
Q2, 60.23 to 77.13	88	39.8	15.9	13.6	39.7
Q3, 77.14 to 97.9	88	33.0	13.6	13.6	39.8
Q4, 98.7 to 174	88	26.1	14.8	6.8	52.3
P-value				0.11	
Screen time, hours/week, quartiles					
Q1, 2 to 20.5	92	28.3	16.3	12.0	43.5
Q2, 21 to 30.5	90	30.0	15.6	11.1	43.3
Q3, 31 to 40.5	88	37.5	19.3	6.8	36.4
Q4, 41 to 104	90	31.1	16.7	13.3	38.9
P-value				0.87	
Ever smoked cigarettes					

	N (Row Total)	Normal HOMA-IR at T1 and T2 N=115	Transition from normal to IR N=61	Transition from IR to normal N=40	IR at T1 and T2 N=146
No	298	30.9	14.8	10.1	44.3
Yes	59	35.6	27.1	15.3	22.0
P-value				0.007	
Average Calorie intake, kcal					
Q1, 663 to 1688	89	23.6	14.6	9.0	52.8
Q2, 1690 to 2169	91	31.9	14.3	17.6	36.3
Q3, 2173 to 2812	89	38.2	16.9	7.9	37.1
Q4, 2817 to 5964	89	31.5	22.5	9.0	37.1
P value				0.10	
Change in waist circumference, cm					
Q1, median=-14.9 to 2.8	92	39.1	7.6	16.3	37.0
Q2, median=3 to 6.25	90	37.8	14.4	13.3	34.4
Q3, median=6.3 to 9.34	91	30.8	13.2	7.7	48.4
Q4, median=9.35 to 25.1	89	19.1	32.6	6.7	41.6
P-value				<0.0001	
Change in BMI, kg/m <sup>2</sup>					
Q1, median=-6.3 to 0.26	92	44.6	8.7	13.0	33.7
Q2, median=0.27 to 1.1931	91	34.1	13.2	16.5	36.3
Q3, median=1.1933 to 2.20	91	28.6	19.8	8.8	42.9
Q4, median=2.21 to 9.79	88	19.3	26.1	5.7	48.9
P-value				0.001	
Change in body fat percentage					
Q1, median=-22 to -1.8	91	34.1	6.6	16.5	42.9
Q2, median=-1.79 to 1.19	90	34.4	12.2	13.3	40.0
Q3, median=1.20 to 3.80	89	33.7	18.0	9.0	39.3
Q4, median=3.90 to 20	89	25.8	30.3	5.6	38.2
P-value				0.005	
Change in triceps skinfold, mm					
Q1, median=-16 to -1	101	42.6	9.9	14.9	32.7
Q2, median=-0.7 to 1.5	87	34.5	16.1	12.6	36.8
Q3, median=2 to 5	89	25.8	18.0	4.5	51.7
Q4, median=5.5 to 21	84	22.6	25.0	10.7	41.7
P-value				0.006	

<sup>1</sup>Values represent row percentages

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



**Table 3.**

Unadjusted associations between baseline sleep characteristics and 2-year changes in HOMA-IR from baseline (T1) to follow-up (T2)

Baseline sleep characteristics	Normal HOMA-IR at T1 and T2 N=115	Transition from normal to IR N=61	Transition from IR to normal N=40	IR at T1 and T2 N=146
Sleep duration				
Recommended sleep duration	34.7 <sup>1</sup>	16.8	11.2	37.2
Within 1 hour below recommendation	33.0	15.5	7.8	43.7
1 hour below recommendation	20.6	19.1	15.9	44.4
P-value		0.35		
Sleep duration variability over 7 days				
<1 hour SD	32.0	12.4	11.3	44.3
1–2 hour SD	28.4	18.4	10.5	42.6
>2 hour SD	40.0	18.7	12.0	29.3
P-value		0.34		
Sleep midpoint				
Tertile 1– 12:55 AM to 2:58 AM	36.2	12.6	11.0	40.2
Tertile 2– 2:59 AM to 4:33 AM	27.3	17.2	10.2	45.3
Tertile 3– 4:34 AM to 7:59 AM	31.8	21.5	12.2	34.6
P-value		0.40		
Social jetlag				
Within 1 hour, weekend to weekday	30.1	18.1	13.0	38.9
>1 hour later on weekend	31.3	13.5	8.3	46.9
>2 hours later on weekend	36.6	18.3	9.9	35.2
P-value		0.60		

<sup>1</sup>Values represent row percentages

**Table 4.**

Adjusted associations between baseline sleep characteristics and 2-year changes in HOMA-IR from baseline (T1) to follow-up (T2)

Baseline sleep characteristics	Transition to IR compared to normal HOMA-IR at both, <sup>1</sup> OR (95% CI)	Transition from IR to normal compared to normal HOMA-IR at both, <sup>1</sup> OR (95% CI)	IR at both time points compared to normal HOMA-IR at both, <sup>1</sup> OR (95% CI)
Sleep duration			
Recommended sleep duration	Reference	Reference	Reference
Within 1 hour below recommendation	1.43 (0.64, 3.20)	0.87 (0.33, 2.33)	1.20 (0.65, 2.24)
1 hour below recommendation	2.74 (1.01, 7.43)	2.83 (0.98, 8.22)	2.00 (0.90, 4.47)
P, trend <sup>2</sup>	0.05	0.10	0.11
Sleep duration variability over 7 days			
<1 hour SD	Reference	Reference	Reference
1–2 hour SD	1.49 (0.6, 3.37)	0.98 (0.41, 2.35)	1.17 (0.4, 2.14)
>2 hour SD	1.02 (0.39, 2.8)	0.68 (0.23, 2.00)	0.62 (0.29, 1.33)
P, trend <sup>2</sup>	0.99	0.48	0.28
Sleep midpoint			
Tertile 1– 12:55 AM to 2:58 AM	Reference	Reference	Reference
Tertile 2– 2:59 AM to 4:33 AM	1.82 (0.80, 4.17)	1.24 (0.49, 3.10)	1.64 (0.89, 3.05)
Tertile 3– 4:34 AM to 7:59 AM	2.63 (1.03, 6.72)	1.54 (0.54, 4.44)	1.64 (0.79, 3.41)
P, trend <sup>2</sup>	0.04	0.42	0.15
Social jetlag			
Within 1 hour, weekend to weekday	Reference	Reference	Reference
>1 hour later on weekend	0.69 (0.31, 1.52)	0.62 (0.25, 1.56)	1.13 (0.62, 2.04)
>2 hours later on weekend	0.77 (0.34, 1.73)	0.62 (0.24, 1.65)	0.73 (0.37, 1.43)
P, trend <sup>2</sup>	0.43	0.26	0.50

<sup>1</sup>From a multinomial logistic regression model adjusted for age, sex, and baseline puberty status; sleep duration and midpoint were mutually adjusted

<sup>2</sup>Obtained by replacing the categorical predictor with a variable representing ordinal categories of each sleep category