



## Guest Editorial

## Hot topics in muscle metabolism and adaptation



This special issue highlights comprehensive reviews and researches on topics in fundamentally important areas of skeletal muscle physiology.<sup>1–6</sup> The most important adaptive response of skeletal muscle to exercise, particularly resistant exercise, is increased protein synthesis, which is critical for promoting muscle hypertrophy and/or preventing atrophy. This process plays an essential role in promoting and maintaining skeletal muscle contractile and metabolic functions.

The review by Tinline-Goodfellow et al.<sup>2</sup> elegantly reviewed the literature on the regulation of muscle protein synthesis with a focus on the mechanistic target of the rapamycin (mTOR) complex 1 (mTORC1) pathway. The authors not only provided a comprehensive review of findings in support of the indispensable role of mTORC1 in promoting muscle protein synthesis in response to growth factors, nutrients and mechanical stimuli, but also brought our attention to the role of skeletal muscle periphery region, including recruitment of mTORC1 to this region, in promoting protein synthesis. They dived deep into the possible molecular mechanisms, linking the spatial specificity of skeletal muscle anabolism to substrate availability, signaling regulation as well as protein translation. This review provides a contemporary understanding of the beauty of the regulatory system for skeletal muscle mass control in our body.

Islam and Gillen's review<sup>3</sup> dealt with another equally important area of skeletal muscle physiology: exercise-enhanced insulin sensitivity and glycemic control. They focused on the exciting development of two similar exercise modes, high-intensity interval training (HIIT) and sprint interval training (SIT). These two exercise modes have recently gained much attention due to their short duration and high efficacy in achieving many of the health benefits of traditional moderate-intensity continuous training. The article presented unequivocal evidence that HIIT and SIT either in an acute or chronic setting are equally, if not more, effective in promoting insulin sensitivity in both healthy individuals and patients with metabolic syndrome. The authors provided a balanced review of the evidence in supporting the role of some molecular regulators, but correctly pointed to the lack of the scientific evidence of causal links. The review also raised a question about blunted glycemic benefits of HIIT and/or SIT in females and the need for more studies to improve the understanding of how intense interval training improves insulin sensitivity.

Finally, Koopmans et al.<sup>1</sup> brought us to another level of complexity of muscle adaptation with a focus on the role of myonuclei. The review provided a much-needed detailed description of the findings of how myonuclei, resident or satellite cell-derived, undergo transcriptional, epigenetic, mobility and morphological regulations in response exercise

stimuli in vivo. Intriguingly, the review presented evidence supporting that myonuclei may have the memory of past training adaptations. The authors postulated that this memory is likely due to epigenetic modification either by DNA methylation or myomiR modification.

These articles presented the most updated evidence in the research field regarding skeletal muscle adaptation to exercise training dealing with three fundamentally important questions related to the control of protein synthesis, glucose metabolism and gene regulation in myonuclei, hopefully provoking more questions and stimulating more research in exercise-induced adaptations in skeletal muscle.

## References

- Koopmans PJ, Zwetsloot KA, Murach KA. Going nuclear: molecular adaptations to exercise mediated by myonuclei. *Sports Med Health Sci.* 2023;5(1):2–9. <https://doi.org/10.1016/j.smhs.2022.11.005>.
- Tinline-Goodfellow CT, Lees MJ, Hodson N. The skeletal muscle fiber periphery: a nexus of mTOR-related anabolism. *Sports Med Health Sci.* 2023;5(1):10–19. <https://doi.org/10.1016/j.smhs.2022.11.004>.
- Islam H, Gillen JB. Skeletal muscle mechanisms contributing to improved glycemic control following intense interval exercise and training. *Sports Med Health Sci.* 2023; 5(1):20–28. <https://doi.org/10.1016/j.smhs.2023.01.002>.
- Katz A. The role of glycogen phosphorylase in glycogen biogenesis in skeletal muscle after exercise. *Sports Med Health Sci.* 2023;5(1):29–33. <https://doi.org/10.1016/j.smhs.2022.11.001>.
- Liu M, Yang Y, Shao H, Liu J, Niu M, Fu L. Globular adiponectin ameliorates insulin resistance in skeletal muscle by enhancing the LKB1-mediated AMPK activation via SESN2. *Sports Med Health Sci.* 2023;5(1):34–41. <https://doi.org/10.1016/j.smhs.2022.08.001>.
- Jeon S, Sontag SA, Herda TJ, Trevino MA. Chronic training status affects muscle excitation of the vastus lateralis during repeated contractions. *Sports Med Health Sci.* 2023;5(1):42–49. <https://doi.org/10.1016/j.smhs.2022.12.005>.

Zhen Yan\*

Center for Exercise Medicine Research, Fralin Biomedical Research Institute  
at Virginia Tech Carilion, Roanoke, VA, 24016, USA  
Department of Human Nutrition, Foods, and Exercise, College of Agriculture  
and Life Sciences, Virginia Tech, Roanoke, VA, 24016, USA

David A. Hood  
Muscle Health Research Centre, School of Kinesiology and Health Science,  
York University, 4700 Keele St., Toronto, ON, M3J 1P3, Canada

\* Corresponding author. Center for Exercise Medicine Research, Fralin  
Biomedical Research Institute at Virginia Tech Carilion, Roanoke, VA,  
24016, USA.

E-mail address: [zhenyan1@vtc.vt.edu](mailto:zhenyan1@vtc.vt.edu) (Z. Yan).

<https://doi.org/10.1016/j.smhs.2023.01.004>

Available online 24 January 2023

2666-3376/© 2023 Chengdu Sport University. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).