



OPEN ACCESS

APPROVED BY
Frontiers in Cardiovascular Medicine,
Frontiers Media SA, Switzerland

*CORRESPONDENCE
Luigi P. Badano
✉ luigi.badano@unimib.it

SPECIALTY SECTION
This article was submitted to Structural
Interventional Cardiology, a section of the
journal Frontiers in Cardiovascular Medicine

RECEIVED 20 February 2023
ACCEPTED 21 February 2023
PUBLISHED 13 March 2023

CITATION
Gavazzoni M, Heilbron F, Badano LP, Radu N,
Cascella A, Tomaselli M, Perelli F, Caravita S,
Baratto C, Parati G and Muraru D (2023)
Corrigendum: The atrial secondary tricuspid
regurgitation is associated to more favorable
outcome than the ventricular phenotype.
Front. Cardiovasc. Med. 10:1169907.
doi: 10.3389/fcvm.2023.1169907

COPYRIGHT
© 2023 Gavazzoni, Heilbron, Badano, Radu,
Cascella, Tomaselli, Perelli, Caravita, Baratto,
Parati and Muraru. This is an open-access
article distributed under the terms of the
[Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/).
The use, distribution or reproduction in other
forums is permitted, provided the original
author(s) and the copyright owner(s) are
credited and that the original publication in this
journal is cited, in accordance with accepted
academic practice. No use, distribution or
reproduction is permitted which does not
comply with these terms.

Corrigendum: The atrial secondary tricuspid regurgitation is associated to more favorable outcome than the ventricular phenotype

Mara Gavazzoni¹, Francesca Heilbron¹, Luigi P. Badano^{1,2*},
Noela Radu^{2,3}, Andrea Cascella², Michele Tomaselli¹,
Francesco Perelli², Sergio Caravita^{1,4}, Claudia Baratto¹,
Gianfranco Parati¹ and Denisa Muraru^{1,2}

¹Department of Cardiology, Istituto Auxologico Italiano, IRCCS, Milan, Italy, ²Department of Medicine and Surgery, University of Milano-Bicocca, Milan, Italy, ³University of Medicine and Pharmacy Carol Davila Bucharest, Emergency University Hospital Bucharest, Bucharest, Romania, ⁴Department of Management, Information and Production Engineering, University of Bergamo, Dalmine, Italy

KEYWORDS

atrial secondary tricuspid regurgitation, atrial fibrillation, heart failure hospitalizations, secondary tricuspid regurgitation, prognosis

A Corrigendum on

The atrial secondary tricuspid regurgitation is associated to more favorable outcome than the ventricular phenotype

By Gavazzoni M, Heilbron F, Badano LP, Radu N, Cascella A, Tomaselli M, Perelli F, Caravita S, Baratto C, Parati G, Muraru D. (2022) Front. Cardiovasc. Med. 9:1022755. doi: 10.3389/fcvm.2022.1022755

In the published article, there was an error in affiliation (2) for the following 3 authors: Mara Gavazzoni, Michele Tomaselli, Claudia Baratto. The affiliation (2) “Department of Medicine and Surgery, University of Milano-Bicocca, Milan, Italy” does not apply to these three authors. The only affiliation for these authors should be (1): “Department of Cardiology, Istituto Auxologico Italiano, IRCCS, Milan, Italy”, as displayed in the new version of the paper.

The authors apologize for this error and state that this does not change the scientific conclusions of the article in any way. The original article has been updated.