



Robotic operations in urgent general surgery: a systematic review

Alexander Reinisch¹ · Juliane Liese² · Winfried Padberg² · Frank Ulrich¹

Received: 18 March 2022 / Accepted: 6 May 2022 / Published online: 21 June 2022
© The Author(s) 2022

Abstract

Robotically assisted operations are the state of the art in laparoscopic general surgery. They are established predominantly for elective operations. Since laparoscopy is widely used in urgent general surgery, the significance of robotic assistance in urgent operations is of interest. Currently, there are few data on robotic-assisted operations in urgent surgery. The aim of this study was to collect and classify the existing studies. A two-stage, PRISMA-compliant literature search of PubMed and the Cochrane Library was conducted. We analyzed all articles on robotic surgery associated with urgent general surgery resp. acute surgical diseases of the abdomen. Gynecological and urological diseases so as vascular surgery, except mesenterial ischemia, were excluded. Studies and case reports/series published between 1980 and 2021 were eligible for inclusion. In addition to a descriptive synopsis, various outcome parameters were systematically recorded. Fifty-two studies of operations for acute appendicitis and cholecystitis, hernias and acute conditions of the gastrointestinal tract were included. The level of evidence is low. Surgical robots in the narrow sense and robotic camera mounts were used. All narrow-sense robots are nonautonomous systems; in 82%, the Da Vinci[®] system was used. The most frequently published emergency operations were urgent cholecystectomies (30 studies, 703 patients) followed by incarcerated hernias (9 studies, 199 patients). Feasibility of robotic operations was demonstrated for all indications. Neither robotic-specific problems nor extensive complication rates were reported. Various urgent operations in general surgery can be performed robotically without increased risk. The available data do not allow a final evidence-based assessment.

Keywords General surgery · Minimally invasive surgery · Robotic surgery · Emergency surgery · Urgent surgery

Introduction

Laparoscopic interventions are established in general surgery for urgent, acute diseases of the abdominal organs. For acute appendicitis and cholecystitis, laparoscopic operations are the standard of care and are recommended in guidelines [1, 2]. Many surgical departments also operate on

gastrointestinal perforations, incarcerated hernias and bowel obstruction laparoscopically on a regular basis.

However, robotic surgery has been an integral part of general surgery for over two decades and is becoming increasingly widespread. From 2012 to 2018, an increase in general surgical robotic interventions by more than a factor of 8 was described, reaching up to 15.1% of all general surgical operations in the USA [3]. This raises the question of the significance of robotic surgery in urgent surgery. Recently, this led to the publication of a position paper by the World Society of Emergency Surgery (WSES) [4].

We can distinguish two groups of robots as follows: surgical robots (SRs) in a narrower sense and robotic camera mounts (RCMs). SRs in the stricter definition are nonautonomous systems, which are controlled by a surgeon via a console. The main representative of this group of robots is the Da Vinci[®] system from Intuitive Surgery Inc. (Sunnyvale, CA, USA) since the products from Computer Motion (ZEUS, AESOP; Santa Barbara, CA, USA) were discontinued after Computer Motion and Intuitive merged in 2003.

Alexander Reinisch and Juliane Liese have contributed equally to this work.

Winfried Padberg and Frank Ulrich have contributed equally to this work.

✉ Alexander Reinisch
Alexander_reinisch@hotmail.com

¹ Department of General, Visceral and Oncologic Surgery, Wetzlar Hospital and Clinics, Forsthausstr. 1, 35578 Wetzlar, Germany

² Department of General, Visceral, Thoracic, Transplant and Pediatric Surgery, University Hospital Giessen, Giessen, Germany

Recently, the Da Vinci[®] system has faced competition from other manufacturers (e.g., CMR Surgical Versius[®], Cambridge, UK; Hugo[®], Medtronic, Dublin, Ireland; and Microhand S, China; Dexter[®], Distalmotion, Switzerland).

A number of advantages are accredited to robotic surgical systems as follows: systems such as the Da Vinci[®] system should help to overcome the disadvantages of laparoscopic surgery, such as physiological tremors and restricted degrees of freedom. RCMs, such as Soloassist[®] II (AKTORMed, Barbing, Germany), promise the liberation of the physician acting as a surgical assistant. However, these operations are very similar to “classic” laparoscopy. All robotic systems also advertise improved ergonomics and optimized visualization. Whether these properties lead to advantages over traditional laparoscopy is the subject of debate. All robotic systems are theoretically predisposed for technical malfunctions that may cause harm to the patient. Several publications report on better outcomes of robotic surgery, e.g., conversion rates, morbidity (including postoperative ileus) and postoperative stay in colorectal surgery [5, 6]. However, almost all publications examine elective operations. Against the background and known advantages of laparoscopic operations in urgent operations, the possible benefit of surgical robots in urgent general surgery must be examined and discussed. The aim of this review is to analyze and classify the available data on this topic.

Methods

Literature search

Two of the authors (AR and JL) independently searched PubMed (1980-present) and the Cochrane Library (1980-present). The systematic review was performed according to the preferred reporting items for systematic reviews and meta-analysis (PRISMA) guidelines [7].

A two-stage analysis was conducted. First, a preliminary search and screening of 500 results was performed on October 26th and 27th, 2020, to elaborate the principles of the further analysis. The results of this first search indicated that the number of publications of interest was very limited. Therefore, we decided to include case reports and case series in the systematic analysis. Furthermore, this first search showed a massive heterogeneity of data. To give the reader a full overview of the topic, we decided not to attempt a meta-analysis since a meta-analysis would have limited the useable studies to only a few. A strict systematic review or a meta-analysis would necessarily have meant that most urgent surgical diseases could not have been analyzed. The primary goal of this work, however, was to cover robotic-assisted operations in urgent surgery as broadly as possible, which was only possible through a partially descriptive evaluation.

The second definitive search was run on December 31st, 2021.

The following search terms were used in combination with “robotic” and “robotic surgery”: adhesiolysis, appendectomy, appendicitis, bowel obstruction, cholecystectomy, cholecystitis, diverticular disease, diverticulitis, hernia, incarceration, perforation, peritonitis, ulcer, urgency and emergency (Appendix 1).

The search terms were partially truncated to include as many grammatical variables as possible.

Inclusion criteria

All studies, case reports and case series describing urgent general and visceral surgical procedures in connection with robotic surgery were included. Manuscripts were included in which typical general or visceral surgical operations were described; inclusion was not decided by which specialty the operation performed. If study populations were published multiple times, the more recent publication was included.

Exclusion criteria

Publications that did not report the original data were excluded, as were those with overlapping study populations (see above). Urological, gynecological or vascular surgery emergency interventions were excluded. Reports on the thoracoscopic robotic treatment of diaphragmatic hernias were also excluded. If the full text was not available, the study was also excluded.

Evaluation

If the specified information was extractable, complications were classified according to the Clavien–Dindo classification, whenever applicable [8]. In all other cases and biliary complications in cholecystectomies, the complications were mentioned separately.

Outcomes of interest

All included manuscripts were examined with a focus on the following factors: (a) primary objective of the study and technical aspects, (b) complications (see above) and whether these complications could be related to the use of the robot, (c) pros and cons of the use of the robotic operation, (d) financial aspects, (e) factors related to the acute/urgent situation and (f) further outcomes of interest.

Results

The literature search resulted in a total of 3072 (+5, see below) findings, of which 219 were eliminated (duplicates, letters, replies, guidelines, etc.). A total of 2853 papers were analyzed. The review of the references of the analyzed full texts led to the inclusion of five additional studies. A total of 2645 publications were excluded since the title and/or abstract did not meet the inclusion criteria. The full texts of 208 studies met the criteria and were further analyzed. No unpublished studies were obtained. The full-text analysis revealed that 155 of these studies did not contain data of interest for this review, and one full text was unavailable. Three similar case reports were combined into a case series [9–11]. In total, 52 studies were included, five of which were evaluated for more than one indication (Fig. 1) [12–16]. Apart from one paper on a spleen hematoma, all reports on urgent robotic operations could be assigned to four classic general surgical fields: appendectomies, cholecystectomies, hernias (partly with intestinal obstruction) and gastrointestinal procedures [15].

Of the included papers, 15 were case reports or case series. None of the studies were randomized, and 24 were controlled. Six studies were prospective, and the remaining studies were retrospective [17–22]. Only four retrospective and none of the prospective studies explicitly examined urgent robotic operations, and two of these were controlled studies [23–26].


Two studies focused on the use robotic camera mounts (RCM), and they included 302 urgently operated patients

[14, 21]. All other 50 studies examined the use of surgical robots in the narrow sense (SR). In these studies, 955 urgently operated patients were included. A total of 655 of these patients (68.6%) were treated using the Da Vinci® system (41 studies, 82% of the SR studies). Hosein et al. and Gangemi et al. did not specify the type of SR, and the 261 patients included in their studies accounted for 27.4% of the SR patients [27, 28]. An alternative SR was utilized in only 39 operations (3.6%).

Complication rates were listed in all but one study, but with varying degrees of accuracy [29]. Some studies reported complications only above a certain degree of severity, i.e., Clavien–Dindo grade ≥ 3 . It should be noted that only 22 of the 52 studies differentiated whether the complication occurred in the subgroup of emergency robotic procedures, and 12 of these were case reports. In most of the included studies, only the complication rate of the entire robotic group was listed. In addition, there was no strict distinction between intra- and postoperative complications, while the follow-up periods differed greatly.

Since urgent operations were neither an interest, end-point nor variable in the majority of the studies, evidence grading, e.g., the Newcastle–Ottawa Quality Assessment Scale, cannot be applied meaningfully. Regarding urgent operations, the levels of evidence for almost all included studies must be assessed as low, which was true particularly for case series and case reports. One paper was excluded because it was a duplicate publication of a more or less identical patient group [26].

Fig. 1 PRISMA-Flow diagram showing the literature search and the study selection with 52 relevant studies enrolled in this review



† Three similar case reports by one author were combined into a case series.

Appendectomies

Seven papers reported on urgent appendectomies using robotic surgery in a total of 196 patients (Table 1). The vast majority of patients ($n = 185$) were analyzed in two studies, both using RCM. Three more publications were case reports or case series. As a result, only 11 patients were urgently appendectomized using an SR. No complications or conversions were reported.

The three case reports/case series specified closure of the appendix stump (sutures), and the other studies did not provide any information on this. There were no reports of robotic single-incision appendectomies.

Cholecystectomies

Thirty publications reported on urgent robotic cholecystectomies, none of which were randomized (Table 2). Three studies were prospectively controlled, and 14 were retrospectively controlled. Of the uncontrolled studies, three were prospective, and six were retrospective. Four case reports or case series have been summarized in the table for better clarity [13, 30–32].

A total of 804 urgent cholecystectomies were performed with robotic assistance, 101 of them with an RCM. The remaining 703 patients underwent surgical intervention with an SR. The Da Vinci[®] system was used in 546 (77.7%) patients; the robot type was not reported for 130 patients. One study explicitly analyzed urgent operations as the main interest [23].

Eleven publications reported cholecystectomies using the single-incision robotic technique (SIRC).

The reports of complications differed greatly, as did the follow-up periods (0–6 months). Kubat et al. showed 12% complications in urgent cholecystectomies, including one (1.5%) biliary tract injury [23]. The highest reported complication rate was 20% (including seromas), and 9 studies stated that no complications occurred. The incidence of Clavien–Dindo grade ≥ 3 complications varied between 1.7 and 4.2%. Biliary tract problems (injury, leakage) were demonstrated in three studies, each with one patient. Hernias represented a particular complication of single-site operations; these were described in three studies, with an incidence of 0.9–6.5%. Conversion to a laparoscopic procedure or to open surgery was reported in 7 studies, and two studies indicated that conversion was necessary due to inflammation [33, 34]. In one study, conversions were an exclusion criterion [35]. Six authors reported a prolonged operation time and one reported a decreased operation time in the robotic group compared to laparoscopic cholecystectomies. One study reported higher costs for robotically operated patients [36].

Gastrointestinal procedures

Urgent gastrointestinal operations were investigated in 12 studies (Table 3). There were three studies in this group of topics that specifically investigated urgent operations: Anderson et al., Beltzer et al. and, most recently, Robinson et al. [24, 37, 38]. The publication by Ohmura et al. on the use of an RCM evaluated for appendicitis and cholecystitis also reports on the operation of 16 perforations in the upper gastrointestinal tract [14]. Three very similar case reports of Kudsi et al. were combined into a case series [9–11].

It remains unclear how many of the 60 robotically operated patients reported by Beltzer et al. were treated in an emergency situation. For this review, we accepted two of the 60 patients with diverticulitis type IIc of classification of diverticular disease (CDD) since they undoubtedly belonged to the group of urgent operations [39]. The study did not distinguish the complications between urgent and elective operations. The overall complication rate was 30%, with percentages of 8.3% for Clavien–Dindo grade 3b complications and 6.7% for anastomotic leakage. One fatality (Clavien–Dindo grade 5), and the need for conversion in another case were described [37]. Anderson et al. listed a complication rate of 20%, though with no further specifications ($n = 1$) [24]. Robinsons' primary focus was on a typical urgent operation: perforated gastrojejunal ulcers. He reported noninferiority to the laparoscopic approach but dramatically higher costs for robotic operations. None of the case reports/case series reported any complications.

Hernias/miscellaneous

SRs are commonly used in hernia surgery; however, the urgency of these operations arises from the incarceration of the hernia, which is often associated with intestinal obstruction. We identified eight reports of urgent robotic operations for hernias (Table 4). Four studies retrospectively analyzed urgent hernia operations, two of which were controlled studies [25, 40, 41]. In addition, we identified a database analysis that included urgent robotic hiatal hernia operations [27]. A rare indication for urgent robotic surgery was the operation of posttraumatic splenic bleeding reported by Giulianotti et al. [15]. No reports on the use of an SR in adhesive intestinal obstruction were found.

For urgent robotic hernia operations, the published complication rates were very heterogeneous; the database analysis of Hosein et al. reported one death (0.1%) and an overall complication rate of 2% [27]. Muysoms et al. reported 3.5% minor and 0.2% major complications [41]. Kudsi et al. published a complication rate of 36.6% (23% minor and 13% major complications); however, this rate was significantly lower than that in the open surgery subgroup of Clavien–Dindo grade ≥ 3 complications [26]. A further

Table 1 Appendicitis

References	Study design		Outcome			Further notes			
	Design	Study period	<i>n</i> (urgent, robotic)	<i>n</i> (non-urgent)	Primary study objective		Robot	Differentiated urgent vs. non-urgent	Complications conversions
Cadière et al. [16]	R-NRNC	3/1997–2/2001	1	–	Feasibility	Da Vinci	No	No complications	See also <i>cholecystitis</i>
Keikar et al. [22]	P-NRNC	3/2019–4/2019	4	145	Feasibility	Versius	No	No conversions	
Kibar et al. [51]	CR	n.d	1	26	–	Da Vinci	Yes	No complications	Appendicovesical fistula
Mittal et al. [21]	P-NRNC	n.d	22	–	Feasibility/usefulness	FreeHand RCM	No	No complications	Liberation of the surgical assistant
Ohmura et al. [14]	R-NRC	12/2014–3/2017	163	3	feasibility	Soloassist RCM	No	“No device-related complications”	See also <i>cholecystitis</i> and <i>hollow organs</i>
Yi et al. [12]	CS	3/2014	2	783	Feasibility	Micro Hand S	Yes	No conversions	See also hollow organs
Yi et al. [13]	CS	4/2014–5/2014	3	1	Feasibility	Micro Hand S	No	No conversions	See also <i>cholecystitis</i> and <i>hollow organs</i>

CR Case report, CS Case series, n.d. not documented, P-NRNC prospective, non-randomized, non-controlled, R-NRC retrospective, non-randomized, controlled, R-NRNC retrospective, non-randomized, non-controlled, RCM Robotic camera mount

Table 2 Cholecystitis

References	Study design			Outcome			Further notes	
	Design	Study period	<i>n</i> (urgent, robotic) <i>n</i> (non-urgent) <i>n</i> (other)	Primary study objective	Robot	Differentiated urgent vs. non-urgent		Complications conversions
Ayloo et al. [52]	R-NRC	9/2005–6/2012	45 134 147	MIRC vs. MILC	Da Vinci	No	1.7% CD ≥ 3; 1.7% CD < 3” no conversions	3 conversions in MILC; OT longer in MIRC
Balachandran et al. [53]	R-NRC	10/2011–7/2014	76 339 263	SIRC vs. MILC	Da Vinci	No	1% ileus 0.2% bile leakage 2.9% to MILC, 3.2% to open	6.5% hernias in SIRC LOS shorter SIRC
Bibi et al. [33]	R-NRNC	6/2012–1/2013	31 71 –	Safety of SIRC	Da Vinci	No	4% CD < 3 3.9% conversions	50% of conversions due to inflammation
Buzad et al. [17]	P-NRC	1/2012–5/2012	2 18 10	SIRC vs. SILC	Da Vinci	No	No complications	No differences in OT and costs
Cadière et al. [16]	R-NRNC	3/1997–2/2001	4 44 98	Feasibility	Da Vinci	No	No conversion 25% (<i>n</i> = 1) CD 2	See also <i>appendicitis</i>
Chung et al. [29]	R-NRC	8/2013–1/2015	7 63 70	SILC vs. MILC	Da Vinci	No	no conversions n.d.	2.8% 30 d readmission
Daskalaki et al. [54]	R-NRNC	7/2011–2/2013	28 156 –	ICG Cholangiography	Da Vinci	No	1.4% conversion to open	cholangiography > 94%
Gangemi et al. [28]	R-NRC	2008–2015	130 546 289	Risk factors for conversion	n.d.	N.d.	1% CD ≥ 3; 2.2% CD < 3	no conversions
Giulianotti et al. [15]	R-NRNC	10/2000–11/2002	7 45 123	Descriptive; RC vs. LC	Da Vinci	No	0.15% (<i>n</i> = 1) conversions 1.9% (<i>n</i> = 1) complications n.d.	n.d.
Gonzalez et al. [55]	R-NRC	1/2012–9/2012	20 146 335	SILC vs. SIRC vs. SPIDER	Da Vinci	No	1.9% (<i>n</i> = 1) conversions 1.2% CD ≥ 3; 0.6% CD < 3	OT for SIRC longer than SILC/SPIDER incisions)

Table 2 (continued)

References	Study design			n		Primary study objective	Robot	Outcome		Further notes
	Design	Study period	n (urgent, robotic)	n (non-urgent)	n (other)			Differentiated urgent vs. non-urgent	Complications conversions	
Honaker et al. [56]	R-NRC	3/2013–2/2014	1	17	–	MIRC vs. MILC	Da Vinci	No	No complications	LOS shorter MIRC
Jeong Jang et al. [57]	R-NRNC	4/2019–8/2020	2 (5)	40	72 (69)	Feasibility of SIRC	Da Vinci	No	No conversions 1.35% CD 1	3 acute cholecystitis diagnosed intraoperatively
Kalteis et al. [58]	R-NRC	4/2002–11/2004	20	52	72	Safety and effectiveness	AESOP	No	No conversions 1.4% CD < 3	Liberation of the surgical assistant
Konstantinidis et al. [18]	P-NRNC	3/2011–7/2011	4	41	–	Feasibility and effectiveness	Da Vinci	No	2.2.% CD 3b, 5.3% < CD 3	20% intraoperative gallbladder-ruptures
Kornprat et al. [19]	P-NRC	2001–2006	2	18	–	MIRC vs. MILC	Zeus	No	No complications	OT longer in MIRC
Kubat et al. [23]	R-NRC	5/2012–8/2013	74	26	76	Urgent vs. non-urgent MIRC	Da Vinci	Yes	No conversions 12% complications in urgent, incl. 1.5% (n=1) BD injury 1.5% (n=1) conversions	LOS longer in urgent, overall complications + SSI no differences
Kulaylat et al. [36]	R-NRC	1/2015–12/2018	10	69	–	SIRC/MIRC vs. SILC/MILC	Da Vinci	No	“No differences in complications”	OT longer in robotic, costs higher in robotic
Li et al. [42]	R-NRC	8/2013–12/2015	17	220	61	Safety, resources	Da Vinci	No	No conversions 3.8% < CD 3 (20% in MILC)	1.9% conversion rate in MILC OT, LOS and costs higher in SIRC
Mattei et al. [59]	R-NRNC	2013–?	1	367	19	Feasibility in pediatric patients	Da Vinci	No	No conversions 20% seromas (CD < 3)	OT longer, LOS shorter
Ohmura et al. [14]	R-NRC	12/2014–3/2017	101	172	–	Feasibility	Soloassist RCM	No	No conversions “No device-related complications”	LOS shorter; liberation of surgical assistant
					848				No conversions	See also <i>appendix</i> and <i>hollow organs</i>

Table 2 (continued)

References	Study design		Primary study objective		Robot	Outcome		Further notes
	Design	Study period	<i>n</i> (urgent, robotic)	<i>n</i> (non-urgent) <i>n</i> (other)		Differentiated urgent vs. non-urgent	Complications conversions	
Rosales-Velderrain et al. [60]	P-NRNC	3/2013–5/2015	4	10	Da Vinci	No	7% (<i>n</i> = 1) seroma No conversions	
Schertz et al. [35]	R-NRC	8/2013–4/2018	3/6	101/99	Da Vinci	No	1 enterotomy (MIRC), 2.9% hernias (SIRC)	OT, LOS in SIRC shorter
Su et al. [61]	R-NRC	2/2014–9/2015	10	41	Da Vinci	No	No complications No conversions	OT in urgent longer Pain lower in SIRC
Svoboda et al. [20]	P-NRC	11/2012–2/2014	159	112	Da Vinci	No	0.9% (<i>n</i> = 1), incisional hernia no conversions	OT longer in obese patients
Tao et al. [62]	R-NRC	1/2006–2/2020	13	161	Da Vinci	No	12.3% overall in RC No conversion	RC vs. LC CD 1 >, CD 2 <, CD 3 equal, no CD IV
Vidovszky et al. [34]	P-NRNC	1/2012–1/2013	13	82	Da Vinci	No	4.2% CD 3; 2.1% < CD 3 8.4%; 1 due to inflammation	One disruption of DC 5 inflammations diagnosed intra-operatively
Case Series Case Reports	CS/CR	2001–2021	11	–	Da Vinci (3), Microhand S (1)	No	No complications No conversions	Bustos et al. [31]; Hanisch et al. [30]; Milone et al. [32]; Yi et al. [13]

Only statistically significant differences listed

BMI Body mass index, *CD* Clavien-Dindo classification of complications [8], *CR* Case report, *CS* Case series, *DC* Ductus cysticus, *LC* laparoscopic cholecystectomy, *LOS* length of stay, *MILC* Multi-incision laparoscopic cholecystectomy, *MILC* Multi-incision robotic cholecystectomy, *n.d.* not documented, *OT* Operation Time, *P-NRC* prospective, non-randomized, controlled, *P-NRNC* prospective, non-randomized, non-controlled, *RC* Robotic cholecystectomy, *R-NRC* retrospective, non-randomized, controlled, *R-NRNC* retrospective, non-randomized, non-controlled, *RCM* robotic camera mount, *SI* Single-Incision, *SILC* Single-Incision laparoscopic cholecystectomy, *SIRC* Single-Incision robotic cholecystectomy

Table 3 Gastrointestinal procedures

References	Study design		Outcome			Further notes			
	Design	Study period	<i>n</i> (urgent, robotic)	<i>n</i> (non-urgent) <i>n</i> (other)	primary study objective		Robot	Differentiated urgent vs. non-urgent	Complications conversions
Anderson et al. [24]	R-NRC	2/2015–2/2017	6	–	Urgent subtotal colectomy: robotic vs. Laparoscopic	Da Vinci	Yes	20% (<i>n</i> =1) CD 2 No conversions	OT longer in robotic 2 dockings
Beltzer et al. [37]	R-NRC	10/2013–11/2018	2	58 46	Diverticular disease: robotic vs. Laparoscopic	Da Vinci	No	15% cd ≥ 3; 15% < 3 1.7% (<i>n</i> =1) conversions	Length of postop ileus in robotic group shorter
Felli et al. [63]	CR	n.d	1	–	Hemicolectomy for hemorrhagic cancer	Da Vinci	Yes	No complications No conversion	Double-barreled ileocolostomy; oncologic resection
Guerra et al. [64]	CR	n.d	1	–	Acute large bowel malignant obstruction	Da Vinci	Yes	No complications No conversions	Oncological resection of a splenic flexure tumor
Kudsi et al. [9–11]	3 CR	2019–2020	3	–	3 colon resections	Da Vinci	Yes	No complications No conversions	All with hand sutured anastomosis
Ohmura et al. [14]	R-NRC	12/2014–3/2017	16	–	Feasibility	Soloassist RCM	No	“No device-related complications” No conversions	LOS shorter liberation of surgical assistant
Pedraza et al. [65]	CR	n.d	1	933	Iatrogenic colonic perforation;	Da Vinci	Yes	No complications	Primary repair
Robinson et al. [38]	R-NRC	2015–2019	24	–	Perforated gastroduodenal ulcer	Da Vinci	Yes	8.3%, median cd-score 2.2 No conversions	Complications similar to laparoscopic group
Sudan et al. [66]	CR	n.d	1	–	Duodenal stump insufficiency	Da Vinci	Yes	No complications	5 d after BPD-DS, biliary peritonitis
Sun et al. [67]	CR	3/2014	1	–	Gastric perforations	Microhand S	Yes	No complications	
Yi et al. [12]	CS	3/2014	1	–	Repair of gastric perforation	Microhand S	No	No complications	See also appendicitis and cholecystitis
Yi et al. [13]	CS	4/2014–5/2014	1	–	Repair of gastric perforation	Microhand S	No	No complications	Withdrawal of 2 patients

Only statistically significant differences listed

BPD biliopancreatic diversion with duodenal switch, *CD* Clavien-Dindo classification of complications [8], *CDD* classification of diverticular disease [39], *CR* Case report, *CS* Case series, *LOS* length of stay, *n.d.* not documented, *OT* Operation time, *R-NRC* Retrospective, non-randomized, controlled

Table 4 Hernias/Miscellaneous

References	Study design		Outcome				Further notes		
	Design	Study period	n (urgent, robotic)	n (non-urgent) n (control)	Primary study objective	Robot		Differentiated urgent vs. non-urgent	Complications
Bou-Ayash et al. [25]	R-NRNC	2/2013–5/2020	19	–	Incarcerated inguinal hernia	Da Vinci	Yes	5.3% CD IVa (n=1; hypercarbia); 10.6% CD <3	
Cubas et al. [68]	CR	n.d	1	–	Incarcerated morgagni hernia	Da Vinci	Yes	No conversions No complication	
Ceccarelli et al. [69]	CS	12/2009–12/2019	3	–	Strangulated hiatal hernias	Da Vinci	Yes	No conversion 33% (n=1) CD 3	
Giulianotti et al. [15]	R-NRNC	10/2000–11/2002	1	2 lap 192	Feasibility; here: posttraumatic spleen hematomia	Da Vinci	No	No conversions No complications	One CD V in non-urgent patients
Hosein et al. [27]	database query	2015–2017	131	704 1488 open 6774 lap	Hiatal hernia repair: robotic vs. Lap. Vs. Open	n.d	No	2% overall complications; 0.1% (n=1) CD V n.d	More urgent and severe ill cases in open; robotic less complication than lap/open
Muhonen et al. [70]	CS	n.d	1	– 2 lap	Incarcerated paraduodenal hernia	Da Vinci	Yes	No complications No conversions	
Muysoms et al. [41]	R-NRNC	1/2016–12/2019	8	396 272	Robotic vs. Lap. Inguinal hernia	Da Vinci	No	3.5% CD II, 0.2% CD IIIb No conversions	Shorter hospital stay, higher costs in robotic group
Kudsi et al. [40]	R-NRNC	2013–2019	34	–	Incarcerated hernia robotic vs. Open	Da Vinci	Yes	23.3% < CD 3; 13.3% ≥ CD 3; 3.2% recurrence No conversions	OT shorter in open, more CD ≥ 3 in open, more SSI in open IPOM & TAPP
Smith et al. [71]	CR	n.d	1	–	Incarcerated inguinal hernia	Da Vinci	Yes	No complications No conversion	TAPP

Only statistically significant differences listed

CD Clavien-Dindo classification of complications, CS Case series, lap laparoscopic, IPOM intraperitoneal only mesh, n.d. not documented, OT operation time, R-NRNC non-randomized, controlled, R-NRNC retrospective, non-randomized, not controlled, SSI surgical site infections, TAPP transabdominal pre-peritoneal hernia repair

interesting outcome was the significantly shorter hospital stay but higher costs and the recurrence rate of 3.2% reported by Myosoms et al. [41]. All studies emphasized that the interventions were technically feasible using an SR.

Discussion

General considerations and assessment of evidence

To date, there is only limited research on urgent robotic operations in general surgery; therefore, the data available are still very limited. Against this background, we consider a review such as ours, which systematically collects and examines the existing data and presents it descriptively, to be as comprehensive as possible and to be valuable and necessary.

Our review gives clear indications that robotic surgery has not yet arrived in urgent general surgery on a larger scale. A randomized controlled trial has not yet been performed, which is not totally unexpected given that randomized trials dealing with emergency and urgent surgery are generally very rare. However, even the nonrandomized studies offer specific data only to a limited extent: only six studies named urgent operations as a main interest or important variable [23–27, 38]. This makes a systematic assessment of the evidence complicated and a meta-analysis practically impossible. It can be stated that there is only a low level of evidence regarding robotic surgery in the context of urgent operations. Nevertheless, we were able to identify and summarize a notable number of publications covering a wide range of diseases.

Case reports and series are sources with limited evidence and are not suitable for a meta-analysis. In the absence of sources with better evidence, these are nevertheless presented in our study, as these reports fulfill the function of documenting the technical feasibility of certain interventions in the sense of a proof of concept.

With the recently published position paper of the WSES, the topic of urgent robotic operations was highlighted for the first time. De'Angelis et al. also conducted an extensive literature search. The number of papers screened was comparable to our study, but de'Angelis et al. used only ten manuscripts for their analysis: five retrospective cohort studies, and five case reports/case series [4]. We decided against reducing the number of publications used through a stricter assessment since we wanted to provide a maximum amount of information about robotic emergency operations. Nevertheless, our presentation goes beyond a purely narrative review, as, under verifiable conditions, we offer the first complete overview of the published data. However, the conclusions the authors of the WSES statement published are very similar to our findings, though we covered a wider field of operations [4]. The combination of de'Angelis et al.

and our work creates, for the first time, a deeper impression of the significance of robotic operations in urgent surgery.

The main indications for robotic emergency interventions were gallbladder, hernias, and gastrointestinal surgery, as well as appendectomies. Some common aspects can be identified. First, there are supposedly higher costs of robotic interventions. Our review shows that robotic operations are significantly more expensive than laparoscopic surgeries [36, 38, 41, 42]. To date, there are no data that demonstrate the amortization of the extra costs by a reduction in the length of stay or complications that stem from the use of an SR. Moreover, the abovementioned studies that showed a cost increase per procedure did not take the considerable acquisition costs of the systems into account. The option of refunding by additional charges for the use of an SR is not possible in all health systems. Second, an often cited counterargument against robotic emergency interventions is the increased time requirements due to longer preparation or operation times. Here, the data were ambiguous and ranged from lengthening to shortening of time intervals with a tendency toward increased operative time. Minimally invasive surgery is primarily intended for stable patients, which of course also applies to robotic operations. Thus, even the moderate increases in operating time described above do not seem to be a contraindication for robotic operations.

Robotic camera mounts (RCM)

RCMs can be classified as robotic surgery only to a limited extent. However, they offer some of the alleged advantages, in particular a stable image and ergonomic advantages. However, there are also might be disadvantages, e.g., complications due to technical malfunctions.

According to this review, this type of surgery seems to be not very widespread, especially in urgent surgery, as only two publications have described the use of these systems [14, 21].

No clear statements about the advantages and disadvantages of RCMs can be drawn from the data presented here, but it can be stated that this type of surgery is undoubtedly more similar to laparoscopic than to robotic surgery. However, it seems important that no RCM-associated complications are reported.

Appendectomies

Appendectomy is a very frequent operation, although there have been very few studies that deal with robotic operations for acute appendicitis. This is of particular interest since appendicitis was one of the diseases that significantly led to the development of modern laparoscopy, initially starting with the confirmation of the diagnosis and the first laparoscopic appendectomy by Semm in 1983 [43]. Today,

laparoscopic appendectomy is the therapy of choice for appendicitis [2].

There may be various reasons for the lack of studies on SR for appendicitis: The presumed high costs of robotic operations have been discussed since the start of robotic surgery [30]. Most likely, a “minor” procedure such as an appendectomy with correspondingly low remuneration is estimated as economically unreasonable and will therefore not become part of clinical routine for robotic surgery. It is also conceivable that the advantages of robotic operations for an often rather simple operation without complex preparation are not regarded as sufficient to implement this technique. An urgent appendectomy with the Da Vinci[®] system was reported in only two cases, while this robot system has been generally used most often. The feasibility of appendectomies with the Da Vinci[®] system was shown by several studies reporting on appendectomies in the context of other operations [44, 45].

An option to make a robotic appendectomy economically worth considering is to overthink the closure of the appendix stump: the use of the stapler or clip applicator for the Da Vinci[®] system causes relevant costs. This could be circumvented by closing the stump with a Roeder loop or suture. However, a current meta-analysis showed that stump closure with staples is superior in terms of postoperative complications [46].

In the overall view, appendectomy via SR is technically possible, but the additional expense compared to laparoscopic appendectomy most likely cannot be justified.

Cholecystectomies

Regarding this indication, the number of published studies is better, although far from satisfactory. However, against the background of more available studies, a case report analysis was less necessary. In addition to the fact that one of these case reports was among the first publications on robotic surgery in general surgery (Hanisch et al.), these reports did not provide any crucial information [30]. The statements regarding the advantages and disadvantages of the robotic approach in urgent cholecystectomies were very heterogeneous without a clear trend.

Remarkably, many studies compared single-incision robotic cholecystectomy (SIRC) with nonrobotic surgical procedures. Therefore, we assume a reporting bias: multi-incision robotic cholecystectomy (MIRC) was primarily examined in studies that were published before 2010, and acute cholecystitis was often an exclusion criterion [19]. However, the scientific perception of robotic cholecystectomy may have changed in two aspects: MIRC appears to be such a standard procedure that it is less examined in studies. At the same time, and based on their increased robotic experience, more researchers have opted to include acute

cholecystectomy in their studies. In our estimation, these interventions are currently often performed as SIRC since this technique is assumed to be more innovative. Furthermore, there is another inaccuracy in the analysis: a number of studies reported robotic surgery for acute cholecystitis, with acute inflammation diagnosed intraoperatively. Therefore, this might not be labeled “urgent surgery” correctly in the narrower sense. The subject of this analysis is not the question of single- vs. multi-incision operations. The currently most up-to-date Practice Guideline on Safe Cholecystectomy votes for MIRC, in particular due to an increased rate of biliary tract injuries in the single-incision group [47]. Regarding the literature analyzed in our study, there was one biliary tract injury during an operation for acute cholecystitis using the SIRC technique. The data on complications and conversions were heterogeneous, as were the statements on the possible advantages and disadvantages of robotic operations. Biliary tract problems were reported in three patients, which resulted in a biliary complication rate of 0.4% among all patients operated on with an SR, which was within the expected range. Except for one study, these complications were not differentiated between urgent and elective operations. Notably, Kulaylat et al. reported an increase in hospital costs of 38% for the robotic procedure compared to laparoscopic cholecystectomy [36]. In summary, at the moment, neither the advantages nor the risks of robotic operations in urgent, acute cholecystectomies can be adequately assessed. However, no clear contraindications for the use of robots in this situation were found in this review.

Gastrointestinal procedures

The advantage of robotic surgery in acute conditions of the gastrointestinal tract, especially perforations, seems obvious: since a robot provides a significant advantage for suturing and tying knots, perforations and ruptures can be closed excellently. Furthermore, robotic operations for benign and malignant diseases of the colon or the upper gastrointestinal tract are clinical routine, with an increasing proportion being performed in the elective setting [48]. Our review shows that for urgent operations, there are hardly any data available. However, our own experience and the case reports/series listed demonstrate that these interventions were definitely possible in an urgent setting.

Beltzer et al. summarized that there was no advantage in using the robot for surgical procedures in diverticular disease [37]. Anderson et al. also did not state any explicit advantages and reported a longer operation time [24]. There are currently no data that propagate the use of SR in this indication group for a better outcome in a population operated urgently.

Robinson et al. presented a study primarily focused on urgent robotic surgery for perforated gastrojejunal perforations

[38]. The authors were able to show that robotic surgery was not inferior to laparoscopic surgery. In addition, they reported two interesting aspects that otherwise hardly received attention: (1) the immediate preparation time in the OR was even shorter than that with laparoscopic operations; and (2) 54% of the operations were performed on weekends or during the evening, night or early morning. This invalidates some of the arguments that have been put forward against robotic emergency operations.

As mentioned above, in regard to surgery of the gastrointestinal tract, the cost disadvantages of SRs matter again: stapler and sealing devices in addition to draping, scissors, forceps, etc., are significant cost factors. Schiergens et al. showed that the use of an SR for an elective sigma resection increases the cost of surgical supplies by more than a factor of 4 compared to open surgery and by more than a factor of 2 compared to laparoscopic surgery. These numbers were very comparable to the cost increase reported by Robison et al. for urgent gastrojejunal ulcers [38, 49].

Hernias/miscellaneous

SRs are used regularly in minimally invasive hernia surgery [3]. This may contribute to the fact that there are quite a few studies on urgent robotic hernia surgery. Even if the evidence cannot be described as satisfactory, the overall picture is similar to those of the other indication groups: robotic interventions are feasible in incarcerated hernias and in urgent situations. The complication and conversion rates were low and comparable to those of nonrobotic, minimally invasive procedures. To date, however, no clear advantages of the robotic technique have been demonstrated.

The situation is different in hiatal hernias: in most cases, hiatal hernias are technically more demanding than inguinal or ventral hernias. The sutures required to reconstruct the hiatus and to create the fundoplication as well as the preparation make this operation ideal for the use of an SR. Current studies showed fewer complications and a shorter hospital stay for robotic hiatus hernia surgery in the elective setting [50]. Even if no data are available for urgent hiatoplasty, it is quite likely that such effects can be observed here.

The report of a splenectomy for bleeding (along with several reports of elective splenectomies) indicated that this operation is also possible with an SR. If the patient's circulatory system remains stable and a minimally invasive procedure is conceivable, the procedure can also be carried out with an SR if the appropriate expertise is available.

Conclusion

A particular value of our work is that it provides a well-founded summary of the existing data for surgeons and researchers who are interested in urgent abdominal robotic surgery. Based on this, the specific studies needed can be initiated in the future. Furthermore, our review may help to establish framework conditions for a register for urgent robotic interventions.

With this review, we provide the most complete overview of the current literature on robotic surgery for urgent general surgical operations. Our analysis of the literature gives the impression that, in particular, robotic cholecystectomies in acute cholecystitis and operations on impacted hernias are feasible and rational and can be carried out without increased risk. Initial studies on urgent robotic operations in the gastrointestinal tract have thus far shown ambivalent results but have been proven technically feasible. High costs remain a significant burden for these operations.

Appendix 1

Search algorithm

```
#1 (“robot*”[MeSH Terms] AND “robot*”[Title/
Abstract]) OR “robotic surgery” OR “robot-assisted” OR
“robot assisted”
#2 #1 AND adhesiolysis
#3 #1 AND (appendicitis OR append*)
#4 #1 AND bowel obstruction
#5 #1 AND (cholecystitis OR cholecystectomy OR
cholecyst*)
#6 #1 AND diverticular disease
#7 #1 AND diverticulitis
#8 #1 AND “emergency”
#9 #1 AND “emergencies”
#10 #1 AND hernia
#11 #1 AND (incarcerated OR incarcerat*)
#12 #1 AND “ischaemia”
#13 #1 AND perforation
#14 #1 AND peritonitis
#15 #1 AND ulc*
#16 #1 AND urgency
#17 #1 AND urgent
#2 OR #3 OR #4 OR #5 OR #6 OR #7 OR #8 OR #9
OR #10 OR #11 OR #12 OR #13 OR #14 OR #15 OR #16
OR #17
```

Author contributions All authors contributed to the study conception and design. Data acquisition was performed by AR and JL, and analysis and interpretation of the data were performed by AR, JL, WP and FU. Critical revision was performed by WP and FU. The first draft of the manuscript was written by AR and JL, and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Funding Open Access funding enabled and organized by Projekt DEAL. The authors declare that no funds, grants or other support was received during the preparation of this manuscript.

Declarations

Conflict of interest Alexander Reinisch declares no financial or non-financial interests to disclose. Juliane Liese declares no financial or nonfinancial interests to disclose. Winfried Padberg declares no financial or nonfinancial interests to disclose. Frank Ulrich declares no financial or nonfinancial interests to disclose.

Ethical approval Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Wakabayashi G, Iwashita Y, Hibi T, Takada T, Strasberg SM, Asbun HJ, Endo I, Umezawa A, Asai K, Suzuki K, Mori Y, Okamoto K, Pitt HA, Han HS, Hwang TL, Yoon YS, Yoon DS, Choi IS, Huang WS, Giménez ME, Garden OJ, Gouma DJ, Belli G, Dervenis C, Jagannath P, Chan ACW, Lau WY, Liu KH, Su CH, Misawa T, Nakamura M, Horiguchi A, Tagaya N, Fujioka S, Higuchi R, Shikata S, Noguchi Y, Ukai T, Yokoe M, Cherqui D, Honda G, Sugioka A, de Santibañes E, Supe AN, Tokumura H, Kimura T, Yoshida M, Mayumi T, Kitano S, Inomata M, Hirata K, Sumiyama Y, Inui K, Yamamoto M (2018) Tokyo guidelines 2018: surgical management of acute cholecystitis: safe steps in laparoscopic cholecystectomy for acute cholecystitis (with videos). *J Hepatobiliary Pancreat Sci* 25(1):73–86. <https://doi.org/10.1002/jhbp.517>
- Gorter RR, Eker HH, Gorter-Stam MA, Abis GS, Acharya A, Ankersmit M, Antoniou SA, Arolfo S, Babic B, Boni L, Bruntink M, van Dam DA, Defoort B, Deijen CL, DeLacy FB, Go PM, Harmsen AM, van den Helder RS, Iordache F, Ket JC, Muysoms FE, Ozmen MM, Papoulas M, Rhodes M, Straatman J, Tenhagen M, Turrado V, Vereczkei A, Vilallonga R, Deelder JD, Bonjer J (2016) Diagnosis and management of acute appendicitis. EAES consensus development conference 2015. *Surg Endosc* 30(11):4668–90. <https://doi.org/10.1007/s00464-016-5245-7>
- Sheetz KH, Claffin J, Dimick JB (2020) Trends in the adoption of robotic surgery for common surgical procedures. *JAMA Netw Open* 3(1):e1918911. <https://doi.org/10.1001/jamanetworkopen.2019.18911>
- de Angelis N, Khan J, Marchegiani F, Bianchi G, Aisoni F, Alberti D, Ansaloni L, Biffi W, Chiara O, Ceccarelli G, Coccolini F, Cicuttin E, D'Hondt M, Di Saverio S, Diana M, De Simone B, Espin-Basany E, Fichtner-Feigl S, Kashuk J, Kouwenhoven E, Leppaniemi A, Beghdadi N, Memeo R, Milone M, Moore E, Peitzmann A, Pessaux P, Pikoulis M, Pisano M, Ris F, Sartelli M, Spinoglio G, Sugrue M, Tan E, Gavriilidis P, Weber D, Kluger Y, Catena F (2022) Robotic surgery in emergency setting: 2021 WSES position paper. *World J Emerg Surg* 17(1):4. <https://doi.org/10.1186/s13017-022-00410-6>
- Ng KT, Tsia AKV, Chong VYL (2019) Robotic versus conventional laparoscopic surgery for colorectal cancer: a systematic review and meta-analysis with trial sequential analysis. *World J Surg* 43(4):1146–1161. <https://doi.org/10.1007/s00268-018-04896-7>
- Tan A, Ashrafian H, Scott AJ, Mason SE, Harling L, Athanasiou T, Darzi A (2016) Robotic surgery: disruptive innovation or unfulfilled promise? A systematic review and meta-analysis of the first 30 years. *Surg Endosc* 30(10):4330–52. <https://doi.org/10.1007/s00464-016-4752-x>
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J, Moher D (2009) The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 339:b2700. <https://doi.org/10.1136/bmj.b2700>
- Dindo D, Demartines N, Clavien PA (2004) Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 240(2):205–213. <https://doi.org/10.1097/01.sla.0000133083.54934.ae>
- Kudsi OY, Gokcal F (2019) Urgent robotic mesocolic excision for obstructing proximal transverse colon cancer—a video vignette. *Colorectal Dis* 21(9):1093–1094. <https://doi.org/10.1111/codi.14714>
- Kudsi OY, Bou-Ayash N (2020) Bleeding sigmoid diverticulosis—urgent stapless totally robotic sigmoidectomy—a video vignette. *Colorectal Dis* 22(9):1205. <https://doi.org/10.1111/codi.15051>
- Kudsi OY, Bou-Ayash N (2020) Caecal volvulus—urgent totally robotic right colectomy—a video vignette. *Colorectal Dis* 22(10):1448–1449. <https://doi.org/10.1111/codi.15068>
- Yi B, Wang G, Li J, Jiang J, Son Z, Su H, Zhu S (2016) The first clinical use of domestically produced Chinese minimally invasive surgical robot system “micro hand S.” *Surg Endosc* 30(6):2649–55. <https://doi.org/10.1007/s00464-015-4506-1>
- Yi B, Wang G, Li J, Jiang J, Son Z, Su H, Zhu S, Wang S (2017) Domestically produced Chinese minimally invasive surgical robot system “micro hand S” is applied to clinical surgery preliminarily in China. *Surg Endosc* 31(1):487–93. <https://doi.org/10.1007/s00464-016-4945-3>
- Ohmura Y, Nakagawa M, Suzuki H, Kotani K, Teramoto A (2018) Feasibility and usefulness of a joystick-guided robotic scope holder (soloassist) in laparoscopic surgery. *Visc Med* 34(1):37–44. <https://doi.org/10.1159/000485524>
- Giulianotti PC, Coratti A, Angelini M, Sbrana F, Cecconi S, Balestracci T, Caravaglios G (2003) Robotics in general surgery: personal experience in a large community hospital. *Arch Surg* 138(7):777–84. <https://doi.org/10.1001/archsurg.138.7.777>
- Cadière GB, Himpens J, Germay O, Izizaw R, Degueudre M, Vandromme J, Capelluto E, Bruyns J (2001) Feasibility of robotic laparoscopic surgery: 146 cases. *World J Surg* 25(11):1467–1477. <https://doi.org/10.1007/s00268-001-0132-2>
- Buzad FA, Corne LM, Brown TC, Fagin RS, Hebert AE, Kaczmarek CA, Pack AN, Payne TN (2013) Single-site robotic

- cholecystectomy: efficiency and cost analysis. *Int J Med Robot* 9(3):365–370. <https://doi.org/10.1002/rcs.1507>
18. Konstantinidis KM, Hirides P, Hirides S, Chrysocheris P, Georgiou M (2012) Cholecystectomy using a novel Single-Site(®) robotic platform: early experience from 45 consecutive cases. *Surg Endosc* 26(9):2687–94. <https://doi.org/10.1007/s00464-012-2227-2>
 19. Kornprat P, Werkgartner G, Cerwenka H, Bacher H, El-Shabrawi A, Rehak P, Mischinger HJ (2006) Prospective study comparing standard and robotically assisted laparoscopic cholecystectomy. *Langenbecks Arch Surg* 391(3):216–221. <https://doi.org/10.1007/s00423-006-0046-4>
 20. Svoboda S, Qaqish TR, Wilson A, Park H, Youssef Y (2015) Robotic single-site cholecystectomy in the obese: outcomes from a single institution. *Surg Obes Relat Dis* 11(4):882–885. <https://doi.org/10.1016/j.soard.2014.11.016>
 21. Mittal R, Sbaih M, Motson RW, Arulampalam T (2020) Use of a robotic camera holder (FreeHand). *Minim Invasive Ther Allied Technol* 29(1):56–60. <https://doi.org/10.1080/13645706.2019.1576052>
 22. Kelkar D, Borse MA, Godbole GP, Kurlekar U, Slack M (2020) Interim safety analysis of the first-in-human clinical trial of the Versius surgical system, a new robot-assisted device for use in minimal access surgery. *Surg Endosc*. <https://doi.org/10.1007/s00464-020-08014-4>
 23. Kubat E, Hansen N, Nguyen H, Wren SM, Eisenberg D (2016) Urgent and elective robotic single-site cholecystectomy: analysis and learning curve of 150 consecutive cases. *J Laparoendosc Adv Surg Tech A* 26(3):185–191. <https://doi.org/10.1089/lap.2015.0528>
 24. Anderson M, Lynn P, Aydinli HH, Schwartzberg D, Bernstein M, Grucela A (2020) Early experience with urgent robotic subtotal colectomy for severe acute ulcerative colitis has comparable perioperative outcomes to laparoscopic surgery. *J Robot Surg* 14(2):249–253. <https://doi.org/10.1007/s11701-019-00968-5>
 25. Bou-Ayash N, Gokcal F, Kudsi OY (2020) Robotic inguinal hernia repair for incarcerated hernias. *J Laparoendosc Adv Surg Tech A*. <https://doi.org/10.1089/lap.2020.0607>
 26. Kudsi OY, Gokcal F, Bou-Ayash N, Chang K (2020) Comparison of midterm outcomes between open and robotic emergent ventral hernia repair. *Surg Innov*. <https://doi.org/10.1177/1553350620971182>
 27. Hosein S, Carlson T, Flores L, Armijo PR, Oleynikov D (2021) Minimally invasive approach to hiatal hernia repair is superior to open, even in the emergent setting: a large national database analysis. *Surg Endosc* 35(1):423–8. <https://doi.org/10.1007/s00464-020-07404-y>
 28. Gangemi A, Danilkowicz R, Bianco F, Masrur M, Giulianotti PC (2017) Risk factors for open conversion in minimally invasive cholecystectomy. *JSLs*. <https://doi.org/10.4293/JSLs.2017.00062>
 29. Chung PJ, Huang R, Policastro L, Lee R, Schwartzman A, Alfonso A, Sugiyama G (2015) Single-site robotic cholecystectomy at an inner-city academic center. *JSLs*. <https://doi.org/10.4293/JSLs.2015.00033>
 30. Hanisch E, Markus B, Gutt C, Schmandra TC, Encke A (2001) Robot-assisted laparoscopic cholecystectomy and fundoplication—initial experiences with the Da Vinci system. *Chirurg* 72(3):286–288. <https://doi.org/10.1007/s001040051307>
 31. Bustos R, Mashbari H, Gangemi A (2019) First report of gallbladder volvulus managed with a robotic approach. *Case Rep Surg* 2019:2189890. <https://doi.org/10.1155/2019/2189890>
 32. Milone M, Vertaldi S, Bracale U, D’Ambra M, Cassese G, Manigrasso M, De Palma GD (2019) Robotic cholecystectomy for acute cholecystitis: three case reports. *Medicine (Baltimore)* 98(30):e16010. <https://doi.org/10.1097/MD.00000000000016010>
 33. Bibi S, Rahnamai-Azar AA, Coralic J, Bayoumi M, Khorsand J, Farkas DT, Prasad LM (2015) Single-site robotic cholecystectomy: the timeline of progress. *World J Surg* 39(10):2386–2391. <https://doi.org/10.1007/s00268-015-3135-0>
 34. Vidovszky TJ, Carr AD, Farinholt GN, Ho HS, Smith WH, Ali MR (2014) Single-site robotic cholecystectomy in a broadly inclusive patient population: a prospective study. *Ann Surg* 260(1):134–141. <https://doi.org/10.1097/SLA.0000000000000295>
 35. Schertz P, Misra S, Livert D, Mulligan J, Rohatgi C (2019) Comparison of intraoperative outcomes between single-incision robotic cholecystectomy and multi-incision robotic cholecystectomy. *Cureus* 11(8):e5386. <https://doi.org/10.7759/cureus.5386>
 36. Kulaylat AN, Richards H, Yada K, Coyle D, Shelby R, Onwuka AJ, Aldrink JH, Diefenbach KA, Michalsky MP (2020) Comparative analysis of robotic-assisted versus laparoscopic cholecystectomy in pediatric patients. *J Pediatr Surg*. <https://doi.org/10.1016/j.jpedsurg.2020.11.013>
 37. Beltzer C, Knoerzer L, Bachmann R, Axt S, Dippel H, Schmidt R (2019) Robotic versus laparoscopic sigmoid resection for diverticular disease: a single-center experience of 106 cases. *J Laparoendosc Adv Surg Tech A* 29(11):1451–1455. <https://doi.org/10.1089/lap.2019.0451>
 38. Robinson TD, Sheehan JC, Patel PB, Marthy AG, Zaman JA, Singh TP (2022) Emergent robotic versus laparoscopic surgery for perforated gastrojejunal ulcers: a retrospective cohort study of 44 patients. *Surg Endosc* 36(2):1573–1577. <https://doi.org/10.1007/s00464-021-08447-5>
 39. Lembecke B (2015) Diagnosis, differential diagnoses, and classification of diverticular disease. *Viszeralmedizin* 31(2):95–102. <https://doi.org/10.1159/000380833>
 40. Kudsi OY, Bou-Ayash N, Chang K, Gokcal F (2020) Perioperative and midterm outcomes of emergent robotic repair of incarcerated ventral and incisional hernia. *J Robot Surg*. <https://doi.org/10.1007/s11701-020-01130-2>
 41. Muysoms F, Vierstraete M, Nachtergaele F, Van Garsse S, Pletinckx P, Ramaswamy A (2021) Economic assessment of starting robot-assisted laparoscopic inguinal hernia repair in a single-centre retrospective comparative study: the EASTER study. *BJS Open*. <https://doi.org/10.1093/bjsopen/zraa046>
 42. Li YP, Wang SN, Lee KT (2017) Robotic versus conventional laparoscopic cholecystectomy: a comparative study of medical resource utilization and clinical outcomes. *Kaohsiung J Med Sci* 33(4):201–206. <https://doi.org/10.1016/j.kjms.2017.01.010>
 43. Semm K (1983) Endoscopic appendectomy. *Endoscopy* 15(2):59–64. <https://doi.org/10.1055/s-2007-1021466>
 44. Hüttenbrink C, Hatiboglu G, Simpfendorfer T, Radtke JP, Becker R, Teber D, Hadaschik B, Pahernik S, Hohenfellner M (2018) Incidental appendectomy during robotic laparoscopic prostatectomy—safe and worth to perform? *Langenbecks Arch Surg* 403(2):265–269. <https://doi.org/10.1007/s00423-017-1630-5>
 45. Akl MN, Magrina JF, Kho RM, Magtibay PM (2008) Robotic appendectomy in gynaecological surgery: technique and pathological findings. *Int J Med Robot* 4(3):210–213. <https://doi.org/10.1002/rcs.198>
 46. Mannu GS, Sudul MK, Bettencourt-Silva JH, Cumber E, Li F, Clark AB, Loke YK (2017) Closure methods of the appendix stump for complications during laparoscopic appendectomy. *Cochrane Database Syst Rev* 11:CD006437. <https://doi.org/10.1002/14651858.CD006437.pub3>
 47. Brunt LM, Deziel DJ, Telem DA, Strasberg SM, Aggarwal R, Asbun H, Bonjer J, McDonald M, Alseidi A, Ujiki M, Riall TS, Hamill C, Moulton CA, Pucher PH, Parks RW, Ansari MT, Connor S, Dirks RC, Anderson B, Altieri MS, Tsamalaidze L, Stefanidis D, Group tPoBDICW (2020) Safe cholecystectomy multi-society practice guideline and state of the art consensus

- conference on prevention of bile duct injury during cholecystectomy. *Ann Surg* 272(1):3–23. <https://doi.org/10.1097/SLA.00000000000003791>
48. Konstantinidis IT, Ituarte P, Woo Y, Warner SG, Melstrom K, Kim J, Singh G, Lee B, Fong Y, Melstrom LG (2020) Trends and outcomes of robotic surgery for gastrointestinal (GI) cancers in the USA: maintaining perioperative and oncologic safety. *Surg Endosc* 34(11):4932–4942. <https://doi.org/10.1007/s00464-019-07284-x>
 49. Schiergens TS, Siemel WGR, Werner J (2020) Implementation of new technologies in the DRG system under economic aspects. *Chirurg* 91(7):561–566. <https://doi.org/10.1007/s00104-020-01184-w>
 50. Soliman BG, Nguyen DT, Chan EY, Chihara RK, Meisenbach LM, Graviss EA, Kim MP (2020) Robot-assisted hiatal hernia repair demonstrates favorable short-term outcomes compared to laparoscopic hiatal hernia repair. *Surg Endosc* 34(6):2495–502. <https://doi.org/10.1007/s00464-019-07055-8>
 51. Kibar Y, Yalcin S, Kopru B, Kaya E, Topuz B, Ebiloglu T (2016) Robot-assisted laparoscopic repair of spontaneous appendicovesical fistula. *J Endourol Case Rep* 2(1):111–113. <https://doi.org/10.1089/cren.2016.0063>
 52. Ayloo S, Roh Y, Choudhury N (2014) Laparoscopic versus robot-assisted cholecystectomy: a retrospective cohort study. *Int J Surg* 12(10):1077–1081. <https://doi.org/10.1016/j.ijssu.2014.08.405>
 53. Balachandran B, Hufford TA, Mustafa T, Kochar K, Sulo S, Khorsand J (2017) A comparative study of outcomes between single-site robotic and multi-port laparoscopic cholecystectomy: an experience from a tertiary care center. *World J Surg* 41(5):1246–1253. <https://doi.org/10.1007/s00268-016-3799-0>
 54. Daskalaki D, Fernandes E, Wang X, Bianco FM, Elli EF, Ayloo S, Masrur M, Milone L, Giulianotti PC (2014) Indocyanine green (ICG) fluorescent cholangiography during robotic cholecystectomy: results of 184 consecutive cases in a single institution. *Surg Innov* 21(6):615–621. <https://doi.org/10.1177/1553350614524839>
 55. Gonzalez AM, Rabaza JR, Donkor C, Romero RJ, Kosanovic R, Verdeja JC (2013) Single-incision cholecystectomy: a comparative study of standard laparoscopic, robotic, and SPIDER platforms. *Surg Endosc* 27(12):4524–31. <https://doi.org/10.1007/s00464-013-3105-2>
 56. Honaker MD, Paton BL, Stefanidis D, Schiffern LM (2015) Can robotic surgery be done efficiently while training residents? *J Surg Educ* 72(3):377–380. <https://doi.org/10.1016/j.jsurg.2014.11.008>
 57. Jeong Jang E, Kang SH, Kim KW (2021) Early outcomes of robotic single site cholecystectomy using the DaVinci Xi® system. *JLSLS*. <https://doi.org/10.4293/JLSLS.2020.00082>
 58. Kalteis M, Pistrich R, Schimetta W, Pölz W (2007) Laparoscopic cholecystectomy as solo surgery with the aid of a robotic camera holder: a case-control study. *Surg Laparosc Endosc Percutan Tech* 17(4):277–282. <https://doi.org/10.1097/SLE.0b013e31806030ae>
 59. Mattei P (2018) Single-site robotic-assisted laparoscopic cholecystectomy in children and adolescents: a report of 20 cases. *Surg Endosc* 32(5):2402–8. <https://doi.org/10.1007/s00464-017-5939-5>
 60. Rosales-Velderrain A, Alkhoury F (2017) Single-port robotic cholecystectomy in pediatric patients: single institution experience. *J Laparoendosc Adv Surg Tech A* 27(4):434–437. <https://doi.org/10.1089/lap.2016.0484>
 61. Su WL, Huang JW, Wang SN, Lee KT (2017) Comparison study of clinical outcomes between single-site robotic cholecystectomy and single incision laparoscopic cholecystectomy. *Asian J Surg* 40(6):424–428. <https://doi.org/10.1016/j.asjsur.2016.03.005>
 62. Tao Z, Emuakhagbon VS, Pham T, Augustine MM, Guzzetta A, Huerta S (2021) Outcomes of robotic and laparoscopic cholecystectomy for benign gallbladder disease in Veteran patients. *J Robot Surg* 15(6):849–857. <https://doi.org/10.1007/s11701-020-01183-3>
 63. Felli E, Brunetti F, Disabato M, Salloum C, Azoulay D, De'angelis N (2014) Robotic right colectomy for hemorrhagic right colon cancer: a case report and review of the literature of minimally invasive urgent colectomy. *World J Emerg Surg* 9:32. <https://doi.org/10.1186/1749-7922-9-32>
 64. Guerra F, Petrelli F, Stortoni P, Patrì A (2021) Acute large bowel malignant obstruction of the splenic flexure: fully robotic resection with central ligation and intracorporeal anastomosis—a video vignette. *Colorectal Dis* 23(1):325. <https://doi.org/10.1111/codi.15384>
 65. Pedraza R, Ragupathi M, Martinez T, Haas EM (2012) Robotic-assisted laparoscopic primary repair of acute iatrogenic colonic perforation: case report. *Int J Med Robot* 8(3):375–378. <https://doi.org/10.1002/rcs.1447>
 66. Sudan R, Desai SS (2012) Emergency and weekend robotic surgery are feasible. *J Robot Surg* 6(3):263–266. <https://doi.org/10.1007/s11701-011-0289-0>
 67. Sun L, Tang S (2015) Perioperative nursing for gastric perforation repair operation performed by the first domestic surgical robot. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 40(4):453–456. <https://doi.org/10.11817/j.issn.1672-7347.2015.04.019>
 68. Cubas R, Garcia M, Mukherjee K (2021) Robotic repair of incarcerated morgagni hernia in an adult on the acute care surgery service. *Rev Fac Cien Med Univ Nac Cordoba* 78(1):91–94. <https://doi.org/10.31053/1853.0605.v78.n1.28299>
 69. Ceccarelli G, Pasculli A, Bugiantella W, De Rosa M, Catena F, Rondelli F, Costa G, Rocca A, Longaroni M, Testini M (2020) Minimally invasive laparoscopic and robot-assisted emergency treatment of strangulated giant hiatal hernias: report of five cases and literature review. *World J Emerg Surg* 15(1):37. <https://doi.org/10.1186/s13017-020-00316-1>
 70. Muhonen J, Hsu M, Strudivant M, Unger A, Dexter D, Giuseppucci P, Esper C (2020) Identification and repair of left-sided paraduodenal hernia using both laparoscopic and robotic techniques. *Case Rep Surg* 2020:7569530. <https://doi.org/10.1155/2020/7569530>
 71. Smith MT, Wong NC, Coleman JA (2021) Inguinal hernia of the distal ureter causing hydronephrosis: a rare case. *Urol Case Rep* 35:101549. <https://doi.org/10.1016/j.eucr.2020.101549>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.