- 7 Pannett B, Coggon D, Acheson ED. A job-exposure matrix for use in population based studies in England and Wales. Br J Ind Med 1985;42:777-83.
- Breslow NE, Day NE. Statistical methods in cancer research. Vol
  The analysis of case-control studies. Lyon: International Agency for Research on Cancer, 1980.
- 9 Heasman MA, Lipworth L. Accuracy of certification of cause of death. London: HMSO, 1966. (General Registrar Office studies on medical and population subjects No 20.)
- 10 Alderson MR. Some sources of error in British occupational mortality data. Br J Ind Med 1972;29:245-54.
- 11 Office of Population Censuses and Surveys. Occupational mortality 1970-72. Decennial supplement for England and Wales. London: HMSO, 1978. (Series DS No 1.)

- 12 Rushton L, Alderson MR. An epidemiological survey of eight oil refineries in Britain. Br J Ind Med 1981;38:225-34.
- 13 Theriault G, Goulet L. A mortality study of oil refinery workers. J Occup Med 1979;21:367-70.
- 14 Waxweiler RJ, Alexander V, Leffingwell SS, Haring M, Lloyd JW. Mortality from brain tumour and other causes in a cohort of petrochemical workers. JNCI 1983;70:75-81.
- 15 Thomas TL, Waxweiler RJ, Crandall MS, et al. Brain cancer among OCAW members in three Texas oil refineries. Ann NY Acad Sci 1982;381:120-9.
- 16 Nicholson WJ, Seidman H, Selikoff IJ, et al. Brain tumours among operating engineers in the chemical and petrochemical industry in Texas and Louisiana. Ann NY Acad Sci 1982;381:172-80.

## Correction

## An industry wide mortality study of chemical workers occupationally exposed to benzene. II Dose response analyses (June 1987)

The author regrets that an incorrect table was submitted. Table 14 (p. 391) should read as follows:

Table 14 Mantel-Haenszel relative risks and extension chi-squares for lymphatic and haematopoietic cancer, leukaemia, non-Hodgkin's lymphoma, and non-Hodgkin's lymphopoietic cancer by maximum peak occupational exposure to benzene, adjusted for age and race

Cause of death (8th ICD)	Maximum peak exposure (ppm)	Observed deaths	Relative risk	Chi-square	p Value
Lymphatic and haematopoietic cancer (200-209)	Non-exposed <25	3	1·00 3·38		
	25 25-100 ≥ 100	7 8	2·96 3·01	2.10	0.148
Leukaemia and aleukaemia (204–207)	Non-exposed	0	3.01		
	<25 25–100	1 2	Undefined	2.91	0.088
	≥100	4	Chachica	271	0 000
Non-Hodgkin's lymphoma (200, 202, 203)	Non-exposed < 25	2	1·00 3·44		
	25–100	3 4	3·44 1·98	0.06	0.807
	≥100	3	1.57		
Non-Hodgkin's lymphopoietic cancer (200, 202-207)	Non-exposed	2	1.00		
	<25 25–100	4 6	4·62 3·32	1.95	0-163
	≥100	ž	3.89	1 75	0 103

(Lymphatic and haematopoietic cancer includes non-Hodgkin's lymphoma, Hodgkin's disease, and leukaemia.)