



## CASE REPORT

# Further characterization of *NFIB*-associated phenotypes: Report of two new individuals

Gemma Marinella<sup>1</sup> | Eugenia Conti<sup>1</sup> | Bianca Buchignani<sup>1,2</sup> | Giada Sgherri<sup>1</sup> |  
Rosa Pasquariello<sup>1</sup> | Flavio Giordano<sup>3</sup> | Paola Cristofani<sup>1</sup> | Roberta Battini<sup>1,2</sup>  |  
Agatino Battaglia<sup>1</sup> 

<sup>1</sup>Department of Developmental Neuroscience, IRCCS Stella Maris Foundation, Calambrone, Pisa, Italy

<sup>2</sup>Department of Clinical and Experimental Medicine, University of Pisa, Pisa, Pisa, Italy

<sup>3</sup>Department of Neurosurgery, Children's Hospital A. Meyer-University of Florence, Firenze, Florence, Italy

## Correspondence

Roberta Battini, Department of Clinical and Experimental Medicine, University of Pisa, Director of Child Neurology and Psychiatric Residency Program, Department of Developmental Neuroscience, Stella Maris Scientific Institute, Via dei Giacinti, 2, 56128 Calambrone, Pisa, Italy.  
Email: [roberta.battini@fsm.unipi.it](mailto:roberta.battini@fsm.unipi.it)

## Funding information

Sarzana Family Donation; Italian Ministry of Health, Grant/Award Number: RC2021

## Abstract

*Nuclear Factor I B (NFIB)* haploinsufficiency has recently been identified as a cause of intellectual disability (ID) and macrocephaly. Here we report on two new individuals carrying a microdeletion in the chromosomal region 9p23-p22.3 containing *NFIB*. The first is a 7-year 9-month old boy with developmental delays, ID, definite facial anomalies, and brain and spinal cord magnetic resonance imaging findings including periventricular nodular heterotopia, hypoplasia of the corpus callosum, arachnoid cyst in the left middle cranial fossa, syringomyelia in the thoracic spinal cord and distal tract of the conus medullaris, and a stretched appearance of the filum terminale. The second is a 32-year-old lady (the proband's mother) with dysmorphic features, and a history of learning disability, hypothyroidism, poor growth, left inguinal hernia, and panic attacks. Her brain magnetic resonance imaging findings include a dysmorphic corpus callosum, and a small cyst in the left choroidal fissure that marks the hippocampal head. Array-based comparative genomic hybridization identified, in both, a 232 Kb interstitial deletion at 9p23p22.3 including several exons of *NFIB* and no other known genes. Our two individuals add to the knowledge of this rare disorder through the addition of new brain and spinal cord MRI findings and dysmorphic features. We propose that *NFIB* haploinsufficiency causes a clinically recognizable malformation-ID syndrome.

## KEYWORDS

brain/spinal cord malformation, developmental delay, intellectual disability, *NFIB*, periventricular nodular heterotopia, syringomyelia

## 1 | INTRODUCTION

The Nuclear Factor I (NFI) family of transcription factors play an important role in normal development of several organs. Three NFI family members (*NFIA*, *NFIX*, *NFIB*) are highly expressed in the brain, and deletions and sequence variations in all of them have been

associated with intellectual disability (ID) and variable structural brain anomalies (Betancourt et al., 2014; Chen et al., 2017; Gronostajski, 2000; Mason et al., 2009; Murtagh et al., 2003).

Deletions of chromosome 1p32-p31 (MIM: 613735) as well as deletions or sequence variations of *NFIA* lead to a variable phenotype characterized by global developmental delay, ID, central nervous

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *American Journal of Medical Genetics Part A* published by Wiley Periodicals LLC.

system malformations (loss of white matter, polymicrogyria and hypoplastic falx cerebri, partial inversion of the hippocampi), macrocephaly, and craniofacial dysmorphisms (Bayat et al., 2017; Campbell et al., 2002; Koehler et al., 2010; Labonne et al., 2016; Lu et al., 2007; Negishi et al., 2015; Nyboe et al., 2015; Rao et al., 2014; Shanske et al., 2004; Zinner & Batanian, 2003). Mutations leading to haploinsufficiency of *NFIX* cause the autosomal dominant Malan syndrome (MIM: 614753), whereas dominant-negative mutations in *NFIX* cause the autosomal dominant Marshall–Smith syndrome (MIM: 602535). Marshall–Smith syndrome is due to exon deletions, indels and splice site variants leading to frameshift downstream of Exon 5; the mutant *NFIX* protein has a preserved DNA binding and dimerization domain, and therefore likely acts in a dominant negative manner. The syndrome is characterized by prenatal overgrowth, advanced bone age, “Bullet”-shaped middle phalanges, failure to thrive, high neonatal mortality, thick eyebrows, prominent eyes, depressed nasal bridge, anteverted nares, choanal atresia, overfolded helix, and micrognathia (Adam et al., 2005; Aggarwal et al., 2017; Malan et al., 2010; Martinez et al., 2015; Schanze et al., 2014). Malan syndrome is characterized by postnatal overgrowth, decrease in height, overgrowth with age, persistent macrocephaly, learning disability, autism and anxiety, hypotonia, brain anomalies, and dysmorphic features (Malan et al., 2010; Yoneda et al., 2012). *NFIX* duplication has also been observed in a few individuals presenting ID, short stature, small head circumference, and minor dysmorphic features (Dolan et al., 2010; Trimouille et al., 2018). Recently, haploinsufficiency of *NFIB* (caused by either deletions of 9p23-p22.2 or *NFIB* sequence variations), has been shown to cause ID with macrocephaly, along with motor delay, hypotonia, behavioral abnormalities, and variable structural brain anomalies mainly involving the corpus callosum (Barrus et al., 2020; Schanze et al., 2018).

Herein we report two individuals (the proband and his mother) carrying a microdeletion in the chromosomal region 9p23-p22.3 containing *NFIB*, assessed at the IRCCS Stella Maris Foundation (Scientific Institute for Research, Hospitalization and Healthcare Stella Maris Foundation), Pisa, Italy. We describe new imaging data and clinical findings that have not been previously reported.

## 2 | CASE REPORTS


Patient 1 is a 7-year 9-month old boy born by caesarean for breech presentation at 37 + 3 weeks of gestation. Apgar was 7/8. His birth weight was 1800 g (<10th centile), length 48 cm (50th centile), OFC 33.5 cm (50th centile). He was found to have a ventricular septal defect (VSD), a patent foramen ovale (PFO), and jaundice which was treated with phototherapy. Developmental delay was obvious from early on; he held his head at 9 months, sat unsupported at 18 months, and walked alone at 26 months; babbling appeared at around 2 years, and first bisyllabic words at 3 years. Presently, at age 7-year 9-month, he pronounces both simple and complex sentences with some phono-articulatory alterations.

VSD and PFO spontaneously resolved. At age 3 months he underwent surgery for a right inguinal hernia, and at age 3 years 7 months for a left congenital hydrocele. With time, his growth remained between 10th and 25th centile. Frequent, severe urinary tract infections were reported from early on.


Array-comparative genomic hybridization (Array-CGH) CytoSure Oligo Arrays OGT-Oxford Gene Technology), performed elsewhere, showed a deletion of the short arm of a Chromosome 9 in the p23p22.3 region sizing ~232 Kb, starting at 14.107.008 pb and ending at 14.339.185 pb (hg19) including most of the gene *NFIB* (Nuclear Factor I/B; *NFIB*, \*600728). The rearrangement was confirmed by fluorescence in situ hybridization (FISH) using bac RP11-1147E20 which maps in 9p23p22.3. Array-CGH on both parents showed the same deletion in the proband's mother.

The child was referred to our institute at 7 years due to psychomotor developmental delay and dysmorphic features. Physical examination showed height 115 cm (10–25th centile), weight 18.8 kg (3rd–10th centile), occipitofrontal circumference (OFC) 53.5 cm (50–98th centile). There were frontal bossing, broad forehead, high anterior hairline, sparse eyebrows, palpebral fissures length of 2.2 cm, anteverted nares, long and smooth philtrum (lip/philtrum 5), thin vermilion of the upper lip, micro-retrognathia, prominent digit pads, winged scapulae, and cervico-dorsal left convex scoliosis (Figure 1). Neuropsychological evaluation showed a receptive and expressive language disorder, emotional behavioral immaturity, learning disability, and borderline intellectual functioning (Wechsler Preschool and Primary Scale of Intelligence-WPPSI III test total intelligence quotient-QIT 77; performance intelligence quotient-PIQ 85; performance intelligence quotient-VIQ 80) (Luiselli et al., 2013). Fragile X testing and a polygraphic video-electroencephalography (video-EEG) recording (performed while the patient was awake and asleep) were normal. Brain and spinal cord magnetic resonance imaging (MRI) at 7 years of age showed two heterotopic nodules beneath the frontal horn of the left and right lateral ventricles, slight hypoplasia of the rostrum and the splenium of the corpus callosum, a small arachnoid cyst in the left middle cranial fossa, and syringomyelia in the thoracic spinal cord and distal tract of the conus medullaris and slightly stretched appearance of the “filum terminale” (Figure 2A–D). On this basis and given the suspected bladder dysfunction, the child underwent a spinal cord detethering by release of filum terminalis.

Patient 2, the proband's mother, is a 32-year-old lady with a history of learning disability, hypothyroidism, poor growth, left inguinal hernia, and panic attacks. Her cranio-facial features were very similar to those observed in the proband (Figure 3). In addition, she showed bilateral short fourth metatarsal with nail dysplasia, and widely spaced toes (Figure 4). Her weight was 61 kg (50–75th centile), height 148.4 cm (<3rd centile), OFC 55.5 cm (50–98th centile). Brain MRI at the age of 32 years showed a dysmorphic corpus callosum, and a small cyst in the left choroidal fissure that marks the hippocampal head (Figure 2E,F). She has a mild ID (total intelligence quotient-QIT 55; verbal comprehension index-VCI 61; perceptual reasoning index-PRI 67; working memory index-WMI 69; processing speed index-PSI



**FIGURE 1** Front (a) and side (b) view of Patient 1 displaying a definite and well recognizable craniofacial phenotype



**FIGURE 2** Patient 1: (a,b) Coronal T1-weighted image shows periventricular nodular heterotopia confined to the walls of the frontal horns bilaterally (white arrow) and a small arachnoid cyst in the left middle cranial fossa (\*). (c) Sagittal T1-weighted image shows slight hypoplasia of the rostrum (head arrow) and the splenium (\*) of the corpus callosum. (d) Sagittal T2 FIESTA shows syringomyelia in the thoracic spinal cord and distal tract of the conus medullaris and slightly stretched appearance of the small terminal. Patient 2: (e) Sagittal T1-weighted image shows a dysmorphic corpus callosum. (f) Coronal T2-weighted image shows a small cyst in the left choroidal fissure (white arrow) that marks the hippocampal head



**FIGURE 3** Front view of Patient 2, the proband's mother



**FIGURE 4** Feet of Patient 2, showing bilateral short fourth metatarsal with nail dysplasia, and widely spaced toes

67, assessed with the Wechsler Adult Intelligence Scale psychometric test).

### 3 | DISCUSSION

We report on two individuals with ID, abnormal brain and spinal cord MRI findings, and dysmorphic features who both had a heterozygous microdeletion of a Chromosome 9 in the p23p22.3 region, sizing ~232 Kb, including most of the gene *NFIB* (*MIM:618286*). The most distinct dysmorphic features in these individuals include broad forehead, high anterior hairline, sparse eyebrows, anteverted nares, long and smooth philtrum, and thin vermilion of the upper lip. Such clinical features are highly similar to those observed in individuals previously

reported by Schanze et al. (2018), and seem to be distinctive enough to allow a clinical recognition of the *NFIB* haploinsufficiency even before the cytogenomic confirmation.

Contrary to what previously observed, our proband did not have macrocephaly and showed poor growth (Schanze et al., 2018). Both had a dysmorphic corpus callosum and small arachnoid cysts, consistent with previously reported MRI findings (Barrus et al., 2020; Schanze et al., 2018). The periventricular nodular heterotopias (PNH) along the frontal horns of the lateral ventricles observed in Proband 1 appear to be very similar to those found in Patient 7 described by Schanze et al. (2018). PNH, most often bilateral and symmetric, have been observed in several conditions. They are thought to result from loss of radial glial cell attachment to disrupted neuroepidyma (Ferland et al., 2009). Classic PNH, occurring near or in the ventricular walls, are associated with mutations in *FLNA* and *ARFGEF2*, and are probably associated with vesicle trafficking and neuroepidymal repair during neuron proliferation and intracellular transport (Ferland et al., 2009). They can occur in isolation or associated with other central nervous system (CNS) anomalies. It might well be that such PNH will prove more common than presently reported, as a number of described individuals have not undergone brain imaging. *NFIB* haploinsufficiency should be considered in the differential diagnosis of individuals with PNH, particularly bearing in mind the low penetrance of the phenotype.

Our Proband 1 also showed previously unreported spinal cord anomalies on MRI, including syringomyelia in the thoracic spinal cord and distal tract of the conus medullaris and slightly stretched appearance of the "filum terminale." Seemingly, bilateral short fourth metatarsal with nail dysplasia, and widely spaced toes, observed in Proband 2, have not been previously reported, although such features might well be secondary to an associated autosomal dominant disorder. As far as we know, no other family members were reported to have similar anomalies.

All reported findings expand the spectrum of structural brain/spinal cord and physical anomalies associated with *NFIB* haploinsufficiency and highlight the value of performing MRI in such individuals.

Minor heart defects similar to those observed in our Proband 1 have also been reported in 5 out of the 17 individuals reported by Schanze et al. (2018).

The Proband 2's milder MRI anomalies may represent the disorder's variable expressivity in this family.

The clinical spectrum of *NFIB*-related conditions will continue to evolve as new individuals are properly studied. Special attention should be given to the neuropsychological data, and brain/spinal MRI findings, including malformations of cortical and spinal cord development. Thereafter, genotype–phenotype correlations can be developed to help physicians provide anticipatory guidance to both the affected individuals and their families in the future.

In conclusion, the reported family adds to the knowledge of this rare disorder through the addition of new brain and spinal cord MRI findings and dysmorphic features, and showing its variable expressivity.

## ACKNOWLEDGMENTS

Research reported in this publication was supported by the project “Study on phenotype-genotype correlations of genetic rare diseases expressing in developmental age with neuropsychiatric symptoms,” funded by Sarzana Family Donation (Roberta Battini, magnetic resonance). Furthermore, it was partially supported by the Italian Ministry of Health (Grant RC2021) and taxpayers' contributions (“5 × 1000”) to IRCCS Stella Maris Foundation for year 2021 (Agatino Battaglia, RB), RC2021-2022, and public grant 5X1000 for IRCCS Fondazione Stella Maris. Open access funding provided by BIBLIOSAN.

## CONFLICT OF INTEREST

The authors declare no conflicts of interest.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

## ORCID

Roberta Battini  <https://orcid.org/0000-0002-7889-1355>

Agatino Battaglia  <https://orcid.org/0000-0002-7128-7606>

## REFERENCES

- Adam, M. P., Hennekam, R. C. M., Keppen, L. D., Bull, M. J., Clericuzio, C. L., Burke, L. W., Ormond, K. E., & Hoyme, H. E. (2005). Marshall-Smith syndrome: Natural history and evidence of an osteochondrodysplasia with connective tissue abnormalities. *American Journal of Medical Genetics*, 137A, 117–124. <https://doi.org/10.1002/ajmg.a.30580>
- Aggarwal, A., Nguyen, J., Rivera-Davila, M., & Rodriguez-Buritica, D. (2017). Marshall-Smith syndrome: Novel pathogenic variant and previously unreported associations with precocious puberty and aortic root dilatation. *European Journal of Medical Genetics*, 60, 391–394. <https://doi.org/10.1016/j.ejmg.2017.04.012>
- Barrus, K., Rego, S., Yip, T., Martin, P. M., Glen, O. A., Van Ziffle, J., & Slavotinek, A. M. (2020). The expanding spectrum of NFIB-associated phenotypes in a diverse patient population – A report of two new patients. *American Journal of Medical Genetics. Part A*, 182, 2959–2963. <https://doi.org/10.1002/ajmg.a.61852>
- Bayat, A., Kirchhoff, M., Madsen, C. G., Roos, L., & Kreiborg, S. (2017). Familial craniofacial abnormality and polymicrogyria associated with a microdeletion affecting the NFIA gene. *Clinical Dysmorphology*, 26, 148–153. <https://doi.org/10.1097/MCD.0000000000000182>
- Betancourt, J., Katzman, S., & Chen, B. (2014). Nuclear factor one B regulates neural stem cell differentiation and axonal projection of corticofugal neurons. *Journal of Comparative Neurology*, 522, 6–35. <https://doi.org/10.1002/cne.23373>
- Campbell, C. G. N., Wang, H., & Hunter, G. W. (2002). Interstitial microdeletion of chromosome 1p in two siblings. *American Journal of Medical Genetics*, 111, 289–294. <https://doi.org/10.1002/ajmg.10595>
- Chen, K. S., Lim, J. W. C., Richards, L. J., & Bunt, J. (2017). The convergent roles of the nuclear factor I transcription factors in development and cancer. *Cancer Letters*, 410, 124–138. <https://doi.org/10.1016/j.canlet.2017.09.015>
- Dolan, M., Mendelsohn, N. J., Pierpont, M. E., Schimmenti, L. A., Berry, S. A., & Hirsch, B. (2010). A novel microdeletion/microduplication syndrome of 19p13.13. *Genetics in Medicine*, 12, 503–511. <https://doi.org/10.1097/GIM.0b013e3181e59291>
- Ferland, R. J., Batiz, L. F., Neal, J., Lian, G., Bundock, E., Lu, J., Hsiao, Y. C., Diamond, R., Mei, D., Banham, A. H., Brown, P. J., Vanderburg, C. R., Joseph, J., Hecht, J. L., Folkerth, R., Guerrini, R., Walsh, C. A., Rodriguez, E. M., & Sheen, V. L. (2009). Disruption of neural progenitors along the ventricular and subventricular zones in periventricular heterotopia. *Human Molecular Genetics*, 18, 497–516. <https://doi.org/10.1093/hmg/ddn377>
- Gronostajski, R. M. (2000). Roles of the NFI/CTF gene family in transcription and development. *Gene*, 249, 31–45. [https://doi.org/10.1016/S0378-1119\(00\)00140-2](https://doi.org/10.1016/S0378-1119(00)00140-2)
- Koehler, U., Holinski-Feder, E., Ertl-Wagner, B., Kunz, J., von Moers, A., von Voss, H., & Schell-Apacik, C. (2010). A novel 1p31.3p32.2 deletion involving the NFIA gene detected by array CGH in a patient with macrocephaly and hypoplasia of the corpus callosum. *European Journal of Pediatrics*, 169, 463–468. <https://doi.org/10.1007/s00431-009-1057-2>
- Labonne, J. D. J., Shen, Y., Kong, I. K., Diamond, M. P., Layman, L. C., & Kim, H. G. (2016). Comparative deletion mapping at 1p31.3-p32.2 implies NFIA responsible for intellectual disability coupled with macrocephaly and the presence of several other genes for syndromic intellectual disability. *Molecular Cytogenetics*, 9, 24. <https://doi.org/10.1186/s13039-016-0234-z>
- Lu, W., Quintero-Rivera, F., Fan, Y., Alkuraya, F. S., Donovan, D. J., Xi, Q., Turbe-Doan, A., Li, Q. G., Campbell, C. G., Shanske, A. L., Sherr, E. H., Ahmad, A., Peters, R., Rilliet, B., Parvex, P., Bassuk, A. G., Harris, D. J., Ferguson, H., Kelly, C., ... Maas, R. L. (2007). NFIA haploinsufficiency is associated with a CNS malformation syndrome and urinary tract defects. *PLoS Genetics*, 3, e80. <https://doi.org/10.1371/journal.pgen.0030080>
- Luiselli, J., Happé, F., Hurst, H., Freeman, S., Goldstein, G., Mazefsky, C., Carter, A. S., Kaufman, A. S., Simmons, E. S., Eernisse, E. R., Tsatsanis, K., Fluit, F., Klein-Tasman, B., Feinstein, A., Kahl, A., Poyau, S., Jones, C. R. G., Volkmar, F. R., Wolf, J. M., ... Newman, D. B. (2013). WPPSI-III – Wechsler preschool and primary scale of intelligence. In F. R. Volkmar (Ed.), *Encyclopedia of Autism Spectrum Disorders* (Third ed.). Springer Nature. [https://doi.org/10.1007/978-1-4419-1698-3\\_101583](https://doi.org/10.1007/978-1-4419-1698-3_101583)
- Malan, V., Rajan, D., Thomas, S., Shaw, A. C., Louis Dit Picard, H., Layet, V., Till, M., Van Haeringen, A., Mortier, G., Nampoothiri, S., Pušeljčić, S., Legeai-Mallet, L., Carter, N. P., Vekemans, M., Munnich, A., Hennekam, R. C., Colleaux, L., & Cormier-Daire, V. (2010). Distinct effects of allelic NFIX mutations on nonsense-mediated mRNA decay engender either a sotos-like or a Marshall-Smith Syndrome. *American Journal of Human Genetics*, 87, 189–198. <https://doi.org/10.1016/j.ajhg.2010.07.001>
- Martinez, F., Marín-Reina, P., Sanchis-Calvo, A., Perez-Aytés, A., Oltra, S., Roselló, M., Mayo, S., Monfort, S., Pantoja, J., & Orellana, C. (2015). Novel mutations of NFIX gene causing Marshall-Smith syndrome or Sotos-like syndrome: One gene, two phenotypes. *Pediatric Research*, 78, 533–539. <https://doi.org/10.1038/pr.2015.135>
- Mason, S., Piper, M., Gronostajski, R. M., & Richards, L. J. (2009). Nuclear factor one transcription factors in CNS development. *Molecular Neurobiology*, 39, 10–23. <https://doi.org/10.1007/s12035-008-8048-6>
- Murtagh, J., Martin, F., & Gronostajski, R. M. (2003). The nuclear factor I (NFI) gene family in mammary gland development and function. *Journal of Mammary Gland Biology and Neoplasia*, 8(2), 241–254. <https://doi.org/10.1023/A:1025909109843>
- Negishi, Y., Miya, F., Hattori, A., Mizuno, K., Hori, I., Ando, N., Okamoto, N., Kato, M., Tsunoda, T., Yamasaki, M., Kanemura, Y., Kosaki, K., & Saitoh, S. (2015). Truncating mutation in NFIA causes brain malformation and urinary tract defects. *Human Genome Variation*, 2, 15007. <https://doi.org/10.1038/hgv.2015.7>
- Nyboe, D., Kreiborg, S., Kirchhoff, M., & Hove, H. B. (2015). Familial craniosynostosis associated with a microdeletion involving the NFIA

- gene. *Clinical Dysmorphology*, 24, 109–112. <https://doi.org/10.1097/MCD.0000000000000079>
- Rao, A., O'Donnell, S., Bain, N., Meldrum, C., Shorter, D., & Goel, H. (2014). An intragenic deletion of the NFIA gene in a patient with a hypoplastic corpus callosum, craniofacial abnormalities and urinary tract defects. *European Journal of Medical Genetics*, 57, 65–70. <https://doi.org/10.1016/j.ejmg.2013.12.011>
- Schanze, D., Neubauer, D., Cormier-Daire, V., Delrue, M. A., Dieux-Coeslier, A., Hasegawa, T., Holmberg, E. E., Koenig, R., Krueger, G., Schanze, I., Seemanova, E., Shaw, A. C., Vogt, J., Volleth, M., Reis, A., Meinecke, P., Hennekam, R. C. M., & Zenker, M. (2014). Deletions in the 3' part of the NFIX gene including a recurrent alu-mediated deletion of exon 6 and 7 account for previously unexplained cases of Marshall-Smith syndrome. *Human Mutation*, 35, 1092–1100. <https://doi.org/10.1002/humu.22603>
- Schanze, I., Bunt, J., Lim, J. W. C., Schanze, D., Dean, R. J., Alders, M., Blanchet, P., Attié-Bitach, T., Berland, S., Boogert, S., Boppudi, S., Bridges, C. J., Cho, M. T., Dobyns, W. B., Donnai, D., Douglas, J., Earl, D. L., Edwards, T. J., Faivre, L., ... Richards, L. J. (2018). NFIB Haploinsufficiency is associated with intellectual disability and macrocephaly. *American Journal of Human Genetics*, 103, 752–768. <https://doi.org/10.1016/j.ajhg.2018.10.006>
- Shanske, A. L., Edelmann, L., Kardon, N. B., Gosset, P., & Levy, B. (2004). Detection of an interstitial deletion of 2q21-22 by high resolution comparative genomic hybridization in a child with multiple congenital anomalies and an apparent balanced translocation. *American Journal of Medical Genetics*, 131A, 29–35. <https://doi.org/10.1002/ajmg.a.30311>
- Trimouille, A., Houcinat, N., Vuillaume, M. L., Fergelot, P., Boucher, C., Toutain, J., Caignec, C. L., Vincent, M., Nizon, M., Andrieux, J., Vanlerberghe, C., Delobel, B., Duban, B., Mansour, S., Baple, E., McKeown, C., Poke, G., Robertshaw, K., Fifield, E., ... Moutton, S. (2018). 19p13 microduplications encompassing NFIX are responsible for intellectual disability, short stature and small head circumference. *European Journal of Human Genetics*, 26, 85–93. <https://doi.org/10.1038/s41431-017-0037-7>
- Yoneda, Y., Saito, H., Touyama, M., Makita, Y., Miyamoto, A., Hamada, K., Kurotaki, N., Tomita, H., Nishiyama, K., Tsurusaki, Y., Doi, H., Miyake, N., Ogata, K., Naritomi, K., & Matsumoto, N. (2012). Missense mutations in the DNA-binding/dimerization domain of NFIX cause Sotos-like features. *Journal of Human Genetics*, 57, 207–211. <https://doi.org/10.1038/jhg.2012.7>
- Zinner, S. H., & Batanian, J. R. (2003). Second reported patient with del(1)(p32.1p32.3) and similar clinical features suggesting a recognizable chromosomal syndrome. *American Journal of Medical Genetics*, 22A(2), 164–167. <https://doi.org/10.1002/ajmg.a.20265>

**How to cite this article:** Marinella, G., Conti, E., Buchignani, B., Sgherri, G., Pasquariello, R., Giordano, F., Cristofani, P., Battini, R., & Battaglia, A. (2023). Further characterization of NFIB-associated phenotypes: Report of two new individuals. *American Journal of Medical Genetics Part A*, 191A:540–545. <https://doi.org/10.1002/ajmg.a.63018>