

Correction to “Photogeneration of Spin Quintet Triplet–Triplet Excitations in DNA-Assembled Pentacene Stacks”

Sarah R. E. Orsborne, Jeffrey Gorman, Leah R. Weiss, Akshay Sridhar, Naitik A. Panjwani, Giorgio Divitini, Peter Budden, David Palecek, Seán T. J. Ryan, Akshay Rao, Rosana Collepardo-Guevara, Afaf H. El-Sagheer, Tom Brown, Jan Behrends, Richard H. Friend,* and Florian Auras*

J. Am. Chem. Soc. **2023**, *145* (9), 5431–5438. DOI: [10.1021/jacs.2c13743](https://doi.org/10.1021/jacs.2c13743)



Cite This: *J. Am. Chem. Soc.* **2023**, *145*, 8271–8271



Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

Pages 5436. In the analysis and discussion of the ESR experiments, the definition of the quintet zero-field-splitting Hamiltonian was mistakenly typed as

$$\frac{\hat{H}_{zfs}}{h} = \mathbf{S}^T \cdot \mathbf{D}_q \cdot \mathbf{S} = D_q (\hat{S}_z^2 - 2\hat{I}_3) + E_q (\hat{S}_z^2 - \hat{S}_z^2)$$

The equation and definition of its terms should read as follows:

$$\frac{\hat{H}_{zfs}}{h} = \mathbf{S}^T \cdot \mathbf{D}_q \cdot \mathbf{S} = D_q (\hat{S}_z^2 - 2\hat{I}) + E_q (\hat{S}_x^2 - \hat{S}_y^2)$$

where \mathbf{S} is the vector of quintet spin operators, \hat{I} is the 5×5 identity matrix, and \hat{S}_i is the spin operator defined along the principal axis $i \in (\hat{x}, \hat{y}, \hat{z})$ of the ZFS tensor \mathbf{D}_q , determined by the symmetry of the underlying spin–spin interactions.

This is also updated in Section K (“Theoretical quintet ($S=2$) zero-field splitting parameters”, page S22) of the [Supporting Information](#).

These corrections do not affect the conclusions presented in the published article.

■ ASSOCIATED CONTENT

SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.3c02230>.

Experimental methods, synthetic procedures, and additional spectroscopic data (updated) ([PDF](#))

Published: March 31, 2023

