



Retrospective Cohort Study

# Causes of gastrointestinal bleeding in children based on endoscopic evaluation at a tertiary care center in Bahrain

Hasan M Isa, Fatema A Alkharsi, Hana A Ebrahim, Kamel J Walwil, Jumana A Diab, Nafelah M Alkowari

**Specialty type:** Gastroenterology and hepatology

**Provenance and peer review:**

Unsolicited article; Externally peer reviewed.

**Peer-review model:** Single blind

**Peer-review report's scientific quality classification**

Grade A (Excellent): 0  
Grade B (Very good): B, B  
Grade C (Good): C  
Grade D (Fair): 0  
Grade E (Poor): 0

**P-Reviewer:** Çolak E, Turkey; Hu B, China

**Received:** December 17, 2022

**Peer-review started:** December 17, 2022

**First decision:** January 12, 2023

**Revised:** January 23, 2023

**Accepted:** March 17, 2023

**Article in press:** March 17, 2023

**Published online:** April 16, 2023



**Hasan M Isa, Fatema A Alkharsi,** Department of Pediatrics, Salmaniya Medical Complex, Manama 26671, Bahrain

**Hasan M Isa,** Department of Pediatrics, Arabian Gulf University, Manama 26671, Bahrain

**Hana A Ebrahim, Kamel J Walwil, Jumana A Diab,** Department of Pediatrics, Royal College of Surgeons in Ireland-Medical University of Bahrain, Busaiteen 15503, Bahrain

**Nafelah M Alkowari,** Department of Pediatrics, King Hamad University Hospital, Muharraq 24343, Bahrain

**Corresponding author:** Hasan M Isa, MBChB, Associate Professor, Department of Pediatrics, Salmaniya Medical Complex, Arabian Gulf University, Manama 26671, Bahrain.

[halfaraj@hotmail.com](mailto:halfaraj@hotmail.com)

## Abstract

### BACKGROUND

Gastrointestinal bleeding (GIB) is a serious health problem worldwide, particularly during childhood. This can be an alarming sign of an underlying disease. Gastrointestinal endoscopy (GIE) is a safe method for the diagnosis and treatment of GIB in most cases.

### AIM

To determine the incidence, clinical presentation, and outcomes of GIB in children in Bahrain over the last two decades.

### METHODS

This was a retrospective cohort review of the medical records of children with GIB who underwent endoscopic procedures in the Pediatric Department at Salmaniya Medical Complex, Bahrain, between 1995 and 2022. Demographic data, clinical presentation, endoscopic findings, and clinical outcomes were recorded. GIB was classified into upper (UGIB) and lower (LGIB) GIB according to the site of bleeding. These were compared with respect to patients' sex, age, and nationality using the Fisher's exact, Pearson's  $\chi^2$ , or the Mann-Whitney U tests.

### RESULTS

A total of 250 patients were included in this study. The median incidence was 2.6/100000 per year (interquartile range, 1.4-3.7) with a significantly increasing

trend over the last two decades ( $P < 0.0001$ ). Most patients were males ( $n = 144$ , 57.6%). The median age at diagnosis was 9 years (5–11). Ninety-eight (39.2%) patients required upper GIE alone, 41 (16.4%) required colonoscopy alone, and 111 (44.4%) required both. LGIB was more frequent ( $n = 151$ , 60.4%) than UGIB ( $n = 119$ , 47.6%). There were no significant differences in sex ( $P = 0.710$ ), age ( $P = 0.185$ ), or nationality ( $P = 0.525$ ) between the two groups. Abnormal endoscopic findings were detected in 226 (90.4%) patients. The common cause of LGIB was inflammatory bowel disease (IBD) ( $n = 77$ , 30.8%). The common cause of UGIB was gastritis ( $n = 70$ , 28%). IBD and undetermined cause for bleeding were higher in the 10–18 years group ( $P = 0.026$  and  $P = 0.017$ , respectively). Intestinal nodular lymphoid hyperplasia, foreign body ingestion, and esophageal varices were more common in the 0–4 years group ( $P = 0.034$ ,  $P < 0.0001$ , and  $P = 0.029$ , respectively). Ten (4%) patients underwent one or more therapeutic interventions. The median follow-up period was two years (0.5–3). No mortality was reported in this study.

### CONCLUSION

GIB in children is an alarming condition, whose significance is increasing. LGIB, commonly due to IBD, was more common than UGIB, commonly due to gastritis.

**Key Words:** Pediatric; Gastrointestinal bleeding; Endoscopy; Causes; Outcome; Bahrain

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** The incidence, clinical presentation, cause, and outcomes of gastrointestinal bleeding (GIB) in children in Bahrain are unknown. We observed a significant increase in the annual incidence of GIB. Lower GIB (LGIB) was more common than upper GIB (UGIB). The most common cause of LGIB was inflammatory bowel disease. The most common cause of UGIB was gastritis. Causes of GIB varied with patient's age and differed from that reported in other countries. No mortality was observed in any patient. These findings are essential to tailor management based on the most common causes and patient age.

**Citation:** Isa HM, Alkharsi FA, Ebrahim HA, Walwil KJ, Diab JA, Alkowari NM. Causes of gastrointestinal bleeding in children based on endoscopic evaluation at a tertiary care center in Bahrain. *World J Gastrointest Endosc* 2023; 15(4): 297-308

**URL:** <https://www.wjgnet.com/1948-5190/full/v15/i4/297.htm>

**DOI:** <https://dx.doi.org/10.4253/wjge.v15.i4.297>

## INTRODUCTION

Gastrointestinal (GI) bleeding (GIB) is a serious health problem worldwide, particularly during childhood. It is defined as any kind of hemorrhage or blood loss in the digestive tract, from the mouth to the anus[1], which can range from nearly undetectable to acute, massive, or life-threatening[1]. It is divided into three main clinical forms: Upper GIB (UGIB), lower GIB (LGIB), and bleeding of obscure origin[2]. UGIB and LGIB are defined based on their relationship with the ligament of Treitz[3]. The source of bleeding in UGIB is proximal to the ligament of Treitz (from the upper part of the esophagus to the duodenojejunal flexure), whereas that of LGIB is distal to the ligament (small bowel and colon) [3]. GIB can also be occult, that is, not visible to the patient or physician, leading to either a positive fecal occult blood test or iron-deficiency anemia[3].

GIB can cause anxiety in children, caregivers, and healthcare providers[4] and can be an alarming sign of an underlying disease[5]. GIB has a variety of causes; a good elicitation of patients' medical history and physical examination can adequately differentiate between macroscopic and microscopic forms of bleeding[5].

GI endoscopy (GIE) is a safe method for the diagnosis and treatment of GIB[4]. GIE can reveal the underlying etiology of GIB in most cases[3]. However, it requires the experience and coordination of the medical team to provide specialized patient care[6].

According to the Italian Society of Pediatric Gastroenterology, the incidence rate of GIB in children is 6.4% [3]. However, there is a paucity of studies regarding the incidence and causes of GIB in the pediatric population in the Middle East. Thus, we aimed to determine the incidence, clinical presentation, causes, and outcomes of GIB in children at the main tertiary hospital in Bahrain over the last two decades. Additionally, we aimed to stratify the causes of GIB based on patient age.

## MATERIALS AND METHODS

### **Study design and setting**

We conducted a retrospective cohort review of all medical records of children diagnosed with GIB in the Pediatric Department at Salmaniya Medical Complex (SMC), Manama, Bahrain, between January 1995 and October 2022. SMC is the only tertiary hospital in Bahrain to where children with GIB are referred for diagnosis and management. Endoscopic procedures were performed in either the endoscopy unit or the main operating theatre. The endoscopy unit in SMC consists of three rooms, in which three pediatric gastroenterology consultants and one chief resident can perform endoscopic procedures. Two types of endoscopic equipment were used: Olympus (PCF-230 and XQ230, Olympus Cooperation, Shinjuku, Tokyo, Japan) and Pentax (EG-2901 and EC-380IF, Pentax Ricoh Imaging Company Ltd, Tokyo, Japan).

### **Study participants**

All children who were admitted to the pediatric department for GIB and underwent GIE were included in the study. Patients who were discharged from the emergency department and those who did not undergo endoscopic procedures were excluded. Patients were classified as having upper, lower, or both GIB according to their presenting symptoms.

### **Data collection**

From 1995 to 2010, the data were retrieved from archived paper-based medical records, while from 2010 to 2022, the data were retrieved from the I-Seha electronic medical records. The following demographic data were collected: Year of presentation, sex, nationality, age at diagnosis, and history of associated chronic diseases.

Presenting symptoms, such as hematemesis, melena, and rectal bleeding (hematochezia), were noted. Additional symptoms, including recurrent vomiting or abdominal pain, chest pain, diarrhea, constipation, weight loss, and history of foreign body or caustic ingestion, were noted. Physical findings such as oral ulcers, pallor, jaundice, perianal fistula or fissure, hemorrhoids, and failure to thrive were recorded. A perianal fistula was defined as a small passage that connects an infected gland inside the anus to an opening on the skin around the anus. A perianal fissure is a tear in the anal mucosa. Failure to thrive (thinness) was defined as a weight for age z-score of < 2 standard deviations according to the World Health Organization growth references[7].

Endoscopy data on the type, upper (UGIE) or lower (LGIE) GIE, and the total number of procedures performed were collected. The causes of GIB were based on endoscopic findings, such as Mallory-Weiss syndrome, esophagitis, esophageal ulcer or varices, foreign body ingestion, gastritis, peptic or duodenal ulcer, gastroenteritis, inflammatory bowel disease (IBD), nodular lymphoid hyperplasia, Meckel's diverticulum, rectal ulcers or polyps, and anal fissures. Data on therapeutic interventions, as well as number of patients who required repeat endoscopic interventions, follow-up duration, and patient outcomes were collected.

### **Statistical analysis**

Data were analyzed using the Statistical Package for the Social Sciences (SPSS) (version 21; IBM Corp. Released 2012. IBM SPSS Statistics for Windows, Armonk, NY: IBM Corp, United States). The annual incidence of GIB was calculated. The 28-year study period was divided into four periods (1995–2001, 2002–2008, 2009–2015 and 2016–2022), which were compared in terms of the mean annual incidence of GIB using a one-way analysis of variance test. Bonferroni post-hoc analysis was used for pairwise multiple comparisons between the four periods. Patient ages were classified into three groups: 0–4, 5–9, and 10–18 years. Categorical variables are presented as frequency and percentage. Continuous variables are presented as median and interquartile range (IQR). UGIB and LGIB were compared with respect to sex, age, and nationality. The causes of GIB were compared according to age group. Fisher's exact test or Pearson's  $\chi^2$  test was used to compare categorical variables, while the Mann-Whitney *U* test was used to compare continuous variables with a skewed distribution. *P* values < 0.05 were considered statistically significant.

### **Ethical approval**

This study was conducted in accordance with the principles of Helsinki Declaration, and it was ethically approved by the Research and Research Ethics Committee, Salmaniya Medical Complex, Government hospitals, Kingdom of Bahrain (IRB number: 6170122, January 17, 2022).

## RESULTS

During the study period, 250 children were admitted to the hospital for GIB and underwent a GIE procedure. All the patients were included in this study. According to the 2020 Bahrain Health Statistics, the total population in Bahrain was 1472204, with 481819 people within the pediatric age group (up to

18 years). The median incidence was 2.6/100000 per year (IQR, 1.4–3.7) with an increasing trend (Figure 1). A significant difference in the mean GIB incidence was found between the four periods, [ $F(3, 24) = 10.280, P < 0.0001$ ]. The mean incidence in 2009–2015 was significantly higher ( $4.9 \pm 1.9$ ) than that in 1995–2001 ( $1.7 \pm 0.8$ ), 2002–2008 ( $1.5 \pm 0.9$ ) and 2016–2022 ( $2.7 \pm 1.9$ ) ( $P = 0.001, P < 0.0001, P = 0.028$ , respectively). The demographic data of the included patients are shown in Table 1. Most of the patients were males ( $n = 144, 57.6\%$ ). The majority were Bahraini ( $n = 224, 89.6\%$ ), while the rest were non-Bahraini ( $n = 24, 9.6\%$ ) [eight (3.2%) were from India, three (1.2%) from Pakistan, two (0.8%) from Iraq, two (0.8%) from Egypt, one (0.4%) from Oman, one (0.4%) from Qatar, one (0.4%) from Yemen, one (0.4%) from Sudan, one (0.4%) from Syria, one (0.4%) from Sweden, one (0.4%) from Bangladesh, one (0.4%) from China, and one (0.4%) from Philippines]. Two patients (0.8%) were of unspecified nationality. The median age at the time of endoscopic diagnosis was 9 years (IQR, 5–11). The most commonly affected age group was the 10–18 years group ( $n = 107, 42.8\%$ ). Thirty-nine (15.6%) patients had one or more associated diseases that were not considered as a direct cause of GIB. The common associated diseases were gastroesophageal reflux disease (GERD) ( $n = 7, 2.8\%$ ) and celiac disease ( $n = 7, 2.8\%$ ), followed by sickle cell disease ( $n = 5, 2\%$ ), and autoimmune hepatitis ( $n = 4, 1.6\%$ ). Other associated diseases were cerebral palsy ( $n = 3, 1.2\%$ ), mental retardation ( $n = 3, 1.2\%$ ), biliary atresia ( $n = 2, 0.8\%$ ), Wilson's disease ( $n = 2, 0.8\%$ ), sclerosing cholangitis ( $n = 2, 0.8\%$ ), chronic liver disease with portal hypertension ( $n = 2, 0.8\%$ ), trisomy 21 ( $n = 2, 0.8\%$ ), Ehlers Danlos syndrome ( $n = 1, 0.4\%$ ), familial Mediterranean fever ( $n = 1, 0.4\%$ ), cholecystitis ( $n = 1, 0.4\%$ ), intestinal worms ( $n = 1, 0.4\%$ ), tracheoesophageal fistula ( $n = 1, 0.4\%$ ), liver cirrhosis ( $n = 1, 0.4\%$ ), Turner's syndrome ( $n = 1, 0.4\%$ ), insulin-dependent diabetes mellitus ( $n = 1, 0.4\%$ ), hydrocephalus ( $n = 1, 0.4\%$ ), glucose-6-phosphate dehydrogenase deficiency ( $n = 1, 0.4\%$ ), congenital heart disease ( $n = 1, 0.4\%$ ), hiatal hernia ( $n = 1, 0.4\%$ ), autoimmune hemolytic anemia ( $n = 1, 0.4\%$ ), food allergy ( $n = 1, 0.4\%$ ) and anorexia nervosa ( $n = 1, 0.4\%$ ).

LGIB was more frequent ( $n = 151, 60.4\%$ ) than UGIB was ( $n = 119, 47.6\%$ ). Out of the 250 patients, 20 (8%) patients underwent both UGIB and LGIB. Children with LGIB were older [median age of 9 years (IQR, 5–12)] than those with UGIB [median age of 8 years (IQR, 4–11)]. This difference was not statistically significant ( $P = 0.185$ ). There was also no significant difference in sex ( $P = 0.710$ ) and nationality ( $P = 0.525$ ) between the two groups (Figure 2).

The clinical presentations are shown in Table 2. The most common presenting symptom was per rectal bleeding ( $n = 151, 60.4\%$ ), followed by hematemesis ( $n = 117, 46.8\%$ ). Some patients presented with more than one symptom. Physical examination was unremarkable in most of the patients ( $n = 218, 87.2\%$ ). However, 32 (12.8%) patients had positive findings on physical examination (Table 2).

Most patients ( $n = 206, 82.4\%$ ) required one GIE; the remaining 44 (17.6%) required more than one GIE. The median number of endoscopies was one, ranging from one to seven. Abnormal endoscopic findings were detected in 226 (90.4%) patients; twenty-four (9.6%) patients had a normal GIE. Examples of patients with positive findings are shown in Figure 3.

The different causes of GIB according to age group are shown in Table 3. The most common cause of LGIB was IBD ( $n = 77, 30.8\%$ ); thirty-four (44.1%) had ulcerative colitis, 33 (42.9%) had Crohn's disease, and 10 (13%) had unspecified IBD. The most common cause of UGIB was gastritis ( $n = 70, 28\%$ ); nonspecific gastritis ( $n = 55, 78.6\%$ ), *Helicobacter pylori* infection ( $n = 14, 20\%$ ), and eosinophilic gastritis ( $n = 1, 1.4\%$ ). In patients with both UGIB and LGIB, the causes of bleeding were gastritis ( $n = 6, 3.5\%$ ), gastroenteritis ( $n = 4, 2\%$ ), ulcerative colitis ( $n = 3, 1.5\%$ ), Crohn's disease ( $n = 2, 1\%$ ), unspecified IBD type ( $n = 1, 0.5\%$ ), esophageal ulcer ( $n = 1, 0.5\%$ ), gastric ulcer ( $n = 1, 0.5\%$ ), duodenal ulcer ( $n = 1, 0.5\%$ ), and esophagitis with gastritis ( $n = 1, 0.5\%$ ).

IBD and unclear cause for bleeding were significantly higher in the 10–18 years group ( $P = 0.026$  and  $P = 0.017$ , respectively), while intestinal nodular lymphoid hyperplasia, ingestion of foreign bodies, and esophageal varices were more frequent in the 0–4 years age group ( $P = 0.034, P < 0.0001$ , and  $P = 0.029$ , respectively). There were no significant differences between other causes according to age groups.

Ten (4%) patients underwent one or more therapeutic interventions [polypectomy ( $n = 4, 1.6\%$ ), injection sclerotherapy ( $n = 3, 1.2\%$ ) where one required two sessions, clipping ( $n = 2, 0.8\%$ ), and banding of esophageal varices and dilatation of esophageal stricture ( $n = 1, 0.4\%$  each)]. Forty-five (18%) patients underwent follow-up endoscopy for disease reassessment. The median follow up period was two years (IQR, 0.5–3). No mortality was reported in this study.


## DISCUSSION

This study revealed that the median incidence of GIB in children was 2.6/100000 per year, which has significantly increased over the last two decades. This increase could be attributed to the overall increase in some of the underlying etiologies, such as IBD. The burden of IBD is rising worldwide[8]. In the Middle East, the incidence of GIB in children is not well established[9,10]. However, in the United States, it accounts for 1% of all pediatric hospitalizations[11]. Romano *et al*[3] reported a higher incidence reaching 6.4% in Italy.

**Table 1** Demographic data of children presented with gastrointestinal bleeding

Variables	n (%)
<b>Sex</b>	
Male	144 (57.6)
Female	106 (42.4)
<b>Nationality</b>	
Bahraini	224 (89.6)
Non-Bahraini	24 (9.6)
Unspecified	2 (0.8)
<b>Age group (yr) (n = 247)</b>	
0-4	58 (23.2)
5-9	82 (32.8)
10-18	107 (42.8)
<b>Type of endoscopy performed</b>	
Upper gastrointestinal endoscopy	98 (39.2)
Lower gastrointestinal endoscopy	41 (16.4)
Both types of gastrointestinal endoscopies	111 (44.4)

Data are presented as numbers and percentages.



**Figure 1** Incidence of gastrointestinal bleeding in children, 1995-2022.

In the current study, most patients who presented with GIB were males ( $n = 144, 57.6\%$ ). This is comparable to several other studies, with a male predominance ranging between 54% and 59.2% [2,6,9,12,13]. However, Kalyoncu *et al* [14], Banisalamah *et al* [15], and Almadi *et al* [16] reported a higher percentage, ranging from 66.8% to 74.2%. The cause of this male predominance in children with GIB remains unclear.

In this study, the median age at the time of endoscopic diagnosis was 9 years (IQR, 5–11), with no significant difference according to the type of bleeding. Similarly, Cleveland *et al* [4] reported a median



**Table 2** Clinical presentation of children with gastrointestinal bleeding

Clinical presentation	Patients' n (%)
Presenting symptom <sup>a</sup>	
Per rectal bleeding	151 (60.4)
Hematemesis	117 (46.8)
Melena	25 (10)
Associated symptom	
Recurrent abdominal pain	59 (23.6)
Diarrhea	23 (9.2)
Constipation	19 (7.6)
Recurrent vomiting	17 (6.8)
Weight loss	10 (4)
Chest pain	8 (3.2)
History of foreign body ingestion	7 (2.8)
Caustic ingestion	1 (0.4)
Nausea	1 (0.4)
Physical finding	
Normal	218 (87.2)
Pallor	11 (4.4)
Perianal fissure	11 (4.4)
Perianal fistula	4 (1.6)
Jaundice	3 (1.2)
Failure to thrive	2 (0.8)
Oral ulcers	1 (0.4)

<sup>a</sup>Some patients presented with more than one symptom. Data are presented as number and percentage.

age of 10.1 years with a mean age of  $9.3 \pm 5.7$  years. However, Gimiga *et al*[6], Rafeey *et al*[10], and Kalyoncu *et al*[14] reported GIB in younger children ( $1.3$ ,  $6.1 \pm 3.9$ , and  $7.6 \pm 2.4$  years, respectively). Nonetheless, the most commonly affected age group in our study was 10–18 years. However, Jafari *et al* [2] and Hassoon *et al*[9] reported that children aged 6–12 years were more frequently affected, with percentages of 31.8% and 24%, respectively. Gimiga *et al*[6] and Zahmatkeshan *et al*[12] reported that the 2–10 and 3–10 years age groups were the most commonly affected, representing 63.7% and 55.1% of their patient samples, respectively. This variation in age at diagnosis could be attributed to the differences in the study settings, inclusion criteria, and site and causes of bleeding in each study.

In the current study, the most common associated diseases were GERD and celiac disease ( $n = 7$ , 2.8%). Attard *et al*[13] reported a higher percentage of GERD in children with GIB (12.2%). Almadi *et al* [16] also reported a high percentage of GERD (38.13%) in adult patients with GIB. Celiac disease can cause occult GIB; it rarely presents as frank GIB[1].

In the present study, LGIB was more frequent than UGIB was, accounting for 60.4% ( $n = 151$ ) and 47.6% ( $n = 119$ ) of cases, respectively. However, Jafari *et al*[2] determined that UGIB was more common than LGIB. Apart from the Jafari *et al*[2] study, all the other published studies tackled either UGIB or LGIB alone. Thus, our findings cannot be compared with those of other studies.

In this study, per rectal bleeding was the most frequent presenting symptom ( $n = 151$ , 60.4%). Gimiga *et al*[6] and Zahmatkeshan *et al*[12] reported similar findings, with hematochezia as the most common symptom in 54.2% and 80.2% of their patients, respectively.

In our study, 117 (46.8%) patients presented with hematemesis. This percentage is higher than those reported by Jafari *et al*[2] (40.7%) and Rafeey *et al*[10] (26.9%). However, Cleveland *et al*[4] and Hassoon *et al*[9]. reported higher percentages of 58.5% and 73.4%, respectively. This might be related to the fact that hematemesis is bright red and usually alarms the child and family to seek early medical advice[9].

In the present study, melena was the third most common symptom. However, Jafari *et al*[2] (17.73%), Cleveland *et al*[4] (20.8%), and Rafeey *et al*[10] (13.42%) reported higher percentages of melena. In

**Table 3 Causes of gastrointestinal bleeding in children in relation to age group**

Causes <sup>a</sup>	Age group (yr)			Total, n (%)	P value <sup>b</sup>
	0-4, n = 58 (23.2)	5-9, n = 82 (32.8)	10-18, n = 107 (42.8)		
Inflammatory bowel disease	10 (17.2)	27 (32.9)	40 (37.4)	77 (30.8)	<b>0.026</b>
Gastritis	16 (27.6)	23 (28)	31 (29)	70 (28)	0.980
Unclear cause (normal)	3 (5.2)	4 (4.9)	17 (15.9)	24 (9.6)	<b>0.017</b>
Gastroenteritis	7 (12.1)	9 (11)	6 (5.6)	22 (8.8)	0.275
Duodenal ulcer	5 (8.6)	4 (4.9)	11 (10.3)	20 (8)	0.397
Esophagitis	4 (6.9)	5 (6.1)	6 (5.6)	15 (6)	0.947
Rectal polyp	6 (10.3)	5 (6.1)	4 (3.7)	15 (6)	0.237
Anal fissure	1 (1.7)	5 (6.1)	5 (4.7)	11 (4.4)	0.461
Peptic ulcer	4 (6.9)	4 (4.9)	1 (0.9)	9 (3.6)	0.114
Colonic ulcers	2 (3.4)	5 (6.1)	2 (1.9)	9 (3.6)	0.305
Rectal ulcer	0 (0)	2 (2.4)	7 (6.5)	9 (3.6)	0.078
Intestinal nodular lymphoid hyperplasia	4 (6.9)	4 (4.9)	0 (0)	8 (3.2)	<b>0.034</b>
Foreign body ingestion	6 (10.3)	1 (1.2)	0 (0)	7 (2.8)	<b>&lt; 0.0001</b>
Esophageal varices	4 (6.9)	0 (0)	2 (1.9)	6 (2.4)	<b>0.029</b>
Esophageal ulcer	2 (3.4)	1 (1.2)	2 (1.9)	5 (2)	0.646
Meckel's diverticulum	1 (1.7)	2 (2.4)	1 (0.9)	4 (1.6)	0.717
Mallory Weiss syndrome	1 (1.7)	2 (2.4)	0 (0)	3 (1.2)	0.291
Hemorrhoids	1 (1.7)	0 (0)	1 (0.9)	2 (0.8)	0.524
Duodenal varices	1 (1.7)	0 (0)	0 (0)	1 (0.4)	0.195
Colonic angiodysplasia	0 (0)	0 (0)	1 (0.9)	1 (0.4)	0.518
CMPA	1 (1.7)	0 (0)	0 (0)	1 (0.4)	0.195

<sup>a</sup>Some patients may have more than one cause.


<sup>b</sup>Pearson  $\chi^2$ . Data are presented as number and percentage. P value < 0.05 was considered statistically significant. CMPA: cow's milk protein allergy.

contrast, Hassoon *et al*[9] reported a lower percentage of 5.2%. This variation might be explained by the fact that melena is usually unnoticeable by the patients and their families, especially in older children; it typically appears after hematemesis, which leads to delayed presentation and diagnosis[9].


Recurrent abdominal pain was the most frequent associated symptom with GIB ( $n = 59$ , 23.6%). This result is comparable to that reported by Gimiga *et al*[6] (26.3%) and Zahmatkeshan *et al*[12] (24.5%), where abdominal pain was the most common accompanying symptom. However, Rafeey *et al*[10] reported a lower percentage (7.4%) of abdominal pain in children over the age of one. Several mechanisms can explain the association between abdominal pain and GIB, depending on the underlying etiology. Peptic or duodenal ulcers can develop due to the loss of the protective mucosal layer, which normally protects against gastric acid[17]. Deep or perforated ulcers irritate the gastric nerves or cause peritonitis, which causes severe abdominal pain[18].

In the current study, physical examination was unremarkable in most of the patients ( $n = 218$ , 87.2%). However, pallor was documented in 11 (4.4%) patients. Pallor might be related to the severity of bleeding and amount of blood loss. Cleveland *et al*[4] found that esophageal varices and duodenal ulcers are the most common conditions leading to anemia. Persistent or recurrent iron-deficiency anemia is a sign of obscure GIB[3]. Zahmatkeshan *et al*[12] reported iron deficiency anemia in 1.4% of the patients. However, Gimiga *et al*[6] reported a much higher incidence of hypochromic microcytic anemia (61.9%).

The causes of GIB in children vary based on the diagnostic approach used (radiological or endoscopic). Specific causes of GIB can be diagnosed based on imaging findings even before endoscopic intervention, such as foreign body ingestion, esophageal varices, intussusception, Meckel's diverticulum, and IBD[19,20]. The role of radiology in the management of children with GIB differs according to patient age and clinical presentation[19]. Radiological imaging is frequently requested after a negative endoscopic evaluation or for undetermined causes or bleeding sites[19]. Abdominal



**Figure 2 Comparison between upper and lower gastrointestinal bleeding according to the age groups.** <sup>1</sup>Pearson  $\chi^2$  Test. UGIB: Upper gastrointestinal bleeding; LGIB: Lower gastrointestinal bleeding. 250 patients had a total of 270 bleeding episodes (20 patients had both UGIB and LGIB).



**Figure 3 Endoscopic findings in children presented with gastrointestinal bleeding.** A: Esophageal varices; B: Rectal polyp; C: Inflamed ulcerated colonic mucosa in patient with ulcerative colitis.

ultrasonography, barium studies, computed tomography, magnetic resonance imaging, nuclear scintigraphy, and selective angiography may play a role in identifying the underlying pathology and exact source of bleeding[19]. In our study, IBD was the main cause of GIB in adolescents, whereas foreign body ingestion and esophageal varices were the main causes if GIB in preschool children. Çolak recently reported a small bowel intussusception caused by Meckel’s diverticulitis in a 10-year-old girl in whom it was diagnosed using radiological images even before endoscopic evaluation[20].

With advancements in the field of medical interventions, endoscopy has become the modality of choice for diagnosing GIB in children[21]. All the patients in this study were diagnosed *via* endoscopic examination. The causes of GIB among the published studies were diverse, which might account for the variation in disease distribution among different countries (Table 4). The most common cause of GIB in this study was IBD. Gimiga *et al*[6] reported ulcerative colitis (22%) as the second-most common cause of GIB. However, Jafari *et al*[2] and Zahmatkeshan *et al*[12] reported IBD to be a rare cause of GIB, accounting for only 4% and 5.8% of the study population, respectively. In patients with IBD, deep ulcers secondary to colitis might be sufficient to disrupt the underlying blood vessels of the inflamed and friable mucosa, causing GIB[11].

The second-most cause of GIB in this study was gastritis, with *Helicobacter pylori* infection accounting for 20% of the cases. However, Jafari *et al*[2] (7.1%), Rafeey *et al*[10] (14.5%), and Hassoon *et al*[9] (19.6%) reported lower percentages of gastritis in their study population. This difference might be due to the high prevalence of *Helicobacter pylori* gastritis in our community (57%)[22]. Duodenal ulcers were noted in 20 (8%) of our patients, which is comparable to that reported by Hassoon *et al*[9] (7.4%), Rafeey *et al*[10] (7.6%), and Cleveland *et al*[4] (8.2%). Duodenal ulcers can be caused by mucosal irritation from infections, such as *Helicobacter pylori* infection, or certain medications, such as



**Table 4 Summary of previous studies of gastrointestinal bleeding in children from neighboring countries and worldwide**

Ref.	Country	n	Age (yr)	Sex	Bleeding site	Two most common symptoms (%)	Two most common causes (%)
Isa <i>et al.</i> , 2023	Bahrain <sup>a</sup>	250	≤ 18	M > F	Both	Per rectal bleeding (60.4); Hematemesis (46.8)	Inflammatory bowel disease (30.8); Gastritis (28)
Jafari <i>et al</i> [2], 2018	Iran	113	< 18	M > F	Both	Hematemesis (40.7); Coffee ground vomitus (38)	Prolapse gastropathy (18.6) for UGIB; Polyps (32.5) for LGIB
Rafeey <i>et al</i> [10], 2013	Iran	447	< 18	M > F	UGIB	Hematemesis (26.85); Melena (13.42)	Erosive esophagitis (40); Gastric erosion (17)
Zahmatkeshan <i>et al</i> [12], 2012	Iran	363	< 18	M > F	LGIB	Hematochezia (80.2); Bloody diarrhea (18.1)	Juvenile polyp (23.1); Lymphoid nodular hyperplasia (18.2)
Hassoon <i>et al</i> [9], 2012	Iraq	58	4 d-18	M > F	UGIB	Hematemesis (58.5); Melena or hematochezia (5.2)	Esophageal varices (39); Gastric erosions (19.6)
Gimiga <i>et al</i> [6], 2015	Romania	118	< 18	M > F	LGIB	Hematochezia (54.2); Rectorrhagia (40.7)	Solitary colorectal polyps (33); Ulcerative colitis (22)
Cleveland <i>et al</i> [4], 2012	USA	158	< 17	M > F	UGIB	Hematemesis (73.4); Melena (20.8)	Prolapse gastropathy syndrome (12.7); gastric erosions/ulcers (10.8)

<sup>a</sup>The present study. UGIB: upper gastrointestinal bleeding; LGIB: lower gastrointestinal bleeding; USA: United states of America.

nonsteroidal anti-inflammatory drugs[23].

Esophagitis occurs due to irritation of the esophageal mucosal lining by gastric acids[11]. In this study, esophagitis was observed in 15 (6%) patients, which is similar to that seen in Khan *et al*[24] (4%) and Hassoon *et al*[9] (4.9%). However, other studies report a higher percentage, ranging between 9.5% and 40%[2,4,10].

Intestinal polyposis is another cause of GIB. In this study, polyps were detected in only 15 (6%) patients. However, Zahmatkeshan *et al*[12], Jafari *et al*[2] and Gimiga *et al*[6], reported polyps as the most common cause of LGIB, representing 23.1%, 32.5%, and 33% of their study population, respectively.

In the present study, 4% of our patients underwent one or more therapeutic interventions. Polypectomy was the commonest (1.6%), followed by injection sclerotherapy (1.2%). Comparably, Lee *et al*[25] reported that 2.3% of their patients underwent polypectomy; however, they reported a higher percentage of injection sclerotherapy (21%).

Like most retrospective studies, this study was limited by missing medical data such as vital signs and hemoglobin levels at GIB presentation. In addition, this was a single-center study that included only patients who were admitted to the hospital and underwent an endoscopic procedure. Subsequently, children with GIB who presented to the pediatric emergency department and were discharged home, such as those with Mallory–Weiss syndrome, and those who were admitted but did not undergo endoscopy, were excluded. Therefore, this study might not reflect the true incidence of GIB in children. Another limitation is that, in 2020, most endoscopic procedures were cancelled due to the coronavirus pandemic. This resulted in a marked reduction in the total number of patients with GIB who underwent endoscopy that year.

Despite these limitations, this study is important because it is the first study from Bahrain to focus on children presenting with GIB. Moreover, this study covers most aspects of GIB, including the incidence, clinical presentation, diagnosis, therapeutic interventions, and outcomes in children. Furthermore, this study covered both types of GIB (upper and lower), which is considered a strength of this study; most previous studies reported only one type of bleeding. The findings of this study are essential for pediatricians or gastroenterologists to tailor their management strategies based on the most frequent causes of GIB according to the patient's age group. Additionally, it forms a strong foundation for future research.

## CONCLUSION

Gastrointestinal bleeding in children is an alarming condition that has increased significantly over the last two decades. LGIB was more common in our study population than UGIB was. Additionally, most of the patients presented with bleeding per rectum. Endoscopic procedures are the best modality for diagnosing this condition. IBD was the most frequent cause of LGIB, whereas gastritis was the most frequent cause of UGIB. The causes of GIB varied according to the patients' age group and were different from those reported in neighboring countries and the rest of the world. The children in our study had particularly good outcomes. Further studies that include GIB identified in an emergency

setting and those that assess the effect of GIB on patients' hemodynamic stability, the need for blood transfusion, and the long-term impact of this condition are needed.

## ARTICLE HIGHLIGHTS

### **Research background**

Gastrointestinal bleeding (GIB) is a serious health problem worldwide, particularly in childhood. The incidence, clinical presentation, and causes of pediatric GIB vary among countries.

### **Research motivation**

Due to limited data on GIB in the pediatric population in the Middle East, we were motivated to study this health problem in Bahrain.

### **Research objectives**

To assess the incidence, clinical presentation, causes, and outcomes of GIB in children at the main tertiary hospital in Bahrain over the last two decades and to stratify the causes of GIB according to the patients' age group.

### **Research methods**

We retrospectively reviewed and collected the demographic data, clinical presentation, endoscopic findings, and outcomes of children with GIB admitted to the Pediatric Department at Salmaniya Medical Complex, Kingdom of Bahrain, from medical records between 1995 and 2022. The causes of GIB were compared according to patient's age at presentation.

### **Research results**

A total of 250 patients with GIB were included in this study. The median incidence was 2.6/100000 per year (interquartile range, 1.4–3.7) with a significantly increasing trend over the last two decades ( $P < 0.0001$ ). Most patients were males ( $n = 144, 57.6\%$ ). The median age at diagnosis was 9 years (IQR, 5–11). Ninety-eight (39.2%) patients required upper gastrointestinal endoscopy alone, 41 (16.4%) required colonoscopy alone, and 111 (44.4%) required both. Lower GIB (LGIB) was more frequent ( $n = 151, 60.4\%$ ) than upper GIB (UGIB) ( $n = 119, 47.6\%$ ). There were no significant differences in sex ( $P = 0.710$ ), age ( $P = 0.185$ ), or nationality ( $P = 0.525$ ) between the two groups. Abnormal endoscopic findings were detected in 226 (90.4%) patients. Inflammatory bowel disease (IBD) was the most common cause of LGIB ( $n = 77, 30.8\%$ ), whereas gastritis was the most common cause of UGIB ( $n = 70, 28\%$ ). IBD and undetermined cause for bleeding were higher in the 10–18 years group ( $P = 0.026$  and  $P = 0.017$ , respectively), while intestinal nodular lymphoid hyperplasia, foreign body ingestion, and esophageal varices were more common in the 0–4 years group ( $P = 0.034$ ,  $P < 0.0001$ , and  $P = 0.029$ , respectively). Ten (4%) patients underwent one or more therapeutic intervention. The median follow-up period by endoscopy was two years (IQR, 0.5–3). No mortality was reported in this study.

### **Research conclusions**

GIB in children is an alarming condition that is increasing significantly. LGIB were more frequent than UGIB. IBD was the most common cause of LGIB, whereas gastritis was the most common cause of UGIB in our children. The cause for GIB varied based on patient age and differed from those reported in neighboring countries and the rest of the world.

### **Research perspectives**

Further studies are needed that include children with GIB from an emergency setting and studies that assess the effect of this bleeding on patients' hemodynamic stability, the need for blood transfusion, and the long-term impact of this condition.

---

## ACKNOWLEDGEMENTS

We gratefully acknowledge all health care providers who provided the needed care for children with gastrointestinal bleeding in the Pediatric Departments at Salmaniya Medical Complex, Kingdom of Bahrain.

## FOOTNOTES

**Author contributions:** Isa HM was the main contributor in study design, literature review, data analysis, drafting manuscript, and over-sight for all phases of the project and the final approval of the version to be published; Alkharsi FA was responsible for literature review, data collection, drafting and revising manuscript; Ebrahim HA, Walwil KJ and Diab JA were responsible for literature review, data collection and revising manuscript; Alkowari NM was responsible for literature review, drafting and revising manuscript; All the authors have read and approved the final manuscript.

**Institutional review board statement:** This study was ethically approved by the Research and Research Ethics Committee, Salmaniya Medical Complex, Government hospitals, Kingdom of Bahrain (IRB number: 6170122, January 17, 2022).

**Informed consent statement:** Consent was not needed as the study was retrospective without exposure to the patients' data.

**Conflict-of-interest statement:** All the Authors have no conflict of interest related to the manuscript.

**Data sharing statement:** Data are available upon reasonable request.

**STROBE statement:** The authors have read the STROBE Statement—checklist of items, and the manuscript was prepared and revised according to the STROBE Statement—checklist of items.

**Open-Access:** This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

**Country/Territory of origin:** Bahrain

**ORCID number:** Hasan M Isa 0000-0001-6022-5576; Fatema A Alkharsi 0000-0002-0387-1587; Hana A Ebrahim 0000-0003-1989-1192; Kamel J Walwil 0000-0003-2810-5106; Jumana A Diab 0000-0002-5968-7252; Nafelah M Alkowari 0000-0002-7691-8067.

**S-Editor:** Liu JH

**L-Editor:** A

**P-Editor:** Liu JH

## REFERENCES

- Bresci G.** Occult and obscure gastrointestinal bleeding: Causes and diagnostic approach in 2009. *World J Gastrointest Endosc* 2009; **1**: 3-6 [PMID: 21160643 DOI: 10.4253/wjge.v1.i1.3]
- Jafari SA, Kiani MA, Kianifar HR, Mansooripour M, Heidari E, Khalesi M.** Etiology of gastrointestinal bleeding in children referred to pediatric wards of Mashhad hospitals, Iran. *Electron Physician* 2018; **10**: 6341-6345 [PMID: 29629057 DOI: 10.19082/6341]
- Romano C, Oliva S, Martellosi S, Miele E, Arrigo S, Graziani MG, Cardile S, Gaiani F, de'Angelis GL, Torroni F.** Pediatric gastrointestinal bleeding: Perspectives from the Italian Society of Pediatric Gastroenterology. *World J Gastroenterol* 2017; **23**: 1328-1337 [PMID: 28293079 DOI: 10.3748/wjg.v23.i8.1328]
- Cleveland K, Ahmad N, Bishop P, Nowicki M.** Upper gastrointestinal bleeding in children: an 11-year retrospective endoscopic investigation. *World J Pediatr* 2012; **8**: 123-128 [PMID: 22573422 DOI: 10.1007/s12519-012-0350-8]
- Saljoughian M.** Gastrointestinal bleeding: An alarming sign. *US Pharm* 2009; **34**: HS12-HS16. Available from: [https://www.researchgate.net/publication/289783639\\_Gastrointestinal\\_bleeding\\_An\\_alarming\\_sign](https://www.researchgate.net/publication/289783639_Gastrointestinal_bleeding_An_alarming_sign)
- Gimiga N, Burlea M, Diaconescu S, Oлару C.** An assessment of the causes of lower gastrointestinal bleeding in a children's hospital in northeastern Romania. *Arch Biol Sci* 2015; **67**: 715-720 [DOI: 10.2298/ABS140917043G]
- WHO Multicentre Growth Reference Study Group.** WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development. Geneva: World Health Organization, 2006
- GBD 2017 Inflammatory Bowel Disease Collaborators.** The global, regional, and national burden of inflammatory bowel disease in 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet Gastroenterol Hepatol* 2020; **5**: 17-30 [PMID: 31648971 DOI: 10.1016/S2468-1253(19)30333-4]
- Hassoon AJ, AlMaeeni AA, Matloub HY.** Upper gastrointestinal bleeding in children. *J Fac Med Baghdad* 2012; **54**: 223-227. Available from: <https://iqjmc.uobaghdad.edu.iq/index.php/19JFacMedBaghdad36/article/download/722/526>
- Rafeey M, Shoaran M, Majidy H.** Diagnostic endoscopy and clinical characteristics of gastrointestinal bleeding in children: a 10-year retrospective study. *Iran Red Crescent Med J* 2013; **15**: 794-797 [PMID: 24616788 DOI: 10.5812/ircmj.7075]

- 11 **Nationwide children's**. GI Bleeding. 2022. Available from: <https://www.nationwidechildrens.org/conditions/gi-bleeding>
- 12 **Zahmatkeshan M**, Fallahzadeh E, Najib K, Geramizadeh B, Haghghat M, Imanieh MH. Etiology of lower gastrointestinal bleeding in children: a single center experience from southern Iran. *Middle East J Dig Dis* 2012; **4**: 216-223 [PMID: 24829660]
- 13 **Attard TM**, Miller M, Pant C, Kumar A, Thomson M. Mortality associated with gastrointestinal bleeding in children: A retrospective cohort study. *World J Gastroenterol* 2017; **23**: 1608-1617 [PMID: 28321162 DOI: 10.3748/wjg.v23.i9.1608]
- 14 **Kalyoncu D**, Urganci N, Cetinkaya F. Etiology of upper gastrointestinal bleeding in young children. *Indian J Pediatr* 2009; **76**: 899-901 [PMID: 19904505 DOI: 10.1007/s12098-009-0195-x]
- 15 **Banisalah AA**, Mraiat ZM. Upper gastrointestinal bleeding in Irbid, Jordan. *Rawal Med J* 2007; **32**: 105-108. Available from: <https://www.semanticscholar.org/paper/Upper-Gastrointestinal-Bleeding-in-Irbid%2C-Jordan-Banisalah-Mraiat/738542c47059fe624e617ca753d5c378ddc8d8fa>
- 16 **Almadi MA**, Almutairdi A, Alruzug IM, Aldarsouny TA, Semaan T, Aldaheer MK, AlMustafa A, Azzam N, Batwa F, Albawardy B, Aljebreen A. Upper gastrointestinal bleeding: Causes and patient outcomes. *Saudi J Gastroenterol* 2021; **27**: 20-27 [PMID: 33047678 DOI: 10.4103/sjg.SJG\_297\_20]
- 17 **Malik TF**, Gnanapandithan K, Singh K. Peptic ulcer disease. 2022. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK534792/>
- 18 **Di Saverio S**, Bassi M, Smericri N, Masetti M, Ferrara F, Fabbri C, Ansaloni L, Ghersi S, Serenari M, Coccolini F, Naidoo N, Sartelli M, Tugnoli G, Catena F, Cennamo V, Jovine E. Diagnosis and treatment of perforated or bleeding peptic ulcers: 2013 WSES position paper. *World J Emerg Surg* 2014; **9**: 45 [PMID: 25114715 DOI: 10.1186/1749-7922-9-45]
- 19 **Racadio JM**, Agha AK, Johnson ND, Warner BW. Imaging and radiological interventional techniques for gastrointestinal bleeding in children. *Semin Pediatr Surg* 1999; **8**: 181-192 [PMID: 10573428 DOI: 10.1016/s1055-8586(99)70025-9]
- 20 **Çolak E**. Meckel' diverticulitis causing small bowel intussusception: A case report. *Pediatr Pract Res* 2021; **9**: 97-99 [DOI: 10.21765/ppjournal.937537]
- 21 **ASGE Standards of Practice Committee**, Fisher L, Lee Krinsky M, Anderson MA, Appalaneni V, Banerjee S, Ben-Menachem T, Cash BD, Decker GA, Fanelli RD, Friis C, Fukami N, Harrison ME, Ikenberry SO, Jain R, Jue T, Khan K, Maple JT, Strohmeyer L, Sharaf R, Dominitz JA. The role of endoscopy in the management of obscure GI bleeding. *Gastrointest Endosc* 2010; **72**: 471-479 [PMID: 20801285 DOI: 10.1016/j.gie.2010.04.032]
- 22 **Zainaldeen HA**, Isa HM. Helicobacter pylori associated gastritis among Bahraini children presenting with abdominal pain. *J Bahrain Med Soc* 2010; **22**: 52-54. Available from: [https://www.researchgate.net/publication/286378046\\_Helicobacter\\_pylori\\_associated\\_gastritis\\_among\\_Bahraini\\_children\\_presenting\\_with\\_abdominal\\_pain](https://www.researchgate.net/publication/286378046_Helicobacter_pylori_associated_gastritis_among_Bahraini_children_presenting_with_abdominal_pain)
- 23 **Barth B**. 7 common causes of pediatric GI bleeding, plus treatment information. 2019. Available from: <https://utswmed.org/medblog/gi-bleeding-children/>
- 24 **Khan MR**, Ahmed S, Ali SR, Maheshwari PK, Jamal MS. Spectrum of upper GI endoscopy in pediatric population at a tertiary care centre in Pakistan. *Open J Pediatr* 2014; **4**: 180-184 [DOI: 10.4236/ojped.2014.43025]
- 25 **Lee WS**, Zainuddin H, Boey CC, Chai PF. Appropriateness, endoscopic findings and contributive yield of pediatric gastrointestinal endoscopy. *World J Gastroenterol* 2013; **19**: 9077-9083 [PMID: 24379634 DOI: 10.3748/wjg.v19.i47.9077]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

