

BMJ Open Social inequalities in aggravating factors of somatic symptom persistence (SOMA.SOC): study protocol for a mixed-method observational study focusing on irritable bowel syndrome and fatigue

Olaf von dem Knesebeck , Rieke Barbek, Anna Christin Makowski

To cite: von dem Knesebeck O, Barbek R, Makowski AC. Social inequalities in aggravating factors of somatic symptom persistence (SOMA.SOC): study protocol for a mixed-method observational study focusing on irritable bowel syndrome and fatigue. *BMJ Open* 2023;**13**:e070635. doi:10.1136/bmjopen-2022-070635

► Prepublication history for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2022-070635>).

Received 29 November 2022
Accepted 12 April 2023



© Author(s) (or their employer(s)) 2023. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

Institute of Medical Sociology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

Correspondence to
Professor Olaf von dem Knesebeck;
o.knesebeck@uke.de

ABSTRACT

Introduction Some studies indicate that persistent somatic symptoms (PSS) are more prevalent among individuals with a low socioeconomic status (SES) and a migration background. However, factors explaining social inequalities in PSS are largely unknown. It is expected that aggravating factors of PSS like illness perception, illness beliefs (health literacy, stigma), illness behaviour and health anxiety may play an important role for this explanation. The SOMA.SOC study will examine social inequalities (according to SES and migration) in factors contributing to symptom persistence in irritable bowel syndrome (IBS) and fatigue.

Methods and analysis The project will collect both quantitative and qualitative data. Quantitative data will be gathered via a representative telephone survey in Germany (N=2400). A vignette design will be used depicting patients varying in sex, condition (IBS/fatigue), occupational status (low/high) and migration (yes/no). In the survey, we will assess public knowledge and beliefs (eg, health literacy), attitudes (stigma) and personal experiences with the condition (eg, somatic symptom burden). Complementary, longitudinal qualitative interviews will be conducted with patients (n=32 at three time points, resulting in N=96 interviews) who will also vary according to sex, condition, occupational status and migration. Patients will be recruited from primary care practices in Hamburg. The interviews will cover origin and development of the condition, coping and help-seeking as well as social interactions and perception of the disease by others (eg, perceived stigma). SOMA.SOC is part of the interdisciplinary SOMACROSS (Persistent SOMAtic Symptoms ACROSS Diseases) research unit.

Ethics and dissemination The study protocol was approved by the Ethics Committee of the Hamburg Medical Association on 25 January 2021 (reference number: 2020–10194- B0-ff). Informed consent will be obtained from all participants. The main findings will be submitted for publication in peer-reviewed journals within 12 months of study completion.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ Social inequalities according to socioeconomic status and migration in a range of aggravating factors (illness perception, symptom beliefs, illness/treatment experiences, illness behaviour and health anxiety) in irritable bowel syndrome (IBS) and fatigue will be explored.
- ⇒ By combining quantitative and qualitative data, we expect a better and more comprehensive understanding of inequalities in aggravating factors of persistent somatic symptoms (PSS).
- ⇒ With the longitudinal design of the qualitative study part, we expect to gain insight into the course of symptoms and changes of experiences over time.
- ⇒ Due to the observational design, causal conclusions cannot be drawn from the study.
- ⇒ As the study is focused on IBS and fatigue, results cannot be generalised to other PSS.

INTRODUCTION

Background

Health inequalities according to socioeconomic status and migration

National and international studies in social epidemiology have consistently shown that socioeconomic status (SES, ie, education, income and/or occupational position) as well as migration and ethnicity are important social determinants of health.^{1,2} Accordingly, people with a low SES generally experience worse health and well-being as well as higher morbidity (including higher risk of chronic disease course) and mortality rates than those with a high SES. Material (eg, material deprivation, unfavourable working, housing and neighbourhood conditions), psychosocial (eg, social isolation, psychosocial stress, coping resources) and behavioural factors (eg, smoking, exercise, alcohol consumption,

nutrition) contribute to the explanation of such health inequalities. Recent studies indicate that differences in knowledge and beliefs about health and illness ('health literacy') are also important for the explanation of health inequalities.^{3,4} However, results on the explanatory contribution of these factors are inconsistent as there is variation according to the disease under study. For example, socioeconomic inequalities of coronary heart disease can largely be explained by differences in classic cardiovascular risk factors (ie, high blood pressure, high cholesterol, diabetes mellitus and cigarette smoking),⁵ while poorer coping styles, ongoing life events, stress exposure and weaker social support are important factors to explain the higher prevalences of mental disorders in lower SES groups.^{6,7}

Although migrants are often, at least initially, relatively healthy compared with the non-migrant population in the host country, available data suggest that they tend to be more vulnerable to certain communicable diseases, occupational health hazards, injuries, poor mental health and maternal and child health problems.^{2,8} Studies furthermore show that there are large differences between migrants in terms of their socioeconomic situation but these differences and their health-related consequences are far from being adequately understood. As migration status and SES reflect different aspects of social inequalities, that are not independent, it is reasonable to explore both aspects and their relation to health simultaneously.⁹

Social inequalities in somatic symptoms

Regarding persistent somatic symptoms (PSS), there are a few studies investigating inequalities according to SES and migration. Using German population survey data, Hinz *et al*¹⁰ found that a low SES is associated with a higher risk for somatic symptoms measured with the Patient Health Questionnaire-15 (PHQ-15).¹¹ This result was confirmed by an analysis of the Gutenberg Health Study conducted in the Rhine-Main Region.¹² Based on the German national Cohort study, Morawa *et al*¹³ showed that persons of Turkish origin report more symptoms according to the PHQ-15 compared with Germans. Moreover, psychiatric patients with a migration background had a significantly higher current symptom load, especially by somatic symptoms, compared with those without migration background.¹⁴

Thus, overall, results suggest that somatic symptoms are more prevalent among individuals with a low SES and a migration background in Germany. Respective empirical studies mostly use sum scales comprising a number of symptoms but magnitude and mechanisms of inequalities may differ depending on symptom under study. Considering specific PSS with high prevalence such as fatigue (about 30% in Germany^{15,16}) or irritable bowel syndrome (IBS, about 10–20% in Germany¹⁷), research on inequalities is scarce, particularly in Germany. Jason *et al*¹⁸ found highest levels of fatigue among minority groups and persons with lower levels of education and occupational status in the USA. A German population

study also revealed increased rates of fatigue among low SES groups.¹⁶ Another study indicates that there are interactions between SES and ethnicity in the association with fatigue.¹⁹ In terms of IBS, there are studies showing higher prevalences in lower SES groups²⁰ but others find a reverse association or no socioeconomic inequalities.^{21–23} There is a paucity of data on differences according to migration status in the epidemiology of IBS.

Associations between social inequalities and aggravating factors of PSS

Aetiological models suggest that psychosocial, behavioural and biomedical factors, as well as their interaction, contribute to the persistence and aggravation of somatic symptom burden.²⁴ Among known psychosocial maintaining or aggravating factors in PSS are health anxiety, treatment experiences and illness beliefs, perception and behaviour. We expect that these factors may partially mediate any differences that might be seen in rates of PSS between those with differential socioeconomic or migration status. However, empirical studies analysing associations between social inequalities and aggravating factors are scarce and thus, there is not much known about the mechanisms that potentially can explain inequalities in PSS.

Studies on illness beliefs examine to what extent an illness is recognised as such and what beliefs about the causes, the course and the treatment of this illness are prevalent among the public. In recent years, these studies have been discussed with reference to the concept of health literacy. Health literacy entails 'the public's knowledge, motivation, and competence to access, understand, appraise, and apply health information in order to make judgments and take decisions in everyday life concerning healthcare'.²⁵ Deficits in health literacy mean that people do not know what they can do for prevention, delay or avoid seeking treatment, view recommended treatments with suspicion or are unsure how to assist those afflicted.²⁶ Limited health literacy has been found among people with a low SES and migrant background.^{4,27,28} For example, a study analysing differences between German and Turkish patients suffering from somatoform disorders showed that Turkish patients believed significantly stronger in supernatural causes of their disease.²⁹ These beliefs reduced the motivation to take up psychotherapy. A lack of health literacy has been found to be associated with negative attitudes towards persons with mental illness,²⁸ which can be conceptualised as public stigma. This entails reactions of the general public towards a group based on stereotypes about this group. In this regard, results suggest that respondents with higher SES desire less social distance from the mentally ill and express more liberal views.³⁰ Research also suggests that public stigma is perceived and often internalised by the persons afflicted. Perceived and internalised stigma have been found to predict worse health outcomes among patients with IBS.³¹ With regards to fatigue, studies report higher levels of perceived stigma when compared with IBS³² and participants report

feelings of estrangement and the belief that others attribute fatigue solely to psychological causes.³³

Only a few studies examined inequalities in symptom or illness perception and health anxiety. In terms of the former, perceived symptom burden seems to be positively associated with education,³⁴ while results on differences according to ethnicity and migration are inconsistent.^{34 35} As for health anxiety, a recent meta-analysis showed a lower risk for health anxiety in people with higher SES.³⁶ Another meta-analysis indicated a higher risk of health anxiety in migrants and ethnic minorities compared with the majority population in North America.³⁷ Regarding illness behaviour, reviews from Germany overall indicate a lower usage of healthcare among migrants, although the results vary in terms of healthcare sector, indicator of healthcare usage and migrant population under study.³⁸ Low SES groups show lower usage regarding medical specialist consultations and prevention services.³⁹

OBJECTIVES AND HYPOTHESES

Against this background, the present study, social inequalities in aggravating factors of somatic symptom persistence (SOMA.SOC), was developed. The objective of the study is to examine social inequalities (according to SES and migration) in factors contributing to symptom persistence in IBS and fatigue. We chose these two specific conditions as they appear relatively frequently in the German population. In terms of IBS, we especially refer to the cardinal symptoms of recurrent abdominal pain and altered bowel movements (as specified in the International Classification of Diseases 10th edition, ICD-10).⁴⁰ Regarding fatigue, different symptoms like emotional and cognitive limitations (eg, lack of motivation and reduced performance) as well as physical issues (eg, muscular weakness) are to be considered (as mentioned in the national guideline¹⁵ and covered in

the ICD-10).⁴⁰ The following aggravating factors will be explored: illness perception, knowledge and beliefs about the symptoms, illness and treatment experiences, illness behaviour and health anxiety.

More specifically the following hypotheses will be explored: (1) SES and migration are associated with factors contributing to the symptom persistence (aggravating factors) of IBS and fatigue. (2) There are differences in public perceptions of and public beliefs about IBS and fatigue according to the SES and migration status of the person expressing the symptoms.

The SOMA.SOC study is part (project 6) of the interdisciplinary research unit Persistent SOMatic symptoms ACROSS diseases—from risk factors to modification (SOMACROSS). SOMACROSS aims to identify disease-overarching and disease-specific biopsychosocial risk factors and mechanisms for the persistence of somatic symptoms.⁴¹ SOMACROSS proposes a biopsychosocial ‘PSS working model’ as a starting point for the investigation of risk factors and aetiological mechanisms, based on the model by Henningsen *et al.*²⁴ The integration of the objectives of the SOMA.SOC project into the working model of SOMACROSS is shown in figure 1 (objectives of the SOMA.SOC study are framed in red). Accordingly, specific focus will be on socio-demographic predisposing factors and psychosocial maintaining/aggravating factors (figure 1).

METHODS AND ANALYSIS

Study design and participants

In order to gain insight into social inequalities in aggravating factors of IBS and fatigue, the project will make use of a mixed methods design by collecting both quantitative and qualitative observational data. For a better understanding of the methods, in the following, these two study parts (population survey and qualitative interviews with




Figure 1 Integration of the study objectives into the working model of the Persistent SOMatic symptoms ACROSS diseases—from risk factors to modification research unit.⁴¹ IBS, irritable bowel syndrome.

Table 1 Factors presented in the vignettes

Factor	Specifications	
Condition	Irritable bowel syndrome (IBS)	Fatigue (Fa)
Sex	Male (M)	Female (F)
Migration history	Migration history (Mh)	No migration history (nM)
Occupational status	High (lawyer, L)	Low (cleaner, C)

patients) will be presented separately. The conditions under study will be conceptualised differently in the two study parts: While we will use vignettes describing a person with typical symptoms of IBS or fatigue in the population survey, patients diagnosed with one of the conditions will be recruited for the qualitative interviews. Data collection started in April 2022 and will be finalised in summer 2024.

Design of the population survey

To explore public knowledge, beliefs, attitudes and experiences regarding PSS in the case of IBS and fatigue, a cross-sectional population survey will be conducted throughout Germany via computer-assisted telephone interviewing. In the beginning of the interview, a case story (vignette) will be presented to the respondents. In the vignette, a person with signs and symptoms indicative of either IBS or fatigue will be described. The vignettes will be varied according to sex (male/female), occupational status (high (lawyer)/low (cleaner)) and migration status (yes/no), resulting in 16 different case stories (see [table 1](#) for an overview of the varied factors and [table 2](#)

Table 2 Vignettes presented in the survey

Vignette no.	Factor combination
1	IBS M Mh L
2	IBS M nM L
3	IBS M Mh C
4	IBS M nM C
5	IBS F Mh L
6	IBS F nM L
7	IBS F Mh C
8	IBS F nM C
9	Fa M Mh L
10	Fa M nM L
11	Fa M Mh C
12	Fa M nM C
13	Fa F Mh L
14	Fa F nM L
15	Fa F Mh C
16	Fa F nM C

For abbreviations of the factor combinations please see [table 1](#).

for resulting vignettes). The vignettes will be designed in cooperation with clinical experts. By combining a vignette design with a survey, social inequalities may be examined from two perspectives: first, as social characteristics of the respondents themselves (please see above, hypothesis 1) and second, of the person described in the vignette (hypothesis 2). In terms of the first hypothesis, associations between social characteristics of the respondents and aggravating factors (eg, illness and treatment experiences, illness behaviour, health anxiety) will be analysed, while for the second hypothesis, differences in perceptions, attitudes and beliefs according to migration and occupational status of the person in the vignette will be examined.

Participants of the population survey

The survey is going to be representative of the resident population in Germany aged 18 years and older, and uses a dual-frame approach, that is, 30% of the gross sample consists of mobile numbers, while 70% will be selected from landline numbers. Thus, there is greater chance of including mobile-only users as well as target groups otherwise hard to reach. Landline numbers will be drawn from all registered private numbers at random. Additional computer-generated numbers will also allow for an ex-directory household. For interviews in households with more than one potential target person, a random selection of the target person is realised using the Kish-Selection-Grid.⁴² Regarding mobile telephone connections, the interview will be conducted with the person answering the phone. To gain a representative sample of the adult population living in Germany, data will be weighted by using an approach that includes a correction for household sizes, selection probabilities and distributions of specific socio-demographic characteristics.⁴³ A total of N=2400 participants will be included in the national telephone survey. The 16 vignettes will be randomly assigned to the respondents (n=150 per vignette). A sample size calculation showed that this number is sufficient to identify small sized differences with a statistical power of 80% and a type I error of 0.05 when comparing two vignettes (hypothesis 2) with regards to categorical outcomes using χ^2 tests. As for continuous outcomes, a sample size of n=150 per vignette allows for the detection of small to medium effect sizes based on t-tests, linear regression models or analyses of variance (statistical power 80%, type I error=0.05). Moreover, based on previous studies with a similar design,⁴⁴⁻⁴⁶ this number of respondents was found to be adequate to detect significant differences in public beliefs between vignettes. Similar surveys with a vignette design resulted in response rates between 48% and 54%.⁴⁴⁻⁴⁶

Design of the qualitative interviews

To complement the population perspective and to analyse social inequalities in factors and mechanisms of symptom persistence among patients with IBS and fatigue, qualitative, semi-structured interviews will be conducted using

a longitudinal design. Longitudinal qualitative research (LQR) is distinct from other qualitative approaches in the way that it incorporates time into the research process. It opens the possibility to make change (or stability) a key focus in analysis.⁴⁷ LQR aims at answering questions regarding lived experiences, for example, during the course of diseases, and how and why these experiences change over time. Against the background of the ongoing process of persistence of somatic symptoms, LQR will be helpful to capture individual courses, narratives and trajectories. Accordingly, interviews will be conducted at three measurement points (at the time of inclusion (t_0), after 6 months (t_1) and after 12 months (t_2)).

Participants of the qualitative interviews

Patients will be recruited from primary care practices in the Hamburg region and will be selected according to condition (IBS/fatigue), sex (male/female), occupational status (high/low) and migration status (yes/no). With respect to the occupational status of patients, the International Socio-Economic Index of Occupational Status according to Ganzeboom *et al*⁴⁸ will be used as a criterion for assigning scores. Scores will be dichotomised to obtain groups of high and low occupational position. Regarding the patients' migration history, we will be adapting the definition of the German Federal Statistical Office. This states that 'A person has a migration background if he or she or at least one parent was not born with German citizenship'.⁴⁹ For recruitment of the patients, the authors will cooperate with the Department of General Practice and Primary Care at the University Medical Center Hamburg-Eppendorf, which established a physicians' research network. Physicians are listed in a database that includes contact details of all primary care physicians residing in Hamburg and its suburbs as well as in adjacent regions (overall about 3000 primary care physicians). The database holds information on contact data, structure of the practice (single vs group practice), physician staff and involvement in recent research projects. Fifty practices will be randomly selected and contacted (figure 2). If interested in participation, the practice will receive detailed study information, a process description and a declaration of consent to participate. Moreover, the participating practices will be provided with all necessary study materials to be handed out to patients (eg, study information, privacy statement, short questionnaire to collect necessary patient data (eg, name, condition, sex, age, occupation, contact information)).

Patients will be selected based on a purposeful sampling procedure according to the four strata mentioned above (condition, sex, occupational status and migration). This procedure was chosen as it is expected to be useful to analyse social inequalities in factors and mechanisms of symptom persistence. Doctors are asked to inform patients who meet the diagnostic criteria (K58.1 (IBS-D), K58.2 (IBS-C), K58.3 (IBS-M), K58.8, R53, F48.0 or G93.3 according to ICD-10)⁴⁰ about the study when they present in the

practice. Those patients who are interested in participation are requested to return a short questionnaire with all relevant information about eligibility including contact details to the project group (either via mail, email or fax). Returned questionnaires will be sorted according to strata (condition, sex, migration status, occupational position). Thereafter, telephone calls will be conducted to provide further information if needed and to arrange an interview appointment. Informed consent will be signed by all participating patients. Participants will be free to conduct the interviews in person or by telephone. To acknowledge their participation and compensate for expenses, each patient will receive €15 for each completed interview (ie, €45 for the three interviews).

Based on experiences in previous projects, we expect a participation rate of at least 50% of the randomly selected practices.⁵⁰ This will lead to around 25 practices taking up patient recruitment. Physicians will receive €50 per patient included in the study. We also estimate the participation rate of the addressed patients at 50%, meaning that the practices will have to contact 96 patients (around 4 patients per practice in 6 months), leading to 48 patients with the required combination of characteristics who supply informed consent to participate. This will allow for a dropout rate of approximately 30% among participants during the course of the study, resulting in $n=32$ patients who will be included in all three measurement points (see figure 2). For each of the 16 combinations (IBS/fatigue), sex (male/female), migration (yes/no) and occupational status (high/low), two semi-structured interviews will be conducted. This sampling procedure will result in an equal number of patients ($n=16$) in terms of condition, sex, occupational status and migration status interviewed at three measurement points. Thus, altogether 96 interviews will be conducted.




Figure 2 Two-stage recruitment process of patients for the qualitative interviews via primary care practices in Hamburg.

Table 3 Instruments intended to use in the population survey

	Construct	Indicator/instrument
(1)	Health literacy	Labelling (recognition of disease); perceived causes; treatability; effectiveness of treatment options
	Illness perception	Illness Perception Questionnaire (brief version) ⁵¹
	Illness belief	Eg, 'Do you think that the person in the vignette has a real disorder?'
(2)	Public stigma	Emotional reactions towards those affected by IBS or fatigue Stereotypes ascribed to persons with IBS or fatigue
	Anticipated stigma	Modified version of the IBS Stigma Scale ⁵²
(3)	Illness behaviour	Eg, 'If you have the feeling of being affected by such complaints, you should get help as soon as possible.' Healthcare usage
	Illness experience	Own affliction; contact to someone affected
	Treatment experience	Treatment experience
	Somatic symptom burden	Somatic Symptom Scale-8 ⁵³
	Health anxiety	Whiteley Index-7 ⁵⁴
(4)	Socio-demographic data	Age, sex, educational attainment, occupational position, income, migration status, marital status

IBS, irritable bowel syndrome.

Assessment and study outcomes

Population survey

The standardised questionnaire used in the telephone survey will cover the following topics: (1) questions referring to knowledge and beliefs about IBS or fatigue, (2) attitudes towards persons affected from IBS or fatigue (stigma), (3) personal experiences with these symptoms and (4) socio-demographic characteristics of respondents (also see [table 3](#)^{51–54}). As for social inequalities, highest educational attainment, occupational position, monthly equivalence household income and migration status (no migration background, first and second generation migrants) will be assessed.⁵⁵ In Germany, about 25% of the general population have a migration background. Accordingly, in our sample (N=2400), we expect about 600 respondents to have a migration background. In terms of knowledge and beliefs about IBS or fatigue, health literacy, illness perceptions and illness beliefs will be measured. Regarding personal experiences with the symptoms, questions about illness behaviour, illness and treatment experiences, somatic symptom burden and health anxiety will be included.

Qualitative patient interviews

Semi-structured interviews will be conducted personally or by telephone (according to patient preference). To this end, a blend of closed and open-ended questions will be used that can be accompanied by follow-up questions to further specify mentioned topics. The precise formulation and order of the questions can be varied. This will allow to cover the main issues with all participants while also offering flexibility in discussing issues pertinent to individuals. The interviews will have a length of about

30 min and will be audio recorded on permission of the patient. A total of three sections are planned for the interview guide:

1. Origin, causes and development of the disease: Here, aspects of symptom/illness perception as well course and severity of symptoms and aspects of health anxiety will be covered.
2. Coping with the illness and help-seeking: In this segment, illness behaviour and experiences, health literacy as well as treatment experiences and treatment expectations will be focused.
3. Social interaction and perception by others: This complex will cover disclosure of illness and (perceived) stigma associated with the disorder.

Data analyses

Population survey

Associations between social characteristics of the respondents and aggravating factors will be analysed by calculating regression models (hypothesis 1). Furthermore, structural equation model approaches will be used to explore to what extent social inequalities in PSS are mediated by the aggravating factors under study and to what extent variance is mediated by other (unconsidered) factors. Differences in public perceptions, attitudes and beliefs according to migration and occupational status of the person in the vignette (hypothesis 2) will be examined using analyses of variance. The power calculation showed that the sample size is sufficient to detect significant main effects (eg, of migration status of the person in the vignette) as well as two-way interactions (eg, between migration and occupational status). In terms of the latter, interaction tests will be performed. Regarding categorical

outcomes (eg, recognition of condition presented in the vignette (yes or no)), χ^2 tests will be applied. Missing data will be imputed if more than 5% of the data are missing. The number of imputations will be chosen dependent on the proportion of missing data.

Qualitative patient interviews

Subject to the consent of the participants, the interviews will be audio recorded. Uniform transcription rules will be established before the start of the analysis, and compliance with these rules will be checked several times during the course of the process. For the analysis of the transcribed interviews, qualitative content analysis (QCA) will be used. A core aspect of QCA is to make the interpretation of text describable and verifiable by using content analysis rules previously formulated on the basis of a work plan. The technique of inductive category formation is particularly suitable for this study design. Inductive categories are developed from the interview material according to a given definition criterion, in order to reach the central aspects of the text. Step by step, categories will be formulated, which are revised within feedback loops eventually reduced to main categories and checked in respect to their reliability.⁵⁶

Mixed methods integration

Although data from both study parts are collected and analysed separately, they are seen as two complementing perspectives on social aspects that contribute to the symptom persistence of IBS and fatigue. In using these different approaches, we expect a better and more comprehensive understanding of inequalities in aggravating factors of PSS and how they are associated with patients' and the public's characteristics. Integration of findings of the two study parts will be feasible as there are overlaps in design (variation of social characteristics) and content of the survey/interview (please see assessment and study outcomes). The process of integrating findings from the two methods will take place at the interpretation stage when both data sets have been analysed separately. Accordingly, findings from the two components of the study will be listed in a 'convergence coding matrix'.⁵⁷ That means, it will be considered where results from the two study parts agree, offer complementary information on the same issue or seem to contradict each other.

Patient and public involvement

None.

ETHICS AND DISSEMINATION

The study is based on data from personal or telephone interviews. Thus, ethical and legal considerations are focused on data protection. We will strictly adhere to data protection regulations. Informed consent will be obtained from all participants. Respondents will be informed that participation in the study is voluntary and that withdrawal is possible at any time. There are no

specific risks for the participants in the study. The study protocol was approved by the Ethics Committee of the Hamburg Medical Association on 25 January 2021 (reference number: 2020–10194-BO-ff), including financial compensations for patients and physicians.

In accordance with the ethics committee approval and the German Research Foundation guidelines for the handling of research data, de-identified quantitative individual data will be made publicly available. Data sharing will follow the FAIR Data Principles (Findable, Accessible, Interoperable and Reusable). The main findings will be submitted for publication in peer-reviewed journals within 12 months of study completion.

Acknowledgements We would like to acknowledge the continued contributions of all colleagues of the Research Unit 5211 'SOMACROSS' (FOR 5211) to the design of this study.

Contributors OvdK and ACM developed the study. OvdK is the principal investigator of the study. RB is research associate in the study. All authors contributed to the refinement of the study protocol, read and approved the final version.

Funding The SOMA.SOC study is supported by the German Research Foundation (Deutsche Forschungsgemeinschaft, DFG) (project number: 445297796, please see <https://gepris.dfg.de/gepris/projekt/460374208?language=en>). It is part of the collaborative research unit 5211 (RU 5211) 'Persistent SOMAtic symptoms ACROSS diseases: from risk factors to modification (SOMACROSS)', also funded by the DFG (spokesperson: Professor Bernd Löwe, MD).

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Olaf von dem Knesebeck <http://orcid.org/0000-0001-6310-5879>

REFERENCES

- Marmot M. Social determinants of health inequalities. *Lancet* 2005;365:1099–104.
- Rechel B, Mladovsky P, Ingleby D, *et al*. Migration and health in an increasingly diverse Europe. *The Lancet* 2013;381:1235–45.
- World Health Organization. Health literacy. The solid facts. 2013. Available: <https://apps.who.int/iris/bitstream/handle/10665/128703/e96854.pdf>
- von dem Knesebeck O, Mnich E, Daubmann A, *et al*. Socioeconomic status and beliefs about depression, schizophrenia and eating disorders. *Soc Psychiatry Psychiatr Epidemiol* 2013;48:775–82.
- Kivimäki M, Shipley MJ, Ferrie JE, *et al*. Best-Practice interventions to reduce socioeconomic inequalities of coronary heart disease mortality in UK: a prospective occupational cohort study. *Lancet* 2008;372:1648–54.
- Lorant V, Deliège D, Eaton W, *et al*. Socioeconomic inequalities in depression: a meta-analysis. *Am J Epidemiol* 2003;157:98–112.
- Muntaner C, Eaton WW, Miech R, *et al*. Socioeconomic position and major mental disorders. *Epidemiol Rev* 2004;26:53–62.
- Razum O, Karrasch L, Spallek J. Migration: a neglected dimension of inequalities in health? *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2016;59:259–65.
- Malmusi D, Borrell C, Benach J. Migration-related health inequalities: showing the complex interactions between gender, social class and place of origin. *Soc Sci Med* 2010;71:1610–9.

- 10 Hinz A, Ernst J, Glaesmer H, *et al.* Frequency of somatic symptoms in the general population: normative values for the patient health questionnaire-15 (PHQ-15). *J Psychosom Res* 2017;96:27–31.
- 11 Kroenke K, Spitzer RL, Williams JBW, *et al.* The patient health questionnaire somatic, anxiety, and depressive symptom scales: a systematic review. *Gen Hosp Psychiatry* 2010;32:345–59.
- 12 Beutel ME, Wiltink J, Ghaemi Kerahrodi J, *et al.* Somatic symptom load in men and women from middle to high age in the gutenber health study-association with psychosocial and somatic factors. *Sci Rep* 2019;9:4610.
- 13 Morawa E, Dragano N, Jöckel K-H, *et al.* Somatization among persons with Turkish origin: results of the pretest of the German national cohort study. *J Psychosom Res* 2017;96:1–9.
- 14 Brandl EJ, Dietrich N, Mell N, *et al.* Clinical and sociodemographic differences between patients with and without migration background in a psychiatric outpatient service. *Psychiatr Prax* 2018;45:367–74.
- 15 Baum E, Donner-Banzhoff N, Maisel P. [Fatigue] [S3 guideline]. 2017.
- 16 Kocalevent RD, Hinz A, Brähler E, *et al.* Determinants of fatigue and stress. *BMC Res Notes* 2011;4:238.
- 17 Federal Ministry of Health. Irritable bowel syndrome. 2020. Available: <https://gesund.bund.de/reizdarmsyndrom>
- 18 Jason LA, Richman JA, Rademaker AW, *et al.* A community-based study of chronic fatigue syndrome. *Arch Intern Med* 1999;159:2129–37.
- 19 Bardwell WA, Burke SC, Thomas KS, *et al.* Fatigue varies by social class in African Americans but not Caucasian Americans. *Int J Behav Med* 2006;13:252–8.
- 20 Spiegel BMR. The burden of IBS: looking at metrics. *Curr Gastroenterol Rep* 2009;11:265–9.
- 21 Canavan C, West J, Card T. The epidemiology of irritable bowel syndrome. *Clin Epidemiol* 2014;6:71–80.
- 22 Lovell RM, Ford AC. Global prevalence of and risk factors for irritable bowel syndrome: a meta-analysis. *Clin Gastroenterol Hepatol* 2012;10:712–721.
- 23 Howell S, Talley NJ, Quine S, *et al.* The irritable bowel syndrome has origins in the childhood socioeconomic environment. *Am J Gastroenterol* 2004;99:1572–8.
- 24 Henningsen P, Zipfel S, Sattel H, *et al.* Management of functional somatic syndromes and bodily distress. *Psychother Psychosom* 2018;87:12–31.
- 25 Sørensen K, Van den Broucke S, Fullam J, *et al.* Health literacy and public health: a systematic review and integration of definitions and models. *BMC Public Health* 2012;12:80.
- 26 Jorm AF. Mental health literacy: empowering the community to take action for better mental health. *Am Psychol* 2012;67:231–43.
- 27 Bostock S, Steptoe A. Association between low functional health literacy and mortality in older adults: longitudinal cohort study. *BMJ* 2012;344:e1602.
- 28 Holman D. Exploring the relationship between social class, mental illness stigma and mental health literacy using British national survey data. *Health (London)* 2015;19:413–29.
- 29 Reich H, Bockel L, Mewes R. Motivation for psychotherapy and illness beliefs in Turkish immigrant inpatients in Germany: results of a cultural comparison study. *J Racial and Ethnic Health Disparities* 2015;2:112–23.
- 30 Angermeyer MC, Dietrich S. Public beliefs about and attitudes towards people with mental illness: a review of population studies. *Acta Psychiatr Scand* 2006;113:163–79.
- 31 Taft TH, Riehl ME, Dowjotas KL, *et al.* Moving beyond perceptions: internalized stigma in the irritable bowel syndrome. *Neurogastroenterol Motil* 2014;26:1026–35.
- 32 Looper KJ, Kirmayer LJ. Perceived stigma in functional somatic syndromes and comparable medical conditions. *J Psychosom Res* 2004;57:373–8.
- 33 Green J, Romei J, Natelson BH. Stigma and chronic fatigue syndrome. *Journal of Chronic Fatigue Syndrome* 1999;5:63–75.
- 34 Luo H, Lindell DF, Jurgens CY, *et al.* Symptom perception and influencing factors in Chinese patients with heart failure: a preliminary exploration. *IJERPH* 2020;17:2692.
- 35 Wilkins SS, Bourke P, Salam A, *et al.* Functional stroke mimics: incidence and characteristics at a primary stroke center in the middle East. *Psychosom Med* 2018;80:416–21.
- 36 Barbek RME, Makowski AC, von dem Knesebeck O. Social inequalities in health anxiety: a systematic review and meta-analysis. *J Psychosom Res* 2022;153:110706.
- 37 Barbek R, Henning S, Ludwig J, *et al.* Ethnic and migration-related inequalities in health anxiety: a systematic review and meta-analysis. *Front Psychol* 2022;13:960256.
- 38 Klein J, von dem Knesebeck O. Inequalities in health care utilization among migrants and non-migrants in Germany: a systematic review. *Int J Equity Health* 2018;17:160.
- 39 Klein J, von dem Knesebeck O. [Social disparities in outpatient and inpatient care: an overview of current findings in Germany]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2016;59:238–44.
- 40 Deutsches Institut für Medizinische Dokumentation und Information. ICD-10-WHO version 2019 [Internet]. n.d. Available: <https://www.dimdi.de/static/de/klassifikationen/icd/icd-10-who/kode-suche/htmlamlt2019/>
- 41 Löwe B, Andresen V, Van den Bergh O, *et al.* Persistent somatic symptoms across diseases — from risk factors to modification: scientific framework and overarching protocol of the interdisciplinary SOMACROSS research unit (RU 5211). *BMJ Open* 2022;12:e057596.
- 42 Kish L. A procedure for objective respondent selection within the household. *Journal of the American Statistical Association* 1949;44:380–7.
- 43 Kolenikov S. Calibrating survey data using iterative proportional fitting (raking). *The Stata Journal* 2014;14:22–59.
- 44 Makowski AC, Mnich EE, Ludwig J, *et al.* Changes in beliefs and attitudes toward people with depression and schizophrenia-results of a public campaign in Germany. *Psychiatry Res* 2016;237:271–8.
- 45 Knesebeck O von dem, Kofahl C, Makowski AC. Differences in depression stigma towards ethnic and socio-economic groups in Germany-exploring the hypothesis of double stigma. *J Affect Disord* 2017;208:82–6.
- 46 von dem Knesebeck O, Löwe B, Lehmann M, *et al.* Public beliefs about somatic symptom disorders. *Front Psychiatry* 2018;9:616.
- 47 Calman L, Brunton L, Molassiotis A. Developing longitudinal qualitative designs: lessons learned and recommendations for health services research. *BMC Med Res Methodol* 2013;13:14.
- 48 Ganzeboom HBG, De Graaf PM, Treiman DJ. A standard international socio-economic index of occupational status. *Social Science Research* 1992;21:1–56.
- 49 destatis. *Population with migration background-results of the microcensus 2019-series 1 section 2.2.* 2020.
- 50 Koens S, Marx G, Gras C, *et al.* Physicians' information seeking behavior in patients presenting with heart failure symptoms-does gender of physician and patient matter? *Patient Educ Couns* 2020;103:2437–42.
- 51 Broadbent E, Petrie KJ, Main J, *et al.* The brief illness perception questionnaire. *J Psychosom Res* 2006;60:631–7.
- 52 Jones MP, Keefer L, Bratten J, *et al.* Development and initial validation of a measure of perceived stigma in irritable bowel syndrome. *Psychol Health Med* 2009;14:367–74.
- 53 Gierk B, Kohlmann S, Kroenke K, *et al.* The somatic symptom scale-8 (SSS-8): a brief measure of somatic symptom burden. *JAMA Intern Med* 2014;174:399–407.
- 54 Fink P, Ewald H, Jensen J, *et al.* Screening for somatization and hypochondriasis in primary care and neurological in-patients: a seven-item scale for hypochondriasis and somatization. *J Psychosom Res* 1999;46:261–73.
- 55 Lampert T, Kroll L, Müters S, *et al.* [Measurement of socioeconomic status in the German health interview and examination survey for adults (DEGS1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2013;56:631–6.
- 56 Mayring P, Brunner E. Qualitative content analysis. In: Buber R, Holzmüller HH, eds. *Qualitative market research: concepts - methods - analyses.* 2007: 669–80.
- 57 O’Cathain A, Murphy E, Nicholl J. Three techniques for integrating data in mixed methods studies. *BMJ* 2010;341:c4587.