

# Complications after endoscopic submucosal dissection for early colorectal cancer (Review)

JIANHUI SUN<sup>1-3\*</sup>, XINYUAN XIE<sup>1-3\*</sup>, YANGCHENG LIU<sup>1,2\*</sup>, XIANGLIN HAO<sup>1-3</sup>,  
GANG YANG<sup>1,2</sup>, DAN ZHANG<sup>1-3</sup> and QIONG NAN<sup>1,2</sup>

<sup>1</sup>Department of Gastroenterology, The First Affiliated Hospital of Kunming Medical University;

<sup>2</sup>Yunnan Provincial Clinical Research Center for Digestive Diseases, Kunming, Yunnan 650032;

<sup>3</sup>Graduate School of Kunming Medical University, Kunming, Yunnan 650500, P.R. China

Received December 15, 2022; Accepted April 11, 2023

DOI: 10.3892/ol.2023.13850

**Abstract.** Colorectal cancer (CRC) is a gastrointestinal malignancy that seriously threatens human life and health, resulting in a heavy disease burden. Endoscopic submucosal dissection (ESD) is widely used in clinical practice and is an effective treatment for early CRC (ECC). Colorectal ESD is a challenging operation, and the incidence of postoperative complications is relatively high because of the thin intestinal wall and limited space for endoscopic operations. Systematic reports on the postoperative complications of colorectal ESD, such as fever, bleeding and perforation, from both China and elsewhere are lacking. In the present review, progress in research on postoperative complications after ESD for ECC is summarized.

## Contents

1. Introduction
2. Fever
3. Bleeding
4. Perforation
5. Post-ESD electrocoagulation syndrome (PEECS)
6. Other complications
7. Summary

## 1. Introduction

Colorectal cancer (CRC) occurs in the colon or rectum, and has the third highest incidence and second highest mortality

rate among types of cancer worldwide (1-3), thus it is associated with a heavy disease burden. According to the 2019 China Cancer Center report, China ranked second in CRC incidence and fourth in CRC mortality in 2015 (4). Early CRC (ECC) refers to CRC confined to the mucosal and submucosal layers, with or without lymph node metastasis (5,6). The 5-year survival rates of patients with ECC are  $\geq 90\%$  (7), whereas the 5-year survival rate of patients with progressive CRC is markedly lower, at  $<10\%$  (8,9). Previous studies have reported that colonoscopy and lesion resection can reduce CRC incidence by 76-90% and mortality by 53% (10,11). The presence of colonic adenomas, particularly progressive adenomas, is associated with significantly increased CRC incidence and mortality, and resection of adenomas significantly reduces CRC incidence [standardized incidence ratio (SIR), 0.24-0.65] and mortality (SIR, 0.26-0.80) (12). Early detection and treatment of cancer are critical to patient prognosis. Current treatments for ECC are primarily surgery, endoscopic submucosal dissection (ESD) and endoscopic mucosal resection (EMR) (13). The advantages and disadvantages of ESD and EMR in different parts of the GI tract are presented in Table I. Surgery was once considered the standard method for treating early-stage gastrointestinal cancer; however, although surgery can completely remove a lesion, it has a number of disadvantages, including the associated trauma, slow recovery and high complication rates. Compared with traditional surgery, ESD preserves normal bowel function, and results in less trauma and faster postoperative recovery (14). Moreover, patients who undergo ESD for the treatment of ECC and precancerous lesions have a higher overall survival rate, with a long-term prognosis similar to that of conventional surgery (15). Compared with EMR, ESD has a relatively higher risk of complications, due to its greater surgical difficulty and longer operative time (16), but can be used to treat larger lesions, reduce the postoperative residual and recurrence rates, and yields a more accurate pathological histology report; therefore, ESD is gradually becoming the main treatment modality for early gastrointestinal cancer and precancerous lesions (17-20) (Fig. 1). Although ESD can achieve relatively good efficacy in the treatment of ECC, the incidence of post-ESD complications, including fever, bleeding, perforation, electrocoagulation syndrome and stricture, is relatively high. The relatively high complication rate

---

*Correspondence to:* Professor Qiong Nan, Department of Gastroenterology, The First Affiliated Hospital of Kunming Medical University, 295 Xichang Road, Wuhua, Kunming, Yunnan 650032, P.R. China

Email: LineWalker692@gmail.com

\*Contributed equally

**Key words:** early colorectal cancer, endoscopic submucosal dissection, postoperative complications

is due to the limits of endoscopic operation and anatomical factors, such as the rich vascularity of the colorectal mucosa, the thin intestinal wall and the substantial curvature of the intestine. Therefore, understanding and controlling the occurrence of common complications after colorectal ESD surgery, related risk factors, and their prevention and control measures, are important topics for clinicians. The present review provides an overview of the occurrence of common postoperative complications after colorectal ESD, related risk factors, and prevention and control measures, to serve as a reference for clinical treatment.

## 2. Fever

*Risk factors for fever.* Fever is common after colorectal ESD, occurring in 46.7-58.3% of cases (21); however, study of fever after colorectal ESD has been somewhat neglected, with investigations focusing primarily on bleeding, perforation or electrocoagulation syndrome (22,23). Tu (24) defined fever as a temperature  $\geq 38.0^{\circ}\text{C}$ , whereas Izumi *et al* (21) defined it as  $>37.0^{\circ}\text{C}$ , which indicates that postoperative fever lacks a unified standard definition. Postoperative fever is related to the release of inflammatory factors (21). ESD is a low-risk surgical operation for infection and, even if perforation occurs, the rate of bacteremia is low and transient (25); therefore, prophylactic antibiotics can generally be dispensed with. Nevertheless, certain studies have reported that prophylactic antibiotics can reduce clinical adverse events, including abdominal pain, diarrhea and fever (26,27). Therefore, prophylactic use of antibiotics after ESD remains controversial.

Risk factors for fever after ESD include age, lesion size, postoperative bleeding or perforation, and surgery duration. Advanced age is often associated with numerous underlying diseases and immune deficiency, which, together with longer ESD surgery duration, increase the risk of postoperative infection (28). Nakanishi *et al* (29) also reported that age was a risk factor for fever in patients following colorectal ESD, as was lesion size (21). Usually ESD resection depth is limited to the mucosal and submucosal layers, but sometimes it penetrates the muscular layer or all layers (30). Deeply infiltrated ulcers can form soon after ESD and the inflammatory response during their healing is another cause of fever (31). Furthermore, deep/large ulcers after ESD are associated with postoperative bleeding and perforation and, if postoperative bleeding requiring re-electrocoagulation occurs, burns to the intestinal wall can cause an inflammatory response in the plasma membrane (32). Complete intraoperative clamping of the wound in ESD can reduce perforation and various adverse events, particularly fever (33), supported by the fact that mucosal defect closure accelerates wound healing (34). Postoperative perforation combined with fever and elevated infection indicators, such as white blood cells, may indicate secondary infection (35), and surgical intervention may be required in severe cases of secondary infection (36,37).

*Fever control measures.* The consensus among Chinese experts is that postoperative antimicrobials should be used for patients with advanced age, extensive underlying disease, large resection area, long operative time, and postoperative bleeding or perforation (38,39). Routine use of antibiotics after ESD

for possible postoperative free abdominal, mediastinal, retroperitoneal or systemic infection is recommended (40). Since post-ESD bleeding or perforation can cause fever, preoperative anticoagulation and antiplatelet drugs should be avoided. Submucosal injection of epinephrine-saline and hemostatic clips can achieve intraoperative hemostasis; to prevent perforation, adequate submucosal injection should be ensured. Cooling treatment should be given for postoperative absorption fever, while for electrocoagulation syndrome, cooling, analgesia and antibiotics are recommended. For intraoperative perforation, timely clamping and antibiotics should be used. In patients with large lesion diameters and long postoperative fasting time, clinicians must be alert for postoperative fever (41). Shortening operation time and reducing serious postoperative complications, such as bleeding and perforation, are important for fever prevention after ESD.

## 3. Bleeding

*Risk factors for bleeding.* Bleeding is a common complication after ESD, and some bleeding associated with the procedure is almost unavoidable, including intraoperative and late postoperative bleeding (Fig. 2). Intraoperative bleeding is defined as active bleeding during ESD. Postoperative delayed bleeding refers to bleeding occurring  $>6$  h after ESD, mainly manifesting as obvious blood in the stool or a decrease in hemoglobin of  $>20$  g/l relative to the preoperative period (42-44), and is an indication for emergency endoscopic hemostasis (45). Bleeding occurs after 0-11.11% of colorectal ESD procedures (42,46-49). Li *et al* (50) reported that most late bleeding after colorectal ESD occurred within 5 days following surgery. Furthermore, the incidence of immediate and delayed postoperative bleeding has been reported to be 0.39% (95% CI, 0.11-1.3%) and 1.8% (95% CI, 1.4-2.4%) in Asian countries, and 3.3% (95% CI, 1.4-7.6%) and 3.9% (95% CI, 2.5-5.8%) in European and North American countries, respectively. Bleeding rates post-ESD are lower in Asia than in Europe (51), which may be related to the fact that ESD was performed relatively earlier and with more advanced techniques in Asian countries.

Risk factors affecting post-ESD bleeding include duration of surgery, lesion size and location, histological type, intraoperative mucosal lift, and preoperative administration of antiplatelet or anticoagulant medications. Yamamoto *et al* (52) reported that operative time, lesion location and histological type were independent risk factors for postoperative bleeding in ESD. Furthermore, the risk of delayed bleeding is higher in cecum lesions than those at other sites (53). Chiba *et al* (54) reported an increased risk of delayed bleeding after ESD when lesions were in the rectum and  $\geq 40$  mm. Inadequate submucosal injection volume and high fibrosis at the lesion site are risk factors for delayed bleeding after colorectal ESD (50,55), as well as a major cause of poor separation of the mucosal layer from the muscular layer. Therefore, more caution is needed for poorly elevated colorectal lesions. A higher risk of postoperative bleeding has been reported in hypertensive patients treated with ESD (56-58); however, other studies reported no correlation between hypertension and postoperative bleeding after ESD (50,59). A previous study reported that the effect of lesion size on delayed bleeding

Table I. Advantages and disadvantages of ESD and EMR at esophagus, stomach and colorectal sites.

|   |  | Site   |  |  |
|---|--|--|--|--|
|   |  | Esophagus  | Gastric  | Colorectal   |
| <b>A, ESD</b>   |  |  |  |  |
| Advantages or disadvantages   | All  |  |  |  |
| Advantages  | 1. High rate of whole resection and radical resection<br>2. Low local recurrence rate.                         | First-line therapy for early-stage esophageal cancer.  | 1. Preferred for early gastric cancer.<br>2. Enables accurate pathological assessment and curative resection.  | 1. The primary treatment option for early colorectal cancer.<br>2. Enables accurate pathological assessment and curative resection.  |
| Disadvantages   | 1. High operational requirements.<br>2. High surgical difficulty.<br>3. High surgical trauma.<br>4. High cost. | 1. A small operating space.<br>2. High risk of postoperative complications, including bleeding, perforation and stenosis.  | High risk of postoperative bleeding and perforation.   | 1. Anatomical factors, such as the thin intestinal wall, the relatively free part of the intestinal segment and the small operating space of the intestinal cavity determine that the operation is more difficult than that of the upper gastrointestinal tract.<br>2. There is a marked risk of postoperative bleeding, perforation and stricture.<br>3. Only routinely performed by a few large centers. |
| <b>B, EMR</b>   |  |  |  |  |
| Advantages or disadvantages   | All  |  |  |  |
| Advantages  | 1. Simple surgical operation.<br>2. Low surgical trauma.<br>3. Lower cost.                                     | 1. Lesion diameter $\leq 10$ mm, superficial gastric tumor, choice of whole block resection.<br>2. Low incidence of postoperative bleeding, perforation and stenosis.                        | 1. UL0 cT1a, differentiated carcinoma, lesion diameter $\leq 20$ mm, able to be resected in its entirety.<br>2. Low incidence of postoperative bleeding and perforation. | 1. Superficial intestinal tumor, $\leq 20$ mm in length and diameter, can be removed whole.<br>2. Low incidence of postoperative bleeding, perforation and stricture.  |
| Disadvantages   | Local residual, high risk of recurrence  | 1. Early esophageal adenocarcinomas after EMR resection, ablation therapy is recommended after resection.<br>2. Partial, incomplete resection, making assessing the radical effect difficult | Partial incomplete excision of local residuals or EPMR, which interferes with accurate pathological assessment.  | Partial incomplete excision of local residuals or EPMR, which interferes with accurate pathological assessment.  |
| UL0, no ulcer and ulcer scar; cT1a, intramucosal carcinoma; EPMR, endoscopy piecemeal mucosal resection; ESD, endoscopic submucosal resection; EMR, endoscopic mucosal resection. |  |  |  |  |




Figure 1. Patient 1. Images were obtained from the Gastrointestinal Endoscopy Center of The First Affiliated Hospital of Kunming Medical University (Kunming, China). Endoscopic submucosal dissection. (A) Laterally spreading tumors located in the rectum. (B) Dissection procedure: Peeling of submucosa and lesions. (C) Post-dissection trauma. (D) A dissected lesion.




Figure 2. Patient 2. Images were obtained from the Gastrointestinal Endoscopy Center of The First Affiliated Hospital of Kunming Medical University (Kunming, China). Complicated bleeding after endoscopic mucosal dissection. (A) Laterally spreading tumors located in the rectum with (B) bleeding during dissection. (C) Trauma after dissection. (D) Complicated bleeding 1 week after endoscopic submucosal dissection. (E) Postoperative bleeding and (F) following repeated endoscopic hemostasis.

was not statistically significant (60); however, some scholars take the opposite view (59). Previous reports stated that antiplatelet agents should be discontinued before ESD treatment (56,61,62), unless patients are at high risk of combined thromboembolism; however, certain studies disagree (47,59). Therefore, risk factors for postoperative bleeding after ESD to date are controversial, and further research is needed.

*Measures to prevent and treat bleeding.* Most bleeding from small vessels after ESD can be stopped by electrocoagulation (14). Bleeding from larger vessels requires hemostasis by metal clip closure combined with electrocoagulation (14).

During ESD, effective identification of submucosal vessels, timely electrocoagulation and careful treatment of trauma are essential. Furthermore, local cleaning with ice-cold saline containing norepinephrine and hemostatic drugs, such as hemagglutinin, can achieve a hemostatic effect and effectively reduce postoperative bleeding risk (63), while prophylactic clamping of trauma reduces delayed bleeding risk after colorectal ESD (64) by reducing irritation from intestinal contents and accelerating incision healing. Therefore, for larger colorectal lesions, clamping of trauma is beneficial to promote incisional healing and prevent delayed bleeding. Emergency management measures related to postoperative




Figure 3. Flowchart for emergency treatment of bleeding or perforation after ESD. ESD, endoscopic submucosal dissection.

bleeding in ESD are shown in Fig. 3. Huang *et al* (65) reported that patients with colorectal lesions who received traction ring-assisted ESD experienced significantly less intraoperative bleeding than those who did not .

#### 4. Perforation

**Risk factors for perforation.** The risk of perforation is elevated after colorectal ESD because the muscular layer of the intestinal wall is thinner than that in the gastric wall (51,66). The main goal of ESD treatment is complete resection without perforation (67) and reported perforation rates after colorectal ESD are 0-10.7% (44,68). Perforations include intraoperative and delayed perforations. Intraoperative perforation (69) is

defined as an intraoperative, endoscopically confirmed bowel wall defect, most of which can be successfully clamped endoscopically. Delayed perforation (70) is defined as perforation occurring after ESD, with postoperative peritonitis, and can be confirmed by abdominal X-ray or computed tomography (CT) (Fig. 4). Most delayed perforations occur within 14 h of surgery (44) and, although incidence is very low, when it occurs, spillage of bowel contents causes severe peritonitis, which often requires surgery (71,72), indirectly illustrating the importance of preoperative bowel cleansing. Similar to postoperative bleeding, immediate and delayed perforation rates were reported to be 3.8% (95% CI, 3.1-4.6%) and 0.18% (95% CI, 0.08-0.42%) in Asian countries, and 6.6% (95% CI, 4.6-9.4%) and 1.2% (95% CI, 0.29-4.6%) in European




Figure 4. Patient 3. Images were obtained from the Gastrointestinal Endoscopy Center of The First Affiliated Hospital of Kunming Medical University (Kunming, China). Complicated perforation after endoscopic mucosal dissection. (A) A polyp located in the hepatic flexure of the transverse colon. (B) Submucosal injection. (C) Trauma after dissection and (D) closure of the trauma with titanium clips. (E) Abdominal computed tomography scan 12 h after surgery which suggested perforation and (F) a large volume of free gas was seen in the abdominal cavity and under the diaphragm.

and North American countries, respectively. These findings indicated that the incidence of postoperative colorectal ESD perforation is lower in Asia than in Europe (51).

Risk factors influencing perforation after colorectal ESD include duration of surgery, lesion size, lesion location and submucosal fibrosis. Su *et al* (73) reported that the perforation rate in patients with colorectal lesions >4 cm in diameter (7.04%) was higher than that in patients with lesion diameter <4 cm (5.2%), there was a significant difference between the two groups. Hong *et al* (74) reported that lesion size, submucosal fibrosis and duration of surgery were associated with postoperative perforation after colorectal ESD. Another study reported tumor size, site, submucosal fibrosis and operator experience as potential risk factors for postoperative perforation (75). Furthermore, a Japanese study reported that longer operative times were more likely to result in postoperative perforation of the colorectal ESD (52). Prolonged operative time is related to various factors, such as operator experience and the difficulty of the ESD procedure. Another study reported a higher incidence of perforation following left hemicolectomy (73). Further studies are needed in the future regarding the relationship between the lesion site and postoperative perforation of colorectal ESD.

**Preventive measures for perforation.** Perforation post-colorectal ESD is a serious complication that can cause extensive peritonitis, infectious shock and even be life-threatening. Adequate

preoperative preparation, such as water fasting and bowel cleansing, is necessary. The colorectal wall is relatively weak and thermal damage from excessive electrocoagulation may lead to perforation; therefore, avoiding excessive intraoperative electrocoagulation can reduce postoperative perforation incidence (72,76). Adequate submucosal injection reduces perforation due to disruption of the muscularis during dissection caused by electrocoagulation. Furthermore, timely perforation closure is closely related to prognosis and can reduce peritonitis occurrence (77). If intraoperative perforation occurs, it can generally be detected and smaller perforations can be closed with metal clips. For larger perforations, metal clips combined with nylon rope wrap, alongside fasting and gastrointestinal decompression, the use of intravenous antibiotics and other means, may be required; generally, the treatment is effective (78). Surgical laparoscopic exploration and repair are preferable in cases of delayed perforation that cannot be completely closed endoscopically or for cases in which conservative treatment is ineffective (14) (Fig. 3).

### 5. Post-ESD electrocoagulation syndrome (PEECS)

**Risk factors for PEECS.** PEECS refers to a group of inflammatory response syndromes, including limited peritonitis, which can be caused by electrocoagulation damage to the intestinal wall during ESD (72,79). The reported incidence of PEECS is 4.8-14.2% (22,80-83). Typical clinical manifestations of

PEECS include abdominal pain, limited peritoneal irritation, fever and chills, among others. Laboratory tests show elevated leukocytes and C-reactive protein, and abdominal X-ray or computed tomography scans reveal no signs of perforation. PEECS does not usually require surgical treatment, but can result in delayed perforation in severe cases (84). Based on analysis of relevant domestic and international literature, the mechanism underlying PEECS remains unclear.

Research on risk factors associated with PEECS has gradually increased in recent years. Age, lesion diameter, combined malignancy and lesion location have been reported as independent risk factors for PEECS development (80). Female sex, lesion site, lesion size and submucosal fibrosis have also been reported to be associated with PEECS occurrence (22,85,86). Sun *et al* (81) reported a higher incidence of PEECS after colonic ESD than rectal ESD, and that larger lesions were associated with a higher incidence of PEECS. Notably, Yamasaki *et al* (87) reported that line-assisted complete clip closure was effective in reducing PEECS incidence. This subject warrants further study in the future.

*Prevention and control measures for PEECS.* Compared to bleeding and perforation, PEECS are less severe and have a better prognosis, but there remains a risk of delayed perforation. Once diagnosed, immediate treatment, such as fasting, keeping the bowels open and intravenous administration of broad-spectrum antibiotics, can alleviate symptoms; however, in rare cases where conservative treatment is ineffective or the condition worsens, the possibility of delayed perforation should be considered. If intestinal perforation is confirmed, immediate surgical repair is required. Li *et al* (84) reported that placement of anal canal drainage and decompression after ESD reduced the incidence of postoperative PEECS, possibly because of a reduced risk of infection, and delayed perforation, due to reduced exposure of the surgical wound to intestinal bacteria and fecal matter. A study in the United States reported that prophylactic antibiotics reduced the risk of PEECS and effectively relieved abdominal pain (88). Furthermore, prolonging mucosal augmentation time at the lesion may reduce the risk of intestinal wall damage caused by electrocoagulation, thus reducing the risk of PEECS (89).

## 6. Other complications

*Stenosis.* Stenosis is a less frequent complication after colorectal ESD, generally defined as the inability of a normal endoscope to pass successfully through the postoperative bowel lumen. A study by Hayashi *et al* (90) reported that the incidence of colorectal stenosis was 0.49% (4/822 cases) and that of postoperative stenosis was 11% (2/18) for 90-100% circumferential lesions. Another study (91) reported that the stenosis rate after rectal ESD was 19.7% (12/61), with that of stenosis after total circumferential resection 71.4% (5/7), and that of stenosis after circumferential >90% lesion resection 43.8% (7/16). The results of these studies suggested that circumferential resection of large lesions is a risk factor for postoperative stenosis. General endoscopic balloon dilatation treatment may improve postoperative stenosis to a certain extent, while local hydrocortisone enemas may also help to prevent postoperative stenosis (92).

*Fulminant necrotizing fasciitis (Fournier's syndrome).* Due to its anatomical location, perforation of the lower rectum after surgery can easily cause mediastinal or subcutaneous emphysema, and it is challenging to confine the infected lesion to a single location. Therefore, such lesions can easily spread along the loose tissue to the buttocks and abdomen, causing multiple muscle necrosis and inflammation of the surrounding fascia. Such fulminant necrotizing fasciitis (Fournier's syndrome) is rare clinically, and has few reports to date. Once present, the condition can easily cause sepsis and diffuse intravascular coagulation with serious consequences. Fournier's syndrome is exceptionally aggressive, with reports of mortality rates as high as 20-40%. Therefore, prompt treatment with broad-spectrum antibiotics and surgery should be used following detection (44).

## 7. Summary

In summary, ESD has both advantages and disadvantages for the treatment of ECC, and reducing the occurrence of postoperative complications is an important safeguard for the successful implementation of colorectal ESD surgery. Therefore, endoscopists should strictly grasp the indications for ESD surgery, understand the relevant risk factors affecting postoperative complications, and take immediate and effective measures to intervene once they occur, which will effectively improve the safety of colorectal ESD treatment. Furthermore, the clinical application of colorectal ESD could be further expanded, so that more patients with early-stage colorectal cancer can benefit from ESD technology in the future.

## Acknowledgements

Not applicable.

## Funding

Funding for the present study was received from The Kunming Medical Joint Special Project-Surface Project (grant no. 202101AC070802) and The Yunnan Provincial Clinical Medical Research Center for Digestive Diseases-Research and Application of Difficult and Critical Digestive Diseases in Yunnan Province (grant no. 202002AA100205).

## Availability of data and materials

Not applicable.

## Authors' contributions

JS was responsible for research design, conceptualization and writing the first draft. XX, DZ and XH were responsible for conceptualization, organization, investigation and research design. YL and GY were responsible for supervision, conceptualization and revision of the manuscript. QN was responsible for research supervision and revision of the manuscript. Data authentication is not applicable. All authors contributed to the article, and read and approved the final manuscript.

### Ethics approval and consent to participate

The publication of the images in the figures was approved by the Ethics Committee of The First Affiliated Hospital of Kunming Medical University (Kunming, China).

### Patient consent for publication

Written informed consent for publication of the images in the figures was obtained from the patients.

### Competing interests

The authors declare that they have no competing interests.

### References

- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A and Bray F: Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 71: 209-249, 2021.
- Chinese Society Of Clinical Oncology CSCO Diagnosis And Treatment Guidelines For Colorectal Cancer Working Group: Chinese society of clinical oncology (CSCO) diagnosis and treatment guidelines for colorectal cancer 2018 (English version). *Chin J Cancer Res* 31: 117-134, 2019.
- Brenner H, Kloor M and Pox CP: Colorectal cancer. *Lancet* 383: 1490-1502, 2014.
- Zheng RS, Sun KX, Zhang S, Zeng HM, Zou XN, Chen R, Gu XY, Wei WW and He J: Analysis of malignant tumor prevalence in China in 2015. *Chin J Oncol* 41: 19-28, 2019 (In Chinese).
- McGill SK, Soetikno R and Kaltenbach T: Optical diagnosis of early colorectal cancer: Riding the highs and lows of the Japanese narrow-band imaging expert team classification. *Gastrointest Endosc* 86: 710-712, 2017.
- Shin JW, Han KS, Hyun JH, Lee SJ, Kim B, Hong CW, Kim BC, Sohn DK, Chang HJ, Kim MJ, *et al*: Risk of recurrence after endoscopic resection of early colorectal cancer with positive margins. *Endoscopy* 50: 241-247, 2018.
- Courtney RJ, Paul CL, Carey ML, Sanson-Fisher RW, Macrae FA, D'Este C, Hill D, Barker D and Simmons J: A population-based cross-sectional study of colorectal cancer screening practices of first-degree relatives of colorectal cancer patients. *BMC Cancer* 13: 13, 2013.
- Yao L, Shen XF and Chen LG: Analysis of risk factors for late bleeding after transendoscopic submucosal dissection for colorectal cancer. *Zhejiang Trauma Surgery* 27: 476-477, 2022.
- Jiang YQ: Clinical study of endoscopic submucosal dissection for early colorectal cancer and precancerous lesions. South China University, 2021.
- Miller KD, Sauer AG, Ortiz AP, Fedewa SA, Pinheiro PS, Tortolero-Luna G, Martinez-Tyson D, Jemal A and Siegel RL: Cancer statistics for Hispanics/Latinos, 2018. *CA Cancer J Clin* 68: 425-445, 2018.
- Wang C: Analysis of risk factors for late bleeding after endoscopic mucosal resection of colorectal polyps. China Medical University, 2021.
- Nishihara R, Wu K, Lochhead P, Morikawa T, Liao X, Qian ZR, Inamura K, Kim SA, Kuchiba A, Yamauchi M, *et al*: Long-term colorectal-cancer incidence and mortality after lower endoscopy. *N Engl J Med* 369: 1095-1105, 2013.
- Bai Y, Yang F, Ma D and Zou W: Guidelines for early colorectal cancer screening and endoscopic management in China (Beijing, 2014). *Chin J Gastrointestinal Endoscopy* 32: 341-360, 2015.
- Zhou PH, Cai MY and Yao LQ: Expert consensus on the treatment of gastrointestinal mucosal lesions by endoscopic submucosal dissection. *Chin J Gastrointestinal Surg*: 1083-1086, 2012.
- Wang H, Lu PX, Zhang DX, Han HC, Zhong YS and Yao LQ: Efficacy of endoscopic treatment or surgery in patients with early colorectal cancer and precancerous lesions. *J Pract Med* 35: 1639-1643, 2019.
- Yao DH: Advances in the clinical application of endoscopic submucosal dissection. *Southwest Defense Med* 20: 215-217, 2010.
- Li P, Wang CJ, Chen GY and Xu CQ: Chinese consensus on screening and diagnosis and treatment of early stage esophageal squamous cell carcinoma and precancerous lesions (Beijing, 2015). *Chin J Gastrointest Endoscopy* 33: 3-18, 2016.
- Expert group of the major project of Beijing Municipal Commission of Science and Technology 'Research on the standardization of early gastric cancer treatment'. Expert consensus opinion on standardized endoscopic resection of early gastric cancer (2018, Beijing). *Chin J Gastrointestinal Endoscopy* 36: 381-392, 2019.
- Yamamoto Y, Kikuchi D, Nagami Y, Nonaka K, Tsuji Y, Fujimoto A, Sanomura Y, Tanaka K, Abe S, Zhang S, *et al*: Management of adverse events related to endoscopic resection of upper gastrointestinal neoplasms: Review of the literature and recommendations from experts. *Dig Endosc* 31 (Suppl 1): S4-S20, 2019.
- Komeda Y, Watanabe T and Kudo M: Requirement of additional surgery after non-curative endoscopic submucosal dissection for early colorectal cancer. *J Invest Surg* 34: 895-896, 2021.
- Izumi K, Osada T, Sakamoto N, Kodani T, Higashihara Y, Ritsuno H, Shibuya T, Nagahara A, Ogihara T, Kikuchi K and Watanabe S: Frequent occurrence of fever in patients who have undergone endoscopic submucosal dissection for colorectal tumor, but bacteremia is not a significant cause. *Surg Endosc* 28: 2899-2904, 2014.
- Ito S, Hotta K, Imai K, Yamaguchi Y, Kishida Y, Takizawa K, Kakushima N, Tanaka M, Kawata N, Yoshida M, *et al*: Risk factors of post-endoscopic submucosal dissection electrocoagulation syndrome for colorectal neoplasm. *J Gastroenterol Hepatol* 33: 2001-2006, 2018.
- Shichijo S, Takeuchi Y, Shimodate Y, Yamashina T, Yamasaki T, Hayashi T, Hirasawa K, Fukunaga S, Yamaguchi S, Asai S, *et al*: Performance of perioperative antibiotics against post-endoscopic submucosal dissection coagulation syndrome: A multicenter randomized controlled trial. *Gastrointest Endosc* 95: 349-359, 2022.
- Tu JF: Study of risk factors and causes of fever after endoscopic submucosal dissection of gastrointestinal tumors. Zhejiang University, 2016.
- Deng CQ and Feng L: Clinical study of bacteremia and endotoxemia after endoscopic mucosal dissection. *Southwest Military Med* 18: 514-516, 2016.
- Zhang QS, Han B, Xu JH, Gao P and Shen YC: Antimicrobial prophylaxis in patients with colorectal lesions undergoing endoscopic resection. *World J Gastroenterol* 21: 4715-4721, 2015.
- La Regina D, Mongelli F, Fasoli A, Lollo G, Ceppi M, Saporito A, Garofalo F, Giuseppe MD and di Tor Vajana AF: Clinical adverse events after endoscopic resection for colorectal lesions: A meta-analysis on the antibiotic prophylaxis. *Dig Dis* 38: 15-22, 2020.
- Liu L, Shen XJ, Yue M, Zhou SQ, Wang SW and Zhu JY: Safety of gastric endoscopic mucosal dissection and risk factors associated with complications in elderly patients. *Chin J Gerontol* 38: 3910-3912, 2018.
- Nakanishi T, Araki H, Ozawa N, Takada J, Kubota M, Imai K, Onogi F, Ibuka T, Shiraki M, Shimizu M and Moriwaki H: Risk factors for pyrexia after endoscopic submucosal dissection of gastric lesions. *Endosc Int Open* 2: E141-E147, 2014.
- Zhou PH, Chen WF and He MJ: Standardized endoscopic diagnosis and treatment of early gastric cancer. *Chin J Pract Surg* 34: 604-607, 2014.
- Zhang LL and Yang Z: Advances in the pharmacological treatment of artificial ulcers after endoscopic mucosal dissection. *J Clin Military Med* 45: 1093-1095, 2017.
- Cha JM, Lim KS, Lee SH, Joo YE, Hong SP, Kim TI, Kim HG, Park DI, Kim SE, Yang DH and Shin JE: Clinical outcomes and risk factors of post-polypectomy coagulation syndrome: A multi-center, retrospective, case-control study. *Endoscopy* 45: 202-207, 2013.
- Harada H, Suehiro S, Murakami D, Nakahara R, Ujihara T, Shimizu T, Miyama Y, Katsuyama Y, Hayasaka K and Tounou S: Clinical impact of prophylactic clip closure of mucosal defects after colorectal endoscopic submucosal dissection. *Endosc Int Open* 5: E1165-E1171, 2017.
- Osada T, Sakamoto N, Ritsuno H, Murakami T, Ueyama H, Matsumoto K, Shibuya T, Ogihara T and Watanabe S: Closure with clips to accelerate healing of mucosal defects caused by colorectal endoscopic submucosal dissection. *Surg Endosc* 30: 4438-4444, 2016.



35. Huang R, Zhang LH, Zhang RC, Luo H, Wang XP, Tao Q, *et al*: Analysis of risk factors associated with fever after endoscopic submucosal dissection. *Chin J Gastrointest Endosc* 31: 72-75, 2014.
36. Suzuki H, Oda I, Sekiguchi M, Abe S, Nonaka S, Yoshinaga S, Nakajima T and Saito Y: Management and associated factors of delayed perforation after gastric endoscopic submucosal dissection. *World J Gastroenterol* 21: 12635-12643, 2015.
37. Matsuda Y, Kataoka N, Yamaguchi T, Tomita M, Sakamoto K and Makimoto S: Delayed esophageal perforation occurring with endoscopic submucosal dissection: A report of two cases. *World J Gastrointest Surg* 7: 123-127, 2015.
38. Li P, Wang YJ, Chen GY and Xu CQ: Chinese consensus on screening and diagnosis and treatment of early stage esophageal squamous cell carcinoma and precancerous lesions (2015-Beijing). *Chin J Pract Intern Med* 36: 20-33, 2016.
39. Yang F, Ma D and Zou DW: Expert recommendations on peri-operative medication for endoscopic submucosal dissection of gastric mucosal lesions (2015, Suzhou). *Chin J Intern Med* 54: 905-908, 2015.
40. Zhou PH, Zhong YS, Li QL and Qi ZP: Expert consensus on endoscopic diagnosis and treatment of gastrointestinal submucosal tumors in China (2018 edition). *Chin J Pract Surg* 38: 840-850, 2018.
41. Ren PY, Yin YQ and Gao YH: Progress in the study of complications of colorectal endoscopic submucosal dissection. *J Gastroenterol Hepatol* 30: 341-345, 2021.
42. Xu K, Jin HL, Ding X and Tong C: The value and safety of endoscopic submucosal dissection for early colorectal cancer. Application value and safety assessment of endoscopic submucosal dissection for early colorectal cancer. *Chin J Endosc* 24: 17-22, 2018.
43. Tanaka S, Kashida H, Saito Y, Yahagi N, Yamano H, Saito S, Hisabe T, Yao T, Watanabe M, Yoshida M, *et al*: Japan gastroenterological endoscopy society guidelines for colorectal endoscopic submucosal dissection/endoscopic mucosal resection. *Dig Endosc* 32: 219-239, 2020.
44. Tanaka S, Kashida H, Saito Y, Yahagi N, Yamano H, Saito S, Hisabe T, Yao T, Watanabe M, Yoshida M, *et al*: JGES guidelines for colorectal endoscopic submucosal dissection/endoscopic mucosal resection. *Dig Endosc* 27: 417-434, 2015.
45. Kim DH and Lim SW: Analysis of delayed postpolypectomy bleeding in a colorectal clinic. *J Korean Soc Coloproctol* 27: 13-16, 2011.
46. Chen J and Yu HG: Clinical analysis of endoscopic submucosal dissection for early colorectal cancer and its precancerous lesions. *Chin J Gastrointest Endoscopy* 33: 151-154, 2016.
47. Arimoto J, Higurashi T, Chiba H, Misawa N, Yoshihara T, Kato T, Kanoshima K, Fuyuki A, Ohkubo H, Goto S, *et al*: Continued use of a single antiplatelet agent does not increase the risk of delayed bleeding after colorectal endoscopic submucosal dissection. *Dig Dis Sci* 63: 218-227, 2018.
48. Milano RV, Viale E, Bartel MJ, Notaristefano C and Testoni PA: Resection outcomes and recurrence rates of endoscopic submucosal dissection (ESD) and hybrid ESD for colorectal tumors in a single Italian center. *Surg Endosc* 32: 2328-2339, 2018.
49. Tanaka S, Terasaki M, Hayashi N, Oka S and Chayama K: Warning for unprincipled colorectal endoscopic submucosal dissection: Accurate diagnosis and reasonable treatment strategy. *Dig Endosc* 25: 107-116, 2013.
50. Li X, Yu HG, Li SQ and Zhu XY: Analysis of risk factors for bleeding after colorectal endoscopic submucosal dissection. *Med Rev* 25: 1026-1029+1035, 2019.
51. Akindoye E, Kumar N, Aihara H, Nas H and Thompson CC: Colorectal endoscopic submucosal dissection: A systematic review and meta-analysis. *Endosc Int Open* 4: E1030-E1044, 2016.
52. Yamamoto K, Shimoda R, Ogata S, Hara M, Ito Y, Tominaga N, Nakayama A, Sakata Y, Tsuruoka N, Iwakiri R and Fujimoto K: Perforation and postoperative bleeding associated with endoscopic submucosal dissection in colorectal tumors: An analysis of 398 lesions treated in Saga, Japan. *Intern Med* 57: 2115-2122, 2018.
53. Suzuki S, Chino A, Kishihara T, Urugami N, Tamegai Y, Suganuma T, Fujisaki J, Matsuura M, Itoi T, Gotoda T, *et al*: Risk factors for bleeding after endoscopic submucosal dissection of colorectal neoplasms. *World J Gastroenterol* 20: 1839-1845, 2014.
54. Chiba H, Ohata K, Tachikawa J, Arimoto J, Ashikari K, Kuwabara H, Nakaoka M, Goto T and Nakajima A: Delayed bleeding after colorectal endoscopic submucosal dissection: When is emergency colonoscopy needed? *Dig Dis Sci* 64: 880-887, 2019.
55. Youk EG, Sohn DK, Hong CW, Lee SD, Han KS, Kim BC, Chang HJ and Kim MJ: Early outcomes of endoscopic submucosal dissection for colorectal neoplasms according to clinical indications. *Dis Colon Rectum* 59: 403-410, 2016.
56. Harada H, Nakahara R, Murakami D, Suehiro S, Nagasaka T, Ujihara T, Sagami R, Katsuyama Y, Hayasaka K, Tounou S and Amano Y: The effect of anticoagulants on delayed bleeding after colorectal endoscopic submucosal dissection. *Surg Endosc* 34: 3330-3337, 2020.
57. Li R, Cai S, Sun D, Shi Q, Ren Z, Qi Z, Li B, Yao L, Xu M, Zhou P and Zhong Y: Risk factors for delayed bleeding after endoscopic submucosal dissection of colorectal tumors. *Surg Endosc* 35: 6583-6590, 2021.
58. Jaruvongvanich V, Prasitlumkum N, Assavapongpaiboon B, Suchartlikitwong S, Sanguankeo A and Upala S: Risk factors for delayed colonic post-polypectomy bleeding: A systematic review and meta-analysis. *Int J Colorectal Dis* 32: 1399-1406, 2017.
59. Ogasawara N, Yoshimine T, Noda H, Kondo Y, Izawa S, Shinmura T, Ebi M, Funaki Y, Sasaki M and Kasugai K: Clinical risk factors for delayed bleeding after endoscopic submucosal dissection for colorectal tumors in Japanese patients. *Eur J Gastroenterol Hepatol* 28: 1407-1414, 2016.
60. Terasaki M, Tanaka S, Shigita K, Asayama N, Nishiyama S, Hayashi N, Nakadoi K, Oka S and Chayama K: Risk factors for delayed bleeding after endoscopic submucosal dissection for colorectal neoplasms. *Int J Colorectal Dis* 29: 877-882, 2014.
61. Seo M, Song EM, Cho JW, Lee YJ, Lee BI, Kim JS, Jeon SW, Jang HJ, Yang DH, Ye BD and Byeon JS: A risk-scoring model for the prediction of delayed bleeding after colorectal endoscopic submucosal dissection. *Gastrointest Endosc* 89: 990-998.e2, 2019.
62. Xu YQ, Lin ZR, Zhong SS, Lin XL and Wei L: Meta-analysis of risk factors for late bleeding after endoscopic resection of colorectal masses. *Chin Elect J Colorectal Dis* 9: 377-386, 2020.
63. Wang T, Wang DN, Liu WT, Zheng ZQ, Chen X, Fang WL, Li S, Liang L and Wang BM: Hemostatic effect of topical hemocoagulase spray in digestive endoscopy. *World J Gastroenterol* 22: 5831-5836, 2016.
64. Ogiyama H, Tsutsui S, Murayama Y, Maeda S, Satake S, Nasu A, Umeda D, Miura Y, Tominaga K, Horiki M, *et al*: Prophylactic clip closure may reduce the risk of delayed bleeding after colorectal endoscopic submucosal dissection. *Endosc Int Open* 6: E582-E588, 2018.
65. Huang XY, Ji M, Li P, Niu YL, Meng FD, Yu L, Wang YJ, Zhao HY, Lv FJ, Zhai HH and Zhang ST: New traction method (traction ring) for colorectal endoscopic submucosal dissection. *J Dig Dis* 23: 318-323, 2022.
66. ASGE Technology Committee; Maple JT, Dayyeh BK, Chauhan SS, Hwang JH, Komanduri S, Manfredi M, Konda V, Murad FM, Siddiqui UD and Banerjee S: Endoscopic submucosal dissection. *Gastrointest Endosc* 81: 1311-1325, 2015.
67. Imai K, Hotta K, Ito S, Yamaguchi Y, Kishida Y, Yabuuchi Y, Yoshida M, Kawata N, Tanaka M, Kakushima N, *et al*: A risk-prediction model for en bloc resection failure or perforation during endoscopic submucosal dissection of colorectal neoplasms. *Dig Endosc* 32: 932-939, 2020.
68. Du CQ, Yuan WZ, Zhao MW, Jia W, Mi J, Yuan YD, *et al*: Efficacy and safety analysis of endoscopic mucosal dissection for colorectal intramucosal cancer and precancerous lesions. *Chin J Endoscopy* 25: 73-77, 2019.
69. Nakajima T, Saito Y, Tanaka S, Iishi H, Kudo SE, Ikematsu H, Igarashi M, Saitoh Y, Inoue Y, Kobayashi K, *et al*: Current status of endoscopic resection strategy for large, early colorectal neoplasia in Japan. *Surg Endosc* 27: 3262-3270, 2013.
70. Iwatsubo T, Takeuchi Y, Yamasaki Y, Nakagawa K, Arai M, Ohmori M, Iwagami H, Matsuno K, Inoue S, Nakahira H, *et al*: Differences in clinical course of intraprocedural and delayed perforation caused by endoscopic submucosal dissection for colorectal neoplasms: A retrospective study. *Dig Dis* 37: 53-62, 2019.
71. Toyonaga T, Man-I M, East JE, Nishino E, Ono W, Hirooka T, Ueda C, Iwata Y, Sugiyama T, Dozaiku T, *et al*: 1,635 Endoscopic submucosal dissection cases in the esophagus, stomach, and colorectum: Complication rates and long-term outcomes. *Surg Endosc* 27: 1000-1008, 2013.
72. Hirasawa K, Sato C, Makazu M, Kaneko H, Kobayashi R, Kokawa A and Maeda S: Coagulation syndrome: Delayed perforation after colorectal endoscopic treatments. *World J Gastrointest Endosc* 7: 1055-1061, 2015.

73. Su H, Wang HH, Liu L, Cheng T, He YQ and Jin P: Comparative analysis of endoscopic submucosal dissection of early colorectal cancer and precancerous lesions of different diameters. *Chin J Gastrointestinal Endoscopy*: 339-343, 2019.
74. Hong SN, Byeon JS, Lee BI, Yang DH, Kim J, Cho KB, Cho JW, Jang HJ, Jeon SW, Jung SA and Chang DK: Prediction model and risk score for perforation in patients undergoing colorectal endoscopic submucosal dissection. *Gastrointest Endosc* 84: 98-108, 2016.
75. Mizushima T, Kato M, Iwanaga I, Sato F, Kubo K, Ehira N, Uebayashi M, Ono S, Nakagawa M, Mabe K, *et al*: Technical difficulty according to location, and risk factors for perforation, in endoscopic submucosal dissection of colorectal tumors. *Surg Endosc* 29: 133-139, 2015.
76. Kawashima K, Hikichi T, Fujiwara T, Gunji N, Nakamura J, Watanabe K, Katakura K and Ohira H: Delayed perforation after endoscopic submucosal dissection for mucosal colon cancer: A conservatively treated case. *Fukushima J Med Sci* 64: 157-162, 2018.
77. Kang DU, Choi Y, Lee HS, Lee HJ, Park SH, Yang DH, Yoon SM, Kim KJ, Ye BD, Myung SJ, *et al*: Endoscopic and clinical factors affecting the prognosis of colorectal endoscopic submucosal dissection-related perforation. *Gut Liver* 10: 420-428, 2016.
78. Wang QE, Zhuang C, Zhang Y, Chen JP and Xu F: Clinical application of single-clamp endoscopic purse-string suturing in intestinal perforation. *Chin J Gastrointestinal Endoscopy*: 601-603, 2019.
79. Kim ER and Chang DK: Management of complications of colorectal submucosal dissection. *Clin Endosc* 52: 114-119, 2019.
80. Wang MZ, Tan SY, Luo HS, Li M, Wu PB, Guo F and Shu Y: Risk factors for electrocoagulation syndrome after endoscopic submucosal dissection of colorectal lesions. *Chinese J Gen Practitioners* 15: 698-701, 2016.
81. Sun D, Qi ZP, Zhong YS, Shi Q, Cai SL and Li B: Clinical features and risk factors of electrocoagulation syndrome after colorectal endoscopic submucosal dissection. *Chin J Pract Surg* 38: 662-665, 2018.
82. Arimoto J, Higurashi T, Kato S, Fuyuki A, Ohkubo H, Nonaka T, Yamaguchi Y, Ashikari K, Chiba H, Goto S, *et al*: Risk factors for post-colorectal endoscopic submucosal dissection (ESD) coagulation syndrome: A multicenter, prospective, observational study. *Endosc Int Open* 6: E342-E349, 2018.
83. Hong MJ, Kim JH, Lee SY, Sung IK, Park HS and Shim CS: Prevalence and clinical features of coagulation syndrome after endoscopic submucosal dissection for colorectal neoplasms. *Dig Dis Sci* 60: 211-216, 2015.
84. Li B, Zhou PH, Yao LQ, Xu MD, Ren Z and Shi QQ: Analysis of the efficacy of endoscopic submucosal dissection for post-operative anal canal drainage and decompression of colorectal mucosal lesions. *Chin J Pract Surg* 37: 802-805, 2017.
85. Yamashina T, Takeuchi Y, Uedo N, Hamada K, Aoi K, Yamasaki Y, Matsuura N, Kanesaka T, Akasaka T, Yamamoto S, *et al*: Features of electrocoagulation syndrome after endoscopic submucosal dissection for colorectal neoplasm. *J Gastroenterol Hepatol* 31: 615-620, 2016.
86. Jung D, Youn YH, Jahng J, Kim JH and Park H: Risk of electrocoagulation syndrome after endoscopic submucosal dissection in the colon and rectum. *Endoscopy* 45: 714-717, 2013.
87. Yamasaki Y, Takeuchi Y, Kato M, Uedo N and Ishihara R: Line-assisted complete closure of large gastric mucosal defects by use of multiple clip-and-line technique. *VideoGIE* 1: 49-50, 2016.
88. Lee SP, Sung IK, Kim JH, Lee SY, Park HS, Shim CS and Ki HK: A randomized controlled trial of prophylactic antibiotics in the prevention of electrocoagulation syndrome after colorectal endoscopic submucosal dissection. *Gastrointest Endosc* 86: 349-357.e2, 2017.
89. Yamasaki Y, Takeuchi Y, Iwatsubo T, Kato M, Hamada K, Tonai Y, Matsuura N, Kanesaka T, Yamashina T, Arao M, *et al*: Line-assisted complete closure for a large mucosal defect after colorectal endoscopic submucosal dissection decreased post-electrocoagulation syndrome. *Dig Endosc* 30: 633-641, 2018.
90. Hayashi T, Kudo SE, Miyachi H, Sakurai T, Ishigaki T, Yagawa Y, Toyoshima N, Mori Y, Misawa M, Kudo T, *et al*: Management and risk factor of stenosis after endoscopic submucosal dissection for colorectal neoplasms. *Gastrointest Endosc* 86: 358-369, 2017.
91. Ohara Y, Toyonaga T, Tanaka S, Ishida T, Hoshi N, Yoshizaki T, Kawara F, Lui KL, Tepmalai K, Damrongmanee A, *et al*: Risk of stricture after endoscopic submucosal dissection for large rectal neoplasms. *Endoscopy* 48: 62-70, 2016.
92. Ma MX and Bourke MJ: Complications of endoscopic polypectomy, endoscopic mucosal resection and endoscopic submucosal dissection in the colon. *Best Pract Res Clin Gastroenterol* 30: 749-767, 2016.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.