




# Validation of the Arabic version of the Eating Attitude Test in Lebanon: a population study

Chadia Haddad<sup>1,2</sup>, Chloe Khoury<sup>3</sup>, Pascale Salameh<sup>4,5</sup>, Hala Sacre<sup>5</sup>, Rabih Hallit<sup>6</sup>, Nelly Kheir<sup>3</sup>, Sahar Obeid<sup>1,5,7,\*</sup> and Souheil Hallit<sup>5,6,\*</sup> 

<sup>1</sup>Research and Psychology Departments, Psychiatric Hospital of the Cross, P.O. Box 60096, Jal Eddib, Lebanon: <sup>2</sup>INSERM, Univ. Limoges, CHU Esquirol, IRD, U1094 Tropical Neuroepidemiology, Institute of Epidemiology and Tropical Neurology, GEIST, Limoges, France: <sup>3</sup>Faculty of Science, Lebanese University, Fanar, Lebanon: <sup>4</sup>Faculties of Pharmacy and Medicine, Lebanese University, Beirut, Lebanon: <sup>5</sup>INSPECT-LB: Institut National de Santé Publique, Epidemiologie Clinique et Toxicologie-Liban, Beirut, Lebanon: <sup>6</sup>Faculty of Medicine and Medical Sciences, Holy Spirit University of Kaslik (USEK), Jounieh, Lebanon: <sup>7</sup>Faculty of Arts and Sciences, Holy Spirit University of Kaslik (USEK), Jounieh, Lebanon

Submitted 26 February 2020: Final revision received 16 July 2020: Accepted 27 July 2020: First published online 8 September 2020

## Abstract

**Objectives:** To validate an Arabic version of the Eating Attitude Test (EAT-26) and identify factors (such as depression, stress, anxiety and body dissatisfaction) that might be associated with disordered eating among a sample of the Lebanese population.

**Design:** Cross-sectional study.

**Setting:** All Lebanese governorates.

**Participants:** A total of 811 participants randomly selected participated in this 5-month study (January–May 2018).

**Results:** The EAT-26 scale items converged over a solution of six factors that had an eigenvalue over 1, explaining a total of 60.07% of the variance (Cronbach's  $\alpha = 0.895$ ). The prevalence of disordered eating attitudes was 23.8%. Higher EAT-26 scores (disordered eating attitudes) were significantly associated with higher depression ( $\beta = 0.325$ ), higher emotional eating ( $\beta = 0.083$ ), daily weighing ( $\beta = 3.430$ ), higher physical activity ( $\beta = 0.05$ ), starving to reduce weight ( $\beta = 4.94$ ) and feeling pressure from TV/magazine to lose weight ( $\beta = 3.95$ ).

**Conclusions:** The Arabic version of EAT-26 can be a useful instrument for screening and assessing disordered eating attitudes in clinical practice and research. Some factors seem to be associated with more disordered eating attitudes among participants for whom psychological counseling may be needed. Yet, our findings are considered preliminary, and further studies are warranted to confirm them.

**Keywords**  
Eating  
Attitudes  
Depression  
Emotional eating  
Arabic  
Lebanon

'Eating Attitude' is a term used globally to describe the beliefs, thoughts, feelings and behaviours related to food<sup>(1)</sup>. People develop positive or negative attitudes leading to actions consistent with cognitions and emotions towards food. Diverse approaches to food might have consequences on general well-being, and psychological components of eating attitudes might be important determinants of health.

Rapid changes in food choices and behaviours have occurred lately, with people desiring to stay slim<sup>(2–5)</sup>. Unhealthy eating attitudes, related to abnormal preoccupations with food and weight, such as continuous concerns about controlling and losing weight, food restriction and amplified anger and bad temper when hungry, had quickly

emerged and resulted in severe eating disorders<sup>(6,7)</sup>. Eating disorders, defined as abnormal or disturbed eating behaviours, are highly encountered and include harmful eating attitudes with severe distress or concerns about body weight or shape<sup>(8)</sup>. According to many studies, the prevalence of unfavourable eating attitudes varies between 10 and 16% in America<sup>(9,10)</sup>, 3–13.7% in Europe<sup>(11)</sup> and 4–30% in Asia<sup>(2,5,12,13)</sup>. Arab countries, especially Lebanon, are affected by westernisation and tend to adopt western traditions in everything, including eating attitudes. Thus, Lebanese people (mostly women) favour fasting or restrained eating to lose weight<sup>(6)</sup>.

Psychological factors and emotional status play very important roles in food choices and behaviours<sup>(14)</sup>. Negative emotions could have an impact on the behaviour

Chadia Haddad and Chloe Khoury are the first co-authors.  
Sahar Obeid and Souheil Hallit are the last co-authors.

\*Corresponding authors: Email souheilhallit@hotmail.com; saharobeid23@hotmail.com

© The Author(s), 2020. Published by Cambridge University Press on behalf of The Nutrition Society



of individuals by reducing their actions, which usually involves avoidance of the stimulus or the activity<sup>(14)</sup>. Emotional regulation can affect the eating attitude as it plays an influential role in daily food intake and may cause what is known as 'emotional overeating'<sup>(15)</sup>. Furthermore, stress (often coupled with depression and anxiety when chronic) and anxiety are also associated with disordered eating<sup>(16,17)</sup>. Indeed, studies have shown that anxiety appears much earlier than eating disorders, making it a cause of binge eating<sup>(18)</sup>.

Moreover, anxiety showed some association with fasting, especially 'social appearance anxiety', where people get anxious about what society may think of their appearance. Studies have shown that a higher frequency of use of social media resulted in high rates of eating disorders: pressure may lead to body dissatisfaction, lowered self-esteem and urge to become extremely skinny, especially in women<sup>(6,7,19)</sup>. Consequently, the frequency of restrained eating is increasing worldwide; people stop eating voluntarily to lose weight and become skinny<sup>(17)</sup>, without knowing that this will increase the food intake after some time, leading to binge eating again.

Furthermore, studies have demonstrated that intense physical activity is linked to eating disorders<sup>(20,21)</sup>. An excessive workout is often used as an approach to cope with emotional conditions, either by reinforcing mood or by avoiding destructive thoughts related to not exercising<sup>(22)</sup>. On the other hand, people with food addiction are less active and prefer spending their leisure time sitting rather than walking or doing any physical activity<sup>(23)</sup>. Also, education level has an impact on food behaviours and attitudes<sup>(24-26)</sup>: uneducated persons tend to consume large amounts of junk, high energetic content food<sup>(26)</sup>, while more educated people read and analyse health facts and choose healthier lifestyles<sup>(24)</sup>.

In Lebanon, there is a need to assess the risk of eating-related diseases and eating attitudes using appropriate tools; one of the most used is the Eating Attitude Test (EAT-26)<sup>(27)</sup>. The EAT-26 was designed to diagnose anorexia nervosa but has been found useful in non-clinical settings to screen for disordered eating attitudes, particularly atypical preoccupation with food and weight<sup>(27)</sup>. This test was validated in Mexico as a secondary prevention tool to modify public health programmes<sup>(28)</sup>, and in Spain<sup>(29,30)</sup>, where it is known as Disorder Eating Attitude Scale. It was also validated as a very sensitive method in Italy<sup>(31)</sup>, as a sensitive and specific useful tool in non-western countries<sup>(32)</sup> and as the most reliable and commonly used self-report tool for eating disorders worldwide. Additionally, the EAT-26 was validated in Arabic among a representative sample of female students in Riyadh, Saudi Arabia<sup>(33)</sup>. However, it was not validated in the Lebanese dialect and among the general population; hence, the importance of its validation to use it as a screening tool for disordered eating attitudes.

Therefore, the primary objective of the study is to validate the Arabic version of the EAT-26 and identify factors

(such as depression, stress, anxiety, emotion regulation and body dissatisfaction) that might be associated with disordered eating among a sample of the Lebanese population.

## Methods

### Participants

This cross-sectional study conducted between January and May 2018, enrolled 811 community dwelling participants using a proportionate random sample from all Lebanese governorates (Beirut, Mount Lebanon, North, South and Bekaa). Each governorate is divided into Caza (stratum). Two villages were randomly selected from the list of villages provided by the Central Agency of Statistics in Lebanon. Participants were randomly selected from each village. Prior to participation, individuals were briefed on the study objectives and methodology and were assured of the anonymity of their participation. No financial reward was given to the participants, and they had the right to accept or refuse to participate in the study. Those who agreed to participate were asked to read and sign a written consent form.

All participants above 18 years of age were eligible to participate. Excluded were those who refused to fill out the questionnaire and those suffering from cognitive impairment as reported by a family member. Data collection was performed through personal interviews with participants by a trained study-independent clinical psychologist, who clinically evaluated the level of psychiatric illness to exclude participants with psychiatric problems. The same methodology was used in previous papers from this project<sup>(34-36)</sup>.

### Minimal sample size calculation

According to Comrey and Lee<sup>(37)</sup>, ten observations are needed for each scale item for the validation process. Since the EAT includes twenty-six items, a minimal sample of 260 participants was deemed necessary.

### Procedure

Study-independent trained clinical psychologists collected data by performing personal interviews with the participants that required approximately 60 min. Completed questionnaires were collected back by the interviewer and sent for data entry. During the data collection process, the anonymity of the participants was guaranteed.

The full sample was randomly divided into two separate samples (406 for the first sample and 405 for the second sample). To ensure the validity of the results from sample 1, the score of the EAT scale used in sample 1 was tested on another sample (sample 2).

### Questionnaire

The questionnaire used during the interview was in Arabic, the native language of Lebanon. The first part assessed the socio-demographic details of the participants. The BMI was



calculated as follows: weight (kg) divided by the square of the height (metres), as self-reported by each participant. Consumption of alcohol, caffeinated beverages and tobacco was categorised into dichotomous variables (yes/no). The physical activity index is a frequently used indicator of physical activity at the population level. This index is based on responses to a series of questions about the intensity, frequency and duration of participation in physical activity during leisure time. The Total Physical Activity Index was calculated by multiplying the intensity, duration and frequency of daily activity<sup>(38)</sup>. The education level was categorised into four groups: primary, complementary, secondary and university; however, when stratifying over education, the four groups were reduced to two, low level of education (illiterate, primary and complementary) and high level of education (secondary, university).

The second part of the questionnaire consisted of the perception of eating habits among participants. The questions were selected from previous articles about eating disorders<sup>(39–41)</sup>. The final part included the following scales.

#### *Eating Attitude Test*

The EAT is used to assess disordered food attitude<sup>(42)</sup>. The questionnaire comprises twenty-six questions each with six response options, varying from infrequently/almost never/never (0) to always (3). The total score is calculated by summing all questions answers and can vary from 0 to 78. A score of 20 or above indicates possible disordered food attitudes<sup>(43)</sup>. In the current study, the Cronbach's  $\alpha$  was 0.908.

#### *Body dissatisfaction subscale of the Eating Disorder Inventory second version*

In the present study, body dissatisfaction subscale of Eating Disorder Inventory second version was used to measure body disturbance. The body dissatisfaction subscale contains nine items assessing levels of dissatisfaction with their overall body shape as well as specific body parts. A four-point Likert scale was used ranging from 0 (sometimes, rarely and never) to 3 (always). Higher scores indicate more body dissatisfaction<sup>(44)</sup>. In the current study, the Cronbach's  $\alpha$  was 0.779.

#### *Perceived Stress Scale*

The questions in the Perceived Stress Scale ask about feelings and thoughts during the last month<sup>(45)</sup>. The Perceived Stress Scale is a ten-item scale, with answers ranging from never (0) to almost always (4). Items 4, 5, 7 and 8 are reversed items. The total score is calculated by summing the ten items with higher scores indicating more perceived stress<sup>(45)</sup>. In the current study, the Cronbach's  $\alpha$  was 0.709.

#### *Hamilton Anxiety Rating Scale*

The Hamilton Anxiety Rating Scale, validated in Lebanon<sup>(46)</sup>, consists of fourteen symptom-defined elements and targets

both psychological and somatic symptoms. Each item is scored on a basic numeric scoring of 0 (not present) to 4 (severe). The total score ranging from 0 to 56, with higher scores indicating higher anxiety<sup>(47)</sup>. In the current study, the Cronbach's  $\alpha$  was 0.912.

#### *Hamilton Depression Rating Scale*

The Hamilton Scale for Depression, validated in Lebanon<sup>(48)</sup>, is used to evaluate the severity of depression in patients who are already diagnosed as depressed. The total score is based on the sum of the first seventeen items only. Higher scores indicated higher depression<sup>(49)</sup>. In the current study, the Cronbach's  $\alpha$  was 0.879.

#### *Emotion Regulation Questionnaire (ERQ)*

The Emotion Regulation Questionnaire is used to measure respondents' tendency to regulate their emotions in two ways: (1) cognitive reappraisal and (2) expressive suppression. The cognitive reappraisal facet is a way of managing and controlling attention and cognitively changing the meaning of emotionally stimulating stimuli<sup>(50)</sup>. It is considered a healthy emotion regulation strategy<sup>(50)</sup>. The expressive suppression involves inhibition of emotional expressive behavior, thereby changing the emotional impact of a situation<sup>(51)</sup>. It is considered a less healthy emotion regulation strategy<sup>(51)</sup>. A ten-item scale ranging from 1 (strongly disagree) to 7 (strongly agree). Items 1, 3, 5, 7, 8 and 10 make up the cognitive reappraisal facet and items 2, 4, 6 and 9 make up the expressive suppression facet. Each facet's scoring is kept separate. The higher the scores, the greater the use of the emotion regulation strategy<sup>(52)</sup>. In the current study, the Cronbach's  $\alpha$  values for the cognitive reappraisal facet and the expressive suppression facet were 0.744 and 0.732, respectively.

The emotional eating, restrained eating, orthorexia nervosa and binge eating scales were used to test the convergent validity of the EAT scale.

#### *Emotional Eating Scale*

The Emotional Eating Scale, validated in Lebanon<sup>(53)</sup> is a twenty-five-item scale with three-factor analytically derived subscales: anger, anxiety and depression. Participants rate the extent to which certain feelings lead to the urge to eat using a five-point Likert scale ranging from 0 (no desire to eat) to 4 (an overwhelming urge to eat). Higher scores indicate a reliance on using food to help managing emotions<sup>(54)</sup>. In the current study, the Cronbach's  $\alpha$  was 0.957.

#### *Dutch Restrained Eating Scale*

Recently, Dutch Restrained Eating Scale, validated in Lebanon<sup>(55)</sup>, is composed of ten items scored using a five-point Likert scale, varying from 1 (never) to 5 (always). By dividing the total items score by the total number of items, the score for this scale was acquired. A greater score would show higher degree of restrained eating<sup>(56)</sup> (Cronbach's  $\alpha = 0.928$ ).



### *Orthorexia Nervosa Scale – ORTO-15*

The Orthorexia Nervosa Scale, validated in Lebanon<sup>(57)</sup>, is a measure instrument comprising fifteen multiple-choice items. Orthorexia Nervosa Scale is a self-reported questionnaire<sup>(58)</sup> with a four-point Likert scale (never, sometimes, often and always). Lower scores would indicate higher levels of orthorexia tendencies and behaviours. In the current study, the Cronbach's  $\alpha$  was 0.822.

### *Binge Eating Scale*

The Binge Eating Scale, validated in Lebanon<sup>(59)</sup>, was originally developed to identify binge eaters within an obese population<sup>(60)</sup>. It does not specify a time frame and presents a series of differently weighted statements for each item, from which respondents select the statement that best describes their attitudes and behaviours. This yields a continuous measure of binge eating pathology of 0–46. The severity of binge eating was divided into three categories as follows: < 17 as non-binging, between 18 and 26 as moderate binging and  $\geq 27$  as severe binging<sup>(61)</sup>. The Binge Eating Scale has good test-retest reliability ( $r = 0.87$ ,  $P < 0.001$ ). In the current study, the Cronbach's  $\alpha$  was 0.862.

### **Forward and back translation procedure**

Forward translation was first conducted by a single bilingual translator, whose native language is Arabic and fluent in English. An expert committee formed by healthcare professionals and a language professional verified the Arabic-translated version. A backward translation was then performed by a native English speaker translator, fluent in Arabic and unfamiliar with the concepts of the scales. The back-translated English questionnaire was subsequently compared with the original English one, by the expert committee, aiming to discern discrepancies and to solve any inconsistencies between the two versions. The process of forward-back translation was repeated until all ambiguities disappeared.

### **Statistical analysis**

SPSS software version 25 was used to conduct data analysis. A descriptive analysis was done using the counts and percentages for categorical variables and mean and SD for continuous measures. We checked the distribution normality for the EAT scale using the Shapiro Wilk test. Since the assumption of normality was not normally distributed, the comparison of means was performed using the non-parametric tests (Kruskal–Wallis and Mann–Whitney tests). Spearman correlation was used for the linear correlation between continuous variables. For categorical variables, the  $\chi^2$  and Fisher exact tests were used.

A stepwise linear regression was conducted, taking the EAT score as the dependent variable. All variables that showed a  $P < 0.1$  in the bivariate analysis were considered important variables to be entered in the model in order to

eliminate potentially confounding factors as much as possible. Afterwards, other stepwise linear regressions taking the EAT as the dependent variable but stratified over education level were conducted (stratification analysis).

Two different methods were used to confirm the EAT questionnaire construct validity. First, a factor analysis was run using the principal component analysis technique, run on sample 1. Since the extracted factors were found to be significantly correlated, the promax rotation technique was used. To ensure the model's adequacy, the Kaiser–Meyer–Olkin measure of sampling adequacy and Bartlett's test of sphericity were calculated. Factors with an eigenvalue higher than one were retained. Moreover, Cronbach's  $\alpha$  was recorded for reliability analysis for each scale.

Second, a confirmatory factor analysis was carried out in sample 2. To assess the structure of the instrument, the maximum likelihood method for discrepancy function was used. Several goodness-of-fit indicators were reported: relative chi-square ( $\chi^2/\text{df}$ ), root mean square error of approximation, goodness of fit index and the adjusted goodness of fit index. The goodness of fit index was calculated by the value of  $\chi^2/\text{df}$  (cut-off values < 2–5). The root mean square error of approximation tests the fit of the model to the covariance matrix. As a guideline, values of < 0.05 indicate a close fit and values below 0.11 an acceptable fit. The goodness of fit index and adjusted goodness of fit index are  $\chi^2$ -based calculations independent of df. The recommended thresholds for acceptable values are  $\geq 0.90$ <sup>(62)</sup>. A  $P < 0.05$  was considered significant.

## **Results**

Out of 1000 questionnaires distributed, 811 (81.1%) were completed and collected. The socio-demographic characteristics of the participants in the first sample are summarised in Table 1. The results from sample 1 showed that disordered eating attitudes was found in 95 (23.8%) (95% CI 0.152–0.323) participants, whereas 305 (76.2%) participants had appropriate eating attitudes (95% CI 0.715–0.810). No significant difference was found between samples 1 and 2 for all the variables (socio-demographic variables, smoking, alcohol and carbonated beverages intake, and physical activity).

### **Exploratory factor analysis on sample 1**

Out of all the items of EAT-26 scale, none of the items was removed since no one over-correlated to each other ( $r > 0.9$ ), or had a low loading on factors (< 0.3) or even had a low communality (< 0.3). The factor analysis for the EAT scale was run over half of the original sample (total  $n$  403). The EAT-26 scale items converged over a solution of six factors that had an eigenvalue over 1 (factor 1 = restrained eating attitude, factor 2 = anorexic attitude,



**Table 1** Socio-demographic characteristics of the study sample

	Total		Sample 1		Sample 2		P
	Frequency (%)		Frequency (%)		Frequency (%)		
	n	%	n	%	n	%	
Gender							
Male	270	33.5	131	3.4	139	34.6	0.518
Female	536	66.5	273	67.6	263	65.4	
Marital status							
Single	533	67.0	263	66.1	270	67.8	0.395
Married	230	28.9	120	30.2	110	27.6	
Widowed	11	1.4	7	1.8	4	1.0	
Divorced	22	2.8	8	2.0	14	3.5	
Education level							
Primary	24	3.1	11	2.8	13	3.3	0.760
Complementary	61	7.8	34	8.7	27	6.9	
Secondary	125	15.9	60	15.3	65	16.6	
University	574	73.2	288	73.3	286	73.1	
Monthly income							
No income	340	45.1	168	44.4	172	45.7	0.140
<1000 USD	247	32.8	121	32.0	126	33.5	
1000–2000 USD	117	15.5	56	14.8	61	16.2	
>2000 USD	50	6.6	33	8.37	17	4.5	
Smoking							
Yes	246	30.8	127	31.7	119	29.8	0.572
No	554	69.2	274	68.3	280	70.2	
Alcohol							
Yes	32	4.2	12	3.1	20	5.4	0.124
No	724	95.8	372	96.9	352	94.6	
Caffeinated beverages							
Yes	721	90.0	362	90.5	359	89.5	0.646
No	80	10.0	38	9.5	42	10.5	
Practicing sport activities							
Yes	490	62.4	248	63.6	242	61.3	0.501
No	295	37.6	142	36.4	153	38.7	
Age (in years)							0.354
Mean	27.59		27.96		27.19		
SD	11.76		11.83		11.68		
BMI (kg/m <sup>2</sup> )							0.182
Mean	24.35		24.10		24.60		
SD	5.28		4.98		5.55		

The  $\chi^2$  and Fisher exact tests were used to evaluate the association between categorical variables, and the Mann–Whitney test was used to evaluate the association between the age, body mass index (BMI) and the sample group.

factors 3 and 4 = cycle of bulimic crises, factor 5 = impact of others' perception and factor 6 = choice of food quality and time spent on meals; total variance explained = 60.07%, Kaiser-Meyer-Olkin = 0.911, Bartlett's test of sphericity  $P < 0.001$  and Cronbach's  $\alpha = 0.895$ ) According to the promax rotated matrix, the components are summarised in Table 2.

Another factor analysis was run by forcing a rotation over a three-factor solution instead of six; three items (2, 18 and 26) were removed, leaving a total of 23 items (total variance explained = 48.73%, Kaiser-Meyer-Olkin = 0.904, Bartlett's test of sphericity  $P < 0.001$ ). According to the promax rotated matrix, the components are summarised in Table 3.

### Confirmatory factor analysis on sample 2

A confirmatory factor analysis was run on sample 2, using the structure obtained in sample 1. The following results

were obtained: the maximum likelihood  $\chi^2 = 723$  and  $df = 300$ , which gave a  $\chi^2/df = 2.4$ . For non-centrality fit indices, the Steiger-Lind root mean square error of approximation was 0.13 (0.131–0.142). Moreover, the Joreskog goodness of fit index equaled 0.767 and adjusted goodness of fit index equaled 0.711.

### Bivariate analysis conducted on the whole sample

A significantly higher mean of EAT score was found in participants following a diet (18.85 *v.* 11.66,  $P < 0.001$ ), exercising (16.66 *v.* 12.06,  $P < 0.001$ ), vomiting (25.83 *v.* 12.49,  $P < 0.001$ ), taking diet pills (23.43 *v.* 12.74,  $P < 0.001$ ), starving to lose weight (21.26 *v.* 11.84,  $P < 0.001$ ) in the last 30 d and weighing daily (20.09 *v.* 12.49,  $P = 0.005$ ) compared with those who do not follow these eating habits. Also, a significantly higher mean of EAT score was found in participants who received comments from their family members about losing weight (16.75 *v.* 12.48,  $P < 0.001$ ), those who

**Table 2** Promax rotated matrix of the eating attitude eating scale items

Items	Factor loading					
	1	2	3	4	5	6
Factor 1: restrained eating attitude ( $\alpha = 0.785$ )						
6. Aware of the energy content of foods that I eat	0.852					
16. Avoid foods with sugar in them	0.790					
17. Eat diet foods	0.742					
7. Particularly avoid food with a high carbohydrate content (i.e., bread, rice, potatoes, etc.)	0.734					
5. Cut my food into small pieces	0.516					
19. Display self-control around food	0.405					
Factor 2: anorexic attitude ( $\alpha = 0.811$ )						
11. Am preoccupied with a desire to be thinner		0.831				
12. Think about burning up energies when I exercise		0.797				
1. Am terrified about being overweight		0.733				
14. Am preoccupied with the thought of having fat on my body		0.719				
10. Feel extremely guilty after eating		0.489				
2. Avoid eating when I am hungry		0.434				
Factors 3 and 4: cycle of bulimic crises ( $\alpha = 0.710$ and $0.807$ )						
3. Find myself preoccupied with food			0.852			
4. Have gone on eating binges where I feel that I may not be able to stop			0.843			
9. Vomit after I have eaten			0.482			
18. Feel that food controls my life			0.478			
21. Give too much time and thought to food				0.721		
22. Feel uncomfortable after eating sweets				0.684		
24. Like my stomach to be empty				0.609		
23. Engage in dieting behaviour.				0.589		
25. Have the impulse to vomit after meals				0.491		
Factor 5: impact of others' perception ( $\alpha = 0.734$ )						
13. Other people think that I am too thin					0.837	
8. Feel that others would prefer if I ate more					0.819	
20. Feel that others pressure me to eat					0.709	
Factor 6: choice of food quality and time spent on meals ( $\alpha = 0.432$ )						
26. Enjoy trying new rich foods						0.611
15. Take longer than others to eat my meals						0.479
Percentage of variances explained	32.06	7.41	6.73	5.28	4.39	4.20

have been insulted (21.84 *v.* 12.79,  $P < 0.001$ ), those who have been in a bad relationship (15.35 *v.* 13.16,  $P = 0.027$ ), had a family history of eating disorders (17.18 *v.* 12.82,  $P = 0.001$ ) and felt pressure from media to lose weight (19.68 *v.* 12.45,  $P < 0.001$ ) compared with those who do not agree with these statements.

In addition, a higher body dissatisfaction ( $r = 0.135$ ), higher restrained eating ( $r = 0.229$ ), higher anxiety ( $r = 0.285$ ), higher depression ( $r = 0.302$ ), higher physical activity ( $r = 0.092$ ), higher binge eating ( $r = 0.277$ ) and higher emotional eating ( $r = 0.143$ ) were significantly but weakly associated with higher EAT scores. However, higher orthorexia scores (lower orthorexia tendencies and behaviours) ( $r = -0.256$ ) were significantly associated with lower EAT scores (more appropriate eating) (Tables 4 and 5).

#### **Multivariable analysis conducted on the whole sample**

The results of a first linear regression, taking the EAT score as the dependent variable and the socio-demographic characteristics only as independent variables, showed that secondary ( $\beta = -5.87$ ) and university level of education

( $\beta = -5.26$ ) were significantly associated with lower EAT score (appropriate eating attitudes) (Table 6, model 1).

The results of a second stepwise linear regression, considering the EAT score as the dependent variable and the scales and the opinion about eating habits as independent variables, showed that higher depression ( $\beta = 0.325$ ), higher emotional eating ( $\beta = 0.083$ ), daily weighing ( $\beta = 3.430$ ), higher physical activity (0.05), starving to reduce weight ( $\beta = 4.94$ ) and receiving pressure from TV/magazine to lose weight ( $\beta = 3.95$ ) were significantly associated with higher EAT scores (disordered eating attitudes) (Table 6, model 2).

#### **Stratification over education level conducted on the whole sample**

A first linear regression, taking the EAT-26 as the dependent variable and the socio-demographic characteristics as independent variables, stratified over education level, showed that low income ( $\beta = 11.21$ ) was significantly associated with higher EAT-26 scores (disordered eating attitudes) in the low-educated group. Higher BMI ( $\beta = 0.24$ ) and intermediate income ( $\beta = 3.86$ ) were significantly associated with higher EAT-26 scores (disordered eating

**Table 3** Promax rotated matrix of the eating attitude eating scale items using a forced three-factor solution

Items	Factor loading		
	1	2	3
Factor 1: restrained eating attitude ( $\alpha = 0.852$ )			
14. Am preoccupied with the thought of having fat on my body	0.789		
11. Am preoccupied with a desire to be thinner	0.748		
12. Think about burning up energies when I exercise	0.675		
17. Eat diet foods	0.646		
24. Like my stomach to be empty	0.607		
1. Am terrified about being overweight	0.603		
23. Engage in dieting behaviour	0.585		
16. Avoid foods with sugar in them	0.541		
22. Feel uncomfortable after eating sweets	0.495		
6. Aware of the energy content of foods that I eat	0.438		
2. Avoid eating when I am hungry	0.417		
Factor 2: anorexic attitude ( $\alpha = 0.785$ )			
4. Have gone on eating binges where I feel that I may not be able to stop		0.783	
3. Find myself preoccupied with food		0.706	
9. Vomit after I have eaten		0.676	
25. Have the impulse to vomit after meals		0.676	
21. Give too much time and thought to food		0.587	
10. Feel extremely guilty after eating		0.530	
Factors 3 ( $\alpha = 0.742$ )			
8. Feel that others would prefer if I ate more			0.734
13. Other people think that I am too thin			0.720
20. Feel that others pressure me to eat			0.588
15. Take longer than others to eat my meals			0.525
19. Display self-control around food			0.523
5. Cut my food into small pieces			0.489
Percentage of variances explained	32.68	8.27	7.70

**Table 4** Spearman correlation between Eating Attitude Test (EAT) and other scales

	EAT score	
	Correlation coefficient	P
Body dissatisfaction score	0.135	<0.001
Restrained Eating Scale	0.229	<0.001
Perceived Stress Scale	0.068	0.055
Anxiety	0.285	<0.001
Depression	0.302	<0.001
Emotion regulation cognitive reappraisal facet	-0.044	0.239
Emotion regulation expression suppression facet	0.042	0.271
Emotional eating	0.143	<0.001
Orthorexia nervosa	-0.256	<0.001
Binge eating scale	0.277	<0.001
Physical activity index	0.092	0.035

Numbers in bold indicate significant *P*-values. Pearson correlation coefficient was used to evaluate the association between the EAT score and the continuous variables.

attitudes), whereas increase age ( $\beta = -0.11$ ) was significantly associated with lower EAT-26 scores (appropriate eating attitudes) in the high-educated group.

A second linear regression analysis taking the EAT-26 as the dependent variable and the scales and the opinion about eating habits as independent variables, stratified over education level, showed that starving to reduce weight ( $\beta = 14.00$ ) was significantly associated with higher EAT-26 scores (disordered eating attitudes) in the low-educated group. Daily weighting ( $\beta = 3.17$ ), higher depression

( $\beta = 0.30$ ), receiving pressure from the TV/magazine to lose weight ( $\beta = 4.24$ ) and physical activity ( $\beta = 0.07$ ) were significantly associated with higher EAT scores (disordered eating attitudes) in the high-educated group (Table 7).

## Discussion

To the best of our knowledge, this is the first study to assess factors associated with the eating attitude among a

**Table 5** Bivariate analysis of the factors associated with the Eating Attitude Test (EAT) score

		EAT score		
		Mean	SD	P
Education level	Primary	13.29	12.38	0.169
	Complementary	16.73	15.93	
	Secondary	12.23	11.19	
Monthly income	University	13.39	12.20	0.402
	No income	13.06	11.53	
	<1000 USD	13.37	13.31	
	1000–2000 USD	15.74	14.71	
Dieted to lose weight (past 30 d)	>2000 USD	11.71	10.77	<0.001
	No	11.66	11.44	
Exercised to lose weight (past 30 d)	Yes	18.85	14.43	<0.001
	No	12.06	12.03	
Vomited or taken laxatives to lose weight (past 30 d)	Yes	16.66	13.59	<0.001
	No	12.49	11.84	
Taken diet pills to lose weight (past 30 d)	Yes	25.83	15.23	<0.001
	No	12.74	11.99	
Starving to lose weight (past 30 d)	Yes	23.43	15.39	<0.001
	No	11.84	11.28	
Daily weighing	Yes	21.26	15.36	<0.001
	No	12.44	11.46	
Receiving comments from the family concerning losing weight	Yes	20.09	16.37	<0.001
	No	12.48	12.20	
Being insulted because of weight	Yes	16.75	13.68	<0.001
	No	12.79	11.78	
Being in a bad romantic relationship	Yes	21.84	17.92	0.027
	Yes	24.35	17.16	
Family history of eating disorders	No	13.16	12.45	0.001
	Yes	15.35	13.56	
Pressure from TV, magazine to lose weight	No	12.82	12.03	<0.001
	Yes	17.18	14.81	
	No	12.45	12.05	
	Yes	19.68	14.52	

Numbers in bold indicate significant P-values; the Mann–Whitney test was used to evaluate the association between the EAT score and dichotomous variables, whereas the Kruskal–Wallis test was used to evaluate the association between the EAT score and the categorical variables.

**Table 6** Multivariable analysis

	Unstandardised $\beta$	Standardised $\beta$	P	CI	
				Lower Bound	Upper Bound
Model 1: Linear regression taking the Eating Attitude Test score as the dependent variable and the socio-demographic characteristics only as the independent variables*					
Secondary level of education compared to illiteracy*	-5.878	-0.167	0.007	-10.182	-1.575
University level of education compared to illiteracy*	-5.266	-0.187	0.009	-9.218	-1.314
Model 2: Linear regression taking the Eating Attitude Test score as the dependent variable and the scales and the behavioural practices of certain eating habits as independent variables†					
Depression	0.325	0.191	<0.001	0.188	0.462
Emotional eating	0.083	0.123	0.003	0.028	0.137
Daily weighing (yes v no*)	3.430	0.104	0.013	0.726	6.134
Physical activity index	0.054	0.095	0.016	0.010	0.098
Starving to reduce weight (yes v no*)	4.942	0.155	<0.001	2.258	7.625
Feeling pressure from TV/magazine to lose weight (yes v no*)	3.955	0.115	0.006	1.151	6.759

\*Reference group; Variables entered in model 1: age, BMI, gender, marital status, monthly income and education level.

†Variables entered in model 2: age, BMI, gender, marital status, monthly income, education level, body dissatisfaction, anxiety, depression, emotional eating, binge eating, physical activity index, dieted to lose weight, exercised to lose weight, vomited or taken laxatives to lose weight, taken diet pills to lose weight, starving to lose weight, daily weighing, receiving comments from the family concerning losing weight, being insulted, being physically abused, being sexually abused, being in a bad romantic relationship, family history of eating disorders, pressure from TV, magazine to lose weight. Nagelkerke  $R^2 = 26.5\%$ .

representative Lebanese population sample. Our results showed that 23.8% of participants had disordered eating attitudes, whereas 76.2% had appropriate eating attitudes and are in line with other studies: a study carried out among

young women from Brazil (2003) showed a clinically significant disturbed eating behaviour in 16.5% of women, with more disordered eating attitudes among those aged 16–19 years and among overweight/obese women<sup>(63)</sup>. The



**Table 7** Stratification over the education level\*

	Low education level (below 12 years of education)			High education level (over 12 years of education)		
	$\beta$	95 % CI	<i>P</i>	$\beta$	95 % CI	<i>P</i>
Model 1: Linear regression taking the Eating Attitude Test score as the dependent variable and the socio-demographic characteristics as the independent variables						
Age	0.18	-0.09, 0.46	0.186	-0.11	-0.20, 0.02	<b>0.009</b>
Body mass index	0.25	-0.22, 0.73	0.295	0.24	0.04, 0.43	<b>0.016</b>
Low income v no income*	11.21	3.35, 19.07	0.006	-1.87	-4.02, 0.27	0.087
Intermediate income v no income*	-3.39	-14.29, 7.50	0.537	3.86	1.20, 6.53	<b>0.005</b>
Model 2: Linear regression taking the Eating Attitude Test score as the dependent variable and the scales and the behavioural practices of certain eating habits as independent variables						
Daily weighting (yes v no*)	10.30	-5.84, 26.46	0.194	3.17	0.41, 5.93	<b>0.024</b>
Pressure from TV, magazine to lose weight (yes v no*)	-0.10	-12.00, 11.80	0.986	4.24	1.37, 7.11	<b>0.004</b>
Starving to reduce weight (yes v no*)	14.00	4.76, 23.24	0.004	3.65	0.82, 6.48	<b>0.012</b>
Depression (HAMD)	0.88	-0.29, 2.05	0.130	0.30	0.14, 0.45	<b>&lt;0.001</b>
Physical activity	0.04	-0.27, 0.36	0.764	0.07	0.03, 0.12	<b>0.001</b>

\*Reference group; Low education level includes illiterate, primary and complementary categories, whereas the high education level includes the secondary and university categories; numbers in bold indicate significant *P*-values.

prevalence of disorder eating attitudes in western countries varies from 0.4 % in Spain<sup>(64)</sup> to 33 % in Australia<sup>(65)</sup>.

In our study, EAT scale items converged over a solution of six factors explaining a total of 60.07 % of the variance and leading to an internal consistency of Cronbach's  $\alpha = 0.895$ . The validation of the EAT in Mexico showed five factors (constituting a total of 46.6 % of the variance): 'dietary restraint', 'bulimia', 'drive of thinness', 'food preoccupation and perceived social pressure' and with an internal consistency of Cronbach's  $\alpha = 0.9$ <sup>(66)</sup>. The Spanish validation in its turn yielded three factors explaining 41 % of the variance, with a reliability coefficient of 0.93<sup>(30)</sup>. This test was also validated in Italy and consisted of four phases: 'Item generation, internal consistency and reliability analysis, normative data generation, and validity analysis', with an internal consistency Cronbach's  $\alpha$  values of 0.90 and 0.93<sup>(67)</sup>. Thereby, Cronbach's  $\alpha$  values show a very small variation between different countries: this is due to the fact that the Cronbach's  $\alpha$  is not a characteristic for the whole test but for the sample took in the test; it might vary if the test is done on a general population or a sample composed of people having a specific characteristic, for example. However, these obtained values were almost similar, meaning these populations share very close eating attitudes.

Receiving pressure from TV or magazines to lose weight, starving to reduce weight (in the past 30 d), higher depression and higher physical activity were associated with higher EAT scores (disordered eating attitudes) in the better educated group. Previous studies revealed that physical activity levels explained the relationship between prudent dietary patterns, depressive symptoms<sup>(68)</sup> and severity of anxiety symptoms<sup>(69)</sup>; specifically, lower scores of the healthy dietary pattern and higher scores of the unhealthy dietary pattern independently predicted increased

depressive symptoms across time without establishing any correlation with educational level. Furthermore, social media exposure was associated with eating pathology, independent of direct media exposure and other cultural exposures. The different variables explored were not directly correlated with the level of education in the literature. Thus, our findings, related to the stratification by level of education, warrant further investigation to better understand the relation between the level of education and the different variables associated at the same time with the score of EAT.

In the current study, higher depression levels were found to be significantly associated with more disordered eating attitudes (higher EAT scores), similarly to other studies<sup>(17,19,70)</sup>, knowing that depression is often due to an anxiety feeling over a long period of time<sup>(71)</sup>. These unwanted feelings are usually accompanied by an intensified need for eating high-sugar food or high-fat food as a self-medication<sup>(17)</sup>. This might be attributed to a greater body concern of women nowadays for having an ideal social media body to look 'beautiful and attractive'<sup>(6)</sup>. In fact, the desire to become similar to TV characters increased the anxiety and depression among women<sup>(19)</sup>, and these unpleasant feelings resulted sometimes in excessive eating<sup>(72)</sup>. Clearly, peer pressure and social media stars have a bad influence on individuals, especially women, to the point where body image dissatisfaction became a 'normative discontent'<sup>(16)</sup>. Studies show that a higher frequency of social media use, especially the image-centric platforms, resulted in greater body dissatisfaction concern<sup>(6)</sup>. Effectively, these social media platforms tend to promote disordered eating behaviours by always encouraging the 'thin ideal' leading to disastrous weight loss tendencies<sup>(7)</sup>, fasting, higher depression, daily weighing, etc.

Therefore, people are tending to adapt starving methods such as fasting, purging, chronic dieting<sup>(19)</sup> or the daily



weighing method, thinking that it would help them lose weight. Although daily weighing might help them manage their weight by always keeping an eye on it<sup>(73)</sup>, starvation for example, will lead to even more weight gain, because our body does not recognise the difference between this procedure and dieting.

In return, emotional eating was proven to be strongly related to disordered eating attitudes, which is consistent with the results of other studies<sup>(15)</sup>. Many factors were evaluated in the literature to highlight the effect of emotions on eating attitudes. Individuals suffering from a lack of attention from parents or friends<sup>(74)</sup>, and sexual and emotional abuse, showed higher levels of eating disorders, due to the compensation of affection in food, and for those people, food is known to be their 'comfort zone'. Nowadays, people are exposed to numerous criticisms which creates among adolescents a huge personal irritation and a lower self-esteem, making it a crucial cause for restrained eating, followed by binge eating or over eating.

High education level had been found to be related with appropriate eating attitudes; thus, a stratification by level of education was found to be important. A low education level was related to unhealthy dietary habits: low-educated people tend to consume dense food, whereas high-educated people have more positive attitudes in choosing foods as they acquire adequate health information that improves ability in choosing the right foods. A study conducted in 2013<sup>(75)</sup> revealed that more educated people were more compliant with dietary guidelines and expressed more frequently positive attitudes towards healthy eating than low-educated people. Our results showed that low income was associated with higher disordered eating behaviour among low-educated groups. Less educated and low-income people have less healthy dietary habits, in part because of their higher priority for price and familiarity and their lower priority for health as a reason for buying food. A Finnish study showed that the level of education is related to individual priorities in the choice of food, where the more educated consume fewer energy-rich foods than those with less education<sup>(76)</sup>. However, the association between the level of education and eating attitudes is not fully understood, and further studies are needed to investigate the attitudes and motives for making food choices.

### **Clinical implications**

Healthcare professionals and dietitians should take into consideration the factors affecting a person's eating attitude before prescribing a diet plan. Reasons differ from one person to another, as for some, eating disorders are a response to a major impact that affected their lives (early separation, stress, depression, anxiety, etc.), while for others, a non-balanced diet occurs following a low self-esteem and a self-objectification, due to the major influence of social media and TV programmes. Therefore, healthcare professionals should not only be responsible for prescribing

trendy diet plans but also change their approach with patients and understand the reasons behind such disordered attitudes towards eating, then adapt the diet plan accordingly.

### **Limitations**

An information bias might have occurred during the data collection since it was based on a self-report questionnaire. Our data could not be generalised to the entire population, since the group of well-educated female participants was more represented; this led us to conduct the stratified analysis and show differential results between high-*v.* low-educated subgroups. We would assume some kind of self-selection bias: maybe from each randomly selected household, younger and more educated individuals (e.g., students) participated in the study more often than older and less educated individuals. One additional limitation is the absence of an evaluation tool for obsessive-compulsive disorder and anorexic behaviours. It will be interesting to evaluate these behaviours among persons with disordered eating attitudes.

### **Conclusion**

The current study found a high prevalence of disordered eating attitudes in the Lebanese population. Yet, the obtained results are considered preliminary, and further research should be conducted to prove it. Disordered eating attitudes and disorders will remain a controversial issue with respect to their relationship with psychological health and mental disorders. Future studies should point out at the importance of raising the awareness about social media, peer pressure and psychological influence on eating attitudes, since these are the main causes of eating disorders nowadays.

### **Acknowledgements**

*Acknowledgements:* The authors would like to thank Marwan Akel, Karl Honein and Maria Akiki for their help in the data entry and all the participants who helped us during this project. *Financial support:* None. *Conflict of interest:* The authors have no conflicts of interest to report. *Authorship:* S.O. and S.H. conceived and designed the survey. C.H., S.H. and P.S. were involved in the statistical analysis and data interpretation. C.H. and C.K. wrote the manuscript. H.S., R.H. and N.K. reviewed the manuscript. All authors read the manuscript, critically revised it for intellectual content and approved the final version. *Ethics of human subject participation:* The current study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving

research study participants were approved by the Psychiatric Hospital of the Cross Ethics and Research Committee. Written informed consent was obtained from all subjects/patients. *Consent for publication:* All authors gave their consent for publication. *Availability of data and material:* The authors do not have the right to share any data information as per their institutions policies.

## References

- Alvarenga Mdos S, Scagliusi FB & Philippi ST (2010) Development and validity of the disordered eating attitude scale (DEAS). *Percept Mot Skills* **110**, 379–395.
- Fatima W & Ahmad LM (2018) Prevalence of disordered eating attitudes among adolescent girls in Arar city, Kingdom of Saudi Arabia. *Health Psychol Res* **6**, 7444.
- Kazim A, Almarzooqi M & Karavetian M (2017) The prevalence and determents of eating disorders among Emirati female students aged 14–19 years in Ajman, UAE. *J Food Nutr Disord* **6**, 2.
- Rauof M, Ebrahimi H, Jafarabadi MA *et al.* (2015) Prevalence of eating disorders among adolescents in the northwest of Iran. *Iran Red Crescent Med J* **17**(10), e19331. doi: 10.5812/ircmj.19331.
- Saleh RN, Salameh RA, Yhya HH *et al.* (2018) Disordered eating attitudes in female students of An-Najah national university: a cross-sectional study. *J Eat Disord* **6**, 16.
- Griffiths S, Murray SB, Krug I *et al.* (2018) The contribution of social media to body dissatisfaction, eating disorder symptoms, and anabolic steroid use among sexual minority men. *Cyberpsychol Behav Soc Netw* **21**, 149–156.
- Cavazos-Rehg PA, Krauss MJ, Costello SJ *et al.* (2019) “I just want to be skinny.”: a content analysis of tweets expressing eating disorder symptoms. *PLoS One* **14**, e0207506.
- American Psychiatric Association, DSM-5 Task Force (2013) *Diagnostic and Statistical Manual of Mental Disorders: DSM-5™*, 5th ed. USA: American Psychiatric Publishing, Inc. doi: 10.1176/appi.books.9780890425596.
- Jones JM, Bennett S, Olmsted MP *et al.* (2001) Disordered eating attitudes and behaviours in teenaged girls: a school-based study. *CMAJ* **165**, 547–552.
- Vilela JE, Lamounier JA, Dellaretti Filho MA *et al.* (2004) Eating disorders in school children. *J Pediatr (Rio J)* **80**, 49–54.
- Preti A, Girolamo G, Vilagut G *et al.* (2009) The epidemiology of eating disorders in six European countries: results of the ESEMeD-WMH project. *J Psychiatr Res* **43**, 1125–1132.
- Musaiger AO, Al-Mannai M, Tayyem R *et al.* (2013) Risk of disordered eating attitudes among adolescents in seven Arab countries by gender and obesity: a cross-cultural study. *Appetite* **60**, 162–167.
- Yu J, Lu M, Tian L *et al.* (2015) Prevalence of disordered eating attitudes among university students in Wuhu, China. *Nutr Hosp* **32**, 1752–1757.
- Babic-Zielińska E (2006) Role of psychological factors in food choice—a review. *Pol J Food Nutr Sci* **15**, 379–384.
- Gianini LM, White MA & Masheb RM (2013) Eating pathology, emotion regulation, and emotional overeating in obese adults with binge eating disorder. *Eat Behav* **14**, 309–313.
- Farrow C, Haycraft E & Meyer C (2011) Similarities between eating attitudes among friendship groups in childhood: the moderating role of child anxiety. *J Pediatr Psychol* **36**, 1144–1152.
- Yau YH & Potenza MN (2013) Stress and eating behaviors. *Minerva Endocrinol* **38**, 255–267.
- Levinson CA & Rodebaugh TL (2016) Clarifying the prospective relationships between social anxiety and eating disorder symptoms and underlying vulnerabilities. *Appetite* **107**, 38–46.
- Barcaccia B, Balestrini V, Salianni AM *et al.* (2018) Dysfunctional eating behaviors, anxiety, and depression in Italian boys and girls: the role of mass media. *Braz J Psychiatry* **40**, 72–77.
- Shroff H, Reba L, Thornton LM *et al.* (2006) Features associated with excessive exercise in women with eating disorders. *Int J Eat Disord* **39**, 454–461.
- Dalle Grave R, Calugi S & Marchesini G (2008) Compulsive exercise to control shape or weight in eating disorders: prevalence, associated features, and treatment outcome. *Compr Psychiatry* **49**, 346–352.
- Taranis L & Meyer C (2011) Associations between specific components of compulsive exercise and eating-disordered cognitions and behaviors among young women. *Int J Eat Disord* **44**, 452–458.
- Li JTE, Pursey KM, Duncan MJ *et al.* (2018) Addictive eating and its relation to physical activity and sleep behavior. *Nutrients* **10**(10), 1428.
- Li J & Powdthavee N (2015) Does more education lead to better health habits? evidence from the school reforms in Australia. *Soc Sci Med* **127**, 83–91.
- Pilecki MW, Kowal M, Woronkiewicz A *et al.* (2014) The socioeconomic status and family context of eating attitudes and dietary behaviours of children in polish primary schools. *Arch Psychiatr Psychother* **1**, 5–13.
- Finger JD, Tylleskär T, Lampert T *et al.* (2013) Dietary behaviour and socioeconomic position: the role of physical activity patterns. *PLoS One* **8**, e78390.
- Garner DM, Olmsted MP, Bohr Y *et al.* (1982) The eating attitudes test: psychometric features and clinical correlates. *Psychol Med* **12**, 871–878.
- Arellano JR, Torres M, Rivera C *et al.* (2009) Abnormal eating attitudes in Mexican female students: a study of prevalence and sociodemographic-clinical associated factors. *Eat Weight Disord* **14**, e42–49.
- dos Santos Alvarenga M, Francischi R, Fontes F *et al.* (2010) Adaptation and validation of the Spanish version of the disordered eating attitude scale (DEAS). *Perspectivas en Nutrición Humana* **12**, 11–23.
- Castro J, Toro J, Salamero M *et al.* (1991) The eating attitudes test: validation of the Spanish version. *Evaluación psicológica* **7**(2), 175–189.
- Dotti A & Lazzari R (1998) Validation and reliability of the Italian EAT-26. *Eat Weight Disord* **3**, 188–194.
- Nasser M (1986) The validity of the eating attitude test in a non-Western population. *Acta Psychiatr Scand* **73**, 109–110.
- al-Subaie A, al-Shammari S, Bamgboye E *et al.* (1996) Validity of the Arabic version of the eating attitude test. *Int J Eat Disord* **20**, 321–324.
- Haddad C, Obeid S, Akel M *et al.* (2019) Correlates of orthorexia nervosa among a representative sample of the lebanese population. *Eat Weight Disord* **24**(3), 481–493. doi: 10.1007/s40519-018-0631-x.
- Haddad C, Zakhour M, Akel M *et al.* (2019) Factors associated with body dissatisfaction among the lebanese population. *Eat Weight Disord* **24**(3), 507–519. doi: 10.1007/s40519-018-00634-z.
- Sfeir E, Haddad C, Salameh P *et al.* (2019) Binge eating, orthorexia nervosa, restrained eating, and quality of life: a population study in lebanon. *Eat Weight Disord*. doi: 10.1007/s40519-019-00831-4.
- Comrey AL & Lee HB (2013) *A First Course in Factor Analysis*. England: Psychology Press.
- Weary-Smith KA (2007) *Validation of the Physical Activity Index (PAI) as a Measure of Total Activity Load and Total Kilocalorie Expenditure During Submaximal Treadmill Walking*. Pennsylvania, USA: University of Pittsburgh.



39. Salafia EHB, Jones ME, Haugen EC *et al.* (2015) Perceptions of the causes of eating disorders: a comparison of individuals with and without eating disorders. *J Eat Disord* **3**, 32.
40. Zullig KJ, Matthews-Ewald MR & Valois RF (2016) Weight perceptions, disordered eating behaviors, and emotional self-efficacy among high school adolescents. *Eating Behav* **21**, 1–6.
41. Bosi ATB, Çamur D & Güler Ç (2007) Prevalence of orthorexia nervosa in resident medical doctors in the faculty of medicine (Ankara, Turkey). *Appetite* **49**, 661–666.
42. Garner DM & Garfinkel PE (1979) The eating attitudes test: an index of the symptoms of anorexia nervosa. *Psychol Med* **9**, 273–279.
43. Garner DM, Olmsted MP, Bohr Y *et al.* (1982) The eating attitudes test: psychometric features and clinical correlates. *Psychol Med* **12**, 871–878.
44. Garner DM (1991) *Eating Disorder Inventory-2: Professional Manual*. Odessa: Psychological Assessment Resources.
45. Cohen S, Kamarck T & Mermelstein R (1983) A global measure of perceived stress. *J Health Soc Behav* **24**, 385–396.
46. Hallit S *et al.*, *Clin Epidemiol Global Health*, doi: 10.1016/j.cegh.2020.03.028.
47. Thompson E (2015) Hamilton rating scale for anxiety (HAM-A). *Occup Med (Lond)* **65**, 601.
48. Obeid S, Hallit CAE, Haddad C *et al.* (2018) Validation of the Hamilton Depression Rating Scale (HDRS) and socio-demographic factors associated with Lebanese depressed patients. *Encephale* **44**(5), 397–402. doi: 10.1016/j.encep.2017.10.010.
49. Hamilton M (1960) A rating scale for depression. *J Neurol Neurosurg Psychiatry* **23**, 56.
50. Gross JJ (2002) Emotion regulation: affective, cognitive, and social consequences. *Psychophysiology* **39**, 281–291.
51. Gross JJ & Levenson RW (1993) Emotional suppression: physiology, self-report, and expressive behavior. *J Pers Soc Psychol* **64**, 970.
52. Gross JJ & John OP (2003) Individual differences in two emotion regulation processes: implications for affect, relationships, and well-being. *J Pers Soc Psychol* **85**, 348.
53. Rahme C, Obeid S, Sacre H *et al.* (2020) Emotional eating among Lebanese adults: scale validation, prevalence and correlates. *Eat Weight Disord*. doi: 10.1007/s40519-020-01001-7.
54. Arnow B, Kenardy J & Agras WS (1995) The Emotional Eating Scale: the development of a measure to assess coping with negative affect by eating. *Int J Eat Disord* **18**, 79–90.
55. Saade S, Hallit S, Haddad C *et al.* (2019) Factors associated with restrained eating and validation of the Arabic version of the restrained eating scale among an adult representative sample of the Lebanese population: a cross-sectional study. *J Eat Disord* **7**, 24.
56. Van Strien T, Frijters JE, Van Staveren WA *et al.* (1986) The predictive validity of the Dutch restrained eating scale. *Int J Eat Disord* **5**, 747–755.
57. Haddad C, Hallit R, Akel M *et al.* (2019) Validation of the Arabic version of the ORTO-15 questionnaire in a sample of the Lebanese population. *Eat Weight Disord* **5**(4), 951–960. doi: 10.1007/s40519-019-00710-y.
58. Donini LM, Marsili D, Graziani MP *et al.* (2004) Orthorexia nervosa: a preliminary study with a proposal for diagnosis and an attempt to measure the dimension of the phenomenon. *Eat Weight Disord* **9**, 151–157.
59. Zeidan RK, Haddad C, Hallit R *et al.* (2019) Validation of the Arabic version of the binge eating scale and correlates of binge eating disorder among a sample of the Lebanese population. *J Eat Disord* **7**, 40.
60. Gormally J, Black S, Daston S *et al.* (1982) The assessment of binge eating severity among obese persons. *Addict Behav* **7**, 47–55.
61. Greeno CG, Marcus MD & Wing RR (1995) Diagnosis of binge eating disorder: discrepancies between a questionnaire and clinical interview. *Int J Eat Disord* **17**, 153–160.
62. Marsh HW, Hau K-T & Wen Z (2004) In search of golden rules: comment on hypothesis-testing approaches to setting cutoff values for fit indexes and dangers in overgeneralizing Hu and Bentler's (1999) findings. *Struct Equat Model* **11**, 320–341.
63. Nunes MA, Barros FC, Anselmo Olinto MT *et al.* (2003) Prevalence of abnormal eating behaviours and inappropriate methods of weight control in young women from Brazil: a population-based study. *Eat Weight Disord* **8**, 100–106.
64. Ballester Ferrando D, De Gracia Blanco M, Patino Maso J *et al.* (2002) Eating attitudes and body satisfaction in adolescents: a prevalence study. *Actas Esp Psiquiatr* **30**, 207–212.
65. Grigg M, Bowman J & Redman S (1996) Disordered eating and unhealthy weight reduction practices among adolescent females. *Prev Med* **25**, 748–756.
66. Alvarez-Rayón G, Mancilla-Díaz J, Vázquez-Arévalo R *et al.* (2004) Validity of the Eating Attitudes Test: a study of Mexican eating disorders patients. *Eat Weight Dis-Stud Anorexia, Bulimia Obes* **9**, 243–248.
67. Schindler A, Mozzanica F, Monzani A *et al.* (2013) Reliability and validity of the Italian Eating Assessment Tool. *Ann Otol Rhinol Laryngol* **122**, 717–724.
68. Jacka FN, Cherbuin N, Anstey KJ *et al.* (2014) Dietary patterns and depressive symptoms over time: examining the relationships with socioeconomic position, health behaviours and cardiovascular risk. *PLoS One* **9**, e87657.
69. Bittencourt SA, Lucena-Santos P, Moraes JF *et al.* (2012) Anxiety and depression symptoms in women with and without binge eating disorder enrolled in weight loss programs. *Trends Psychiatry Psychother* **34**, 87–92.
70. Costarelli V & Patsai A (2012) Academic examination stress increases disordered eating symptomatology in female university students. *Eat Weight Disord-Stud Anorexia, Bulimia Obes* **17**, e164–e169.
71. Cole DA, Peeke LG, Martin JM *et al.* (1998) A longitudinal look at the relation between depression and anxiety in children and adolescents. *J Consult Clin Psychol* **66**, 451–460.
72. Gan W, Nasir MM, Zalilah M *et al.* (2011) Disordered eating behaviors, depression, anxiety and stress among Malaysian university students. *Coll Stud J* **45**, 296–310.
73. Vanwormer JJ, French SA, Pereira MA *et al.* (2008) The impact of regular self-weighing on weight management: a systematic literature review. *Int J Behav Nutr Phys Act* **5**, 54.
74. Tasca GA & Balfour L (2014) Attachment and eating disorders: a review of current research. *Int J Eat Disord* **47**, 710–717.
75. Le J, Dallongeville J, Wagner A *et al.* (2013) Attitudes toward healthy eating: a mediator of the educational level-diet relationship. *Eur J Clin Nutr* **67**, 808–814.
76. Kontinen H, Sarlio-Lähteenkorva S, Silventoinen K *et al.* (2013) Socio-economic disparities in the consumption of vegetables, fruit and energy-dense foods: the role of motive priorities. *Public Health Nutr* **16**, 873–882.