



Review Article

Socio-economic and cultural disparities in diet among adolescents and young adults: a systematic review

Lucille Desbouys^{1,*}, Caroline Méjean², Stefaan De Henauw³ and Katia Castetbon¹

¹Research Center in Epidemiology, Biostatistics and Clinical Trials, School of Public Health, Université libre de Bruxelles (ULB), CP598, Route de Lennik 808, 1070 Brussels, Belgium; ²MOISA, Université de Montpellier, INRA, CIRAD, CIHEAM-IAMM, Montpellier SupAgro, Montpellier, France; ³Department of Public Health, Ghent University, Ghent, Belgium

Submitted 14 September 2018: Final revision received 2 May 2019: Accepted 22 May 2019: First published online 30 August 2019

Abstract

Objective: To explore dietary differences according to socio-economic and sociocultural characteristics of adolescents and young adults.

Design: A systematic review was conducted.

Setting: The main search source was MEDLINE, consulted between January 2012 and March 2017. Quality of selected studies was assessed based on dietary measurement method, sample selection, socio-economic indicator choice and statistical modelling.

Participants: Cross-sectional and longitudinal studies, assessing relationships between socio-economic status and dietary intake (patterns, scores and food groups) in the 10- to 40-year-old general population of high-income countries, were selected.

Results: Among the 7250 reports identified, forty were selected, seventeen of which were of high quality; their conclusions, related only to adolescents, were combined and presented. The most favourable dietary patterns, higher dietary scores, greater consumption of fruits, vegetables and dairy products, and lower consumption of sugary sweetened beverages and energy-dense foods, were associated with better parental socio-economic status, particularly in terms of higher education. Migrant status was associated with plant-based patterns, greater consumption of fruits and vegetables and of sugary sweetened beverages and energy-dense foods. For the other food groups, and for young adults, very few high-quality studies were found.

Conclusions: The socio-economic gradient in adolescent diets requires confirmation by higher-grade studies of a wider set of food groups and must be extended to young adult populations. Future nutritional interventions should involve the most vulnerable adolescent populations, taking account of socio-economic status and migration.

Keywords

Diet
Nutrition
Socio-economic factors
Adolescent
Young adult

Dietary risk was shown to be responsible for more than one-third of deaths worldwide in 2013⁽¹⁾. Nutritional behaviour has thus been targeted by the WHO so as to reduce the current increase in non-communicable diseases⁽²⁾. At each life stage, a balanced, diversified diet is necessary. Adolescence is one of the most crucial stages in life, requiring specific nutrition⁽³⁾. Adolescence and early adulthood correspond to key transition periods for acquisition of health behaviours (e.g. tobacco and alcohol consumption, diet-related habits, physical activity

and sleep, etc.) that otherwise might later provoke non-communicable diseases⁽⁴⁾. Important changes in health behaviour may occur during this period, while previously acquired habits may be strengthened^(5–7).

In Europe and the USA, socio-economic disparities in mortality and morbidity rates, as well as in perceived health, are widening^(8–11). Nutritional issues are also involved^(12–15). A reference literature review focusing on diet disparities concluded that, in adult populations in industrialized countries, a socio-economic gradient existed⁽¹⁶⁾.

*Corresponding author: Email lucille.desbouys@ulb.ac.be



Indeed, consumption of whole grains, fresh fruits and vegetables and low-fat dairy products increased with socio-economic status (SES), while that of less healthy products such as refined grains and added fats decreased. In a more recent expert report comprising a comprehensive literature review on socio-economic diet disparities, conclusions pertaining to adults also tended to converge towards a socio-economic gradient, despite studies heterogeneously available according to food group⁽¹⁷⁾. Only twenty European studies combining children and adolescents were identified. They came to diverging conclusions, mainly based on dietary behaviour (e.g. weekly daily breakfast frequency) rather than on quantitative amounts of food eaten. Other recent reviews involving specific food groups or populations gave scattered information, and included only children^(18–20) or else did not make a distinction between children and adolescents⁽²¹⁾. Overall, maternal education was shown to be a strong determinant of a child's dietary quality⁽¹⁸⁾. Lower parental SES has been related to higher consumption of sugar-sweetened beverages (SSB), while children of married couples or cohabitating parents may have lower SSB consumption⁽¹⁹⁾. Finally, fruit and vegetable consumption by low-income children differed according to their race/ethnicity⁽²¹⁾. For the other food groups, available information was insufficient for drawing evidence-based conclusions. And, to our knowledge, no study has specifically focused on diet disparities in young adults.

Education, employment and income, the three components that generally characterise SES in research, are responsible for major health disparities⁽²²⁾. Although closely related, they are not interchangeable⁽²³⁾, and may even influence pathways leading to health inequalities⁽²²⁾. Moreover, individual characteristics (age, sex, generation, family conditions, etc.) may interact with SES characteristics and should therefore be taken into account so as to better interpret observed gradients⁽¹⁷⁾. Dietary disparities have also been studied via less common indicators, such as place of living, ethnicity and migration background, which were assimilated as socio-economic and cultural indicators^(17,19). In addition, nutrition-related characteristics like BMI and physical activity might also be included in statistical modelling that explores diet disparities, but their role in potential over-adjustment needs clarification. Indeed, inter-relationships between all these indicators require careful interpretation of observed dietary disparities according to SES characteristics.

However, information available on such disparities during adolescence and early adulthood is scattered. Although conclusions have tended to indicate a social gradient for certain food groups, specificities of life-stage disparities have not been thoroughly addressed and their identification could be relevant for developing targeted interventions. To our knowledge, no recent work has systematically updated available information on diet disparities focusing on adolescence and young adulthood,

and oriented towards a wide set of socio-economic factors, including migratory characteristics. The aim of the current systematic review was thus to explore how diet (overall and by food group) differs according to socio-economic and cultural characteristics of adolescents and young adults from high-income countries.

Methods

Search strategy

A systematic review of the literature according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines⁽²⁴⁾ was conducted between December 2016 and March 2017. Targeted studies sought to examine individual diet according to social, economic and cultural characteristics as their primary or secondary objective. The Population, Intervention, Comparison, Outcomes and Study design (PICOS) inclusion and exclusion criteria are presented in Table 1. A relatively large range of ages was targeted (10–40 years) in order to include those studies examining the general population and which analysed subgroups of adolescents and young adults.

In order to follow up previously published reviews, articles published between 1 January 2012 (the endpoint of the most recently updated review⁽¹⁷⁾) and 31 March 2017 were searched for in MEDLINE®. A controlled vocabulary from the Medical Subject Headings (MeSH) was used to build a syntax (see online supplementary material) according to keywords encountered in the articles selected in previous works^(16,17). MeSH keywords relative to diet were: 'Diet', 'Food' (without tree explosion), 'Fruits', 'Vegetables', 'Dairy Products', 'Nutrition surveys', 'Feeding behavior' (without tree explosion), 'Food preferences' and 'Nutrition'. MeSH keywords concerning the social, economic or cultural factors were: 'Socioeconomic factors', 'Risk factors', 'Ethnic groups', 'Family', 'Family characteristics', 'Health status', 'Human migration' and 'Residence characteristics'. Geographic keywords were added: 'Europe', 'Canada', 'United States', 'Australia' and 'New Zealand'. Asia was not included due to specific dietary habits (types of food, dietary patterns). Since recently published articles may not be referenced in MEDLINE according to the MeSH thesaurus, the review was completed by a free search, covering the latest year and using a similar vocabulary. No language restriction was used, so as to obtain a maximum of available information. In fact, no full texts in any language other than English were finally selected. Finally, references cited in literature reviews published on similar topics^(15,18–21,25–32) were searched for via MEDLINE, examined and added to the corpus if relevant.

Selection process

PRISMA guidelines⁽²⁴⁾ were used to present the flow selection process (Fig. 1). Titles were independently screened

**Table 1** PICOS (Population, Intervention, Comparison, Outcomes and Study design) criteria for inclusion and exclusion of studies in the systematic review

Population	General population, 10–40 years, living in Europe, the USA, Canada, Australia or New Zealand Results specifically presented in adolescent (i.e. 10–17-year-old) and/or young adult (i.e. 18–40-year-old) subgroups Excluded: patients, elderly, infants or pre-school children, pregnant or lactating women, overweight or obese persons, those participating in a diet programme, persons with eating disorders, specific ethnic groups (e.g. Inuit), low-income countries or geographic areas such as Asia
Intervention	Not applicable
Comparison	Of participants, their parents, their household: • Socio-economic status: education level, income, occupation, employment status • Family structure: matrimonial status, parenthood, sibling(s), household size • Cultural aspects and migration status: country of origin, language spoken, migration background Excluded: socio-economic status of a geographic area, a school or another non-individual level
Outcome	Diet assessed by usual intake or food frequency, in terms of food groups, food patterns and diet scores Excluded outcomes: energy, macro- and micronutrient intakes, eating behaviour (meal frequency, breakfast skipping, takeaway or fast-food consumption) and diet assessed through biomarkers
Study design	Cross-sectional Longitudinal: description of cohort at baseline or at follow-up point

by two investigators, while abstracts and full texts were read by one investigator. All full texts were available through academic resources, except for two, which were obtained after electronic contact with authors.

Reasons for record exclusion are presented in the PRISMA flowchart (Fig. 1). Among 140 abstracts assessed for eligibility, ninety-five full texts were excluded: sixty-one because results were not specifically presented for adolescents or young adults, but for a broader age range, and fourteen because diet description covered only nutrients or diet behaviour (e.g. fast foods, breakfast frequency, etc.).

Information was extracted according to a previously established reading grid, which included the following items: name of first author, year of publication, study objectives, country/countries or region, data collection period, study design, sampled population (i.e. national, student, etc.), number of participants included in diet analysis according to socio-economic and cultural factors, age range, diet collection method, diet outcome, socio-economic and cultural status variables, and main results concerning associations between diet and socio-economic or cultural status and adjustment variables.

Quality assessment

Appropriate methods and the quality of each included study were assessed using a set of criteria (Fig. 2). First, to verify risk of information bias, diet collection methods were examined: repeated 24 h recalls, FFQ including a sufficient number of food items (i.e. at least several tens of items) and diet records were considered a valid method for food intake data collection^(33,34). Studies based on other types of questionnaires (short FFQ, diet history and single 24 h recall, for example) were considered to be of lesser quality, were not described in detail and were not tabulated.

Risk of selection bias was investigated by examining the sampling method; attention was paid primarily to sample size and scope. When a small sample was studied (fewer than 500 participants), or when only a call for volunteers or convenience sampling was used, the quality of the methods was considered 'low'. Moreover, if the study population was highly specific (e.g. one year of school grade in one city), the study was considered to be of poor quality.

Accuracy of the exposure measurement was then assessed by the relevance of the socio-economic categories chosen (sufficient number of categories making possible a potential gradient, i.e. minimum of three categories, adapted to the population under study) and reliability of the index when such a composite SES was used (e.g. based on both education level and occupation status).

Finally, we focused on analysis modelling, i.e. appropriateness of the final model, and whether potential confounding factors and mediators (i.e. BMI, physical activity, screen time, age, gender, place of living) were identified and accurately integrated into the model. Factors possibly causing confounding results, either concomitant or as mediators in the relationship between SES and diet, are numerous and differently involved depending on the context. Therefore, the objective was to identify potentially over-adjusted models or inappropriate choices of adjustment variables. If no multivariate analysis was found in the article, univariate results were considered, as well as stratification options.

Analysis process

A narrative synthesis, completed by detailed tables, is presented herein. Given the small number and heterogeneity of selected reports, findings concerning young adults (18–40 years old), food groups such as meat, fish and eggs, starchy foods and legumes, water and low-calorie drinks, fat, pulses, nuts and alcoholic drinks, along with disparities according to rural or urban living environment, are not presented.

Results were sorted by type of diet outcome: patterns, diet scores and food groups (vegetables and fruits, dairy foods, SSB, salty and sweet energy-dense foods). For each, socio-economic indicators related to education, occupation,




Fig. 1 Flowchart showing selection of reports included in the systematic review using PRISMA guidelines (PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses; PICOS, Population, Intervention, Comparison, Outcomes and Study design)

income level, migration status and family structure were presented when available. Names of dietary patterns have been quoted as named in the original articles. Only results of high-quality studies have been detailed in summary tables. Those with lower quality have been added as complementary information in the text. In the tables, studies have been arranged in alphabetical order by first author's name.

Results

Among 7250 records identified after removing duplicates, forty met inclusion criteria (Fig. 1). Among the forty selected studies, seventeen were considered of satisfactory quality and have been presented in detail and tabulated. The main reason for lower quality was lack of accuracy concerning diet outcome measurement.




Fig. 2 Criteria used to assess the methodological quality of studies included in the systematic review

Indeed, twenty-two studies of poor quality used a short FFQ, a single 24 h recall or dietary history.

Dietary patterns

In total, six reports corresponding to five studies presented *a posteriori* dietary patterns. Among them, three studies (four reports) were considered of good quality (Table 2) and two of lower quality^(35,36).

Different categories of dietary patterns were identified and considered according to their potential health benefits or disadvantages. Methods used were cluster analyses^(35,37–39) and principal component analyses^(36,40). Pattern content varied according to the context. ‘Healthy’^(37,38), ‘Mediterranean’⁽⁴⁰⁾, ‘vegetarian’⁽³⁹⁾ and ‘dairy product’⁽³⁷⁾ patterns were identified. Such healthy patterns were confronted with less favourable profiles^(37–39). ‘Western’⁽⁴⁰⁾ and ‘traditional’⁽³⁸⁾ pattern compositions strongly depended on the context: they differed from healthier profiles by their high content in meat, potatoes, bread and cereals, and might also include energy-dense and ultra-processed products. Overlaps between healthy and traditional patterns were also described, creating ‘western and Mediterranean’⁽⁴⁰⁾ and ‘traditional/health conscious’⁽³⁹⁾ patterns, with the latter considered as ‘fairly healthy’.

Among the four dietary pattern studies of good quality (Table 2), in three out of three studies examining education level, patterns considered as healthy were associated with higher parental education levels, especially maternal^(37,39), for girls only in one study⁽⁴⁰⁾. In three out of these three studies analysing occupation, healthy patterns were related to higher parental occupation position⁽³⁷⁾, in girls only in one study⁽⁴⁰⁾, and were observed more frequently when the adolescents’ mothers were unemployed, in comparison to working mothers in a third study⁽³⁹⁾. In all these studies, less favourable patterns were associated with lower parental education^(37,39) (in girls only in one study)⁽⁴⁰⁾. The ‘western’ profile was related to a lower parental occupation⁽⁴⁰⁾, and ‘snacks/sugared drinks’ were more frequent among working mothers of adolescents⁽³⁹⁾. Moreover in a fourth study, tracking of healthy or unfavourable patterns at three time points was correlated with higher and lower maternal education level, respectively⁽³⁸⁾. Results were consistent in studies using less accurate diet measurement methods^(35,36), but slightly discordant when the SES index based on parental education, occupation and income was examined in Germany: the ‘western’ pattern was associated with higher parental SES, while the reverse was observed for the ‘traditional and western’ profile⁽³⁶⁾.

Table 2 Dietary patterns according to socio-economic and cultural characteristics of adolescents* (four reports)

First author, year, reference	Population, design, time of collection, country	Age (years)	n	Diet collection method	Exposure variables (number of categories)	Main associations	Adjustments
Araujo (2015) ⁽³⁷⁾	EPITeen study, public and private schools of Porto, cohort at baseline, 2003–2004, Portugal	13	1489	FFQ	<ul style="list-style-type: none"> Parental educ. (in years, 4) Parental educ. (in years, 4) 	<ul style="list-style-type: none"> Higher % of higher educ. in 'Healthier' and 'Dairy products' patterns Higher educ. increase odds to be in 'Healthier' and 'Dairy products' patterns and decrease odds to be in 'Fast-food and sweets' pattern 	<p>None</p> <p>Sex, regular practice of sports, leisure-time activities, TV watching on weekend, fried-food consumption, BMI and mother's BMI</p>
Bibiloni (2012) ⁽⁴⁰⁾	Balearic Islands, cross-sectional, 2007–2008, Spain	12–17	1231	24hR (x2), FFQ	<ul style="list-style-type: none"> Living with both parents (2) Mother's occup. (3) Parental educ. (3) Parental occup. (3) 	<ul style="list-style-type: none"> NS Higher % of white collar in 'Healthier' and 'Dairy products' patterns 'Western' pattern asso. with low educ. (girls) and 'Mediterranean' pattern asso. with medium and high educ. (girls) 'Western' pattern asso. with low occup. and 'Mediterranean' pattern asso. with high occup. (girls) 	<p>None</p> <p>In univariate analysis only. NS when adjusted for age group, number of daily meals and snacks, media screen time, sleep time, physical activity, body composition, desire to change weight and all SES variables</p>
Northstone (2013) ⁽³⁸⁾	ALSPAC study, cohort at follow-up age 7, 10 and 13 years, 1998, 2001 and 2004, UK	7, 10 and 13	6837, 6972 and 5661	Record (3 d)	<ul style="list-style-type: none"> Ethnicity (2) Maternal educ. (3) 	<ul style="list-style-type: none"> Staying in 'processed' cluster asso. with being non-White (v. White) Staying in 'healthy' cluster at 3 time points asso. with higher maternal educ. and staying in 'processed' cluster asso. with lower maternal educ. 	<p>Sex, ethnicity, maternal age, maternal smoking and all SES variables</p>
Northstone (2014) ⁽³⁹⁾	ALSPAC study, cohort at follow-up age 13 years, 2004, UK	13	3951	FFQ	<ul style="list-style-type: none"> Housing tenure (3) Ethnicity (2) Maternal educ. (5) Mother has a partner (2) Mother in employment (2) Older sibling (3) Younger siblings (3) 	<ul style="list-style-type: none"> NS 'Snack/sugared drinks' pattern asso. with being White (v. non-White) and 'vegetarian' pattern asso. with being non-White (v. White) 'Traditional/health conscious' and 'vegetarian' pattern asso. with higher maternal educ. and 'processed' and 'snack/sugared drinks' patterns asso. with lower maternal educ. NS 'Traditional/health conscious' pattern asso. with being unemployed (v. working) and 'snack/sugared drinks' pattern asso. with working (v. being unemployed) 'Processed' and 'snack/sugared drinks' patterns asso. with presence of two or more older or younger siblings 	<p>Sex, maternal age and all SES variables</p>

EPITeen, Epidemiological Health Investigation of Teenagers in Porto; ALSPAC, Avon Longitudinal Study of Parents and Children; 24hR, 24 h recall; educ., education; occup., occupation; asso., associated; TV, television; SES, socio-economic status.

*Details on risk of bias assessment are not presented since only studies of good quality are tabulated.



Ethnicity was explored in the Avon area of the UK at 13-year follow-up⁽³⁹⁾: the 'vegetarian' pattern was associated with being 'non-white' in comparison with the 'white' group in this predominantly white population. On the other hand, the unfavourable 'snacks and sugared drinks' profile was more frequent among white than among non-white adolescents. Nevertheless, non-white adolescents were more likely to remain in the 'processed' pattern when they were tracked over time, according to a second report concerning the same cohort⁽³⁸⁾. Finally, in one study regarding family structure indicators, 'snacks and sugared drinks' and 'processed' patterns were pointed out as being more frequent in families with more siblings⁽³⁹⁾.

Scores

Eleven selected reports, corresponding to ten studies, analysed *a priori* diet scores in adolescent populations. Five studies (six reports) were considered of good quality (Table 3) and five of lower quality, and were not tabulated^(41–45). One study (two reports) of good quality was conducted in low-socio-economic areas^(46,47).

Different types of scores adapted to adolescents were used, measuring the compliance with a nationally recommended diet^(44,48,49) or to a Mediterranean diet^(46,47,50,51). All these scores were calculated from consumed amounts of several predefined food groups, ranging from seven to sixteen groups. The Diet Quality Index for Adolescents was used in one study⁽⁴⁸⁾: in addition to compliance with recommendations, this score takes into account diet diversity, dietary balance and meal frequency.

Among studies of good quality (Table 3), in five out of five studies, the diet score of adolescents was higher when the parental education level was higher^(48,49), especially maternal education^(46,47,51). A similar trend according to parental occupation was observed in two out of three studies^(48,51), while occupation was not significantly associated in the third⁽⁴⁹⁾. In addition, the diet score was higher when the SES index based on parental education and occupation was higher in the only study that explored such an index⁽⁵⁰⁾. The relationship of diet with income was explored in three studies: among students in Greek areas with low SES, adherence to a Mediterranean diet was positively associated with family affluence⁽⁴⁷⁾ and was higher when the father had an income⁽⁴⁶⁾. Household income was not associated with diet score in the third study⁽⁴⁹⁾.

In the high-quality study examining migration among Greek students attending schools from low-SES areas, adherence to a Mediterranean diet was higher if the mother was a native Greek⁽⁴⁶⁾. Similar trends were pointed out in two studies of lesser quality, showing healthier diet when participants were natives compared with migrants⁽⁴¹⁾ and when they were first- or second-generation migrants compared with the third generation⁽⁴⁴⁾.

Food groups

Twenty-six selected reports, corresponding to twenty-two different studies, described adolescent diets using food groups. Eight studies (nine reports) were considered of good quality (Tables 4–7), including five reports that focused on one or several specific food groups^(49,52–55) and four reports that covered almost all main food groups and subgroups^(50,56–58). The other fourteen studies (seventeen reports) were considered of lower quality^(41,59–74).

Fruits and vegetables

The 'vegetable' group was not defined in most reports^(50,54,56–58); in others^(49,55), it was composed of raw, frozen, canned and cooked vegetables. The 'fruit' group composition was less homogeneous: some included 100% fruit juice⁽⁵⁶⁾, all types of fruit juice⁽⁵⁸⁾, dried fruits⁽⁵⁶⁾ or only fresh⁽⁴⁹⁾ or whole fruits⁽⁵⁷⁾, while some did not define composition^(50,54,55). One report showed analyses of grouped fruits and vegetables⁽⁵⁸⁾. Fruit and vegetable consumption was generally higher when SES indicators were more favourable and none of the selected studies showed an inverse association (Table 4).

Four studies of good quality analysed the association between parental education and vegetable intake. In two studies, and after various adjustments, adolescents with more highly educated parents daily consumed more vegetables^(54,55). In one study, vegetable intake did not vary according to parental education level after adjustment for sex, age and energy intake⁽⁵⁶⁾. Nevertheless, in the fourth study, the highest intake category was associated with higher parental education for boys, after adjustment for sex- and age-recommended amounts of vegetables⁽⁴⁹⁾. In addition, these four studies all showed higher fruit intake and daily consumption when parental education was higher^(49,54–56). Moreover, studies of lower quality showed positive associations between parental education and fruit and vegetable consumption frequency^(60,67).

Two studies investigated the association between vegetable intake and household income/wealth, but found no statistical association^(49,56). In one of three studies investigating fruit consumption, daily fruit intake was higher when household income and wealth levels were higher, after adjustment for age, sex and energy intake in one study⁽⁵⁶⁾, whereas in another study⁽⁴⁹⁾ dichotomized fruit intake was not associated with household income after various adjustments. In a third study, total and whole fruit intake was higher when the family income-to-poverty ratio was higher, whereas 100% fruit juice intake was not associated with family income⁽⁵²⁾. In five out of seven lower-quality studies of fruit and vegetable consumption according to the Family Affluence Scale (FAS) or food insecurity, higher daily consumption was associated with higher FAS^(61,62,65,70,71). Another of these studies also showed that adolescents with a decreasing or increasing poverty level over time consumed less fruits and vegetables than adolescents with a stable non-poor trajectory⁽⁵⁹⁾.

Table 3 Diet scores according to socio-economic or cultural characteristics of adolescents* (six reports)

First author, year, reference	Population, design, time of collection, country	Age (years)	<i>n</i>	Diet collection method	Score	Exposure variables (number of categories)	Main associations	Adjustments
Beghin (2014) ⁽⁴⁸⁾	HELENA study, cross-sectional, 2006–2007, eight European countries	12.5–17.5	1768	24hR (× 2)	DQI-AM	<ul style="list-style-type: none"> Parental educ. (3) Parental occup. (3) 	<ul style="list-style-type: none"> Higher score when educ. higher (Northern Europe) Higher score when occup. higher 	Sex, age and energy intake
Finger (2015) ⁽⁴⁹⁾	KiGGs study, cross-sectional, 2003–2006, Germany	11–17	6359	FFQ	HuSKY	<ul style="list-style-type: none"> Parental educ. (3) Parental occup. (3) Household income (tertiles) 	<ul style="list-style-type: none"> Higher score when educ. higher NS NS 	Age, region, leisure time, media use, total energy expenditure, BMI-for-age, perceived weight status and all SES variables
Grosso (2013) ⁽⁵⁰⁾	Secondary schools of Sicily, cross-sectional, 2010–2011, Italy	13–16	1135	FFQ	KIDMED index	<ul style="list-style-type: none"> SES index (parental educ. and occup., 3) 	<ul style="list-style-type: none"> Higher score when SES higher 	Sex, BMI, physical activity and all SES variables
Kastorini (2016) ⁽⁴⁶⁾	DIATROFI, schools in areas of low SES, intervention study at baseline and after intervention, 2012–2013, Greece	3–18	3941	FFQ	KIDMED index	<ul style="list-style-type: none"> Maternal educ. (3) Mother's country of birth (2) Paternal income source (yes/no) 	<ul style="list-style-type: none"> Higher score when maternal educ. higher Higher score when mother born in Greece than in another country Higher score when presence of paternal income 	Age, sex, food insecurity, time of collection (before v. after) and all SES variables
Ozen (2015) ⁽⁵¹⁾	Balearic Islands, cross-sectional, 2007–2008, Spain	12–17	1691	24hR (× 2), FFQ	Mediterranean diet score	<ul style="list-style-type: none"> Parental educ. (3) Parental occup. (3) 	<ul style="list-style-type: none"> Low adherence to score asso. with low maternal educ. (v. high) among non-functional food consumers Low adherence to score asso. with medium paternal work status (v. high) among non-functional food consumers 	Age, sex, BMI, physical activity, chronic diseases and all SES variables
Yannakoulia (2016) ⁽⁴⁷⁾	DIATROFI, schools in areas of low SES, cross-sectional, 2012–2013, Greece	3–18	11 717	FFQ	KIDMED index	<ul style="list-style-type: none"> Parental educ. (3) FAS (3) 	<ul style="list-style-type: none"> Higher score when educ. higher Higher score when maternal educ. higher Higher score when FAS higher 	Age, sex, sedentary and sports activities and all SES variables

HELENA, Healthy Lifestyle in Europe by Nutrition in Adolescence; KiGGs, German Health Interview and Examination Survey for Children and Adolescents; 24hR, 24 h recall; DQI-AM, Diet Quality Index for Adolescents; HuSKY, Healthy Nutrition Score for Children and Youth; KIDMED, Mediterranean Diet Quality Index for Children and Adolescents; educ., education; occup., occupation; SES, socio-economic status; FAS, Family Affluence Scale; asso., associated.

*Details on risk of bias assessment are not presented since only studies of good quality are tabulated.


Table 4 Vegetable and fruit consumptions according to socio-economic or cultural characteristics of adolescents* (eight reports)

First author, year, reference	Population, design, time of collection, country	Age (years)	n	Diet collection method	Intake or frequency of consumption	Exposure variables (number of categories)	Association	Adjustments
Drewnowski (2015) ⁽⁵²⁾	NHANES, repeated cross-sectional, 2007–2010, USA	14–19	1834	24hR (× 2)	• % of population having total fruit intake < 1.5 cup-equiv./d	• Race/ethnicity (4)	• Lower % of non-Hispanic-Black and 'other Hispanic' than non-Hispanic-White	None
					• Whole fruit intake (cup-equiv./d)	• Family PIR (3)	• Higher % of population when income lower	None
Drouillet-Pinard (2017) ⁽⁵⁶⁾	INCA2 study, cross-sectional, 2006–2007, France	11–17	881	Record (7 d)	• 100 % fruit juice intake (cup-equiv./d)	• Family PIR (3)	• Higher when income higher	None and for sex and race/ethnicity
					• Vegetable intake (g/d)	• Family PIR (3)	• NS	None and for sex and race/ethnicity
Finger (2015) ⁽⁴⁹⁾	KiGGs study, cross-sectional, 2003–2006, Germany	11–17	6359	FFQ	• Fruit intake (g/d)	• Parental occup. (4)	• NS	Age, sex and energy intake
					• Vegetable high or low intake (ratio of g/d intake divided by age- and sex-recommended amount)	• Parental educ. (3)	• NS	
Grosso (2013) ⁽⁵⁰⁾	Secondary schools of Sicily, cross-sectional, 2010–2011, Italy	13–16	1135	FFQ	• Fruit intake (g/d)	• Household income (tertiles)	• NS	Age, sex, BMI, physical activity, place of living and SES
					• Vegetable intake (g/d)	• Parental educ. (3)	• NS	

Disparities in diet among adolescents

Table 4 (Continued)

First author, year, reference	Population, design, time of collection, country	Age (years)	n	Diet collection method	Intake or frequency of consumption	Exposure variables (number of categories)	Association	Adjustments
Grosso (2013) ⁽⁵⁴⁾	Secondary schools of Sicily, cross-sectional, 2010–2011, Italy	13–16	1135	FFQ	<ul style="list-style-type: none"> Vegetable daily consumption Fruit daily consumption 	<ul style="list-style-type: none"> Parental educ. (3) Parental occup. (3) Child educ. (2) Parental educ. (3) Parental occup. (3) 	<ul style="list-style-type: none"> Higher when educ. higher Higher when skilled professions NS Higher when educ. higher Higher when skilled professions 	Age, sex, BMI, daily eat between meals, weekly breakfast, lunch and dinner with parents, influences on food choice and all SES variables
Harris (2015) ⁽⁵⁷⁾	GINIplus study, cohort at follow-up age 10 and 15 years, 2005–2008 and 2010–2013, Germany	10–15	1232	FFQ	<ul style="list-style-type: none"> Change v. tracking of vegetable intake over time (% of energy intake) Change v. tracking of fruit intake over time (% of energy intake) 	<ul style="list-style-type: none"> Child educ. (2) Parental educ. (2) Family income (tertiles) Parental educ. (2) Family income (tertiles) 	<ul style="list-style-type: none"> NS NS NS NS NS 	Age at baseline, baseline energy intake, diet changes, study centre, study intervention arm, pubertal onset, BMI, screen time and all SES variables
Lehto (2015) ⁽⁵⁵⁾	PROGREENS study, cross-sectional, 2009, ten European countries	11	479 to 1218	FFQ	<ul style="list-style-type: none"> Vegetable daily consumption Fruit daily consumption 	<ul style="list-style-type: none"> Parental educ. (2) Parental educ. (2) 	<ul style="list-style-type: none"> Higher when educ. higher (FI, DE, GR, IS, NO, PT, SL) Higher when educ. higher (BG, GR, IS, NO, PT) 	Sex and age
Llull (2015) ⁽⁵⁸⁾	Balearic Islands, cross-sectional, 2007–2008, Spain	12–17	1231	FFQ	<ul style="list-style-type: none"> Vegetable daily consumption Fruit daily consumption Fruit and vegetable daily consumption 	<ul style="list-style-type: none"> Birthplace (4) Birthplace (2) Length of time living in Balearic Islands (4) Birthplace (4) Birthplace (2) Length of time living in Balearic Islands (4) Birthplace (3) Birthplace (2) 	<ul style="list-style-type: none"> Higher for Latin America than Balearic Islands Higher for non-Mediterranean Lower when length of time higher Higher for Latin America than Balearic Islands NS NS Higher for Latin America than Balearic Islands Higher for non-Mediterranean 	Sex and age

NHANES, National Health and Nutrition Examination Survey; INCA2, second French national cross-sectional dietary survey; KiGGS, German Health Interview and Examination Survey for Children and Adolescents; 24hR, 24 h recall; equiv., equivalents; PIR, poverty income ratio; occup., occupation; educ., education; SES, socio-economic status; FI, Finland; DE, Germany; GR, Greece; IS, Iceland; NO, Norway; PT, Portugal; SL, Slovenia; BG, Bulgaria.
 *Details on risk of bias assessment are not presented since only studies of good quality are tabulated.



Table 5 Dairy food consumption according to socio-economic or cultural characteristics of adolescents* (five reports)

First author, year, reference	Population, design, time of collection, country	Age (years)	n	Diet collection method	Intake or frequency of consumption	Exposure variables (number of categories)	Association	Adjustments
Drouillet-Pinard (2017) ⁽⁵⁶⁾	INCA2 study, cross-sectional, 2006–2007, France	11–17	881	Record (7 d)	Milk, yoghurt and cheese intakes (g/d)	<ul style="list-style-type: none"> • Parental occup. (4) • Parental educ. (3) • Household income (tertiles) • Household wealth index (tertiles) • Global SES index (all SES indicators combined, tertiles) 	<ul style="list-style-type: none"> • NS • Yoghurt intake higher when educ., income, wealth and SES higher 	Age, gender and energy intake
Gopinath (2014) ⁽⁵³⁾	Sydney Childhood Eye study, cohort at baseline (age 12 years) and at follow-up age 17 years, 2004–2005 and 2009–2011, Australia	12–17	634	FFQ	Intake ≥ 3.5 servings/d five years later and maintaining consumption above the median over time	<ul style="list-style-type: none"> • Parental educ. at baseline • Ethnicity • Parental occup. 	<ul style="list-style-type: none"> • Higher for tertiary qualifications • NS • NS 	None
Grosso (2013) ⁽⁵⁰⁾	Secondary schools of Sicily, cross-sectional, 2010–2011, Italy	13–16	1135	FFQ	Intake (g/d)	<ul style="list-style-type: none"> • SES index (parental educ. and occup., 3) 	<ul style="list-style-type: none"> • NS 	Age, gender, BMI, physical activity, place of living and SES
Harris (2015) ⁽⁵⁷⁾	GINIplus study, cohort at follow-up age 10 and 15 years, 2005–2008 and 2010–2013, Germany	10–15	1232	FFQ	Change v. tracking of intake over time (% of energy intake)	<ul style="list-style-type: none"> • Parental educ. (2) • Family income (tertiles) 	<ul style="list-style-type: none"> • NS • NS 	Age at baseline, baseline energy intake, diet changes, study centre, study intervention arm, pubertal onset, BMI, screen time and all SES variables
Llull (2015) ⁽⁵⁸⁾	Balearic Islands, cross-sectional, 2007–2008, Spain	12–17	1231	FFQ	Daily consumption	<ul style="list-style-type: none"> • Birthplace (4) • Birthplace (2) • Length of time living in Balearic Islands (4) 	<ul style="list-style-type: none"> • NS • NS • NS 	Gender and age

INCA2, second French national cross-sectional dietary survey; occup., occupation; educ., education; SES, socio-economic status.

*Details on risk of bias assessment are not presented since only studies of good quality are tabulated.

Table 6 Sugar-sweetened beverage consumption according to socio-economic or cultural characteristics of adolescents* (four reports)

First author, year, reference	Population, design, time of collection, country	Age (years)	n	Diet collection method	Intake or frequency of consumption	Exposure variables (number of categories)	Association	Adjustments
Drouillet-Pinard (2017) ⁽⁵⁶⁾	INCA2 study, cross-sectional, 2006–2007, France	11–17	881	Record (7 d)	Intake (g/d)	<ul style="list-style-type: none"> • Parental occup. (4) • Parental educ. (3) • Household income (tertiles) • Household wealth index (tertiles) • Global SES index (all SES indicators combined, tertiles) 	<ul style="list-style-type: none"> • NS • Higher when educ. lower • NS • Higher when wealth lower • Higher when SES lower 	Age, gender and energy intake
Grosso (2013) ⁽⁵⁰⁾	Secondary schools of Sicily, cross-sectional, 2010–2011, Italy	13–16	1135	FFQ	Intake (g/d)	<ul style="list-style-type: none"> • SES index (parental educ. and occup., 3) 	<ul style="list-style-type: none"> • Higher when SES lower 	Age, gender, BMI, physical activity, place of living and SES
Harris (2015) ⁽⁵⁷⁾	GINIplus study, cohort at follow-up age 10 and 15 years, 2005–2008 and 2010–2013, Germany	10–15	1232	FFQ	Change v. tracking of intake over time (% of energy intake)	<ul style="list-style-type: none"> • Parental educ. (2) • Family income (tertiles) 	<ul style="list-style-type: none"> • NS • NS 	Age at baseline, baseline energy intake, diet changes, study centre, study intervention arm, pubertal onset, BMI, screen time and all SES variables
Llull (2015) ⁽⁵⁸⁾	Balearic Islands, cross-sectional, 2007–2008, Spain	12–17	1231	FFQ	Daily consumption	<ul style="list-style-type: none"> • Birthplace (4) • Birthplace (2) • Length of time living in Balearic Islands (4) 	<ul style="list-style-type: none"> • Higher for Latin America and other countries than Balearic Islands • Higher for non-Mediterranean • Higher when length of time lower 	Gender and age

INCA2, second French national cross-sectional dietary survey; occup., occupation; educ., education; SES, socio-economic status.

*Details on risk of bias assessment are not presented since only studies of good quality are tabulated.



Table 7 Salty and sweet energy-dense food consumption according to socio-economic or cultural characteristics of adolescents* (five reports)

First author, year, reference	Population, design, time of collection, country	Age (years)	n	Diet collection method	Intake or frequency of consumption	Exposure variables (number of categories)	Association	Adjustments
Drouillet-Pinard (2017) ⁽⁵⁶⁾	INCA2 study, cross-sectional, 2006–2007, France	11–17	881	Record (7 d)	Stewed fruit/fruit in syrup, dairy desserts, cakes and pastries, confectionery, pizza and sandwich intakes (g/d)	<ul style="list-style-type: none"> • Parental occup. (4) • Parental educ. (3) • Household income (tertiles) • Household wealth index (tertiles) • Global SES index (all SES indicators combined, tertiles) 	<ul style="list-style-type: none"> • Dairy desserts higher and cakes and pastries lower when occup. lower • Dairy desserts higher when educ. lower • NS • NS • Cakes and pastries lower when SES lower 	Age, gender and energy intake
Finger (2015) ⁽⁴⁹⁾	KiGGs study, cross-sectional, 2003–2006, Germany	11–17	6359	FFQ	High or low energy-dense food intake (ratio of g/d intake divided by age- and sex-recommended amount)	<ul style="list-style-type: none"> • Parental educ. (3) • Parental occup. (3) • Household income (tertiles) 	<ul style="list-style-type: none"> • Higher when educ. lower • Higher when occup. lower • Higher when income lower (boys) 	Age, region, media use, total energy expenditure, familial leisure activity, BMI-for-age, perceived weight status and all SES variables
Grosso (2013) ⁽⁵⁰⁾	Secondary schools of Sicily, cross-sectional, 2010–2011, Italy	13–16	1135	FFQ	Intakes of fast foods, snacks and sweets (g/d)	<ul style="list-style-type: none"> • SES index (parental educ. and occup., 3) 	<ul style="list-style-type: none"> • NS 	Age, gender, BMI, physical activity, place of living and SES
Harris (2015) ⁽⁵⁷⁾	GINIplus study, cohort at follow-up age 10 and 15 years, 2005–2008 and 2010–2013, Germany	10–15	1232	FFQ	Sugar-sweetened food intake: change v. tracking over time (% of energy intake)	<ul style="list-style-type: none"> • Parental educ. (2) • Family income (tertiles) 	<ul style="list-style-type: none"> • NS • NS 	Age at baseline, baseline energy intake, diet changes, study centre, study intervention arm, pubertal onset, BMI, screen time and all SES variables
Llull (2015) ⁽⁵⁸⁾	Balearic Islands, cross-sectional, 2007–2008, Spain	12–17	1231	FFQ	Sweets and pastries daily consumption	<ul style="list-style-type: none"> • Birthplace (4) • Birthplace (2) • Length of time living in Balearic Islands (4) 	<ul style="list-style-type: none"> • Sweets higher for Latin America than Balearic Islands • Sweets higher for non-Mediterranean • Higher when length of time lower 	Gender and age

INCA2, second French national cross-sectional dietary survey; KiGGs, German Health Interview and Examination Survey for Children and Adolescents; occup., occupation; educ., education; SES, socio-economic status.

*Details on risk of bias assessment are not presented since only studies of good quality are tabulated.

Vegetable intake was not associated with parental occupation in two studies^(49,56), but in one of these⁽⁵⁶⁾, fruit intake was higher when parental occupational status was higher. Higher daily consumption of vegetables and fruits was associated with parental skilled professions, after various adjustments⁽⁵⁴⁾. Moreover, fruit intake was higher when the global SES index was higher in one study⁽⁵⁶⁾, while it was not associated in another⁽⁵⁰⁾. Vegetable intake was not associated with the overall SES level in these two studies.

Nor was there an association of tracking or change in vegetable and fruit intake over time with parental education or family income in the only study that examined this aspect⁽⁵⁷⁾.

For sociocultural characteristics, fruit and vegetable consumption differed according to birthplace⁽⁵⁸⁾ and ethnic origins⁽⁵²⁾ highly specific to each study context. The first study showed that fruit and vegetable consumption was generally higher for migrants from distant countries and more recent migrants than for natives⁽⁵⁸⁾. In a US sample in the second study, a lower proportion of non-Hispanic Blacks and 'other Hispanics' daily consumed smaller amounts of total fruits than non-Hispanic Whites⁽⁵²⁾. In three out of four studies of lower quality, consumption of fruits and vegetables also differed according to migration status⁽⁴¹⁾ and ethnic origin^(63,72).

Dairy

Most reports defined the 'dairy' group as being composed of milk, yoghurt and cheese^(50,57,58). Some reports also included dairy drinks⁽⁵⁶⁾, flavoured milk, smoothies and milkshakes⁽⁵³⁾ in this group. Some studies indicated higher dairy intake associated with more favourable SES, but overall findings were not consistent (Table 5). Among three studies, one showed that yoghurt intake was higher when parental education, income, wealth and overall SES index were higher, after adjusting for age, sex and energy intake⁽⁵⁶⁾. However, in that study, the studied dairy product intake was never associated with parental occupation, and milk and cheese consumption were not associated with any SES indicator. In the other two studies, dairy intake was higher when parents had tertiary qualifications, but was not associated with occupation⁽⁵³⁾ or SES index (parental occupation and education levels)⁽⁵⁰⁾. Neither changing nor tracking dairy intake over time was associated with parental education or income in the only study concerned⁽⁵⁷⁾.

Neither of two studies examining the association between dairy consumption and ethnicity showed a significant association^(53,58). Among two studies of lesser quality, one described higher consumption of dairy products for breakfast among Spanish adolescents than among other nationalities⁽⁴¹⁾. The other described a proportion of adolescents consuming whole or skimmed milk that differed according to ethnicity, with fewer non-Hispanic Blacks consuming such dairy products⁽⁷²⁾.

Sugar-sweetened beverages

The SSB group was defined throughout the reports as sugary, soft and diet drinks^(50,56,58). In one study, it was also composed of fruit and vegetable juices⁽⁵⁷⁾. SSB drinking, explored in two studies, was higher when parental education^(50,56), household wealth⁽⁵⁶⁾ and global SES⁽⁵⁶⁾ were lower, after various adjustments (Table 6). However, SSB intake was not associated with parental occupation or household income⁽⁵⁶⁾. Four out of five studies of lower quality were rather consistent with each other, showing more frequent SSB consumption when parental education⁽⁶⁰⁾ and FAS⁽⁶²⁾ were lower and when poverty level indicators were higher.

One study carried out in the Balearic Islands explored diet according to birthplace. SSB consumption was higher for adolescents born in Latin America and other foreign countries than for natives, and also higher for those of non-Mediterranean than of Mediterranean origin⁽⁵⁸⁾. Moreover, it was higher when the length of time living in the Balearic Islands was lower. Three out of four lower-quality studies showed significant differences between ethnic groups^(68,69,74).

Change in or tracking of SSB intake over time was not associated with parental education or family income⁽⁵⁷⁾. A lower-quality study of SSB intake decline over time reported differences according to ethnicity, but this was not statistically tested⁽⁷³⁾.

Salty and sweet energy-dense foods

In this group, studies included informal meals generally composed of fatty, salty and sweet snacks and fast foods, without defining a threshold of energy density. Other studies also included soft drinks⁽⁴⁹⁾ or stewed fruits and fruits in syrup⁽⁵⁶⁾. One study focused only on sweet and fatty snacks⁽⁵⁷⁾, and another on sweets and pastries⁽⁵⁸⁾. Amounts of energy-dense foods consumed by adolescents were globally higher when socio-economic characteristics were less favourable, but such findings were not systematically retrieved (Table 7). Two studies out of three showed higher intake when parental education^(49,56), occupation^(49,56) and household income and wealth^(49,56) were lower, after various adjustments. However, an exception was seen: cake and pastry intake was lower when occupational status and global SES were lower⁽⁵⁶⁾. Intakes of stewed fruits, fruits in syrup, confectionery, pizza, sandwiches, fast foods and sweets were otherwise not associated with SES-related indicators^(50,56). Studies of lower quality mainly showed higher consumption of sweets when FAS^(62,70) and parental education⁽⁶⁷⁾ were lower, and higher daily consumption of energy-dense and nutrient-poor snacks when SES was lower, but such associations were not statistically tested⁽⁷⁴⁾.

Only one study in the Balearic Islands explored the associations according to birthplace. Latin American and, more generally, non-Mediterranean adolescents had higher sweets consumption than natives, and sweets and pastry



consumption was higher when the length of time living on the Balearic Islands was lower⁽⁵⁸⁾. One study of lesser quality showed higher sweets and fast-food consumption among adolescents of nationalities other than Spanish⁽⁴¹⁾.

Change in or tracking of sugar-sweetened food intake over time was not associated with education or income⁽⁵⁷⁾. A decline in sweet and salty snack intake over time was observed among Black adolescents with a healthy weight, but ethnic differences were not tested in that study⁽⁷³⁾.

Discussion

Our objective was to update overall knowledge of socio-economic and cultural disparities in dietary patterns, scores and food group consumption by adolescents and young adults. Recent literature on diet disparities has been abundant, but when focusing on this life period, the quality of the studies appears highly variable and available information is scattered. Among adolescents, however, evidence and consistent findings were sufficient to conclude that higher dietary scores and healthier patterns were associated with higher parental education and occupational status, while less favourable patterns were associated with lower SES. Such findings therefore confirmed, at least in part, that a favourable social status is generally associated with a healthier diet.

Regarding food groups, the most substantial bibliographic corpus concerned fruits and vegetables. Such consumption was consistently associated with higher education. In addition, fruit consumption was somewhat higher when household income and wealth were higher. Despite a smaller number of conclusive high-quality studies, SSB, energy-dense food and dairy product consumption were globally associated with SES: SSB and energy-dense food consumption were higher when SES indicators were less favourable, while dairy intake tended to be higher when SES was more favourable. However, available information regarding other groups was very scarce. In addition to SES-related indicators, ethnic and migration disparities were pointed out in several studies but proved to be highly specific to each country and geographic area.

Overall, conclusions are limited due to the heterogeneity of the populations, diet outcome and socio-economic and cultural indicators in question. Thus, it would not have been feasible to carry out a meta-analysis, nor to explore potential publication bias. Moreover, use of a quantitative scale assessing methodological quality would have been too restrictive. Nevertheless, quality assessment was used, making possible a selection based on objective criteria adapted to the diversity of the publications.

Diversity of methods

Most studies using scores were adapted to recommendations dedicated to adolescents, leading to conclusions that could be compared. Since they depended on the population and context of the study, dietary patterns differed from

one study to another, making findings between some countries not directly comparable. However, consistent conclusions were generally drawn. The advantage of describing disparities according to food groups lies in being able to identify specific associated indicators. Food choice and consumption mechanisms may also differ according to food group⁽¹⁶⁾. The issue is to consequently adapt dietary recommendations based on such findings. However, diet collection and description methods differed across studies and thus, for some food groups, it was difficult to draw conclusions.

Statistical models and adjustments were highly variable between studies. Adjustments for sex, age and total energy intake (scores, food groups) enabled taking account of differences in requirements. BMI was sometimes used as an adjustment variable, limiting interpretation, since it may be both a consequence of an unhealthy diet and a reason for adopting a balanced diet. Some authors also chose to adjust for other nutrition-related behaviour (e.g. physical activity, screen time) in order to identify potential confounders, and thus over-adjustment was probable. In some models, identification of the true role of adjustment variables was challenging. Nutrition-related behaviour variables may have been mediators, logically weakening the association between SES and diet. Some adjustment variables were also presented as confounding factors; however, although they were influenced by SES, they could not substitute for that variable in the relationship with diet.

Mechanisms of disparities

Dietary disparities among adolescents overwhelmingly involved inequalities in parental education, particularly maternal. Education is linked to health literacy, i.e. the ability to appropriate health and nutrition information and to generate dietary behaviour that would provide long-term benefits⁽⁷⁵⁾. However, occupation and income were not systematically associated with diet. Income is directly related to financial accessibility to food, and occupation may influence food intake partly via the workplace culture and social networks⁽⁷⁶⁾. Moreover, it has been clearly established that education is a determinant of occupation and income, and that these three indicators are involved in diet disparities, but differently, according to the SES indicator^(16,77,78). In addition, reliability and availability of some SES indicators were insufficient to draw clear conclusions. Nevertheless, the present review shows that parental education was a more systematic determinant of diet than occupation or income. In terms of public health policies, it again emphasizes that nutritional information should be adapted to different education backgrounds and integrated into early education, targeting mothers or caregivers.

According to the food group in question: (i) either all socio-economic indicators were associated (e.g. fruits); (ii) only some SES characteristics were associated (e.g. vegetables, dairy, SSB); or (iii) these associations were contradictory

(e.g. in the case of energy-dense foods). Such disparities within a food group have been described previously; authors have suggested exploring causal mechanisms involved, such as biological (possibly related to higher palatability and lower satiety provided by such energy-dense foods) or behavioural components (accessibility and affordability)⁽¹⁶⁾. For instance, SSB consumption was determined by lower parental financial status (along with less schooling). Indeed, SSB are financially and physically accessible products, often associated with positive values through sports marketing, on the one hand, and time spent in front of screens and sedentary behaviour, on the other⁽¹⁹⁾.

In addition to the main SES indicators, ethnicity and migration status were often associated with diet, but findings appeared to be related to the general background. In some studies carried out in the USA^(52,72), Australia⁽⁵³⁾ and the UK^(38,39), ethnicity was explored mainly as a reflection of SES. In other Mediterranean^(46,58), American⁽⁴⁴⁾ and Canadian⁽⁷¹⁾ studies, parental place of birth, migratory generation and length of time living in the host country were studied. The migration background was thus also explored under the angle of dietary habit acquisition and acculturation^(79–81). For instance, in the general adolescent Balearic Islands population, it was difficult to distinguish effects related to acculturation from those related to SES, since SES indicators were not examined⁽⁵⁸⁾. Nevertheless, higher consumption of SSB and energy-dense foods in recent adolescent migrants may be due to their increased accessibility; furthermore, higher vegetable consumption may be related to culture-specific dietary habits. In addition, variations according to country of origin and stage of nutritional transition should be taken into account.

Conclusions

Based on the present review, findings on dietary patterns and scores, along with fruit and vegetable consumption in adolescents, consistently confirmed the socio-economic gradient observed in adults. However, overall conclusions were much more limited for several food groups and warrant further examination. In addition, high-quality studies remain necessary, especially in terms of reliable dietary and socio-economic evaluations. Sampling of both the general adolescent population and potentially at-risk subgroups such as migrants should also be more carefully examined. Finally, diet in young adults has thus far been poorly described and needs to be concomitantly evaluated so as to improve our understanding of changes in socio-economic and cultural disparities during this transition period.

Nevertheless, the present review, consistent with wide dietary disparities among adolescents, underlines the importance of developing interventions targeted to this age group. Future public health programmes must take the socio-economic gap into account, addressing

nutritional intervention towards both populations as a whole, with the most vulnerable being the adolescent population. Indeed, such initiatives should seek to improve literacy by involving caregivers and taking account of the migration background and associated food culture. Although its long-term sustainability requires confirmation, an improvement in dietary habits during adolescence may continue into adulthood and could contribute to a reduction in non-communicable disease inequalities.

Acknowledgements

Acknowledgements: The authors thank Valérie Durieux for her technical guidance on the literature search and Jerri Bram for English editing. *Financial support:* This work was supported by the French Community of Belgium, as part of the 'Actions de Recherche Concertée' funding programme. The French Community of Belgium had no role in the design, analysis or writing of this article. *Conflict of interest:* None. *Authorship:* L.D. and K.C. formulated the research question and designed the study; L.D. carried out the research; L.D. and K.C. analysed the data; L.D. wrote the paper; K.C., C.M. and S.D.H. edited the article. *Ethics of human subject participation:* Not applicable.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1368980019002362>.

References

1. GBD 2013 Risk Factors Collaborators (2015) Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* **386**, 2287–2323.
2. World Health Organization (2014) Global status report on noncommunicable diseases. <http://www.who.int/nmh/publications/ncd-status-report-2014/en/> (accessed January 2018).
3. Das JK, Salam RA, Thornburg KL *et al.* (2017) Nutrition in adolescents: physiology, metabolism, and nutritional needs. *Ann N Y Acad Sci* **1393**, 21–33.
4. Sawyer SM, Afifi RA, Bearinger LH *et al.* (2012) Adolescence: a foundation for future health. *Lancet* **379**, 1630–1640.
5. Winpenny EM, Penney TL, Corder K *et al.* (2017) Changes in consumption of added sugars from age 13 to 30 years: a systematic review and meta-analysis of longitudinal studies. *Obes Rev* **18**, 1336–1349.
6. Craigie AM, Lake AA, Kelly SA *et al.* (2011) Tracking of obesity-related behaviours from childhood to adulthood: a systematic review. *Maturitas* **70**, 266–284.
7. Nelson MC, Story M, Larson NI *et al.* (2008) Emerging adulthood and college-aged youth: an overlooked age for weight-related behavior change. *Obesity (Silver Spring)* **16**, 2205–2211.



8. Gallo V, Mackenbach JP, Ezzati M *et al.* (2012) Social inequalities and mortality in Europe – results from a large multinational cohort. *PLoS One* **7**, e39013.
9. Mackenbach JP, Stirbu I, Roskam AJ *et al.* (2008) Socioeconomic inequalities in health in 22 European countries. *New Engl J Med* **359**, e14.
10. Centers for Disease Control and Prevention (2013) CDC health disparities & inequalities report – U.S. 2013: some key factors that affect health and lead to health disparities in the United States. http://stacks.cdc.gov/view/cdc/20865/cdc_20865_DS1.pdf (accessed January 2018).
11. Sommer I, Griebler U, Mahlknecht P *et al.* (2015) Socioeconomic inequalities in non-communicable diseases and their risk factors: an overview of systematic reviews. *BMC Public Health* **15**, 914.
12. Wu S, Ding Y, Wu F *et al.* (2015) Socio-economic position as an intervention against overweight and obesity in children: a systematic review and meta-analysis. *Sci Rep* **5**, 11354.
13. Barriuso L, Miqueleiz E, Albaladejo R *et al.* (2015) Socioeconomic position and childhood-adolescent weight status in rich countries: a systematic review, 1990–2013. *BMC Pediatr* **15**, 129.
14. El-Sayed AM, Scarborough P & Galea S (2012) Socioeconomic inequalities in childhood obesity in the United Kingdom: a systematic review of the literature. *Obes Facts* **5**, 671–692.
15. Chung A, Backholer K, Wong E *et al.* (2016) Trends in child and adolescent obesity prevalence in economically advanced countries according to socioeconomic position: a systematic review. *Obes Rev* **17**, 276–295.
16. Darmon N & Drewnowski A (2008) Does social class predict diet quality? *Am J Clin Nutr* **87**, 1107–1117.
17. Expertise Collective (2014) Inégalités sociales de santé en lien avec l'alimentation et l'activité physique. In *Position socioéconomique et alimentation*, pp. 131–187. Paris: Les éditions Inserm.
18. Emmett PM, Jones LR & Northstone K (2015) Dietary patterns in the Avon Longitudinal Study of Parents and Children. *Nutr Rev* **73**, 207–230.
19. Mazarello Paes V, Hesketh K, O'Malley C *et al.* (2015) Determinants of sugar-sweetened beverage consumption in young children: a systematic review. *Obes Rev* **16**, 903–913.
20. Inskip H, Baird J, Barker M *et al.* (2014) Influences on adherence to diet and physical activity recommendations in women and children: insights from six European studies. *Ann Nutr Metab* **64**, 332–339.
21. Di Noia J & Byrd-Bredbenner C (2014) Determinants of fruit and vegetable intake in low-income children and adolescents. *Nutr Rev* **72**, 575–590.
22. Glymour MM, Avendano M & Kawachi I (2014) Social network epidemiology. In *Socioeconomic Status and Health*, 2nd ed., pp. 17–62 [LF Berkman, I Kawachi and MM Glymour, editors]. Oxford: Oxford University Press.
23. Galobardes B, Lynch J & Smith GD (2007) Measuring socioeconomic position in health research. *Br Med Bull* **81–82**, 21–37.
24. Moher D, Liberati A, Tetzlaff J *et al.* (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* **6**, e1000097.
25. Fekete C & Weyers S (2016) Social inequalities in nutrition: evidence, causes and interventions. *Bundesgesundheitsbl* **59**, 197–205.
26. Johnson BJ, Hendrie GA & Golley RK (2016) Reducing discretionary food and beverage intake in early childhood: a systematic review within an ecological framework. *Public Health Nutr* **19**, 1684–1695.
27. Kamphuis CBM, Jansen T, Mackenbach JP *et al.* (2015) Bourdieu's cultural capital in relation to food choices: a systematic review of cultural capital indicators and an empirical proof of concept. *PLoS One* **10**, e0130695.
28. McGill R, Anwar E, Orton L *et al.* (2015) Are interventions to promote healthy eating equally effective for all? Systematic review of socioeconomic inequalities in impact. *BMC Public Health* **15**, 457.
29. deMestral C, Mayen A-L, Petrovic D *et al.* (2017) Socioeconomic determinants of sodium intake in adult populations of high-income countries: a systematic review and meta-analysis. *Am J Public Health* **107**, e1–e12.
30. Novakovic R, Cavelaars A, Geelen A *et al.* (2014) Socioeconomic determinants of micronutrient intake and status in Europe: a systematic review. *Public Health Nutr* **17**, 1031–1045.
31. McCormack LA & Meendering J (2016) Diet and physical activity in rural vs urban children and adolescents in the United States: a narrative review. *J Acad Nutr Diet* **116**, 467–480.
32. Zarnowiecki DM, Dollman J & Parletta N (2014) Associations between predictors of children's dietary intake and socioeconomic position: a systematic review of the literature. *Obes Rev* **15**, 375–391.
33. Yuan C, Spiegelman D, Rimm EB *et al.* (2017) Validity of a dietary questionnaire assessed by comparison with multiple weighed dietary records or 24-hour recalls. *Am J Epidemiol* **185**, 570–584.
34. Willett W (2013) *Nutritional Epidemiology*, 3rd ed. Oxford/New York: Oxford University Press.
35. Fernandez-Alvira JM, de Bourdeaudhuij I, Singh AS *et al.* (2013) Clustering of energy balance-related behaviors and parental education in European children: the ENERGY-project. *Int J Behav Nutr Phys Act* **10**, 5.
36. Richter A, Heidemann C, Schulze MB *et al.* (2012) Dietary patterns of adolescents in Germany – associations with nutrient intake and other health related lifestyle characteristics. *BMC Pediatr* **12**, 35.
37. Araujo J, Teixeira J, Gaio AR *et al.* (2015) Dietary patterns among 13-y-old Portuguese adolescents. *Nutrition* **31**, 148–154.
38. Northstone K, Smith ADAC, Newby PK *et al.* (2013) Longitudinal comparisons of dietary patterns derived by cluster analysis in 7- to 13-year-old children. *Br J Nutr* **109**, 2050–2058.
39. Northstone K, Smith AD, Cribb VL *et al.* (2014) Dietary patterns in UK adolescents obtained from a dual-source FFQ and their associations with socio-economic position, nutrient intake and modes of eating. *Public Health Nutr* **17**, 1476–1485.
40. Bibiloni MDM, Martinez E, Llull R *et al.* (2012) Western and Mediterranean dietary patterns among Balearic Islands' adolescents: socio-economic and lifestyle determinants. *Public Health Nutr* **15**, 683–692.
41. Arriscado D, Muros JJ, Zabala M *et al.* (2014) Factors associated with low adherence to a Mediterranean diet in healthy children in northern Spain. *Appetite* **80**, 28–34.
42. Bibiloni MDM, Pons A & Tur JA (2016) Compliance with the Mediterranean Diet Quality Index (KIDMED) among Balearic Islands' adolescents and its association with socioeconomic, anthropometric and lifestyle factors. *Ann Nutr Metab* **68**, 42–50.
43. Handeland K, Kjellefold M, Wik Markhus M *et al.* (2016) A diet score assessing Norwegian adolescents' adherence to dietary recommendations – development and test–retest reproducibility of the score. *Nutrients* **8**, 467.
44. Martin MA, van Hook JL & Quiros S (2015) Is socioeconomic incorporation associated with a healthier diet? Dietary patterns among Mexican-origin children in the United States. *Soc Sci Med* **147**, 20–29.
45. Papadaki S & Mavriaki E (2015) Greek adolescents and the Mediterranean diet: factors affecting quality and adherence. *Nutrition* **31**, 345–349.
46. Kastorini CM, Lykou A, Yannakoulia M *et al.* (2016) The influence of a school-based intervention programme regarding

- adherence to a healthy diet in children and adolescents from disadvantaged areas in Greece: the DIATROFI study. *J Epidemiol Community Health* **70**, 671–677.
47. Yannakoulia M, Lykou A, Kastorini CM *et al.* (2016) Socio-economic and lifestyle parameters associated with diet quality of children and adolescents using classification and regression tree analysis: the DIATROFI study. *Public Health Nutr* **19**, 339–347.
 48. Beghin L, Dauchet L, de Vriendt T *et al.* (2014) Influence of parental socio-economic status on diet quality of European adolescents: results from the HELENA study. *Br J Nutr* **111**, 1303–1312.
 49. Finger JD, Varnaccia G, Tylleskar T *et al.* (2015) Dietary behaviour and parental socioeconomic position among adolescents: the German Health Interview and Examination Survey for Children and Adolescents 2003–2006 (KiGGS). *BMC Public Health* **15**, 498.
 50. Grosso G, Marventano S, Buscemi S *et al.* (2013) Factors associated with adherence to the Mediterranean diet among adolescents living in Sicily, Southern Italy. *Nutrients* **5**, 4908–4923.
 51. Ozen AE, Bibiloni MM, Murcia MA *et al.* (2015) Adherence to the Mediterranean diet and consumption of functional foods among the Balearic Islands' adolescent population. *Public Health Nutr* **18**, 659–668.
 52. Drewnowski A & Rehm CD (2015) Socioeconomic gradient in consumption of whole fruit and 100 % fruit juice among US children and adults. *Nutr J* **14**, 3.
 53. Gopinath B, Flood VM, Burlutsky G *et al.* (2014) Pattern and predictors of dairy consumption during adolescence. *Asia Pac J Clin Nutr* **23**, 612–618.
 54. Grosso G, Marventano S, Nolfo F *et al.* (2013) Personal eating, lifestyle, and family-related behaviors correlate with fruit and vegetable consumption in adolescents living in Sicily, Southern Italy. *Int J Vitam Nutr Res* **83**, 355–366.
 55. Lehto E, Ray C, Te Velde S *et al.* (2015) Mediation of parental educational level on fruit and vegetable intake among school-children in ten European countries. *Public Health Nutr* **18**, 89–99.
 56. Drouillet-Pinard P, Dubuisson C, Bordes I *et al.* (2017) Socio-economic disparities in the diet of French children and adolescents: a multidimensional issue. *Public Health Nutr* **20**, 870–882.
 57. Harris C, Flexeder C, Thiering E *et al.* (2015) Changes in dietary intake during puberty and their determinants: results from the GINIplus birth cohort study. *BMC Public Health* **15**, 841.
 58. Llull R, Bibiloni M, Pons A *et al.* (2015) Food consumption patterns of Balearic Islands' adolescents depending on their origin. *J Immigr Minor Health* **17**, 358–366.
 59. Kakinami L, Gauvin L, Seguin L *et al.* (2014) Persistent and occasional poverty and children's food consumption: evidence from a longitudinal Quebec birth cohort. *J Epidemiol Community Health* **68**, 987–992.
 60. Ahmadi N, Black JL, Velazquez CE *et al.* (2015) Associations between socio-economic status and school-day dietary intake in a sample of grade 5–8 students in Vancouver, Canada. *Public Health Nutr* **18**, 764–773.
 61. Fismen A-S, Smith ORF, Torsheim T *et al.* (2014) A school based study of time trends in food habits and their relation to socio-economic status among Norwegian adolescents, 2001–2009. *Int J Behav Nutr Phys Act* **11**, 115.
 62. Fismen A-S, Smith ORF, Torsheim T *et al.* (2016) Trends in food habits and their relation to socioeconomic status among Nordic adolescents 2001/2002–2009/2010. *PLoS One* **11**, e0148541.
 63. Meijerink FJ, van Vuuren CL, Wijnhoven HAH *et al.* (2016) Seven-year time trends in energy balance-related behaviours according to educational level and ethnic background among 14-year-old adolescents. *Public Health Nutr* **19**, 777–787.
 64. Minges KE, Chao A, Nam S *et al.* (2015) Weight status, gender, and race/ethnicity: are there differences in meeting recommended health behavior guidelines for adolescents? *J Sch Nurs* **31**, 135–145.
 65. Moore GF & Littlecott HJ (2015) School- and family-level socioeconomic status and health behaviors: multilevel analysis of a national survey in Wales, United Kingdom. *J Sch Health* **85**, 267–275.
 66. Pettigrew S, Jongenelis M, Chapman K *et al.* (2015) Factors influencing the frequency of children's consumption of soft drinks. *Appetite* **91**, 393–398.
 67. Pitel L, Madarasova Geckova A, Reijneveld SA *et al.* (2013) Socioeconomic differences in adolescent health-related behavior differ by gender. *J Epidemiol* **23**, 211–218.
 68. Richmond TK, Spadano-Gasbarro JL, Walls CE *et al.* (2013) Middle school food environments and racial/ethnic differences in sugar-sweetened beverage consumption: findings from the Healthy Choices study. *Prev Med* **57**, 735–738.
 69. Sundborn G, Utter J, Teevale T *et al.* (2014) Carbonated beverages consumption among New Zealand youth and associations with BMI and waist circumference. *Pac Health Dialog* **20**, 81–86.
 70. Voracova J, Sigmund E, Sigmundova D *et al.* (2016) Family affluence and the eating habits of 11- to 15-year-old Czech adolescents: HBSC 2002 and 2014. *Int J Environ Res Public Health* **13**, 1034.
 71. Attorp A, Scott JE, Yew AC *et al.* (2014) Associations between socioeconomic, parental and home environment factors and fruit and vegetable consumption of children in grades five and six in British Columbia, Canada. *BMC Public Health* **14**, 150.
 72. Dodd AH, Briefel R, Cabili C *et al.* (2013) Disparities in consumption of sugar-sweetened and other beverages by race/ethnicity and obesity status among United States schoolchildren. *J Nutr Educ Behav* **45**, 240–249.
 73. Bleich SN & Wolfson JA (2015) Trends in SSBs and snack consumption among children by age, body weight, and race/ethnicity. *Obesity (Silver Spring)* **23**, 1039–1046.
 74. Larson N, Story M, Eisenberg ME *et al.* (2016) Secular trends in meal and snack patterns among adolescents from 1999 to 2010. *J Acad Nutr Diet* **116**, 240–250.e2.
 75. Galobardes B, Shaw M, Lawlor DA *et al.* (2006) Indicators of socioeconomic position (part 1). *J Epidemiol Community Health* **60**, 7–12.
 76. Galobardes B, Morabia A & Bernstein MS (2001) Diet and socioeconomic position: does the use of different indicators matter? *Int J Epidemiol* **30**, 334–340.
 77. Méjean C, Si Hassen W, Lecossais C *et al.* (2016) Socio-economic indicators are independently associated with intake of animal foods in French adults. *Public Health Nutr* **19**, 3146–3157.
 78. Si Hassen W, Castetbon K, Cardon P *et al.* (2016) Socioeconomic indicators are independently associated with nutrient intake in French adults: a DEDIPAC study. *Nutrients* **8**, 158.
 79. Brown AGM, Houser RF, Mattei J *et al.* (2018) Diet quality among US-born and foreign-born non-Hispanic blacks: NHANES 2003–2012 data. *Am J Clin Nutr* **107**, 695–706.
 80. Pillen H, Tsourtos G, Coveney J *et al.* (2017) Retaining traditional dietary practices among Greek immigrants to Australia: the role of ethnic identity. *Ecol Food Nutr* **56**, 312–328.
 81. Osei-Kwasi HA, Powell K, Nicolaou M *et al.* (2017) The influence of migration on dietary practices of Ghanaians living in the United Kingdom: a qualitative study. *Ann Hum Biol* **44**, 454–463.