



Ready-to-eat cereal fortification: a modelling study on the impact of changing ready-to-eat cereal fortification levels on population intake of nutrients

Jessica D Smith^{1,*}, Neha Jain² and Regan L Bailey³

¹Bell Institute of Health and Nutrition, General Mills, Inc., James Ford Bell Technical Center, 9000 Plymouth Avenue N, Golden Valley, MN 55427 USA; ²General Mills India Pvt. Ltd, Mumbai, India; ³Department of Nutrition Science, Purdue University, West Lafayette, IN, USA

Submitted 13 March 2019: Final revision received 2 August 2019: Accepted 19 August 2019: First published online 20 January 2020

Abstract

Objective: Ready-to-eat (RTE) cereal is an important source of nutrients in the American diet. Recent regulatory changes to labelling requirements may impact the fortification of RTE cereal. We used an evidence-based approach to optimize the fortification of RTE cereal considering current dietary patterns and nutrition policy.

Design: A US modelling study of cross-sectional data from the National Health and Nutrition Examination Survey (NHANES) 2013–2014. The percentage of the population below the Estimated Average Requirement (EAR) and above the Upper Tolerable Intake Level (UL) was modelled under three scenarios: baseline, zero fortification and optimized fortification.

Setting: USA.

Participants: Toddlers aged 1–3 years, *n* 559; children aged 4–12 years, *n* 1540; adolescents aged 13–18 years, *n* 992; and adults aged ≥ 19 years, *n* 576.

Results: Comparing current with optimized fortification, nutrient/100 g RTE cereal decreased for vitamin A, thiamin, riboflavin, niacin, vitamin B₆, folic acid, vitamin B₁₂, Ca and Fe (by 2–82%). The amount of vitamins C and D increased (by 13 and 50%, respectively). Among RTE cereal eaters, these changes resulted in modest increases in the percentage of the population aged ≥ 1 year below the EAR (+0.5 to +11.5 percentage points). Decreases were observed in the percentage of the population above the UL.

Conclusions: Fortification of RTE cereal can be optimized to provide key nutrients and minimize the percentage of the population below the EAR and above the UL. Dietary intake modelling is useful to ensure that RTE cereal continues to help the population meet their nutrient needs.

Keywords

Ready-to-eat cereal
Fortification
Dietary reference intakes
National Health and Nutrition
Examination Survey

Voluntary fortification of foods has historically been a successful strategy in reducing nutritional inadequacy in the USA⁽¹⁾. Shifting dietary patterns, particularly emerging nutritional risks from both nutrient inadequacies and, potentially, excess intakes of energy and some nutrients, require that food manufacturers periodically re-examine their fortification policies. Public health efforts to reduce the risk of dietary inadequacy include encouraging dietary diversity, fortification and targeted programmes with dietary supplements⁽²⁾. Indeed, current US fortification practices play a substantial role in reducing the prevalence of intakes below the Estimated Average Requirement

(EAR), the Dietary Reference Intake (DRI) threshold for population nutritional adequacy, across all age groups^(3,4).

The DRI, developed by the US Food and Nutrition Board of The National Academies, provide nutrient reference values for over forty nutrient substances and serve as the scientific basis for the development of food guidelines in the USA and Canada^(5–7). The DRI are comprised of a set of values including: (i) the EAR, which is used to assess population nutritional adequacy; (ii) the RDA, which is the average daily level of intake sufficient to meet the nutrient requirements of nearly all healthy people; and (iii) the Tolerable Upper Intake Level (UL), which is

*Corresponding author. Email jessica.smith@genmills.com



the maximum daily intake unlikely to cause adverse health effects.

The DRI can be used to assess population-level nutrient intakes and form the basis for the Daily Value (DV) recommendations by the US Food and Drug Administration (FDA)⁽¹⁾. The DV are used for food labelling in the USA and provide information on how much of a nutrient, relative to the daily requirement for a 8368 kJ (2000 kcal) diet, is in a serving of food on the Nutrition Facts Panel of packaged food products. The FDA regulates the contents of the Nutrition Facts Panel including the reference amount customarily consumed (RACC), which is used as the basis for the labelled serving size, and the DV. Recently, the FDA updated the requirements of the Nutrition Facts Panel including the RACC for some foods and the DV for several nutrients under Nutrition Label Reform (NLR). The NLR regulation was finalized in 2016 and large companies are required to comply with the regulation by 1 January 2020. Both the changes in RACC and the changes in DV will impact fortified nutrients in foods: companies will either need to change the amount of nutrients added per 100 g of food to maintain their current labelled percentage of the DV (%DV) values or change the labelled %DV if products are not reformulated.

Greater ready-to-eat (RTE) cereal consumption (compared with low/no consumption) is associated with higher nutrient intakes in the diets of US^(4,8) and European^(9,10) children and adolescents. RTE cereal consumption has been associated with enhanced intakes of many vitamins and minerals in adults and children, either directly through its consumption^(8,11) or through its co-consumption with milk⁽¹²⁾. Serum folate and vitamin B₁₂ concentrations were higher⁽¹³⁾ in children who consume RTE cereals⁽¹⁴⁾ and RTE cereal consumption in older adults was associated with higher intakes of fibre, whole grains, and several vitamins and minerals⁽¹⁵⁾. While from foods alone few adults exceed the UL, some children and adolescents (aged 2–18 years) exceed the UL for folic acid (3%), Cu (3%), vitamin A (5%) and Zn (13%), especially among children using dietary supplements⁽¹⁶⁾. Recent data from the Feeding and Infants and Toddlers Study (FITS 2016) suggest that usual intakes from diet and supplements of >30% of infants and toddlers were above the UL for vitamin A and Zn^(17,18).

Because RTE cereal is a major source of nutrients in the American diet, periodic re-evaluation of the fortification approach of RTE cereal should be undertaken to ensure that nutrients within RTE cereal are still meeting public health needs for both children and adults. Furthermore, changing the nutrient content of RTE cereal in response to NLR may impact population-level intakes of several key nutrients, changing the percentage that is consuming either below recommendations (i.e. below the EAR) or above the UL. Therefore, the purpose of the current analysis was to use dietary modelling to test an optimized fortification strategy for RTE cereal that balances the

amount of nutrients to not unduly increase the likelihood of contributing towards dietary intakes above the UL while retaining the public health benefit of reducing the risk of nutritional inadequacy.

Methods

Survey design and participants

The National Health and Nutrition Examination Survey (NHANES) is a nationally representative, cross-sectional series of surveys that samples the resident, non-institutionalized, civilian US population, collected by the National Center for Health Statistics of the Centers for Disease Control and Prevention⁽¹⁹⁾. Data from the 2013 to 2014 NHANES cycle were used for the current analyses. Written informed consent was obtained for all participants or proxies and the NHANES survey protocols were approved by the Research Ethics Review Board at the National Center for Health Statistics.

NHANES data are collected in multiple ways. First, participants are asked to complete an in-person household interview, during which demographic data are collected through a computer-assisted personal interview. Second, participants are asked to attend a health examination in a mobile examination centre and complete an in-person 24 h dietary recall (24HR)⁽²⁰⁾. The final data collection occurs via telephone when a second 24HR is collected; this occurs approximately 3–10 d after the health examination, with an emphasis on collecting both weekday and weekend dietary intakes. Both 24HR are collected as part of the US Department of Agriculture's What We Eat in America (WWEIA) using the validated Automated Multiple-Pass Method^(21,22).

The analytic sample included 8167 participants aged ≥1 year who had complete 24 h dietary intake data. Results were also examined by age group: toddlers aged 1–3 years, *n* 559; children aged 4–12 years, *n* 1540; adolescents aged 13–18 years, *n* 992; and adults aged ≥19 years excluding those pregnant or lactating (*n* 100), *n* 5076. The age groups of 1–3 years and ≥4 years were purposefully chosen because of the FDA's prescribed RACC for foods on a category basis differ for these population age groups. RACC reflect the regulatory serving size that food manufacturers must utilize to calculate the labelled serving size on their products^(23,24). We defined RTE cereal eaters as those participants who reported eating any quantity of RTE cereal (see definition of RTE cereal below) on their first 24HR.

Dietary data

The US Department of Agriculture's Food and Nutrient Database for Dietary Studies (FNDDS) is used to calculate the nutrient values of all reported foods and beverages within NHANES⁽²⁵⁾. Individual foods and beverages within



FNDDS are grouped according to the WWEIA Food Categories, which includes two categories for RTE cereal ('ready-to-eat cereal, lower sugar' and 'ready-to-eat cereal, higher sugar') which we combined. We included all FNDDS food codes within the WWEIA RTE cereal categories in the current analysis. Nutrients from dietary supplements were not analysed in this food optimization project based on the rationale that an underlying premise of the 2015–2020 Dietary Guidelines for Americans is that 'nutritional needs should be met primarily from foods'⁽⁶⁾. The scope of this project included nutrients that are typically added to RTE cereal, including the B-vitamins (thiamin, riboflavin, niacin, vitamin B₆, folic acid, vitamin B₁₂), vitamins A, D and C, and Ca, Fe and Zn^(26–28).

Ready-to-eat cereal fortification models

We considered the amount of nutrient in RTE cereal per 100 g of cereal and adjusted the level of fortification to reflect: (i) the change in fortification according to the three scenarios below; and (ii) the changes to the DV as required by the FDA's NLR (see online supplementary material, Supplemental Table S1) and the changes in the RACC for RTE cereal as part of NLR⁽²⁴⁾. Compared with previous DV, the NLR DV for some nutrients decreased (thiamin, riboflavin, vitamin B₆, vitamin B₁₂, Zn), increased (vitamin D, vitamin C, Ca), changed units (vitamin A, niacin, folate) or did not change (Fe).

For labelling purposes, there are three categories of RTE cereal based on their density (g/cup), each with a different RACC: (i) light-weight cereal which weighs <20 g per cup; (ii) medium-weight cereal which weighs 20 g or more but less than 43 g per cup and also includes high-fibre cereals containing 28 g or more fibre per 100 g; and (iii) heavy-weight cereals, weighing 43 g or more per cup or biscuit-type cereal. The RACC for two categories of RTE cereal changed under NLR: the RACC for medium-weight cereal changed from 30 to 40 g and the RACC for heavy-weight RTE cereal changed from 55 to 60 g. The RACC for light-weight cereal remained 15 g. This change in RACC will influence the 'density' of nutrients per 100 g of cereal if the labelled %DV of nutrients does not change. For example, if a cereal fortifies at 20 %DV for Fe at a 30 g RACC, it will contain a greater absolute amount of Fe per 100 g (approximately 15 mg per 100 g RTE cereal) than a 40 g RACC that also labels at 20 %DV of Fe (approximately 9 mg per 100 g RTE cereal).

For the present study, we assumed that the entire nutrient content of RTE cereal of those vitamins and minerals listed above were from fortification. The fortification scenarios and amount of nutrients per 100 g for the three fortification models are presented in Table 1^(24,29).

Baseline

For the baseline scenario we used the nutrient content of RTE cereal as reported in FNDDS 2013–2014. Table 1

Table 1 Current and proposed levels of fortification of US ready-to-eat (RTE) cereal

Nutrient	Baseline fortification, pre-NLR DV and RACC*		Zero fortification†		Optimized fortification, post-NLR DV and RACC‡	
	%DV	Amount/100 g RTE cereal‡	%DV	Amount/100 g RTE cereal	%DV	Amount/100 g RTE Cereal§
Vitamin A	10	1667 IU	0	0 IU	10	225 µg RAE (750 IU)
Vitamin C	10	20 mg	0	0 mg	10	22.5 mg
Vitamin D	10	133 IU	0	0 IU	10	5 µg (200 IU)
Thiamin	25	1.25 mg	0	0 mg	20	0.6 mg
Riboflavin	25	1.42 mg	0	0 mg	10	0.33 mg
Niacin	25	16.7 mg	0	0 mg	10	4 mg NE
Vitamin B ₆	25	1.67 mg	0	0 mg	20	0.85 mg
Folic acid	50	667 µg	0	0 mg DFE	20	200 µg DFE (120 µg)¶
Vitamin B ₁₂	25	5 µg	0	0 mg	20	1.2 µg
Ca	10	333 mg	0	0 mg	10	325 mg
Fe	25	15 mg	0	0 mg	20	9 mg
Zn	25	12.5 mg	0	0 mg	20	5.5 mg

NLR, Food and Drug Administration's Nutrition Label Reform⁽²⁴⁾; DV, Daily Value; RACC, reference amount customarily consumed; %DV, percentage of the Daily Value; RAE, retinol activity equivalents; NE, niacin equivalents; DFE, dietary folate equivalents.

*Current fortification levels are a typical representation of most RTE cereals; however, not all RTE cereals are fortified to these precise levels. In the baseline condition, we show results for the percentage of the population below the Estimated Average Requirement and above the Tolerable Upper Intake Level based on current intakes of cereal and the most recent version of the US Department of Agriculture's National Nutrient Database.

†These are modelled changes to the fortification scheme of RTE cereals. In modelling these fortification levels, we assumed only a decrease in nutrients. That is, we did not model an increase in nutrients among RTE cereals fortifying below these levels the baseline levels.

‡Pre-NLR RACC for RTE cereals are 30 g for medium-weight cereals and 55 g for heavy-weight cereals⁽²⁴⁾. The DV are reported in the units in which they appeared in the regulations. The DV for several nutrients (i.e. vitamin A, vitamin D, niacin, and folic acid) changed units under NLR regulation. The conversion to the previous DV unit is provided for clarity.

§Post-NLR RACC for RTE cereals are 40 g for medium-weight cereals and 60 g for heavy-weight cereals⁽²⁴⁾. The DV are reported in the units in which they appeared in the regulations. The DV for several nutrients (i.e. vitamin A, vitamin D, niacin, and folic acid) changed units under NLR regulation. The conversion to the previous DV unit is provided for clarity.

¶IRTE cereals that qualify for the Special Supplemental Nutrition Program for Women, Infants, and Children are fortified at higher Fe level; that is, 45 %DV⁽²⁹⁾.

¶¶We assumed all folate in RTE cereal is the folic acid form when converting µg DFE to µg.



reports the typical contemporary fortification practices representative of most of the current US RTE cereal marketplace; however, it should be noted that not all RTE cereals contain these levels of fortification.

Zero fortification

We assumed for the B-vitamins (thiamin, riboflavin, niacin, vitamin B₆, folic acid, vitamin B₁₂), vitamins A, D, and C, and Ca, Fe and Zn that the entire content of RTE cereal was from fortification and none was naturally occurring. Therefore, we assumed under this model that RTE cereal did not contribute to the daily intake of these nutrients.

Optimized fortification

A set of alternative %DV for RTE cereal was developed that considered several sources of information including: (i) current labelled %DV; (ii) NLR changes to the DV; (iii) recommendations within the Dietary Guidelines for Americans (e.g. nutrients of public health concern); and (iv) current nutrient intakes. In some cases, we tested the impact of several different fortified levels of nutrients before selecting the 'optimized' value (data not shown). We considered a fortification level for a nutrient 'optimized' if it decreased the percentage of the population above the UL or the percentage of the population below the EAR for under-consumed nutrients by a meaningful amount (i.e. 5 percentage points (pp)). Here we are showing one set of alternative fortification levels that represent the balance between decreasing the percentage of the population below the EAR and above the UL. Fortification levels of nutrients were adjusted according to the levels in Table 1: we lowered the %DV for B-vitamins, Fe and Zn and the %DV remained the same for vitamin A, vitamin C, vitamin D and Ca. We accounted for the changes in RACC and DV under NLR when calculating the amount per 100 g of RTE cereal. We only modelled a decrease in nutrient content per 100 g RTE cereal: for the RTE cereal FNDDS food codes that either contained no fortification or fortification levels lower than those in our optimized scenario, we did not increase the amount of nutrient. By only modelling a decrease in fortification, we estimated the impact of fortification changes among those RTE cereals that currently fortify at typical levels (Table 1) while excluding those cereals that either do not currently fortify (such as organic RTE cereals) or fortify at levels below most RTE cereals.

Statistical analysis

Statistical analyses were performed using the statistical software package SAS version 9.3 and procedures that account for the complex survey design and sampling weights necessary to make the NHANES data reflective of the US population. Before the diet can be characterized relative to DRI, usual or long-term estimates are needed that are adjusted for the known random measurement error in self-reported diet^(30–32). For this analysis, the National

Cancer Institute method^(33,34) was used to produce the mean and percentiles of usual nutrient intakes. Covariates in the National Cancer Institute usual intake models included whether the 24HR was collected on a weekday or weekend day and the interview method (in-person or telephone) of the 24HR.

Using usual intakes, we calculated the percentage of the population meeting or exceeding the DRI for twelve micronutrients. To calculate the percentage of the population meeting their nutrient requirements, we used the EAR, which represents the median requirements of a nutrient and is used to estimate the needs of a group or population. The percentage of the population below the EAR was calculated for vitamin A, vitamin C, vitamin D, thiamin, riboflavin, niacin, vitamin B₆, vitamin B₁₂, folate, Ca and Zn using the cut-point method. Because the distribution of requirements is skewed for Fe, we used the probability method to determine the prevalence of the population below the EAR.

The percentage of the population above the UL was also determined for the nine of the twelve nutrients with an established UL: vitamin A, vitamin C, vitamin D, niacin, vitamin B₆, folate, Ca, Fe and Zn. The UL for vitamin A was established only for the retinol form of the vitamin; for folate it was established only for folic acid; and for niacin, it was established only for niacin from fortified foods or supplements. Therefore, in calculating the percentage of the population above the UL for these nutrients, we used retinol for vitamin A and folic acid for folate. For niacin, we referred to the FNDDS 2013–2014 which categorizes each food code as being unfortified, fortified or containing fortified ingredients⁽²⁵⁾. For calculating the UL, we considered 0 % of the niacin from unfortified foods, 100 % of the niacin from fortified foods and 75 % of the niacin from foods containing fortified ingredients. We used 75 % of niacin for foods containing fortified ingredients as a conservative estimate of the amount of fortified niacin in these foods that would error on the side of overestimating the percentage of the population above the UL for niacin. We compared the baseline percentage below the EAR and baseline percentage above the UL for males *v.* females using an approximate χ^2 test. In cases with very low estimates and standard errors, normal approximations would not be appropriate and *P* values were not calculated⁽³⁵⁾. We considered a *P* value of <0.05 to be statistically significant.

Results

Fortification levels of ready-to-eat cereal under the three scenarios

Comparing the current with the optimized scenario, there were no changes in the %DV for vitamin A, vitamin C, vitamin D and Ca; however, due to NLR RACC and DV changes, the absolute amount of nutrient content within



RTE cereal decreased for vitamin A (−55 %) and Ca (−2 %) and increased for vitamin C (+13 %) and vitamin D (+50 %). For thiamin, vitamin B₆ and Zn, the %DV changed from 25 %DV at baseline to 20 %DV for the optimized scenario; however, when also considering the NLR changes, this represented about a 50 % decrease in the absolute amount of these nutrients. Fe had a 5-pp decrease in the %DV and this resulted in a 40 % decrease in the absolute amount of Fe. The largest absolute decreases were seen for vitamin B₁₂, riboflavin, niacin and folic acid (−76 to −82 %) and this was largely attributed to the lower DV set forth under NLR (Table 1).

Prevalence and median amount of cereal consumption

Overall, 24.2 % of the US population aged ≥1 year consume RTE cereal on a given day. Toddlers are the age group with the highest prevalence of RTE cereal consumption (45.5 %) and consumption tends to decline with age: 39.9 % of children reported eating RTE cereal, while 27.3 % of adolescents and 20.3 % of adults did so. However, the median daily consumption of RTE cereal was highest in adults (45.9 g/d) and decreased across age groups, with toddlers having the smallest daily intake of RTE cereal (21.2 g/d; Table 2). Demographic characteristics of the study participants can be found in the online supplementary material, Supplemental Table S2.

Percentage of the population below the Estimated Average Requirement under the three fortification scenarios

We noted a pattern of overall higher percentages of the total population below the EAR than seen among RTE cereal eaters only; however, given that RTE cereal consumers are also part of the entire population, no statistical comparisons were possible (nor was this an

objective of the current analyses; Table 3). Similar to the entire US population, the percentage of RTE cereal eaters aged ≥1 year below the EAR was highest for vitamin D (82.5 %; Table 3). Over one-quarter of RTE cereal eaters aged ≥1 year were below the EAR for vitamin C (28.8 %) and Ca (26.9 %). The percentage of RTE cereal eaters aged ≥1 year consuming below the EAR of the other nutrients, including vitamin A, Zn, B-vitamins and Fe, ranged from 0 to 6 %.

When we modelled removing all fortification from RTE cereal, the percentage of RTE cereal eaters aged ≥1 year who were below the EAR for nutrients increased compared with the baseline scenario, ranging from an increase of 25.1 pp for vitamin A to 1.1 pp for riboflavin. Under the optimized fortification scenario, this increase in the percentage of RTE cereal consumers aged ≥1 year below the EAR from baseline was attenuated compared with the zero fortification model, ranging from an increase of 11.5 pp for vitamin A to 0.5 pp for Fe.

At baseline among RTE cereal consumers, the prevalence of intakes below the EAR differed across age groups and tended to be higher for adolescents and adults compared with toddlers and children for vitamin A, vitamin C, vitamin D and Ca (all $P < 0.0001$; Fig. 1 and online supplementary material, Supplemental Tables S3a and S3b). In the modelled scenarios we saw a similar pattern among all age groups: the percentage of RTE cereal eaters below the EAR was lowest at baseline and highest in the zero-fortification scenario (Fig. 1 and Supplemental Tables S3a and S3b).

We did observe a differential pattern in vitamin D and Ca intakes below the EAR by sex. At baseline, a greater percentage of female RTE cereal eaters aged ≥1 year, compared with males, were consuming below the EAR for vitamin D (93.7 *v.* 71.0 %, $P < 0.0001$) and Ca (39.5 *v.* 16.3 %, $P < 0.0001$). Under the zero-fortification scenario, we saw a greater increase in the percentage of females (increase of 31.5 pp) not consuming adequate

Table 2 Prevalence and amount of ready-to-eat (RTE) cereal consumption among US toddlers, children, adolescents and adults*

	All ages (≥1 year)	Toddlers (1–3 years)	Children (4–12 years)	Adolescents (13–18 years)	Adults (≥19 years)
<i>n</i>	8167	559	1540	992	5076
RTE cereal consumers†					
<i>n</i> ‡	1994	237	589	247	921
%	24.2	45.5	39.9	27.3	20.3
Median daily RTE cereal consumption among RTE cereal consumers† only (g/d)	42.4	21.2	30.8	45.4	45.9

*Data are from the National Health and Nutrition Examination Survey (NHANES) 2013–2014, excluding pregnant and lactating women, using self-reported food intake from the first 24 h dietary recall.

†RTE cereal consumers were identified as those who reported consuming any quantity of RTE cereal in their first 24 h dietary recall in NHANES 2013–2014.

‡Percentages were weighted using the appropriate weights set by NHANES to account for the sampling design, non-response and current demographics of the US population.

Table 3 Percentage of the total population and of ready-to-eat (RTE) cereal eaters below the Estimated Average Requirement (EAR) for nutrients fortified in RTE cereal for the US population aged ≥ 1 year*

Nutrient	Total population aged ≥ 1 year (n8167)					
	Baseline		Zero fortification†		Optimized fortification‡	
	% below EAR	SE (%)	% below EAR	SE (%)	% below EAR	SE (%)
Vitamin A	39.5	1.8	47.2	1.8	44.0	1.8
Thiamin	6.8	0.9	9.5	0.9	8.0	0.9
Riboflavin	3.1	0.5	3.9	0.5	3.7	0.5
Niacin	1.3	0.3	2.2	0.3	1.9	0.3
Vitamin B ₆	8.6	0.8	13.6	0.9	10.8	0.9
Folate	12.1	1.3	19.9	1.1	16.8	1.2
Vitamin B ₁₂	4.7	0.7	7.0	0.7	6.2	0.7
Vitamin C	41.4	1.5	43.4	1.5	42.0	1.5
Vitamin D	93.7	0.7	95.3	0.6	93.8	0.7
Ca	43.3	1.4	44.6	1.3	43.6	1.3
Fe	3.5	0.6	6.1	0.5	4.4	0.5
Zn	16.9	1.5	22.4	1.2	18.6	1.4

Nutrient	RTE cereal eaters‡ aged ≥ 1 year (n 1994)					
	Baseline		Zero fortification†		Optimized fortification‡	
	% below EAR	SE (%)	% below EAR	SE (%)	% below EAR	SE (%)
Vitamin A	5.7	1.0	30.8	2.6	17.2	2.4
Thiamin	0.7	0.6	11.2	1.4	3.7	1.3
Riboflavin	0.0	0.0	1.1	0.5	0.6	0.3
Niacin	0.1	0.1	2.7	1.2	1.3	0.8
Vitamin B ₆	0.3	0.2	12.1	1.9	3.0	0.7
Folate	0.2	0.1	21.3	1.4	8.3	1.1
Vitamin B ₁₂	0.0	0.1	3.3	1.1	1.3	0.6
Vitamin C	28.8	2.0	38.1	2.4	31.6	2.0
Vitamin D	82.5	3.3	92.0	2.2	83.1	3.2
Ca	26.9	2.4	34.3	2.2	28.1	2.3
Fe	0.0	0.1	7.4	1.1	0.5	0.3
Zn	2.7	1.0	20.4	2.7	7.1	1.6

*Data are from the National Health and Nutrition Examination Survey (NHANES) 2013–2014, excluding pregnant and lactating women, using self-reported food intake from the first 24 h dietary recall. Data by age group (toddlers aged 1–3 years, children aged 4–12 years, adolescents aged 13–18 years and adults aged ≥ 19 years) are available in the online supplementary materials.

†For the zero-fortification scenario, we modelled removing all fortification of the nutrients listed in this table. In the optimized fortification scenario, we modelled fortification levels of 10% of the Daily Value for vitamins A, C and D, riboflavin, niacin and Ca, and 20% of the Daily Value for thiamin, vitamins B₆ and B₁₂, folic acid, Fe and Zn.

‡RTE cereal consumers were identified as those who reported consuming any quantity of RTE cereal in their first 24 h dietary recall in NHANES 2013–2014.

folate compared with males (increase of 24.1 pp). This increase was attenuated, although still present, in the optimized fortification scenario (increase compared with baseline of 13.7 pp for females and 9.2 pp for males). Similarly, removing fortification of all other nutrients appeared to have a greater impact on females compared with males. Again, this impact was attenuated in the optimized fortification scenario (online supplementary material, Supplemental Table S4).

When we examined the impact of the modelled fortification scenarios in the total population (including RTE cereal eaters and non-eaters) to estimate the overall population public health impact, we found a similar pattern of results to that observed among RTE cereal eaters only, although the magnitude of the impact of changing cereal fortification was attenuated since only the nutrient intakes of those consuming RTE cereal were

considered (Table 3 and online supplementary material, Supplemental Tables S3a, S3b and S4).

Percentage of the population above the Upper Tolerable Intake Level under the three fortification scenarios

In both the entire US population and among RTE cereal eaters only, the percentage above the UL was highest at baseline for niacin (total population, 3.4%; RTE cereal eaters, 15.9%) followed by Zn (total population, 2.8%; RTE cereal eaters, 9.1%). For all other nutrients, the percentage above the UL at baseline was less than 1% for the total US population; this was similar for RTE cereal eaters, except for retinol and folic acid where $< 5\%$ of RTE cereal eaters were above the UL. When we modelled removing all fortification from RTE cereal, the percentage




Fig. 1 Percentage of US ready-to-eat (RTE) cereal eaters below the Estimated Average Requirement (EAR) for nutrients fortified in RTE cereal for under three scenarios: baseline (□), zero fortification (■) and optimized fortification (▨): (a) toddlers aged 1–3 years, n 237; (b) children aged 4–12 years, n 589; (c) adolescents aged 13–18 years, n 247; (d) adults aged \geq 19 years, n 921. Data are from the National Health and Nutrition Examination Survey (NHANES) 2013–2014. The baseline scenario is the current nutrient intake; for the zero-fortification scenario, we modelled removing all fortification of the nutrients listed in this figure. In the optimized fortification scenario, we modelled fortification levels of 10 % of the Daily Value for vitamins A, C and D, riboflavin, niacin and Ca, and 20 % of the Daily Value for thiamin, vitamins B₆ and B₁₂, folic acid, iron and zinc. Values are percentages with their standard errors indicated by vertical bars

of RTE cereal eaters aged \geq 1 year who were above the UL decreased to $<1\%$ for all nutrients except Zn (2.5 % above the UL); and in the optimized scenario this decreased to $<1.1\%$ for all nutrients, again except for Zn (5.1 % above the UL). Similar patterns were seen for the entire population (Table 4).

Across age groups, 0 % of RTE cereal eaters were above the UL for vitamin C, vitamin D and vitamin B₆ at baseline. Less than 1 % of adults were above the UL for Ca and Fe, and for all other age groups this was 0 %. Toddlers and children appeared to have a higher percentage above the UL for retinol, niacin, folic acid and Zn compared with adolescents and adults, although statistical comparisons were not possible for many nutrients due to very low estimates (Fig. 2 and online supplementary material, Supplemental Tables S5a and S5b). The percentage above the UL decreased in all age groups in the zero fortification and optimized fortification scenarios compared with baseline for retinol,

niacin, folic acid and Zn, sometimes dramatically among RTE cereal eaters; for example, among toddlers, the percentage above the UL for niacin decreased from 63.8 % at baseline to 10.1 % in the optimized fortification scenario. Likewise, the percentage of adolescents above the UL for folic acid decreased from 28.9 % at baseline to 0.4 % in the optimized fortification scenario (Fig. 2 and Supplemental Tables S5a and S5b).

We found that men who consumed RTE cereal had a higher percentage above the UL for Ca (women, 0.1 %; men, 0.9 %; $P=0.01$) and niacin (women, 10.9 %; men, 22.4 %; $P=0.02$) compared with women. Decreases in the percentage above the UL in the zero fortification and optimized fortification scenarios for RTE cereal eaters appeared similar for both women and men except for niacin, with males potentially having a greater decrease from baseline to the optimized scenario than women (men, -20.9 pp; women, -10.3 pp; online supplementary material, Supplemental Table S6).

Table 4 Percentage of the total population and of ready-to-eat (RTE) cereal eaters above the Tolerable Upper Intake Level (UL) for nutrients fortified in RTE cereal for the US population aged ≥ 1 year*

Nutrient	Total population aged ≥ 1 year (n8167)					
	Baseline		Zero fortification†		Optimized fortification‡	
	% above UL	SE (%)	% above UL	SE (%)	% above UL	SE (%)
Retinol	0.9	0.2	0.3	0.1	0.4	0.1
Niacin‡	3.4	0.5	0.6	0.1	0.7	0.2
Vitamin B ₆	0.0	0.0	0.0	0.0	0.0	0.0
Folic acid	0.3	0.1	0.0	0.0	0.0	0.0
Vitamin C	0.0	0.0	0.0	0.0	0.0	0.0
Vitamin D	0.0	0.0	0.0	0.0	0.0	0.0
Ca	0.3	0.0	0.2	0.0	0.3	0.0
Fe	0.0	0.0	0.0	0.0	0.0	0.0
Zn	2.8	0.3	1.4	0.2	2.0	0.2

Nutrient	RTE cereal eaters aged ≥ 1 year (n1994)					
	Baseline		Zero fortification†		Optimized fortification‡	
	% above UL	SE (%)	% above UL	SE (%)	% above UL	SE (%)
Retinol	3.2	0.7	0.6	0.2	1.1	0.3
Niacin‡	15.9	1.5	0.3	0.3	1.0	0.5
Vitamin B ₆	0.0	0.0	0.0	0.0	0.0	0.0
Folic acid	4.0	0.6	0.0	0.0	0.0	0.0
Vitamin C	0.0	0.0	0.0	0.0	0.0	0.0
Vitamin D	0.0	0.0	0.0	0.0	0.0	0.0
Ca	0.5	0.2	0.3	0.1	0.5	0.2
Fe	0.5	0.3	0.0	0.0	0.0	0.0
Zn	9.1	1.2	2.5	0.5	5.1	0.7

*Data are from the National Health and Nutrition Examination Survey (NHANES) 2013–2014, excluding pregnant and lactating women, using self-reported food intake from the first 24 h dietary recall. Data by age group (toddlers aged 1–3 years, children aged 4–12 years, adolescents aged 13–18 years and adults aged ≥ 19 years) are available in the online supplementary material.

†For the zero-fortification scenario, we modelled removing all fortification of the nutrients listed in this table. In the optimized fortification scenario, we modelled fortification levels of 10% of the Daily Value for vitamins A, C and D, riboflavin, niacin and Ca, and 20% of the Daily Value for thiamin, vitamins B₆ and B₁₂, folic acid, Fe and Zn.

‡The UL for niacin is based on fortified or supplemental niacin only. Therefore, we identified fortified foods using the US Department of Agriculture's Food and Nutrient Database for Dietary Studies 2013–2014 and considered 100% of the niacin content of the foods from fortification for fortified foods, 0% of the niacin content of the foods from fortification for unfortified foods and 75% of the niacin content of the foods from fortification for partially fortified foods in calculating the percentage above the UL.

||RTE cereal consumers were identified as those who reported consuming any quantity of RTE cereal in their first 24 h dietary recall in NHANES 2013–2014.

Discussion

'Fortification' is an umbrella term used to describe both mandatory enrichment and voluntary addition of nutrients to foods. 'Enrichment' is the addition of nutrients either to replace nutrients lost in processing (e.g. thiamin, riboflavin, niacin and Fe added to grains) or for a demonstrated public health need⁽³⁶⁾. Despite the role of discretionary fortification (also referred to as 'voluntary fortification') to reduce the risk of nutrient inadequacy for the total population and across different race/ethnic groups and socio-economic status^(38–40), some critics have argued that voluntary fortification practices have the potential to unduly expose the public to higher than recommended levels of nutrients^(38–40), especially when taken within the context of high supplement use^(16,41–43).

Thus, policy makers and food manufacturers have weighed the benefits of preventing nutrient inadequacies with the risk of intakes above the UL⁽⁴⁴⁾. Indeed, in the current American dietary landscape frank micronutrient

deficiencies are rare, but many nutrients of concern and 'shortfall' nutrients have been identified in the American diet⁽⁶⁾. Thus, encouraging more nutrient-balanced eating patterns should be a public health and food industry priority. This concept is closely aligned with the FDA fortification policy that outlines the major tenants that should guide fortification practices. First, contemporary intakes should be demonstrated to be below desirable levels. Second, enough people in the population should consume the food or beverage selected for fortification. Finally, the amount of a nutrient added to foods should provide reasonable assurance against excessive or toxic intakes⁽⁴⁵⁾.

RTE cereal is a good candidate for voluntary fortification because of its foundational place at the centre of the breakfast meal, its ability to provide fibre and whole grains, its affordability, its popularity and its well-established associations with healthier overall diets⁽¹⁰⁾. While some RTE cereals are pre-sweetened, observational data have shown positive associations between RTE cereal and diet quality and inverse associations with




Fig. 2 Percentage of US ready-to-eat (RTE) cereal eaters above the Tolerable Upper Intake Level (UL) for nutrients fortified in RTE cereal under three scenarios: baseline (□), zero fortification (■) and optimized fortification (▨): (a) toddlers aged 1–3 years, *n* 237; (b) children aged 4–12 years, *n* 589; (c) adolescents aged 13–18 years, *n* 247; (d) adults aged ≥19 years, *n* 921. Data are from the National Health and Nutrition Examination Survey (NHANES) 2013–2014. The baseline scenario is the current nutrient intake; for the zero-fortification scenario, we modelled removing all fortification of the nutrients listed in this figure. In the optimized fortification scenario, we modelled fortification levels of 10 % of the Daily Value for vitamins A, C and D, riboflavin, niacin and calcium, and 20 % of the Daily Value for thiamin, vitamins B₆ and B₁₂, folic acid, iron and zinc. Values are percentages with their standard errors indicated by vertical bars

chronic disease outcomes⁽⁴⁶⁾. Therefore, fortification of RTE cereal should be encouraged and the fortification profile of RTE cereal should be based on current nutrition science, current patterns of food intake, consumer preference and government policy.

RTE cereal fortification requires particular consideration because it is consumed by a wide range of age groups from young children to older adults and contributes differently to overall nutrient intakes in these groups. From our results in the NHANES data, among toddlers and children, RTE cereal is consumed by a larger proportion of the population (40 and 46 % on a given day, respectively) but at a lower total amount (21 and 31 g/d, respectively) than by adolescents and adults, where less of the population consumes RTE cereal (20 and 27 % on a given day, respectively) but in higher quantities (~45 g/d). Below we describe how the optimization scenario presented within the present study impacted the micronutrient status for the overall population and within these specific age groups.

Vitamin D and Ca are nutrients of public health concern among all age groups in the USA and vitamin C is an

under-consumed nutrient^(6,47). The DV for all three of these nutrients increased under NLR: for vitamin D from 10 to 20 µg (400 to 800 IU); for Ca from 1000 to 1300 mg; and for vitamin C from 60 to 90 mg. RTE cereal is the third highest dietary contributor to vitamin D intake in children and adolescents, contributing 8–10 % of daily vitamin D intake⁽⁴⁾, and among older adults, contributing 7 %⁽⁴⁸⁾. There are very few food sources of vitamin D and the 2015–2020 Dietary Guidelines for Americans recommends consuming foods fortified with vitamin D, including RTE cereal, as a strategy to achieve higher levels of vitamin D intake. Thus, it is important that RTE cereal continues to be an important source of these nutrients. In our optimized scenario, the %DV remained the same for vitamin D, Ca and vitamin C, although the amount per 100 g did change due to the changing DV and RACC: the amount of vitamin D and vitamin C increased by 50 and 13 %, respectively, and the amount of Ca in RTE cereal remained similar (decreased by 2 % compared with the baseline amount). Indeed, under the optimized scenario, no changes in the prevalence of intakes below



the EAR were observed for toddlers, and a slight (1–2 pp) increase for both Ca and vitamin D among child, adolescent and adult RTE cereal consumers. However, it is important to note that a large percentage of the population is still under-consuming these nutrients although RTE cereal consumers are more likely to consume adequate levels⁽⁸⁾.

Fe is considered a shortfall nutrient among adolescent girls and women⁽⁶⁾. Fe was also identified as a priority nutrient in the 2017 Food and Nutrition Board report for the diets of infants and children participating in the Special Supplemental Nutrition Program for Women, Infants, and Children (WIC). WIC provides supplemental foods, including RTE cereal, to low-income pregnant, breast-feeding and postpartum women and to infants and children up to 5 years of age who are found to be at nutritional risk⁽²⁹⁾. Because RTE cereal is the primary source of added Fe in the diets of US children⁽⁴⁾, the optimized scenario sought to determine the amount of Fe that continued to help children and women meet needs, without potential excess in some other population groups, such as adult men.

Despite modelling a change in Fe content for the entire category of RTE cereal, it is important to consider that certain RTE cereals provide a higher amount of Fe (28 mg/100 g RTE cereal *v.* 9 mg/100 g RTE cereal in the optimized scenario) and these amounts are unlikely to change in a 'real-world' setting due to WIC programme specifications. Fe intake of WIC participants is higher and the percentage below the EAR is lower compared with income-eligible and income-ineligible non-participants, suggesting the important impact and reach of Fe-fortified cereals among infants and young toddlers (6–24 months)⁽¹⁸⁾.

The optimized scenario represents a proposed reduction in Fe of 40% in RTE cereal. When we modelled removing Fe fortification from RTE cereal, the percentage below the EAR increased from 3.5% at baseline for the total population aged ≥ 1 year to 6.1%. Among RTE cereal eaters, this increase was from 0% below the EAR to 7.4%. The optimized scenario attenuated this increase in percentage below the EAR for Fe to 4.4% for the total population and 0.5% for RTE cereal eaters. Our scenarios did not include the use of supplemental Fe, which increases the likelihood of intakes above the UL for Fe, especially in children⁽¹⁶⁾. Therefore, the optimized scenario represents a conservative approach to fortification that limits the impact on the percentage of the population below the EAR.

Previous reports have identified intakes above the UL for vitamin A, niacin, folic acid and Zn as potentially of concern, especially in some US age and sex groups^(16,17,42,43,49,50). Because RTE cereal is a key contributor of all four of these nutrients in children, the optimized scenario sought to balance adequate intakes

among all age groups while lowering the percentage of the population above the UL.

Retinol is the form of preformed vitamin A that the UL is based upon. RTE cereal fortification provides approximately 35 and 17% of vitamin A to the diets of child (2–18 years) and adult (≥ 19 years) consumers, respectively⁽⁸⁾. While few adults exceed the UL for vitamin A from foods alone^(42,51), many young children do (19% of all toddlers in the current study), which could be more of a concern among supplement users. Other sources of preformed vitamin A include liver, fish oils, dairy products and eggs. While the %DV for vitamin A remained the same in the optimized scenario as at baseline, a 55% decrease in the vitamin A content of RTE cereal would occur due to the changing DV and RTE cereal RACC. This decrease in vitamin A content translated to a 60% reduction in the prevalence of RTE cereal consumers exceeding the UL among toddlers in the optimized scenario compared with baseline (33% baseline to 13% in optimized). However, at the same time, we found that there was a large percentage of the total population below the EAR for vitamin A particularly among adolescents and adults (10% for children and approximately 48% for adolescents and adults). Our optimized scenario minimized the increase in the percentage below the EAR when compared with complete removal of vitamin A fortification, highlighting the importance of retaining vitamin A fortification in RTE cereal.

Niacin content in RTE cereal was decreased by 76% from baseline to the optimal fortification level due to a combination of reducing the %DV, a decrease in the DV and an increase in the RACC of RTE cereal. Typical American diets, with or without fortified foods, tend to provide niacin at levels congruent with the DRI⁽³⁾ since niacin is present in a wide variety of foods including beef, poultry and fish. While the primary safety concern of high doses of niacin is flushing, which is largely benign, it is still relevant to consider lowering the niacin content given that 16% of the US population consuming RTE cereal is achieving intakes above the UL. The large reduction of niacin per 100 g of RTE cereal slightly shifted the percentage below the EAR at the population level (~ 0.5 pp) and among RTE cereal consumers (~ 1 pp), but the reduction in intakes above the UL was more substantial, particularly among RTE cereal eaters (i.e. from 16% above UL at baseline to 1% in the optimized scenario).

Folic acid is the synthetic form of the vitamin folate that is used in fortification and dietary supplements and is the only form of folate that is considered for the UL. The role of folic acid has been established for the prevention of neural tube defects and fortification programmes are mandated in over seventy countries worldwide⁽⁵²⁾. Fortification programmes have increased dietary intakes of folic acid⁽⁵³⁾ and blood folate concentrations in the USA⁽⁵⁴⁾ and reduced the incidence of neural



tube birth defects dramatically. We noted only a very small percentage of the total population aged ≥ 1 year above the UL for folic acid (0.3%); even among RTE cereal eaters, only 4% of the population ≥ 1 year of age was above the UL although toddlers (28.9% above the UL) and children (7.6% above the UL) who eat RTE cereal were more likely to exceed the UL. The optimized framework delineated in this project, which modelled an 82% decrease in folate/folic acid in RTE cereal from baseline, represents a balance between reducing the percentage of the population exceeding the UL (particularly for toddlers) while minimizing the impact on the percentage of the population below the EAR. For example, there was still a substantial percentage of the population, particularly for adolescents (30%) and adult (20%) women in the total population, who were below the EAR. The optimized scenario reduces the percentage above the UL for toddlers who eat RTE cereal (to $< 1\%$ above the UL) while minimizing the impact on the total population of adolescent and adult women below the EAR.

We found that very few adolescents and no adults exceeded the UL for Zn from food, although the percentage of toddlers (60%) and children (4%) was higher. However, the percentage below the EAR for Zn was notable in the total population for adolescents (26%) and adults (19%). In the optimized scenario, a 56% decrease in the amount of Zn added to RTE cereal was proposed, substantially lowering the prevalence of intakes above the UL for toddlers (to 45%) and children (to 1.4%) with minimal changes in intakes below the EAR. Among RTE cereal eaters specifically, high Zn intake remained a concern for toddlers (57% above the UL); however, it should be pointed out that when we modelled removing all fortification of RTE cereal, 30% of toddlers who eat RTE cereal would have intakes above the UL. Without Zn fortification some 36% of adolescents and 25% of adults who eat RTE cereal would not meet the EAR, thus it is important to maintain fortification of Zn in RTE cereal. Other food sources of Zn include meat and poultry.

In addition to folic acid and niacin, the remaining B-vitamin contents (i.e. thiamin, riboflavin, B₆ and B₁₂) of RTE cereal were also reduced in the optimized scenario; this is salient as RTE cereal is the primary contributor to added amounts of these nutrients in the diets of children and adolescents⁽⁴⁾. Other sources include meat, poultry and fish (thiamin, riboflavin, vitamin B₆, vitamin B₁₂), enriched grains (thiamin, riboflavin), potatoes (vitamin B₆) and dairy products (riboflavin and vitamin B₁₂). No toddlers or children, and a small fraction of adults, who consume RTE cereal had intakes for these nutrients in the baseline scenario below the EAR, and only a small proportion (2–5%) of adolescents and adults had intakes below the EAR with the optimized scenario. Thiamin,

riboflavin, and vitamin B₁₂ do not have UL established and the percentage above the UL for vitamin B₆ was 0% for all age groups.

The majority of the available DRI were set between 1997 and 2005, while Ca and vitamin D were updated in 2011⁽²⁸⁾. The UL for children have been criticized as having been established with too few available data and are considered to be too low for many nutrients⁽⁵⁵⁾. Indeed, for Zn, approximately 40% of US infants and children exceed the UL from diet alone. While total vitamin A intakes exceeded the UL in a large proportion of infants and toddlers, it should also be noted that substantial numbers exceeded the UL for vitamin A from foods alone. Thus, the UL for vitamin A and Zn should be considered for re-evaluation, since so many children exceed them and there is little evidence of clinically significant health consequences. Across all ages, the UL for niacin has similarly been called into question⁽⁵⁶⁾. The range of intakes between the EAR and the UL is much narrower in children than in adults, making it challenging to achieve adequate intakes in the population without exceeding the UL. Furthermore, many of the DRI values in children were extrapolated from studies and data in adults. While the current DRI guidelines represent the best-known data on this subject and therefore are the best information we have for setting public health policies related to nutrient intakes in these age groups, evaluation of the appropriateness of these cut-offs should be pursued in future research. Ongoing efforts to improve the DRI process with a focus on disease prevention are currently being explored⁽⁵⁷⁾.

A number of strengths and limitations exist within the current analysis. First, we propose a proactive scientific approach to fortification to ensure public health adequacy balanced with potential concerns of intakes above the UL. The use of nationally representative survey data permits estimates of changes in fortification practices that are national in scope and we estimated usual or habitual dietary intakes of Americans to reduce within-person variation in nutrient intakes. Nevertheless, self-reported dietary data are known to contain systematic bias including energy under-reporting^(58,59) and our models assume that reported consumption of RTE cereal is unbiased, which may result in classification bias. The optimized scenario jointly modelled the upcoming changes to the serving sizes (i.e. the RACC) and the DV under NLR, representing the most 'real life' and informed scenario possible in interpreting our findings⁽⁶⁰⁾. However, the proposed framework for optimizing RTE cereals made some assumptions that should be considered in interpreting these data. First, the model assumes all RTE cereals would be changed to match the proposed scenario, which is speculative but also reasonable to assume. None of models represent intakes of nutrients from dietary supplements purposefully to understand

the unique contribution of RTE cereal, but in doing so overestimates the population proportion for intakes below the EAR and underestimates the proportion with intakes above the UL^(16,42,43) for those who use nutrient-containing dietary supplements^(16,42). Because the models only influence RTE consumers, we present results both for RTE cereal eaters only and for the total population, the results of which would represent the overall public health impact of changing RTE cereal fortification and considers the prevalence of RTE cereal consumption.

In summary, the present work represents an attempt to optimize the amounts of vitamins and minerals that are provided in RTE cereal given the important role that RTE cereal plays helping Americans meet their dietary needs, either directly or through its frequent consumption with milk. The most notable feature of the optimized scenario presented is balancing the decrease in the amounts of those nutrients that may contribute to intakes above the UL in some population subgroups, such as toddlers, while continuing to provide key under-consumed nutrients to the general population, of which RTE cereal is an important contributor. Moving forward, we hope that this framework may inform decisions about fortification practices in this and across other food categories, to ensure that food manufacturers are providing products that meet the contemporary needs of various population subgroups and are using an evidence-based approach to informing fortification practices^(36,61).

Acknowledgements

Acknowledgements: The authors would like to acknowledge the contribution of Nort Holschuh, employee of General Mills, Inc., for providing statistical expertise. **Financial support:** Funding of this project was provided by the Bell Institute of Health and Nutrition, General Mills, Inc. J.D.S. and N.J. are employed by General Mills, Inc. and they contributed to the study design, the analysis of the data, the interpretation of findings and the preparation of the manuscript. **Conflict of interest:** J.D.S. and N.J. are employees of General Mills, Inc. R.L.B. is a scientific consultant to the Bell Institute of Health and Nutrition. **Authorship:** J.D.S. contributed to the study design and statistical analysis plan, carried out the study, guided the data analyses and contributed to the writing of the manuscript. N.J. conducted the statistical analysis and contributed to the writing of the manuscript. R.L.B. provided expertise and guidance on the study design and statistical analysis plan and wrote the manuscript. **Ethics of human subject participation:** The NHANES survey protocol and procedures are approved by the US Centers for Disease Control and Prevention's National Center for Health Statistics. This paper is a secondary data analysis of NHANES and no personally identifiable information about individuals was available to the researchers.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1368980019003690>

References

1. Institute of Medicine (2003) *Dietary Reference Intakes: Guiding Principles for Nutrition Labeling and Fortification*. Washington, DC: The National Academies Press.
2. Bailey RL, West KP Jr & Black RE (2015) The epidemiology of global micronutrient deficiencies. *Ann Nutr Metab* **66**, Suppl. 2, 22–33.
3. Fulgoni VL 3rd, Keast DR, Bailey RL *et al.* (2011) Foods, fortificants, and supplements: where do Americans get their nutrients? *J Nutr* **141**, 1847–1854.
4. Berner LA, Keast DR, Bailey RL *et al.* (2014) Fortified foods are major contributors to nutrient intakes in diets of US children and adolescents. *J Acad Nutr Diet* **114**, 1009–1022.e8.
5. Institute of Medicine (2006) *Dietary Reference Intakes: The Essential Guide to Nutrient Requirements*. Washington, DC: The National Academies Press.
6. US Department of Health and Human Services & US Department of Agriculture (2015) 2015–2020 Dietary Guidelines for Americans, 8th ed. <http://health.gov/dietaryguidelines/2015/guidelines/> (accessed August 2019).
7. Health Canada (2019) History of Canada's food guides from 1942 to 2007. <https://www.canada.ca/content/dam/hc-sc/documents/services/food-nutrition/canada-food-guide/resources/evidence/food-nutrients-health-interim-evidence-update-2018/26-18-2165-History%20of%20CFG-EN-06.pdf> (accessed August 2019).
8. Fulgoni VL & Buckley RB (2015) The contribution of fortified ready-to-eat cereal to vitamin and mineral intake in the US population, NHANES 2007–2010. *Nutrients* **7**, 3949–3958.
9. Michels N, De Henauw S, Breidenassel C *et al.* (2015) European adolescent ready-to-eat-cereal (RTEC) consumers have a healthier dietary intake and body composition compared with non-RTEC consumers. *Eur J Nutr* **54**, 653–664.
10. Priebe MG & McMonagle JR (2016) Effects of ready-to-eat cereals on key nutritional and health outcomes: a systematic review. *PLoS One* **11**, e0164931.
11. Balvin Frantzen L, Trevino RP, Echon RM *et al.* (2013) Association between frequency of ready-to-eat cereal consumption, nutrient intakes, and body mass index in fourth- to sixth-grade low-income minority children. *J Acad Nutr Diet* **113**, 511–519.
12. Hill KM, Jonnalagadda SS, Albertson AM *et al.* (2012) Top food sources contributing to vitamin D intake and the association of ready-to-eat cereal and breakfast consumption habits to vitamin D intake in Canadians and United States Americans. *J Food Sci* **77**, H170–H175.
13. Yeung LF, Cogswell ME, Carriquiry AL *et al.* (2011) Contributions of enriched cereal-grain products, ready-to-eat cereals, and supplements to folic acid and vitamin B-12 usual intake and folate and vitamin B-12 status in US children: National Health and Nutrition Examination Survey (NHANES), 2003–2006. *Am J Clin Nutr* **93**, 172–185.
14. Albertson AM, Affenito SG, Bauserman R *et al.* (2009) The relationship of ready-to-eat cereal consumption to nutrient intake, blood lipids, and body mass index of children as they age through adolescence. *J Am Diet Assoc* **109**, 1557–1565.
15. Albertson AM, Wold AC & Joshi N (2012) Ready-to-eat cereal consumption patterns: the relationship to nutrient intake, whole grain intake, and body mass index in an older American population. *J Aging Res* **2012**, 631310.



16. Bailey RL, Fulgoni VL 3rd, Keast DR *et al.* (2012) Do dietary supplements improve micronutrient sufficiency in children and adolescents? *J Pediatr* **161**, 837–842.
17. Bailey RL, Catellier DJ, Jun S *et al.* (2018) Total usual nutrient intakes of US children (under 48 months): findings from the Feeding Infants and Toddlers Study (FITS) 2016. *J Nutr* **148**, 9S, 1557S–1566S.
18. Jun S, Catellier DJ, Eldridge AL *et al.* (2018) Usual nutrient intakes from the diets of US children by WIC participation and income: findings from the Feeding Infants and Toddlers Study (FITS) 2016. *J Nutr* **148**, 9S, 1567S–1574S.
19. Centers for Disease Control and Prevention, National Center for Health Statistics (2018) National Health and Nutrition Examination Survey. NHANES 2013–2014. <https://www.nchs.gov/nchs/nhanes/ContinuousNhanes/Default.aspx?BeginYear=2013> (accessed June 2019).
20. Johnson CL, Dohrmann SM, Burt VL *et al.* (2014) National health and nutrition examination survey: sample design, 2011–2014. *Vital Health Stat 2* issue 162, 1–33.
21. Moshfegh AJ, Rhodes DG, Baer DJ *et al.* (2008) The US Department of Agriculture Automated Multiple-Pass Method reduces bias in the collection of energy intakes. *Am J Clin Nutr* **88**, 324–332.
22. Blanton CA, Moshfegh AJ, Baer DJ *et al.* (2006) The USDA Automated Multiple-Pass Method accurately estimates group total energy and nutrient intake. *J Nutr* **136**, 2594–2599.
23. US Department of Health and Human Services & Food and Drug Administration, Center for Food Safety and Applied Nutrition (2013) A Food Labeling Guide: Guidance for Industry. <https://www.fda.gov/downloads/Food/GuidanceRegulation/GuidanceDocumentsRegulatoryInformation/UCM265446.pdf> (accessed June 2019).
24. US Department of Health and Human Services & Food and Drug Administration (2016) Food Labeling: Serving Sizes of Foods That Can Reasonably Be Consumed at One Eating Occasion; Dual-Column Labeling; Updating, Modifying, and Establishing Certain Reference Amounts Customarily Consumed; Serving Size for Breath Mints; and Technical Amendments. *Fed Regist* **81**, 34000–340467.
25. US Department of Agriculture, Agricultural Research Service (2014) USDA Food and Nutrient Database for Dietary Studies 2013–2014. <https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/> (accessed June 2019).
26. Institute of Medicine (1998) *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline*. Washington, DC: The National Academies Press.
27. Institute of Medicine (2000) *Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium, and Carotenoids*. Washington, DC: The National Academies Press.
28. Institute of Medicine (2011) *Dietary Reference Intakes for Calcium and Vitamin D*. Washington, DC: The National Academies Press.
29. US Department of Agriculture, WIC Works Resource System (2017) WIC Report: Guidelines for Feeding Healthy Infants. <https://wicworks.fns.usda.gov/resources/guidelines-feeding-healthy-infants> (accessed June 2019).
30. Beaton G, Milner J, Corey P *et al.* (1979) Sources of variance in 24-hour dietary recall data: implications for nutrition study design and interpretation. *Am J Clin Nutr* **32**, 2546–2559.
31. Beaton G, Milner J, McGuire V *et al.* (1983) Source of variance in 24-hour dietary recall data: implications for nutrition study design and interpretation. Carbohydrate sources, vitamins, and minerals. *Am J Clin Nutr* **37**, 986–995.
32. Kipnis V, Midthune D, Freedman L *et al.* (2002) Bias in dietary-report instruments and its implications for nutritional epidemiology. *Public Health Nutr* **5**, 915–923.
33. Dodd K, Guenther P, Freedman L *et al.* (2006) Statistical methods for estimating usual intake of nutrients and foods: a review of the theory. *J Am Diet Assoc* **106**, 1640–1650.
34. Tooze J, Midthune D, Dodd K *et al.* (2006) A new statistical method for estimating the usual intake of episodically consumed foods with application to their distribution. *J Am Diet Assoc* **106**, 1575–1587.
35. Cox DR (1982) Combination of data. In *Encyclopedia of Statistical Sciences*, vol. 2, pp. 45–53 [S Kotz and NL Johnson, editors]. New York: Wiley.
36. Dwyer JT, Wiemer KL, Dary O *et al.* (2015) Fortification and health: challenges and opportunities. *Adv Nutr* **6**, 124–131.
37. Malek AM, Newman JC, Hunt KJ *et al.* (2019) Race/ethnicity, enrichment/fortification, and dietary supplementation in the US population, NHANES 2009–2012. *Nutrients* **11**, E1005.
38. Sacco JE, Dodd KW, Kirkpatrick SI *et al.* (2013) Voluntary food fortification in the United States: potential for excessive intakes. *Eur J Clin Nutr* **67**, 592–597.
39. Sacco JE & Tarasuk V (2011) Discretionary addition of vitamins and minerals to foods: implications for healthy eating. *Eur J Clin Nutr* **65**, 313–320.
40. Sacco JE & Tarasuk V (2009) Health Canada’s proposed discretionary fortification policy is misaligned with the nutritional needs of Canadians. *J Nutr* **139**, 1980–1986.
41. Bailey RL (2020) Current regulatory guidelines and resources to support research of dietary supplements in the United States. *Crit Rev Food Sci Nutr* **60**, 298–309.
42. Bailey RL, Fulgoni VL 3rd, Keast DR *et al.* (2012) Examination of vitamin intakes among US adults by dietary supplement use. *J Acad Nutr Diet* **112**, 657–663.e4.
43. Bailey RL, Fulgoni VL 3rd, Keast DR *et al.* (2011) Dietary supplement use is associated with higher intakes of minerals from food sources. *Am J Clin Nutr* **94**, 1376–1381.
44. Bruins MJ, Mugambi G, Verkaik-Kloosterman J *et al.* (2015) Addressing the risk of inadequate and excessive micronutrient intakes: traditional versus new approaches to setting adequate and safe micronutrient levels in foods. *Food Nutr Res* **59**, 26020.
45. Food and Drug Administration (1980) Nutritional quality of foods; addition of nutrients. *Fed Regist* **45**, 6314–6324.
46. Xu M, Huang T, Lee AW *et al.* (2016) Ready-to-eat cereal consumption with total and cause-specific mortality: prospective analysis of 367,442 individuals. *J Am Coll Nutr* **35**, 217–223.
47. Bailey RL, Dodd KW, Goldman JA *et al.* (2010) Estimation of total usual calcium and vitamin D intakes in the United States. *J Nutr* **140**, 817–822.
48. Papanikolaou Y & Fulgoni VL (2017) Grain foods are contributors of nutrient density for American adults and help close nutrient recommendation gaps: data from the National Health and Nutrition Examination Survey, 2009–2012. *Nutrients* **9**, E873.
49. Bailey RL, Dodd KW, Gahche JJ *et al.* (2010) Total folate and folic acid intake from foods and dietary supplements in the United States: 2003–2006. *Am J Clin Nutr* **91**, 231–237.
50. Bailey RL, McDowell MA, Dodd KW *et al.* (2010) Total folate and folic acid intakes from foods and dietary supplements of US children aged 1–13 y. *Am J Clin Nutr* **92**, 353–358.
51. Blumberg JB, Balz FB, Fulgoni VL *et al.* (2017) Impact of frequency of multi-vitamin/multi-mineral supplement intake on nutritional adequacy and nutrient deficiencies in US adults. *Nutrients* **9**, E849.
52. Bailey LB, Stover PJ, McNulty H *et al.* (2015) Biomarkers of nutrition for development – folate review. *J Nutr* **145**, issue 7, 1636S–1680S.
53. Dietrich M, Brown CJ & Block G (2005) The effect of folate fortification of cereal-grain products on blood folate status, dietary folate intake, and dietary folate sources among adult non-supplement users in the United States. *J Am Coll Nutr* **24**, 266–274.



54. Pfeiffer CM, Johnson CL, Jain RB *et al.* (2007) Trends in blood folate and vitamin B-12 concentrations in the United States, 1988–2004. *Am J Clin Nutr* **86**, 718–727.
55. Zlotkin S (2006) A critical assessment of the upper intake levels for infants and children. *J Nutr* **136**, issue 2, 502S–506S.
56. Minto C, Vecchio MG, Lamprecht M *et al.* (2017) Definition of a tolerable upper intake level of niacin: a systematic review and meta-analysis of the dose-dependent effects of nicotinamide and nicotinic acid supplementation. *Nutr Rev* **75**, 471–490.
57. Yetley EA, MacFarlane AJ, Greene-Finestone LS *et al.* (2017) Options for basing Dietary Reference Intakes (DRIs) on chronic disease endpoints: report from a joint US-/Canadian-sponsored working group. *Am J Clin Nutr* **105**, issue 1, 249S–285S.
58. Subar AF, Freedman LS, Tooze JA *et al.* (2015) Addressing current criticism regarding the value of self-report dietary data. *J Nutr* **145**, 2639–2645.
59. Subar AF, Kipnis V, Troiano RP *et al.* (2003) Using intake biomarkers to evaluate the extent of dietary misreporting in a large sample of adults: the OPEN study. *Am J Epidemiol* **158**, 1–13.
60. Murphy MM, Spungen JH, Barraj LM *et al.* (2013) Revising the daily values may affect food fortification and in turn nutrient intake adequacy. *J Nutr* **143**, 1999–2006.
61. Dwyer JT, Woteki C, Bailey R *et al.* (2014) Fortification: new findings and implications. *Nutr Rev* **72**, 127–141.