




Dietary patterns are associated with premenstrual syndrome: evidence from a case-control study

Bahareh MoradiFili¹, Reza Ghiasvand^{1,*} , Makan Pourmasoumi², Awat Feizi³, Farnaz Shahdadian¹ and Zahra Shahshahan⁴

¹Department of Community Nutrition, School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan, Iran; ²Gastrointestinal and Liver Diseases Research Center, Guilan University of Medical Sciences, Rasht, Iran; ³Department of Biostatistics and Epidemiology, School of Public Health, Isfahan University of Medical Sciences, Isfahan, Iran; ⁴Department of Obstetrics and Gynecology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Submitted 4 October 2018: Final revision received 24 April 2019: Accepted 14 May 2019: First published online 15 October 2019

Abstract

Objectives: Premenstrual syndrome (PMS) is a common cyclic psychological and somatic disorder which reduces women's quality of life. Evidence regarding the association between dietary patterns (DPs) and PMS is rare. The study aimed to determine the relationship between dietary patterns and PMS.

Design: The case-control study was conducted among women with confirmed PMS and healthy individuals recruited from healthcare centres.

Setting: Dietary data were collected using a validated semi-quantitative food frequency questionnaire and DPs were derived using principal component analysis. The association between DPs and likelihood of PMS was determined using logistic regression.

Participants: In total, 225 women with PMS and 334 healthy participants aged 20–46 years took part in the study.

Results: Three major DPs were identified: (i) 'western DP' characterized by high intake of fast foods, soft drink, and processed meats; (ii) 'traditional DP' in which eggs, tomato sauce, fruits, and red meat were highly loaded; and (iii) 'healthy DP' high in dried fruits, condiments and nuts. After taking all possible confounders into account, individuals in the highest tertile of the western DP were more likely to experience PMS (odds ratio (OR) = 1.49; 95% CI: 1.01, 3.52), $P < 0.001$, whilst both healthy and traditional DP was inversely associated with the syndrome (OR = 0.31; 95% CI: 0.17, 0.72, $P = 0.02$; OR = 0.33; 95% CI: 0.14, 0.77, $P = 0.01$, respectively).

Conclusion: The western dietary patterns were positively associated with PMS, whilst the healthy and traditional dietary patterns were inversely associated with it. Further longitudinal studies are required to confirm our findings.

Keywords

Dietary patterns
Premenstrual syndrome
Women
Principal component analysis
Case-control study

Premenstrual syndrome (PMS) is a common cyclic psychological and somatic disorder affecting a large number of women, which consistently occurs during the luteal phase of the menstrual cycle⁽¹⁾. It is characterized by emotional, behavioural and physical symptoms and can manifest with a wide variety of symptoms, including depression, mood liability, irritability, breast tenderness and sleep disturbances⁽²⁾. It has been indicated that 47.8% of women are diagnosed with PMS and the highest prevalence has been reported from Iran, approximately 98% (95% CI: 97, 100)⁽³⁾. The high prevalence and great impact on quality

of life of the women affected leaves PMS as an unresolved yet important issue in women's health. The pathology of PMS has not yet been fully-understood. It is proposed that the PMS results from ovulation and may be due to ovarian steroid interactions relating to neurotransmitter dysfunction⁽¹⁾. The potential contributors include central nervous system (CNS) sensitivity to reproductive hormones, genetic background, nutritional indices and psychosocial factors^(4,5). In this regard, nutritional factors are the most modifiable parameters that can be taken into account as a strategy in PMS management.

*Corresponding author: Email ghiasvand@hlth.mui.ac.ir

Little information is available regarding particular nutrients, foods or dietary patterns that might be associated with this syndrome. It seems that the excess or deficiency of certain nutrients might be due to hormonal and neurotransmitter imbalances⁽⁶⁾. Furthermore, certain foods, such as vegetables, and dietary patterns including low-fat and high-fibre diets are related to decreased plasma oestrogen level and the duration of premenstrual symptoms⁽⁷⁾. Furthermore, studies assessing food-intake have revealed that the excess consumption of sweet-tasting food items, fast foods, deep-fried meals, coffee and alcohol are significantly related to the development of PMS⁽⁷⁾.

According to our knowledge, studies regarding the association of empirically derived dietary patterns and the risk of PMS are limited. Conceptually, assessing the association between dietary patterns and the likelihood of PMS is very close to reality, because foods are consumed as part of a whole diet. On the other hand, dietary patterns might be different based on cultures, geography and beliefs of different populations^(8,9). Limited data are available regarding the association between major dietary patterns and risk of PMS in Middle Eastern countries⁽¹⁰⁾. The purpose of the current study was to investigate the association between major dietary patterns and premenstrual syndrome among Iranian women aged 20–45 years in the context of a case-control study.

Methods

Study population

The current case-control study was conducted among women with PMS and healthy women as a control group referring to health centres in 2017. The study population consisted of 225 women with PMS and 334 healthy controls aged 20–45 years. The PMS patients included in the present study were those with confirmed diagnosis of moderate/severe PMS using the Premenstrual Symptoms Screening Tool (PSST), Body Mass Index (BMI) less than 39.9 kg/m², without specific habits such as smoking, alcoholism, any chronic disease, specific diet, medication therapy or vitamins and minerals supplementation. On the other hand, participants who did not respond to more than 35 items of the Food Frequency Questionnaire (FFQ) or reported total daily energy intake outside the range of 3347–17 572 kJ (800–4200 kcal) were excluded from the analysis. The control group consisted of age-matched women without premenstrual symptoms according to the PSST questionnaire. Similar exclusion criteria applied for the PMS case were used for the control group.

Written informed consent was completed for each individual. The study protocol was approved by the regional bioethics committee (Project number: 395518, Ethical approve number: IR.MUI.REC.1395.3.518). This study is reported based on Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines⁽¹¹⁾.

Data collection

Dietary intake assessment

Dietary intake was assessed using a validated, self-administered, semi-quantitative food-frequency questionnaire (FFQ). Participants reported the frequency of 168 different food and beverage items consumed during the last year. The reliability and validity of the FFQ has been previously reported in the Iranian population⁽¹²⁾. Trained nutritionists filled in the FFQ through face-to-face interviews. Participants were requested to report the frequency of usual food items consumed in the preceding year on a daily, weekly or monthly basis, considering the standard portion size for each food item. The frequency of foods consumed was converted to gram per day using standard household measures. Subsequently, energy and macro and micronutrient intake was determined using the US Department of Agriculture food composition database using the Nutritionist IV programme. We categorized the food items into 36 predefined food groups in order to derive the dietary patterns (Table 1).

Case confirmation

In the current study, PMS was confirmed by the Premenstrual Symptoms Screening Tool (PSST) questionnaire. This tool is a simple, user-friendly screening tool with high reliability and validity for identifying women with Premenstrual Syndrome (PMS)⁽¹³⁾ and comprises nineteen items subdivided into two domains: the first domain consists of fourteen items related to premenstrual symptoms, whilst the second domain is composed of five items that assess the functional impact of premenstrual symptoms.

In order to capture the criteria of premenstrual symptoms, the following question was asked 'Do you experience some or any of the following premenstrual symptoms, which start before your period and stop within a few days of bleeding?' The symptoms listed were anger/irritability, anxiety/tension, tearful, depressed mood/hopelessness, decreased interest at work, decreased interest at home, decreased interest in social activities, difficulty in concentrating, fatigue/lack of energy, overeating/food cravings, insomnia, hypersomnia, feeling overwhelmed, and physical symptoms (breast tenderness, headaches, joint/muscle pain, bloating, weight gain). In order to assess the functional impact of the premenstrual symptoms, women were asked 'Have your symptoms interfered with the following five domains: relationship with your family, work efficiency or productivity, relationship with co-workers, social life activities or home responsibilities'⁽¹⁴⁾. Each item was rated according to a 4-point Likert scale (0 = absent; 1 = mild; 2 = moderate; 3 = severe). In order to detect moderate to severe PMS, the following three conditions must have qualified. (1) From items 1 to 4, at least one of them must be moderate or severe. (2) At least four items of 1 to 14 should be moderate or severe. (3) In addition to the

Table 1 Food groups used in the principal component analysis

Food groups	Subgroups
Processed meat	Sausage, Kielbasa
Red meat	Lamp, Beef, Ground meat
Organ meats	Heart, Liver and Kidney, (Beef) Lamb-tongue, (Beef) Lamb-brain, (Beef) Lamb tripe, Meat of sheep head, Leg of lamb
Fish	Fish, Canned fish
Skinless poultry	Chicken no skin
Poultry-skin	Chicken with skin
Eggs	Egg
Low fat dairy	Low fat milk, Without fat milk, Yogurt, Kashk, Cheese
High fat dairy	Milk high fat, Cocoa milk, Chocolate milk, Yogurt-plain-whole milk, Cream, Ice cream, Cheese-cream
Yogurt drink	Doogh*
Soft drink	Soft drink
Tea	Tea
Coffee	Coffee
Fruits	Cantaloupe, Melon, Watermelon, Pear, Apricot, Cherry, Apple, Peach, Nectarine plum, Fig, Grape, Kiwi, Grapefruit, Orange, Persimmon, Tangerine, Pomegranate, Date, Black cherry, Strawberry, Banana, Sweet lemon, Lemon Cranberry, Pineapple, Raisin, Mulberry
Natural juices	Grapefruit juice, Orange juice, Apple juice, Melon juice
Canned fruits	Canned fruits, Canned pineapple
Dried fruits	Dried fig, Dried peach, Dried apricot, Dried mulberry
Vegetables	Pea, String Bean, vegetable, Pumpkin, Squash, Eggplant, Celery, Cucumber, Lettuce, Tomato, Cauliflower, Sweet peppers, Raw, Raw carrot, Cooked carrot, Raw onion, Fried onion, Cabbage, Mushroom, Maize spinach, Cooked spinach, Turnip, Cooked
Nuts	Seeds, Walnuts pistachios, Hazelnuts, Almonds. (Pumpkin, Sunflower, Watermelon)
Legumes	Lentils, Beans, Chick pea, Broad bean, Soya, Bean-mung, Pea
Whole grains	Barbari bread, Sangak bread, Taftoon bread, Cooked barley, Oat
Refined grains	Lavash bread, Baguettes, Toast, Rice, Spaghetti Vermicelli, Pasta, Wheat flour, Biscuits,
Fast foods	Fried potato, Hamburger, Pizza
Mayonnaise	Mayonnaise
Tomato sauce	Tomato sauce
Salty snacks	Crackers, Cheese puffs, Chips
Olive	Olive seed, Olive oil
Sugar-sweets-desserts	Cake, Sugar, Honey, Jam, Sweets, Gaz, Candy Sohan, Chocolate, Caramel, Halva, Donuts,
Hydrogenated fats	Solid oil, Fat animal, Animal oil, Butter, Margarine
Vegetables oils	Vegetable oil
Potato	Cooked potato
Garlic	Garlic
Condiments	Juice lime, Pepper black
Salt	Salt
Pickles	Pickles
Broth	Broth

*Doogh is a cold yogurt-based beverage.

previous conditions, among the last five items, at least one of them must be moderate or severe.

Assessment of other variables

Other required information such as age (years), marital status (married/single or widow), education (lower than diploma/diploma (12 years of formal education)/Basic Sciences/post-graduate education), socio-economic status and medication use were obtained using a pretested questionnaire.

Socio-economic status (SES) of the study participants was evaluated based on the following binary variables: education level of both subjects and the family head (academic/non-academic), job of both subjects and the family head (yes/no), family size (fewer than 4/higher than 4), having a car (yes/no), home status (the owner/tenant), and having foreign travel (yes/no) by using self-reported questionnaire, and then a total score was calculated.

Furthermore, physical activity was examined using the International Physical Activity Questionnaire (IPAQ) and expressed as metabolic equivalent-min/week. The validity and reliability of IPAQ has been previously determined⁽¹⁵⁾. Participants were asked to recall their type of physical activity and its intensity as well as the duration of activity for the past 7 d. To estimate total daily physical activity level, respective metabolic equivalents (MET) values (for instance, walking = 3.3, moderate-intensity activity = 4.0, vigorous-intensity activity = 8.0) were multiplied by duration time (min), and then summed. Data obtained from the physical activity questionnaire were expressed as daily metabolic equivalents (MET-min/d). Afterwards, continuous values were categorized into high, moderate, and mild levels⁽¹⁶⁾.

Participants' weights were measured using a balanced digital scale (SECA, Model no: 710, Germany) to the nearest 0.05 kg. Height was recorded using a wall-mounted tape-meter. Height was measured in a standing position with

a tape measure to the nearest 0.5 cm in the normal position. Participant's weight and height were measured with light clothing and barefoot. Body mass index was calculated as weight (kg) divided by the square of height (m²). Hip circumference was measured with a tape measure to the nearest 1 mm at the widest point between hip and buttock. Waist circumference was also measured to the nearest 1 mm at the midway level between the lower rib margin and the iliac crest using a steel tape measure, and waist-hip ratio (WHR) was estimated by dividing waist circumference (cm) to hip circumference (cm).

Statistical analysis

Exploratory factor analysis with Principal component analysis (PCA) as the extraction method was used for extracting the dietary patterns from 36 food groups consumed. Factor analysis is a statistical method used to describe variability among observed, correlated variables (in the current study, food groups) in terms of a potentially lower number of unobserved variables, called factors (in this study; dietary patterns). Factor analysis searches for such joint variations in response to unobserved latent variables (dietary pattern). The observed variables (food groups) are modelled as linear combinations of the potential factors, plus 'error' terms⁽¹⁷⁾.

Scree plot and eigenvalue greater than 1.5 was used for determining the appropriate number of factors. An orthogonal rotation method – i.e. Varimax – was used for improving the interpretability of extracted factors. Finally, three factors were considered as the major dietary patterns and were labelled based on the interpretation of the highly loaded food groups in each factor. For each participant, factor scores of dietary patterns were calculated by using a regression method in which, by summing the intake of food groups weighted by factor loading, each participant received a factor score for each identified dietary pattern. KMO and Bartlett's tests were used to evaluate the adequacy of sample size and the applicability factor analysis method on our data. Participants were categorized into tertiles based on each dietary pattern's score. To assess the association between dietary patterns and the chance for developing PMS, the binary logistic regression was used in crude and different multivariable adjusted models. An adjustment for age, marital status, education and socioeconomic status (SES) was done in the first model. Further adjustments were conducted for BMI and physical activity in the second model. Energy plus those factors mentioned in the second model were further adjusted in the third model. The first tertile of the dietary patterns scores was considered as the reference category in the logistic regression models.

Quantitative data were compared between case and control groups and across tertiles of dietary patterns using independent t-test, one-way ANOVA and the analysis of covariance (ANCOVA) with Bonferroni correction. The

Chi-square test was also used for comparing categorical data between the case and control groups. Statistical analyses were performed using statistical package for social sciences (SPSS) software version 16.0 (SPSS Inc., Chicago, IL, USA). Also, *P*-values less than 0.05 was considered statistically significant.

Result

Major dietary patterns

The three major dietary patterns identified according to the factor loading are (Table 2): (i) 'western dietary pattern', which was highly loaded by fast foods, soft drinks, processed meats, salt, salty snacks, sugar-sweets-desserts, organ meat, broth, poultry-skin, hydrogenated fats, mayonnaise, high fat dairy, vegetables oil, tea and red meat; (ii) 'traditional dietary pattern', which was high in eggs, tomato sauce, fruits, red meats, refined grains, vegetables, pickles, coffee, natural juices, mayonnaise, skinless poultry, vegetable oils, salty snacks, whole grains and high fat dairy; (iii) 'healthy dietary pattern', which was high in dried fruits, condiments, nuts, canned fruits, legumes, vegetables,

Table 2 Factor loadings matrix for dietary patterns identified by using factor-analysis

Food groups	Western pattern	Healthy pattern	Traditional pattern
Fast foods	0.71	–	–
Soft drink	0.68	–	–
Processed meats	0.67	–	–
Salt	0.54	–	–
Salty snacks	0.52	–	0.22
Sugar-sweets-desserts	0.5	–	–
Organ meats	0.49	–	–
Poultry-skin	0.44	–	0.24
Broth	0.44	–	–
Mayonnaise	0.43	–	0.3
Hydrogenated fats	0.43	–	–
High fat dairy	0.26	–	0.2
Vegetables oils	0.25	–	0.22
Tea	0.2	–	0.18
Red meats	–	–	0.44
Fish	–	0.3	–
Skinless poultry	–	–	0.25
Eggs	–	–	0.57
Yogurt drink	–	0.23	–
Coffee	–	–	0.35
Fruits	–	0.36	0.46
Natural juices	–	0.33	0.33
Canned fruits	–	0.4	–
Dried fruits	–	0.54	–
Vegetables	–	0.39	0.4
Nuts	–	0.48	–
Legumes	–	0.39	–
Whole grains	–	–	0.21
Refined grains	–	–	0.42
Tomato sauce	–	–	0.53
Olive	–	0.47	–
Potato	–	–	0.11
Garlic	–	0.38	–
Condiments	–	0.52	–
Pickles	–	0.26	0.36

Loading factors less than 0.2 are not reported for simplicity.

Table 3 General characteristics of participants with and without premenstrual syndrome

Variables	Controls (n 334)			Cases (n 225)			P-value*
	Mean	SD	%	Mean	SD	%	
Age (year)	28.90	6.92		27.96	6.22		0.09
Body mass index (kg/m ²)	22.04	2.54		24.80	2.57		<0.001
Waist-circumference (cm)	78.41	7.34		85.39	9.04		<0.001
Hip-circumference (cm)	101.06	7.76		109.47	9.90		<0.001
Waist to hip ratio	0.77	0.07		0.78	0.06		0.58
Physical activity (MET-min/week)	1806.97	1459.64		403.18	334.50		<0.001
Socio-economic status (score)	3.12	1.43		3.36	1.40		0.05
Marital status							
Married	180		53.9	86		38.3	
Widowed/Divorced	11		3.3	30		13.3	<0.001
Single	143		42.8	109		48.4	
Education							
Lower than diploma	38		11.4	15		6.7	
High school diploma (12 year of formal education)	100		29.9	52		32.1	0.04
Bachelor of sciences	136		40.7	111		49.3	
Postgraduate	60		18	47		20.9	

Values are presented as mean \pm standard deviation (SD) for continuous and number (%) for categorical variables.

*Resulted from independent samples t-test for quantitative variables and chi-square test for categorical data.

garlic, fruits, natural juices, fish, pickles and yogurt drink. In total, the identified dietary patterns explained 44.8% of dietary food intakes in this population.

General characteristics and dietary intake of participants

According to the exclusion criteria, 559 subjects (225 cases and 334 controls) remained for the current analysis. The general characteristics of the study population are shown in Table 3. Individuals in the case group were more likely to be overweight, single, and highly educated, compared with those in the control group. Also, they had greater waist-circumference and hip-circumference and less physical activity ($P < 0.05$).

The comparison of the demographic and lifestyle factors of the study population in different levels of the three major dietary patterns is summarized in Table 4. Distribution of participants in terms of age, weight, physical activity level, waist, hip, WHR, BMI, SES marital status and education was significantly different across western dietary pattern categories. In addition, there were significant differences in WHR, physical activity, marital status and education in the healthy dietary pattern category. Furthermore, across different levels of the traditional dietary pattern, participants were significantly different according to age, weight, waist, hip, WHR, BMI and marital status ($P < 0.05$). In comparison to those in the lowest tertile, individuals in the highest tertile of the western dietary pattern were younger, and had higher mean BMI, waist-circumference, hip-circumference and less physical activity. Comparison of food items in the healthy dietary pattern showed that individuals in the highest tertile were more likely to have physical activity and less WHR. Furthermore, individuals with the highest adherence to traditional dietary pattern were older, and

had higher BMI, waist-circumference, hip-circumference and WHR ($P < 0.05$). Also, 81.4% of participants with PMS and 18.6% of healthy subjects were adherent to the western dietary pattern ($P < 0.05$).

Table 5 illustrates the comparison of macro and micro-nutrients intake in different categories of major dietary patterns. Greater adherence compared with lower adherence to western dietary pattern was significantly associated with higher intake of energy, fat, saturated fatty acid (SFA), MUFA, PUFA, linoleic fatty acid, linolenic fatty acid, oleic fatty acid and Vitamin E, and lower intake of iron, Calcium, Magnesium, Vitamin B1, Vitamin B6, Folate, Vitamin C, and Vitamin D. In addition, participants in the different tertiles of the healthy dietary pattern had significantly different intakes of energy, fat, MUFA, PUFA, linoleic fatty acid, linolenic fatty acid, oleic fatty acid, DHA, Iron, Calcium, Vitamin E, Vitamin B6, Vitamin C, and Vitamin D.

A similar comparison of nutrients intake in the traditional dietary pattern category showed a prominent difference in terms of energy, fat, SFA, MUFA, PUFA, linoleic fatty acid, linolenic fatty acid, oleic fatty acid, Magnesium, Vitamin E, Vitamin B6, Vitamin C and Vitamin D intake.

The relationship between dietary patterns and PMS

Crude and multivariable adjusted odds ratio with 95% CI are represented in Table 6. In all fitted models, the lowest categories of adherence to dietary patterns (tertile 1) was defined as the reference. Those in the top tertile of adherence to western dietary pattern had significantly, 1.77 times, more chance of developing PMS compared with those in the bottom tertile of the crude model (OR: 1.77; 95% CI: 1.05, 2.93). After adjusting for potential important confounding variables, there was a positive significant

Table 4 General characteristics of the study participants across tertiles of the major dietary patterns

Variable	Western dietary pattern									Healthy dietary pattern									Traditional dietary pattern											
	T1			T2			T3			P*	T1			T2			T3			P*	T1			T2			T3			P*
	Mean	SD	%	Mean	SD	%	Mean	SD	%		Mean	SD	%	Mean	SD	%	Mean	SD	%		Mean	SD	%	Mean	SD	%	Mean	SD	%	
Age (year)	29.80	7.14		28.63	6.78		27.15	5.84		<0.001	29.02	7.05		28.22	6.30		28.33	6.69		0.46	28.03	6.16		27.56	6.68		29.98	6.97		0.001
Weight (Kg)	59.52	8.67		60.36	8.43		64.93	8.16		<0.001	61.27	9.23		61.95	8.64		61.58	8.35		0.75	58.79	8.74		62.53	8.31		63.49	8.49		0.001
Body mass index (kg/m ²)	22.37	2.88		22.77	2.82		24.34	2.58		<0.001	23.18	3.18		23.31	2.75		23	2.70		0.59	22.26	2.82		23.33	2.59		23.89	2.99		<0.001
Waist-circumference (cm)	79.43	8.61		80.68	8.61		83.66	8.59		<0.001	82.28	9.50		81.19	8.65		80.23	8.04		0.10	78.66	8.19		81.31	8.27		83.8	9.10		<0.001
Hip-circumference (cm)	101.34	7.88		102.73	8.68		109.53	9.94		<0.001	103.79	10.19		104.84	9.25		104.98	9.21		0.43	102.74	8.83		105.21	9.65		105.65	9.94		0.007
Waist to hip ratio	0.78	0.08		0.78	0.06		0.76	0.04		0.002	0.79	0.08		0.77	0.06		0.76	0.05		<0.001	0.76	0.07		0.77	0.06		0.79	0.06		<0.001
Physical activity (MET-min/week)	1286.64	1234		1635.53	1421		738.22	1116		<0.001	1103.68	1103		978.45	1181		1577.26	1545		<0.001	1322.23	1279		1239.76	1335		1098.59	1325		0.25
Socio economic status	3.42	1.43		2.85	1.38		3.37	1.39		<0.001	3.01	1.37		3.30	1.52		3.35	1.35		0.05	3.10	1.44		3.24	1.45		3.32	1.37		0.34
Marital status																														
Married	101	55.5	91	49.7	68	37.2	0.007	90	49.5	74	40.4	96	52.5	0.04	97	53.3	65	35.5	98	53.6	0.001									
Widow	13	7.1	14	7.7	13	7.1		13	7.1	20	10.9	7	3.8		14	7.7	11	6	15	8.2										
Single	68	37.4	78	42.6	102	55.7		79	43.4	89	48.6	80	43.7		71	39	107	58.5	70	38.3										
Education																														
Lower than diploma	13	7.1	25	13.7	12	6.6	<0.001	18	9.9	24	13.1	8	4.4	<0.001	22	12.1	10	5.5	18	9.8	0.20									
High school diploma	39	21.4	67	36.6	41	22.4		68	37.4	39	21.3	40	21.9		44	24.2	50	27.3	53	29										
Basic sciences	85	46.7	64	35	96	52.5		71	39	84	45.9	90	49.2		82	45.1	91	49.7	72	39.3										
Post graduate	45	24.7	27	14.8	34	18.6		45	24.7	36	19.7	45	24.6		34	18.7	32	17.5	40	21.9										
Premenstrual syndrome																														
No	146	80.2	146	79.8	34	18.6	<0.001	105	57.7	93	50.8	128	69.9	<0.001	123	67.6	112	61.2	91	49.7	0.002									
Yes	36	19.8	37	20.2	149	81.4		77	42.3	90	49.2	55	30.1		59	32.4	71	38.8	92	50.3										

Values are presented as mean ± standard deviation (SD) for continuous and number (%) for categorical variables.

*Result from Analysis of Variance for quantitative variables and chi-square test for categorical data.



Table 5 Comparison of macro- and micro-nutrients intake across tertiles of major dietary patterns

	Western dietary pattern						P*	Healthy dietary pattern						P*	Traditional dietary pattern						P*
	T1		T2		T3			T1		T2		T3			T1		T2		T3		
	Mean	SE	Mean	SE	Mean	SE		Mean	SE	Mean	SE	Mean	SE		Mean	SE	Mean	SE	Mean	SE	
Energy (kJ)†	8945.27	44.84	9338.69	44.44	11 317.72	44.76	<0.001	9282.20	47.06	9656.67	46.90	10 665.01	46.89	<0.001	8442.77	41.92	9442.74	41.92	11 677.63	42.11	<0.001
Protein (g)	89.05	1.44	81.58	1.41	77.08	1.49	<0.001	79.51	1.45	83.01	1.43	85.14	1.45	0.024	78.42	1.52	83.53	1.44	85.71	1.57	0.005
Carbohydrate (g)	360.07	3.87	354.40	3.78	319.27	3.99	<0.001	352.63	3.97	336.80	3.93	344.27	3.98	0.018	346.47	4.15	332.83	3.93	354.36	4.30	0.001
Fat (g)	67.19	1.65	74.24	1.61	92.35	1.70	<0.001	75.02	1.76	81.56	1.74	77.08	1.77	0.031	79.09	1.83	82.83	1.74	71.93	1.90	<0.001
Saturated fatty acid (g)	21.31	0.83	24.03	0.81	28.73	0.85	<0.001	24.88	0.83	25.97	0.83	23.24	0.84	0.07	24.62	0.87	26.61	0.83	22.86	0.91	0.011
Mono-unsaturated fatty acid (g)	21.14	0.54	23.94	0.52	29.28	0.55	<0.001	23.62	0.57	25.49	0.57	25.26	0.58	0.043	25.78	0.59	26.21	0.56	22.48	0.61	<0.001
Poly-unsaturated fatty acid (g)	15.22	0.54	17.27	0.53	23.26	0.56	<0.001	17.08	0.58	19.94	0.57	18.74	0.58	0.002	18.67	0.60	20.22	0.57	16.88	0.63	0.001
Linoleic fatty acid (g)	18.39	0.56	19.76	0.55	23.33	0.58	<0.001	19.37	0.57	20.68	0.56	21.44	0.57	0.039	20.96	0.60	21.34	0.56	19.19	0.62	0.04
Linolenic fatty acid (g)	12.76	0.53	15.01	0.52	20.71	0.54	<0.001	14.82	0.56	17.30	0.56	16.37	0.56	0.008	16.36	0.59	17.77	0.56	14.37	0.61	<0.001
Eicosapentaenoic acid (g)	0.003	0.002	0.006	0.002	0.007	0.002	0.455	0.007	0.002	0.007	0.002	0.002	0.002	0.167	0.004	0.002	0.006	0.002	0.006	0.002	0.733
Docosahexaenoic acid (g)	0.010	0.004	0.022	0.004	0.011	0.005	0.112	0.023	0.004	0.14	0.004	0.006	0.004	0.017	0.012	0.005	0.012	0.004	0.019	0.005	0.517
Iron (mg)	18.81	0.42	19.42	0.41	16.75	0.44	<0.001	19.30	0.42	17.28	0.41	18.40	0.42	0.003	18.78	0.44	18.16	0.42	18.04	0.46	0.486
Calcium (mg)	1198.16	23.80	1063.90	23.25	736	24.49	<0.001	948.15	26.61	976.73	26.34	1071.81	26.7	0.003	956.45	28.13	990.66	26.68	1049.62	29.15	0.096
Magnesium (mg)	348.21	18.96	399.53	18.53	262.70	19.51	<0.001	348.16	19	317.85	18.81	344.42	19.07	0.463	400.25	19.73	315.62	18.71	294.85	20.44	0.001
Zinc (mg)	11.25	0.28	11.04	0.27	10.35	0.28	0.082	10.89	0.27	10.63	0.27	11.12	0.27	0.46	11.12	0.28	11.01	0.27	10.52	0.30	0.354
Vitamin E (mg)	10.13	0.60	8.61	0.59	17.75	0.62	<0.001	12.39	0.64	13.54	0.64	10.39	0.65	0.003	9.57	0.67	13.82	0.63	12.92	0.69	<0.001
Vitamin B ₁ (mg)	2.06	0.03	1.93	0.03	1.66	0.03	<0.001	1.95	0.03	1.85	0.03	1.84	0.03	0.059	1.87	0.03	1.83	0.03	1.94	0.03	0.106
Vitamin B ₆ (mg)	2.07	0.04	1.89	0.04	1.63	0.04	<0.001	1.73	0.04	1.83	0.04	2.02	0.04	<0.001	1.73	0.04	1.84	0.04	2.01	0.04	<0.001
Folate (mcg)	372.19	9.48	389.17	9.26	318.64	9.75	<0.001	375.03	9.49	345.54	9.39	359.45	9.52	0.087	362.19	9.99	348.23	9.48	369.52	10.35	0.298
Vitamin B ₁₂ (mcg)	4.34	0.97	4.29	0.94	7.12	0.99	0.083	4.16	0.95	6.62	0.94	4.97	0.95	0.175	4.81	1.00	6.41	0.95	4.54	1.03	0.332
Vitamin C (mg)	201.42	8.46	176.80	8.26	88.70	8.70	<0.001	122.62	8.59	142.47	8.51	201.41	8.62	<0.001	144.29	9.30	141.05	8.82	181.27	9.64	0.007
Vitamin D (mcg)	2.74	0.12	1.72	0.12	1.76	0.13	<0.001	1.73	0.12	2.18	0.12	2.03	0.12	0.036	1.58	0.13	2.05	0.12	2.31	0.13	0.001

Values are reported as means ± standard errors (SEs).

*P-values were obtained by using analysis of covariance; adjustment was made for age and energy intake.

†Adjusted for age.

Dietary pattern and premenstrual syndrome

Table 6 Crude and multivariable adjusted odds ratios (95% CI for OR) premenstrual syndrome across tertiles of dietary patterns

	Western dietary pattern						Healthy dietary pattern						Traditional dietary pattern					
	T1	T2	95% CI	T3	95% CI	P for trend	T1	T2	95% CI	T3	95% CI	P for trend	T1	T2	95% CI	T3	95% CI	P for trend
Crude	1	1.02	0.61, 1.71	1.77	1.05, 2.93	<0.001	1	0.96	0.72, 1.29	0.43	0.31, 0.58	<0.001	1	1.32	0.86, 2.03	2.11	1.37, 3.22	0.002
Model 1*	1	1.22	0.69, 2.14	2.39	1.34, 4.26	<0.001	1	1.15	0.75, 1.79	0.54	0.34, 0.85	0.002	1	1.27	0.81, 2.00	2.25	1.43, 3.55	0.001
Model 2†	1	1.43	0.67, 3.05	1.92	1.03, 4.42	<0.001	1	0.93	0.48, 1.80	0.59	0.29, 1.17	0.205	1	0.58	0.29, 1.16	0.96	0.47, 1.97	0.222
Model 3‡	1	1.36	0.63, 2.94	1.49	1.01, 3.52	<0.001	1	0.75	0.38, 1.49	0.31	0.17, 0.72	0.02	1	0.45	0.22, 0.95	0.33	0.14, 0.77	0.017

*Model 1: adjusted for age, marital status, education, and socio-economic status.

†Model 2: adjusted for age, marital status, education, socio-economic status, BMI and physical activity.

‡Model 3: adjusted for age, marital status, education, socio-economic status, BMI, physical activity and energy intake.

relationship observed between the chance of PMS and adherence to western dietary pattern in the first model (OR: 2.39; 95 % CI: 1.34, 4.26), second model (OR: 1.92; 95 % CI: 1.03, 4.42), and after adjustment for all confounding variables (OR: 1.49; 95 % CI: 1.01, 3.52).

On the contrary, individuals in the highest tertile of adherence to healthy dietary pattern were 43 % less likely to have odds of PMS compared with those in the lowest category in the crude model (OR: 0.43; 95 % CI: 0.31, 0.58). This association remained significant even after taking potential confounders into account in model 1 (OR: 0.54; 95 % CI: 0.34, 0.85), and model 3 (OR: 0.31; 95 % CI: 0.17, 0.72). But there was insignificant association in model 2 (OR: 0.59; 95 % CI: 0.29, 1.17).

Furthermore, a similar finding for traditional dietary pattern was shown in the crude model (OR: 2.11; 95 % CI: 1.37, 3.22). In model 1, after adjustment for different confounders, those in the top category had 2.25 times higher odds of PMS (OR: 2.25; 95 % CI: 1.43, 3.55). However, these effects were the reverse in model 2 (OR: 0.96; 95 % CI: 0.47, 1.97) and model 3 after adjustment for all confounding variables (OR: 0.33; 95 % CI: 0.14, 0.77).

Discussion

In the current study three major dietary patterns including western, healthy and traditional dietary habits were identified. Among the dietary patterns obtained, the western dietary pattern – which was characterized by high loading of fast foods, soft drink, processed meats, salt, salty snacks, sugar-sweets-desserts, organ meat, broth, poultry-skin, hydrogenated fats, mayonnaise, high fat dairy, vegetables oil, tea and red meat – was significantly associated with a higher likelihood of PMS. In contrast, we found a significant association between a healthy dietary pattern high in dried fruits, condiments, nuts, canned fruits, legumes, vegetables, garlic, fruits, natural juices, fish, pickles and yogurt drink and PMS in the multi-variable adjusted models. In addition, individuals with highest adherence to the traditional dietary pattern high in eggs, tomato sauce, fruits, red meats, refined grains, vegetables, pickles, coffee, natural juices, mayonnaise, skinless poultry, vegetables oils, salty snacks, whole grains and high fat dairy also had lower odds for PMS.

Determining dietary patterns allows holistic identification of diet and disease relations and provides further information beyond associations found by the analysis of individual foods and nutrients. To our knowledge, studies evaluating the association of dietary patterns and PMS risk are limited. Previous studies have mostly focused on the relationship between particular foods or nutrients and PMS. They have supported the positive relationship between unhealthy food items, which are highly loaded in the western dietary pattern, such as fast foods⁽¹⁸⁾, and sweet foods and beverages and PMS⁽¹⁹⁾. In contrast, certain

healthy food items such as fish might be inversely associated with PMS induced by athletic disturbance⁽²⁰⁾. Likewise, specific nutrients such as omega-3 fatty acids have been positively related to the relief of psychiatric and somatic symptoms of PMS⁽²¹⁾.

The results of the current study with 559 participants demonstrated that habitual dietary patterns might contribute to the risk of PMS, despite the complex and vague pathogenesis of PMS. Our findings are in concordance with the results obtained from a previous study⁽⁷⁾, which found that the western dietary pattern was associated with an increased risk of PMS. However, it did not identify any significant relationship between either the healthy or traditional dietary patterns and PMS. The emerging evidence has found that higher adherence to the western dietary pattern is associated with inadequate nutrient intake; on the other hand, reports have revealed an association between risk of PMS and low level of nutrients^(5,22). The western dietary pattern is also positively related to higher level of inflammatory biomarkers, while healthy diets with high antioxidant content, vitamins and phytochemicals results in reduced inflammatory indices^(23,24). On the other hand, PMS symptoms have been positively associated with some inflammatory markers⁽²⁵⁾. Therefore, the effect of western dietary pattern on the risk of PMS might be linked to the induction of inflammation. In addition, the imbalance of oxidant/antioxidant systems may be due to consumption of an unhealthy diet, which eventually enhances the symptoms of PMS^(26,27). Moreover, adherence to unhealthy dietary patterns might have obesogenic results⁽²⁸⁾, which may contribute to the development of PMS symptoms⁽²⁹⁾.

As mentioned before, the earlier study that evaluated the relationship between dietary patterns and PMS did not show any relationship between both the healthy or traditional dietary patterns and PMS⁽⁷⁾. Participants in the top tertile of the healthy eating plan were 57 % and 46 % less likely to have odds of PMS compared with those in the bottom tertile in the crude and first model, respectively. Interestingly, such associations remained significant in the third model. This finding highlights the beneficial role of healthy food items associated with PMS symptoms.

In the current study, protective association of traditional dietary pattern against PMS was also observed. Although the traditional dietary pattern was shown to be a risk factor for PMS in the crude model, this effect was reversed in the final adjusted model. Therefore, it seems that adjusted factors have a higher contribution to the risk of PMS. Taken together, despite a few unhealthy food items such as red meat, the traditional dietary pattern could be considered as a healthy diet for PMS development.

The current study has several strengths. Although studies examining the association between dietary patterns and PMS are rare, the current study is one of the comprehensive studies with a large number of participants in this field. Second, subjects in this study were selected from individuals referring to healthcare centres, with different



socio-economic status. Thus, variability in women's dietary intakes could cover a wide range of eating habits. In addition, we were able to consider a wide range of confounders for adjustment to reach an independent association in the current study.

There are a number of limitations that should be taken into account. Like all case-control studies, the inherent methodological limitations of such studies make it impossible to draw a causal link between dietary patterns and the risk of PMS. Although case-control studies are efficient in terms of studying the association between diet and disease, the selection and recall biases of such study design must be kept in mind. An additional concern is that women diagnosed with PMS change their diet choices, and this might change the risk estimates. Furthermore, although we assessed energy-adjusted intake of all food groups for extracting dietary pattern scores, the possibility of subject misclassification is unavoidable when using FFQ to assess the long-term dietary intakes.

In conclusion, our results indicated that the western dietary pattern – which includes fast foods, soft drink, processed meats, salt, salty snacks, sugar-sweets-desserts, organ meat, broth, poultry-skin, hydrogenated fats, mayonnaise, high fat dairy, vegetables oil, tea and red meat – might be associated with increased odds of PMS. In addition, a protective association of both healthy and traditional dietary pattern against PMS was observed. This study provides important insights into the association between diet and the risk of PMS, and shows that changing diet quality and avoiding unhealthy diets might have desirable effects in reducing the risk of PMS symptoms. However, further studies are needed to confirm these associations and provide the evidence needed to translate these findings into clinical practice.

Acknowledgements

The authors would like to offer their gratitude all the women who kindly participated in this study. **Financial support:** This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors. **Conflict of interest:** The authors declare no personal or financial conflicts of interest. **Authorship:** B.M.F., R.G., F.S. and A.F. made substantial contributions to the conception, design, acquisition of data, analysis and interpretation of data and drafting of this study. B.M.F., A.F. and Z.S. participated in drafting the article or revising it. R.G. and M.P. gave final approval of the version to be submitted and any revised version. R.G. is guarantor of this work. **Ethics of human subject participation:** This study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects/patients were approved by the ethics committee of Isfahan University of medical sciences (Ethical approval number: IR.MUI.REC.

1395.3.518). Written informed consent was obtained from all subjects/patients.

References

1. Liu Q, Wang Y, van Heck CH *et al.* (2017) Stress reactivity and emotion in premenstrual syndrome. *Neuropsychiatr Dis Treat* **13**, 1597–1602.
2. Naheed B, Kuiper JH, Uthman OA *et al.* (2017) Non-contraceptive oestrogen-containing preparations for controlling symptoms of premenstrual syndrome. *Cochrane Database Syst Rev* **3**, Cd010503.
3. Direkvand-Moghadam A, Sayehmiri K, Delpisheh A *et al.* (2014) Epidemiology of Premenstrual Syndrome (PMS)-A systematic review and meta-analysis study. *J Clin Diagn Res* **8**, 106.
4. Hantsoo L & Epperson CN (2015) Premenstrual dysphoric disorder: epidemiology and treatment. *Curr Psychiatry Rep* **17**, 87.
5. Saeedian Kia A, Amani R & Cheraghian B (2015) The association between the risk of premenstrual syndrome and Vitamin D, calcium, and magnesium status among University students: a case control study. *Health Promot Perspect* **5**, 225–230.
6. Stachowicz M & Lebidzińska A (2016) The effect of diet components on the level of cortisol. *Eur Food Res Technol* **242**, 2001–2009.
7. Farasati N, Siassi F, Koohdani F *et al.* (2015) Western dietary pattern is related to premenstrual syndrome: a case-control study. *Br J Nutr* **114**, 2016–2021.
8. Kant AK, Graubard BI & Schatzkin A (2004) Dietary patterns predict mortality in a national cohort: the National Health Interview Surveys, 1987 and 1992. *J Nutr* **134**, 1793–1799.
9. Hu FB (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* **13**, 3–9.
10. Mirmiran P, Bahadoran Z, Vakili AZ *et al.* (2017) Western dietary pattern increases risk of cardiovascular disease in Iranian adults: a prospective population-based study. *Appl Physiol Nutr Metab* **42**, 326–332.
11. von Elm E, Altman DG, Egger M *et al.* (2008) The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* **61**, 344–349.
12. Mirmiran P, Esfahani FH, Mehrabi Y *et al.* (2010) Reliability and relative validity of an FFQ for nutrients in the Tehran lipid and glucose study. *Public Health Nutr* **13**, 654–662.
13. Hariri FZ, Moghaddam-Banaem L, Siah Bazi S *et al.* (2013) The Iranian version of the Premenstrual Symptoms Screening Tool (PSST): a validation study. *Arch Women Ment Health* **16**, 531–537.
14. Steiner M, Maccougall M & Brown E (2003) The premenstrual symptoms screening tool (PSST) for clinicians. *Arch Women Ment Health* **6**, 203–209.
15. Craig CL, Marshall AL, Sjörström M *et al.* (2003) International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc* **35**, 1381–1395.
16. Noori MA, Ghiasvand R, Maghsoudi Z *et al.* (2016) Evaluation of dietary pattern stability and physical activity in three consecutive generations of women. *Int J Public Health* **61**, 29–38.
17. Bartholomew DJ, Steele F, Galbraith J *et al.* (2008) *Analysis of Multivariate Social Science Data*. New York: Chapman and Hall/CRC.
18. Isik H, Ergol S, Aynioglu O *et al.* (2016) Premenstrual syndrome and life quality in Turkish health science students. *Turk J Med Sci* **46**, 695–701.
19. Rossignol A & Bonnländer H (1991) Prevalence and severity of the premenstrual syndrome. Effects of foods and beverages

- that are sweet or high in sugar content. *J Reprod Med* **36**, 131–136.
20. Takeda T, Imoto Y, Nagasawa H *et al.* (2016) Fish consumption and premenstrual syndrome and dysphoric disorder in Japanese collegiate athletes. *J Pediatr Adolesc Gynecol* **29**, 386–389.
 21. Sohrabi N, Kashanian M, Ghafoori SS *et al.* (2013) Evaluation of the effect of omega-3 fatty acids in the treatment of premenstrual syndrome: “a pilot trial”. *Complement Ther Med* **21**, 141–146.
 22. Serra-Majem L, Bes-Rastrollo M, Roman-Vinas B *et al.* (2009) Dietary patterns and nutritional adequacy in a Mediterranean country. *Br J Nutr* **101 Suppl 2**, S21–28.
 23. Barbaresko J, Koch M, Schulze MB *et al.* (2013) Dietary pattern analysis and biomarkers of low-grade inflammation: a systematic literature review. *Nutr Rev* **71**, 511–527.
 24. Bonaccio M, Cerletti C, Iacoviello L *et al.* (2015) Mediterranean diet and low-grade subclinical inflammation: the Moli-sani study. *Endocr Metab Immune Disord Drug Targets* **15**, 18–24.
 25. Bertone-Johnson ER, Ronnenberg AG, Houghton SC *et al.* (2014) Association of inflammation markers with menstrual symptom severity and premenstrual syndrome in young women. *Hum Reprod* **29**, 1987–1994.
 26. Miller ER, 3rd, Erlinger TP, Sacks FM *et al.* (2005) A dietary pattern that lowers oxidative stress increases antibodies to oxidized LDL: results from a randomized controlled feeding study. *Atherosclerosis* **183**, 175–182.
 27. Duvan CI, Cumaoglu A, Turhan NO *et al.* (2011) Oxidant/antioxidant status in premenstrual syndrome. *Arch Gynecol Obstet* **283**, 299–304.
 28. Jessri M, Wolfinger RD, Lou WY *et al.* (2017) Identification of dietary patterns associated with obesity in a nationally representative survey of Canadian adults: application of a priori, hybrid, and simplified dietary pattern techniques. *Am J Clin Nutr* **105**, 669–684.
 29. Masho SW, Adera T & South-Paul J (2005) Obesity as a risk factor for premenstrual syndrome. *J Psychosom Obstet Gynaecol* **26**, 33–39