

Clinical efficacy of endovascular treatment approach in patients with carotid cavernous fistula: A systematic review and meta-analysis[☆]



Aryoobarzan Rahmatian^a, Shirin Yaghoobpoor^b, Arian Tavasol^b, Komeil Aghazadeh-Habashi^c, Zahra Hasanabadi^d, Matin Bidares^e, Borna Safari-kish^e, Robert M. Starke^f, Evan M. Luther^f, Mohammadreza Hajiesmaeili^g, Fatemeh Sodeifian^b, Tara Fazel^h, Mina Dehghaniⁱ, Reza Ramezan^j, Masood Zangi^k, Niloofar Deravi^b, Reza Goharani^{l,**}, Mobina Fathi^{b,*}

^a Faculty of Medicine, Ilam University of Medical Sciences, Ilam, Iran

^b Student Research Committee, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^c Student Research Committee, Tabriz University of Medical Sciences, Tabriz, Iran

^d Medical Faculty, Qazvin University of Medical Science, Qazvin, Iran

^e Clinical Research Development Center, Najafabad Branch, Islamic Azad University, Najafabad, Iran

^f Department of Neurosurgery, University of Miami, Miami, FL, USA

^g Department of Anesthesiology and Critical Care, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^h School of International Campus, Guilan University of Medical Sciences, Rasht, Iran

ⁱ School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

^j Department of Statistics and Actuarial Science, University of Waterloo, Waterloo, Ontario, Canada

^k Anesthesiology Research Center, Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

^l Department of Anesthesiology and Critical Care, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Keywords:

CCF
Exophthalmos
Trauma
Diplopia
Proptosis
Visual decline
Chemosis

ABSTRACT

Background and objectives: Carotid-cavernous fistulas (CCFs) represent a group of rare, abnormal arteriovenous communications between the carotid arterial system and the cavernous sinuses (CS). CCFs often produce ophthalmologic symptoms related to increased CS pressures and retrograde venous drainage of the eye. Although endovascular occlusion remains the preferred treatment for symptomatic or high-risk CCFs, most of the data for these lesions is limited to small, single-center series. As such, we performed a systematic review and meta-analysis evaluating endovascular occlusions of CCFs to determine any differences in clinical outcomes based on presentation, fistula type, and treatment paradigm.

Method: A retrospective review of all studies discussing the endovascular treatment of CCFs published through March 2023 was conducted using PubMed, Scopus, Web of Science, and Embase databases. A total of 36 studies were included in the meta-analysis. Data from the selected articles were extracted and analyzed using Stata software version 14.

Results: 1494 patients were included. 55.08% were female and the mean age of the cohort was 48.10 years. A total number of 1516 fistulas underwent endovascular treatment, 48.05% of which were direct and 51.95% of which were indirect. 87.17% of CCFs were secondary to a known trauma while 10.18% were spontaneous. The most common presenting symptoms were 89% exophthalmos (95% CI: 78.0–100.0; $I^2 = 75.7\%$), 84% chemosis (95% CI: 79.0–88.0; $I^2 = 91.6\%$), 79% proptosis (95% CI: 72.0–86.0; $I^2 = 91.8\%$), 75.0% bruits (95% CI: 67.0–82.0;

Abbreviations: Carotid-Cavernous Fistulas, (CCFs); Cavernous Sinuses, (CS); Computed Tomography, (CT); Ethylene-Vinyl Alcohol Copolymer, (EVOH) (onyx); Internal Carotid Artery, (ICA); n-Butyl CyanoAcrylate, (n-BCA).

[☆] Approved by ethics committee (IR.SBMU.RETECH.REC.1399.083).

* Corresponding author.

** Corresponding author.

E-mail addresses: rahmatian-a@medilam.ac.ir (A. Rahmatian), sh.yaghoobpoor98@gmail.com (S. Yaghoobpoor), ariantavasol1378@gmail.com (A. Tavasol), komeil.aghazade@gmail.com (K. Aghazadeh-Habashi), z.hasanabadi42@gmail.com (Z. Hasanabadi), matin.bidares@gmail.com (M. Bidares), borna.safari.93@gmail.com (B. Safari-kish), RStarke@med.miami.edu (R.M. Starke), evan.luther@jhsmiami.org (E.M. Luther), mrhajiesmaeili@sbmu.ac.ir (M. Hajiesmaeili), Ftmhsodeifian@yahoo.com (F. Sodeifian), tarafazel7899@gmail.com (T. Fazel), minadehghani2001@gmail.com (M. Dehghani), rramezan@uwaterloo.ca (R. Ramezan), Masood_zangi@yahoo.com (M. Zangi), niloofar.deravi@gmail.com (N. Deravi), r.goharani@sbmu.ac.ir (R. Goharani), Mobina.fathi78@gmail.com (M. Fathi).

<https://doi.org/10.1016/j.wnsx.2023.100189>

Received 18 November 2022; Received in revised form 18 March 2023; Accepted 21 March 2023

2590-1397/© 2023 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

$I^2 = 90.7\%$), 56% diplopia (95% CI: 42.0–71.0; $I^2 = 92.3\%$), 49% cranial nerve palsy (95% CI: 32.0–66.0; $I^2 = 95.1\%$), 39% visual decline (95% CI: 32.0–45.0; $I^2 = 71.4\%$), 32% tinnitus (95% CI: 6.0–58.0; $I^2 = 96.7\%$), 29% elevated intraocular pain (95% CI: 22.0–36.0; $I^2 = 0.0\%$), 31% orbital or pre-orbital pain (95% CI: 14.0–48.0; $I^2 = 89.9\%$) and 24% headache (95% CI: 13.0–34.0; $I^2 = 74.98\%$). Coils, balloons, and stents were the three most used embolization methods respectively. Immediate complete occlusion of the fistula was seen in 68% of cases and complete remission was seen in 82%. Recurrence of CCF occurred in only 35% of the patients. Cranial nerve paralysis after treatment was observed in 7% of the cases.

Conclusions: Exophthalmos, Chemosis, proptosis, bruits, cranial nerve palsy, diplopia, orbital and periorbital pain, tinnitus, elevated intraocular pressure, visual decline and headache are the most common clinical manifestations of CCFs. The majority of endovascular treatments involved coiling, balloons and onyx and a high percentage of CCF patients experienced complete remission with the improvement of their clinical symptoms.

1. Introduction

Carotid-Cavernous Fistulas (CCFs) are abnormal arteriovenous shunts between the carotid system and the cavernous sinuses (CS) resulting in arterialized blood within the venous structures draining the orbit.¹ This frequently causes abnormally increased venous pressures within the CS and orbital veins and can lead to various symptoms such as proptosis, glaucoma, elevated intraocular pressure, cranial neuropathies, and visual decline.² The decision to treat CCFs remains multifactorial and is often based on the severity of clinical symptoms and the angiographic characteristics of the fistula.³ Although other treatment modalities exist, in recent years endovascular occlusion has become the treatment of choice for the majority of CCFs.⁴ However, most of the data for these lesions remains limited to small, single-center series. As such, we performed a systematic review and meta-analysis evaluating endovascular occlusions of CCFs to determine any differences in clinical outcomes based on presentation, fistula type, and treatment paradigm.

2. Methods

2.1. Study selection

The systematic review and meta-analysis were performed by PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines. The following electronic databases were queried: PubMed, Scopus, Web of Science (WoS), and Embase. Relevant articles were identified using the following search syntax ((Endovascular treatment) OR (Interventional Neuroradiology)) AND ((Carotid-Cavernous Fistula Embolization) OR (CCF)). After primary database searching was complete, the reference list of each paper was screened and related citations were extracted.

2.2. Inclusion and exclusion criteria

Only peer-reviewed articles evaluating the clinical efficacy of endovascular treatment in CCFs before March 2023 were included. Exclusion criteria were non-English publications, narrative or systematic reviews and meta-analyses, and studies with a lack of sufficient data to determine CCF etiology or type, pre-treatment symptoms, post-treatment outcomes, or endovascular approaches utilized.

2.3. Data extraction

Two of the authors independently assessed the full text of the included studies and extracted the data. All data regarding demographics of patients (e.g., gender, age, country, and duration of follow-up), etiology of CCF (e.g., trauma or spontaneous), surgical approach to the CCF (e.g., application of coil, balloon, and stent), clinical symptoms of the patients (e.g., proptosis, chemosis, diplopia, bruits, visual decline, orbital or pre-orbital pain and headache) and outcomes following endovascular treatment/observation (e.g. complete or partial remission) were extracted.

2.4. Statistical analysis

Categorical variables were reported as a percentage and continuous variables were reported as means and standard deviation (SD). Regarding demographics data we have reported the percentage of males/females, mean age of patients, mean duration of follow-up, and the number of patients in each country. In terms of etiology, we have reported the percentage of patients with an underlying cause of trauma and patients with spontaneous CCF. In surgical approaches, we have calculated the percentage of the application of each endovascular device including coils, balloons, and stents. We have also assessed the prevalence of clinical symptoms in CCF patients including proptosis, chemosis, diplopia, bruits, visual decline, orbital or pre-orbital pain, and headache. At last, we have evaluated the outcomes of endovascular treatment in CCF by reporting the percentage of patients with either partial or complete remission. Following a Binominal distribution, the variance along with a 95% confidence interval was reported. Thereafter, we combined the incidence of each study using average weight and an inverse association between each study variance and its weight was observed. Heterogeneity was assessed using the I^2 index and Q statistic (α significance level of 10%), and the Freeman-Tukey Double Arcsine Transformation⁵ was used to stabilize the variance. If the I^2 indices were more than 50%, the studies included in the meta-analysis were considered as heterogeneous. To perform the meta-analysis, a random effect model was applied using STATA software (version 14.2). If p was near to 1 or 0, the Meta-prop command was exerted. The ethical competence of this research was approved by the ethics committee of our institution Shahid Beheshti University of Medical Sciences (IR.SBMU.RETECH.REC.1400.477).

3. Results

3.1. Literature search and study selection

This study has been performed based on the PRISMA checklist.⁶ After an initial search of four electronic databases, 1085 papers (813 from PubMed, 102 from Scopus, 84 from Embase, and 86 from Web of Science) were obtained employing the search criteria. After an initial evaluation by the authors, 413 studies were removed due to duplication. Reasons such as unavailable abstracts, case reports, review articles, or non-English studies, excluded 235 further articles. The authors then carefully assessed the full text of each study, and papers were excluded due to lack of sufficient data ($n = 176$), ineligible sample size ($n = 63$), inapplicable approaches ($n = 95$), unavailable full text ($n = 13$), or not reporting the necessary information ($n = 54$). After applying the exclusion and inclusion criteria, a total of 36 studies from 2012 to 2023 were included in this systematic review and meta-analysis (Fig. 1, Table 1).

3.2. Demographics

Our meta-analysis included a total of 1494 patients (634 females (55.08%), 6 studies have not mentioned the gender of the patients) who underwent endovascular treatment of a CCF. The mean age of patients in this meta-analysis was 48.10 years. 9 of these 36 studies were conducted




Fig. 1. Flow diagram of studies identified in the systematic review and meta-analysis.

in China, 5 in the U.S., 2 in Germany, 2 in Italy, 2 in France, 2 in Brazil, 1 in India, 1 in Iran, 1 in Indonesia, 1 in Portugal, 1 in Peru, 1 in Australia, 1 in Turkey, 1 in Vietnam, 1 in Korea, 1 in Taiwan and 1 in Ecuador.

3.3. Etiology

We found that 87.17% of CCFs are caused by trauma, where 10.2% can be considered spontaneous (Table 2).

3.4. Surgical approaches

Endovascular treatment was performed on a total number of 1516 fistulas. In some studies, the type of these fistulas was mentioned. Based on the data available, 54.76% of fistulas were indirect and 45.24% were direct fistulas. A wide range of embolization devices and liquids were used in different studies. Some endovascular treatments consisted of more than one device or liquid. Coils (33.03%) and balloons (23.85%) were the two mostly used devices. The most used liquid embolization method was embolization with Onyx (6.96%). In some cases, these methods were used in combination with each other. In 13.76% of fistulas, several methods were used.

3.5. Clinical symptoms

The outcomes of this analysis show that the most prevalent symptoms of pre-operative CCF treatment were 89% exophthalmos (95% CI: 78.0–100.0; $I^2 = 75.7\%$) (Table 2), 84% chemosis (95% CI: 79.0–88.0; $I^2 = 91.6\%$) (Table 2), 79% proptosis (95% CI: 72.0–86.0; $I^2 = 91.84\%$) (Table 2), 75.0% bruits (95% CI: 67.0–82.0; $I^2 = 90.7\%$) (Table 2), 56% diplopia (95% CI: 42.0–71.0; $I^2 = 92.3\%$) (Table 2), 49% cranial nerve palsy (95% CI: 32.0–66.0; $I^2 = 95.1\%$), 39% visual decline (95% CI: 32.0–45.0; $I^2 = 71.4\%$) (Table 2), 32% tinnitus (95% CI: 6.0–58.0; $I^2 = 96.7\%$) (Table 2), 29% elevated intraocular pain (95% CI: 22.0–36.0; $I^2 = 0.0\%$) (Table 2), 31% orbital or pre-orbital pain (95% CI: 14.0–48.0; $I^2 = 89.9\%$) (Table 2), and 24% headache (95% CI: 13.0–34.0; $I^2 = 74.98\%$) (Table 2) (Fig. 1s).

3.6. Outcomes following endovascular treatment/observation

Most papers reported that the closure of the embolization process was successful. Immediate complete occlusion of fistula after treatment was observed in 75.0% (95% CI: 67.0–83.0; $I^2 = 94.91\%$) of cases. The complete remission rate was 90.0% (95% CI: 86.0–94.0; $I^2 = 90.15\%$). Recurrence of CCF was observed in 35% (95% CI: 5.0–66.0; $I^2 = 99.75\%$) of patients. Cranial nerve paralysis after treatment occurred in 5% (95% CI: 1.0–9.0; $I^2 = 43.08\%$) of patients (Figs. 2s and 3s, Table 2).

3.7. Publication bias

The Begg's funnel plot of included papers is shown in Fig. 4s where no sign of publication bias was detected ($P = 0.161$). Hence, we can conclude that both negative and positive results have been reported (Fig. 4s).

4. Discussion

CCFs are a type of arteriovenous malformation, leads to direct or indirect arteriovenous shunts from the internal or external carotid arteries into the cavernous sinus.⁴¹ This can result in a variety of clinical presentations however most have some ocular involvement at the time of diagnosis.⁴² Classification systems have been introduced to categorize CCFs according to their etiology (traumatic or spontaneous), their hemodynamic status (high or low flow), and their angiographic arterial supply (direct or indirect).^{43–45} Based on an angiographic classification established by Barrow et al.,² CCFs are classified into four types: type A is characterized as high-flow, direct shunts between the CS and the internal carotid artery (ICA). This type of fistula can rarely be a complication of head trauma or be caused by a carotid-cavernous aneurysm rupture. Trauma to the craniofacial region, whether direct or indirect, can weaken the muscle wall of the ICA or result in a laceration that causes a vascular shunt from a high-flow artery system into a low-flow venous sinus, which causes CCF. Type B CCFs are low-flow, indirect, or Dural arteriovenous fistulas (DAVFs) between the meningeal branches of the ICA and the CS. Type C CCFs are shunts between the CS and the external carotid artery (ECA) and, finally, type Ds represent a communication between the CS with both the ICA and ECA. Most direct CCFs will present with a combination of chemosis (90%), proptosis (90%), diplopia (50%), pain (25%), bruit (25%) increased ocular pressure, visual loss, and dysfunction of the trigeminal nerve (up to 50%). Direct CCFs are usually unilateral however bilateral lesions can occur and are often associated with higher morbidity.⁴⁶ Indirect CCFs more frequently occur in post-menopausal women.⁴ Some factors including pregnancy, hypertension, diabetes, sinusitis, atherosclerotic disease, and thrombosis of CS predispose patients to develop indirect CCFs.^{1,2,47} Thrombosis of other veins distant from fistulous communication, large varix of the CS, pseudoaneurysm, and venous drainage into cortical veins are great risk factors of morbidity and mortality in CCF patients. Elevated intracranial pressure, progressive proptosis, declined visual acuity, transient ischemic attacks, and hemorrhage are considered clinical signs and symptoms associated with poorer prognosis.⁴⁸

In this meta-analysis, data extracted from the 36 included studies were analyzed and four major results were obtained. Regarding the etiology of fistula, 87.24% of CCFs were caused by trauma while 10.1% were spontaneous. In the method of intervention, 36.49% of CCF patients were treated by applying a coil as an endovascular device, 25.04% by balloon, and 12.94% by an onyx (Table 2.). The third significant finding of our study was about pre-operative symptoms of CCF patients, which from the most common to the least were identified as exophthalmos, chemosis, proptosis, bruits, diplopia, cranial nerve palsy, visual decline, tinnitus, elevated intraocular pressure, orbital and periorbital pain, and headache with a prevalence of respectively 89%, 84%, 79%, 76%, 52%, 47%, 39%, 32%, 29%, 24% and 19%. Another important finding was that 68% of patients who underwent endovascular treatment-experienced

Table 1
Overview of included studies.

Author	The number of patients	Female	Mean age	Trans arterial route	Transvenous route	Both arterial and venous routes	Etiology		Symptoms			
							Trauma	Spontaneous	Chemosis	Proptosis	Exophthalmos	Bruits
Y. Yu, 2012 ⁷	23	8	36.3	23	0	0	0	23	–	–	–	–
Ying Yu, 2014 ⁸	18	4	37.8	18	0	0	16	2	9	7	–	11
Qinglin Liu, 2021 ⁹	10	3	28	10	0	0	10	0	10	–	6	5
C. M. Wendl, 2017 ¹⁰	14	11	59	–	–	–	5	1	–	–	–	–
Tiago Rodrigues, 2014 ¹¹	38	29	63	5	22	1	–	–	34	32	–	5
Zihuan Zhang, 2021 ¹²	18	9	–	–	–	–	–	–	–	–	–	–
Muhamad Thohar Arifin, 2020 ¹³	31	–	–	13	5	0	21	0	21	21	–	–
Lui's Henrique de Castro-, 2018Afonso	63	–	62.7	2	60	1	–	–	–	–	–	–
CUONG TRAN CHI, 2014 ¹⁴	172	–	–	171	1	0	172	0	–	139	–	170
¹⁵ alioscia derenzis, 2013	13	7	53.7	10	3	0	6	7	9	10	–	4
Hui Guo, 2017 ¹⁶	45	28	53.4	0	45	0	–	–	41	37	–	36
Yin Niu, 2019 ¹⁷	24	–	–	–	–	–	24	0	–	–	–	–
Chao-Bao Luo, 2012 ¹⁸	24	9	39	25	0	0	–	–	24	24	–	24
Xiao-Quan Xu, 2012 ¹⁹	58	15	–	58	0	0	58	0	49	48	–	52
S.Stepha, 2015 ²⁰	60	40	59	–	–	–	–	–	39	46	–	23
Jacob F Baranoski, 2019 ²¹	5	2	47	1	4	0	3	2	4	4	–	–
Luis Henrique de Castro-Afonso, 2017 ²²	62	38	62.7	2	58	1	–	–	–	–	–	–
Bu-Lang Gao, 2017 ²³	188	50	31	–	–	–	188	0	–	–	–	–
Edgar A Samaniego, 2015 ²⁴	7	–	–	7	0	0	5	2	2	–	6	4
Xiang Zhang, 2016 ²⁵	16	4	35.7	17	0	0	17	0	–	–	–	–
André Beer-Furlan1, 2020 ²⁶	7	5	–	0	7	0	–	–	3	5	–	–
Xiaojian Lu, 2014 ²⁷	32	11	32.3	32	0	0	30	2	28	–	28	31
Bekir Sanal, 2018 ²⁸	23	16	61	5	18	1	–	–	17	18	–	–
Jong Kook Rhim, 2018 ²⁹	17	13	64.9	0	34	0	–	–	–	–	–	–
Francesco Briganti, 2013 ³⁰	30	22	51	0	30	0	–	–	16	–	30	–
ALI PASHAPOUR, 2014 ³¹	46	–	36.83	26	10	10	–	–	46	23	–	–
Marcus Ohlsson, 2016 ³²	9	4	35	–	–	–	5	0	–	–	3	4
Arvinda Hanumanthapura, 2013Ramalin ³³ gaia	21	3	31	21	0	0	21	0	20	19	–	21
L. Fernando Gonzalez, 2012 ³⁴	5	4	–	2	3	0	2	3	–	–	–	2
Matthew D. Alexander, 2018 ³⁵	267	209	60.9	73	199	7	–	–	227	221	–	–
Fadi Al Saiegh, 2020 ³⁶	42	24	63.4	8	32	2	7	0	37	31	–	–
Lee J.Holland, 2017 ³⁷	39	29	59	–	–	–	13	21	–	30	–	–
Lorenz Ertl, 2020 ³⁸	33	23	61.9	4	27	2	–	–	28	–	29	–
Andres R. Plasencia, 2012 ³⁹	24	10	37.5	18	5	1	14	8	8	–	–	–
Lun-Xin Liu, 2018 ⁴⁰	10	4	35.6	10	0	0	8	2	10	10	–	10
Total	1494	634 (55.08%)	48.10 (mean)	561	563	26	625	73	682	725	99	402

Symptoms							Method of embolization							
Cranial nerve palsy	Diplopia	Orbital and pre-orbital pain	Tinnitus	Elevated IOP	Visual decline	Headache	Coil	Onyx	Stent	Balloons	nBCA	Covered stent	Glue	Multiple
-	-	6	-	-	8	-	-	23	-	-	-	-	-	-
3	-	-	-	-	-	-	-	-	-	1	-	9	-	18
-	-	-	-	-	-	-	-	-	-	-	-	-	-	11
20	20	6	5	14	19	10	24	-	-	-	-	-	3	1
-	-	-	-	-	-	-	14	-	-	-	-	-	-	5
-	-	-	-	-	-	-	8	2	-	10	-	-	-	-
-	-	-	-	-	-	-	33	5	-	-	3	-	-	22
95	-	-	-	-	67	-	-	-	138	-	-	-	31	-
-	-	-	-	-	-	-	1	-	12	-	-	12	12	-
-	26	-	-	-	19	-	-	-	-	-	-	-	-	-
-	-	-	-	-	-	-	1	-	-	21	-	-	-	2
-	-	-	-	-	-	3	25	-	-	-	-	-	-	-
-	-	-	-	-	21	-	-	-	-	58	-	-	-	-
36	32	19	-	15	19	-	-	-	-	-	-	-	-	-
2	4	-	-	-	-	-	-	-	-	-	-	-	-	5
27	-	-	-	-	-	-	32	6	-	-	3	-	-	21
-	-	-	-	-	-	-	-	-	-	188	-	-	-	-
1	-	-	2	-	3	2	1	-	-	-	-	-	-	6
-	-	-	-	-	-	-	-	-	-	-	-	-	-	17
6	-	-	1	-	-	-	-	7	-	-	-	-	-	-
26	-	-	-	-	20	-	8	-	-	21	-	2	-	1
-	17	10	-	-	9	-	18	-	-	1	-	1	1	3
-	-	-	4	-	-	3	-	-	-	-	-	-	-	-
-	14	-	-	-	-	-	30	-	-	-	-	-	-	-
-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
1	-	-	-	-	-	-	-	-	-	-	-	-	9	-
16	18	-	20	-	-	10	-	-	-	-	-	-	-	21
5	-	-	-	-	-	2	-	1	-	-	-	-	-	4
-	197	-	-	-	106	-	196	7	-	-	4	-	-	20
6	14	-	3	-	8	6	1	38	-	-	-	-	-	3
-	23	24	16	-	-	18	17	-	-	-	-	-	-	5
5	4	2	14	9	22	2	17	2	-	-	-	-	1	13
-	-	-	-	7	6	-	6	-	-	12	2	-	-	2
-	-	-	2	-	1	-	-	-	10	-	-	10	-	-
249	369	67	67	45	328	56	432 (33.03%)	91 (6.96%)	160 (12.23%)	312 (23.85%)	12 (0.92%)	34 (2.60%)	57 (4.36%)	180 (13.76%)

Table 2
Summary of meta-analysis results.

	Number of studies	Pooled Percentage (95% CI)	I ² (%)
Etiology			
Spontaneous	21	10.2%	
Trauma	21	87.17%	
Others	21	2.65%	
Surgical method			
Coil	17	33.03%	
Balloons	8	23.85%	
Stent	3	12.23%	
Gluebrin	6	4.36%	
Onyx	9	6.96%	
nBCA	4	0.92%	
Covered stent	5	2.60%	
Multiple	19	13.76%	
Other	3	2.29%	
Clinical symptoms			
Chemosis	22	84% (79.0%–88.0%)	91.6%
Proptosis	18	79% (72.0%–86.0%)	91.84%
Exophthalmos	5	89% (78.0%–100.0%)	75.7%
Bruits	15	75% (67.0%–82.0%)	90.7%
Cranial nerve palsy	14	49% (32.0%–66.0%)	95.1%
Diplopia	11	56% (42.0%–71.0%)	92.3%
Orbital and pre-orbital pain	6	31% (14.0%–48.0%)	89.9%
Tinnitus	9	32% (6.0%–58.0%)	96.7%
Elevated intraocular pressure	4	29% (22.0%–36.0%)	0.0%
Visual decline	14	39% (32.0%–45.0%)	71.4%
Headache	9	24% (13.0%–34.0%)	74.98%
Surgical approaches			
Direct fistulas	28	48.05%	
Indirect fistulas	28	51.95%	
Trans arterial route	28	56% (39.0%–73.0%)	99.93%
Trans venous route	28	45% (25.0%–65.0%)	99.95%
Both Trans arterial and Transvenous routes	28	0% (0%–1%)	100.0%
Treatment outcome			
Immediate complete occlusion of fistula	20	75% (67.0%–83.0%)	99.3%
Complete remission	17	82% (71%–94%)	99.0%
Recurrence	13	35% (5.0%–66.0%)	99.5%
Cranial nerve paralysis	6	5% (1.0%–9.0%)	1.3%

complete occlusion of the fistula immediately after treatment and 82% of patients experienced complete remission with total improvement of clinical symptoms.

As discussed earlier, we extracted data regarding the etiology of CCF and found that the majority of cases were caused by trauma (87.24%) and spontaneous CCF constituted (10.1%) of cases. This finding confirms the previous literature as follows. Different theories explain the post-traumatic CCF mechanism. Trauma that might be accompanied by increased shear force and bone fracture can lead to a rupture in the carotid artery. However, Helmke et al⁴⁹ reported no fractures in 42 post-traumatic direct CCF cases; thus, the mentioned theory was substituted by another theory that trauma leads to an abrupt rise in the internal carotid artery (ICA) pressure and simultaneous compaction of the distal artery, leading to vessel wall tear and CCF. In addition to penetrating or blunt traumas, iatrogenic damage can also result in CCF, for instance, trans-sphenoidal surgery, carotid angioplasty, etc.⁴⁷ Spontaneous direct CCFs are formed through a cavernous aneurysm rupture or atherosclerotic artery leading to a weakened ICA wall subsequent to predisposing factors including pseudoxanthoma elasticum and Ehlers-Danlos syndrome. Iatrogenic causes such as previous contralateral ICA occlusion through altering the flow dynamics and pressure can play a role in spontaneous aneurysmal tears.^{1,2,46,50–52} The causes of Indirect CCFs are still unknown but there is some evidence supporting congenital origins.^{1,46,53} Factors such as trauma, pregnancy, diabetes, and hypertension are predisposing factors for dural CCF, while the association of trauma with indirect CCF is less prevalent.^{1,2,47} Among the included

studies, Liu LX et al and Zeineddine HA et al^{40,54} reported trauma as cause of 80% of CCF cases, which is also confirmed by our study. However, the Sanal B et al study⁴⁴ demonstrated a higher prevalence of spontaneous causes, which perhaps was due to a higher number of indirect CCF cases than direct ones in that study.

Endovascular intervention technology, which has recently evolved enormously, has provided various options for the treatment of CCFs. Thus, endovascular methods are now counted as the principal therapy in CCFs, or after conservative treatment failure. Regarding direct CCF, the target of therapy is the closure of the rupture between the cavernous sinus and ICA, during which the ICA patency is preserved. This can be achieved by using a detachable balloon to eliminate the fistula trans-arterially, applying coils for the abolition of the cavernous sinus of the ipsilateral site trans-venously or trans-arterially, or positioning a covered stent through the fistula.¹ Embolization may be either trans-arterial or transvenous. If trans-arterial, coils are the optimal choice as embolization agent. Liquid agents such as n-BCA and onyx can also be used. In combination with this agent a balloon might be used to keep the parent vessel safe and prevent embolization agent from migrating to cerebral hemispheres. Both coils and liquid agents can be used for transvenous embolization. Based upon our meta-analysis, the prevalence of patients treated by coils, balloons, and stent were, respectively, 36.49%, 25.08%, and 12.94%. Coils are also used in combination with other embolization agents such as ethylene-vinyl alcohol copolymer (EVOH) (onyx), stent, balloons and *n*-butyl cyanoacrylate (n-BCA) as shown in Table 1. Detachable platinum coils are preferred because apart from simple utilization, whenever they are not placed optimally, they can be removed or adjusted later.⁵⁵ A downside of using balloons is the potential risk of displacement or deflation which may lead to a recurrence of symptoms and rehabilitation of the malformation. In addition, they can result in constant or temporary paralysis of cranial nerves by inducing a mass effect in the cavernous sinus.⁵⁶ Onyx, the most used liquid embolization agent, is used both trans-venously and trans-arterially. Stents have some advantages over coils or balloons, including rapid positioning, no risk of coil decapsulation or herniation, decreased local compaction and mass effects, and not being associated with pseudo-aneurysm formation.⁵⁷ A major disadvantage of stents is that they cannot be applied in acute stages post trauma.⁵⁸ In the studies assessed, transvenous approach was used more than trans-arterial route.

According to the results of our study, the most prevalent pre-operative symptom of CCF was exophthalmos (89%) followed by chemosis, proptosis, bruits, diplopia, cranial nerve palsy, visual decline, tinnitus, elevated intraocular pressure, orbital and periorbital pain, and headache respectively, which is in line with findings from previous studies. Drainage of the anterior part of the orbit gives rise to congestion of the orbital vein and subsequent fluids transudation, the elevation of intraocular pressure, laceration of dilated veins, and impairment in perfusion of the retina.^{4,41,42} Proptosis results from elevated orbital pressure, diplopia from paralysis of cranial nerves, a visual decline from ischemia in the retina or optic nerve, pain from aqueous return decline, and elevation in intraocular pressure. Headache is caused by venous hypertension, hemorrhage, and trigeminal nerve impairment. Direct CCFs generally manifest acutely and progress quickly, which requires immediate action. Perhaps these visual manifestations stem from ischemia in the retina, however, indirect CCFs usually present insidiously.⁴¹ We also extracted data regarding the outcome of endovascular treatments and found that 68% of patients who underwent this treatment achieved immediate occlusion of the fistula after treatment, and 82% reached complete remission with no clinical symptoms remained. This supports the finding of the Phan et al study⁵⁹ which indicated that the success of the endovascular orbital approach in fistula embolization was 89.9% and demonstrated that those few patients who did not improve after the operation, had not received a proper fistula embolization. It is difficult to predict visual outcomes after CCF treatment, but minor disorders are associated with better outcomes. Cases with thrombosis in the superior ophthalmic vein or occlusion of the central retinal vein at the time of

diagnosis showed poorer visual recovery. On the other hand, direct CCF patients manifest poorer vision and benefit from endovascular therapy with greater vision recovery.^{60,61} Most of the included studies' findings were consistent with our results.^{1,44,49,40,54,56,37,62-64}

In summary, in this meta-analysis, we gathered all available relevant evidence about the etiology, endovascular treatment devices, symptoms, and endovascular treatment outcomes. However, there were some potential limitations. Methods used in eligible studies were not similar and we included studies with the most valid methods. In addition, as there was no meta-analysis related to this topic since 2012, we could not compare our findings with previous ones. Further research is required to achieve more definite findings and compare these results to yield a better knowledge regarding different aspects of CCF.

5. Conclusion

CCF, a type of arteriovenous malformation, leads to direct or indirect arteriovenous shunts. Generally, post-traumatic CCFs constitute a significantly higher percentage of CCF cases than spontaneous ones. Exophthalmos, chemosis, proptosis, bruits, diplopia, cranial nerve palsy, visual decline, tinnitus, elevated intraocular pressure, orbital and peri-orbital pain, and headache are the most common clinical manifestations of CCFs. The majority of endovascular treatments involved coiling, balloons and stent and a high percentage of CCF patients experienced complete remission with the improvement of their clinical symptoms.

Credit author statement

Aryobarzan Rahmatian: Conceptualization, Methodology, Writing – original draft. Mobina Fathi: Conceptualization, Writing – original draft, Software, Supervision and editing. Shirin Yaghoobpoor: Writing – original draft. Robert M. Starke: Writing – original draft, reviewing. Evan M. Luther: Writing – original draft, reviewing, reviewing Fatemeh Sodeifian: Writing – original draft. Tara Fazel: Writing – original draft. Mina Dehghani: Writing – original draft, investigating and edition. Reza Ramezan: Writing – original draft. Masood Zangi: Writing – original draft, reviewing. Reza Goharani: Conceptualization, Writing – original draft, Software, Supervision, and editing.

Funding statement

The authors received no financial support for the preparation of this article for publication.

Declarations of competing interest

None.

Acknowledgments

The authors of this study are thankful to the Clinical Research Development Center (CRDC) and Skull Base Research Center of Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran for their support, cooperation, and assistance throughout the period of study.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.wnsx.2023.100189>.

References

- Gemmete JJ, Ansari SA, Gandhi DM. Endovascular techniques for treatment of carotid-cavernous fistula. *J Neuro Ophthalmol*. 2009;29(1):62-71.
- Barrow DL, et al. Classification and treatment of spontaneous carotid-cavernous sinus fistulas. *J Neurosurg*. 1985;62(2):248-256.
- Brzozowski K, et al. Superior ophthalmic vein and ophthalmic artery in immediate evaluation after endovascular treatment of carotid-cavernous fistulas. *Pol J Radiol*. 2019;84:e32.
- Korkmaz B, et al. Endovascular treatment of carotid cavernous sinus fistula: a systematic review. *World J Radiol*. 2013;5(4):143.
- Freeman MF, Tukey JW. *Transformations Related to the Angular and the Square Root*. The Annals of Mathematical Statistics; 1950:607-611.
- Liberati A, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *Ann Intern Med*. 2009;151(4): W-65-W-94.
- Yu Y, et al. Use of onyx for transarterial balloon-assisted embolization of traumatic carotid cavernous fistulas: a report of 23 cases. *AJNR Am J Neuroradiol*. 2012;33(7): 1305-1309.
- Yu Y, et al. Embolization of direct carotid cavernous fistula with Onyx and coils under transarterial balloon protection. *Cardiovasc Intervent Radiol*. 2014;37(3): 679-685.
- Liu Q, et al. Treatment of direct carotid-cavernous fistula with Willis covered stent with midterm follow-up. *Chin Neurosurg J*. 2021;7(1):41.
- Wendl CM, et al. Direct carotid cavernous sinus fistulae: vessel reconstruction using flow-diverting implants. *Clin Neuroradiol*. 2017;27(4):493-501.
- Rodrigues T, et al. Management of dural carotid cavernous fistulas: a single-center experience. *Eur Radiol*. 2014;24(12):3051-3058.
- Zhang Z, et al. A modified treatment through point-to-point coil embolization for direct carotid cavernous to fistula: a single-center result. *Front Neurol*. 2021;12, 639552.
- Thohar Arifin M, et al. Neuro-endovascular intervention in traumatic carotico-cavernous fistulae: a single-center experience. *Int J Gen Med*. 2020;13:917-925.
- Chi CT, et al. Direct traumatic carotid cavernous fistula: angiographic classification and treatment strategies. Study of 172 cases. *Intervent Neuroradiol*. 2014;20(4): 461-475.
- De Renzis A, et al. Balloon-assisted coiling of the cavernous sinus to treat direct carotid cavernous fistula. A single center experience of 13 consecutive patients. *Intervent Neuroradiol*. 2013;19(3):344-352.
- Guo H, et al. Focus on the target: angiographic features of the fistulous point and prognosis of transvenous embolization of cavernous sinus dural arteriovenous fistula. *Intervent Neuroradiol*. 2018;24(2):197-205.
- Niu Y, et al. Detachable balloon embolization as the preferred treatment option for traumatic carotid-cavernous sinus fistula? *Intervent Neuroradiol*. 2020;26(1):90-98.
- Luo CB, et al. Transarterial detachable coil embolization of direct carotid-cavernous fistula: immediate and long-term outcomes. *J Chin Med Assoc*. 2013;76(1):31-36.
- Xu XQ, et al. Follow-up of 58 traumatic carotid-cavernous fistulas after endovascular detachable-balloon embolization at a single center. *J Clin Neuro*. 2013;9(2):83-90.
- Stéphan S, et al. Endovascular treatment of carotid-cavernous fistulae: long-term efficacy and prognostic factors. *J Fr Ophthalmol*. 2016;39(1):74-81.
- Baranoski JF, et al. Flow diverters as a scaffold for treating direct carotid cavernous fistulas. *J Neurointerventional Surg*. 2019;11(11):1129-1134.
- de Castro-Afonso LH, et al. Transvenous embolization of dural carotid cavernous fistulas: the role of liquid embolic agents in association with coils on patient outcomes. *J Neurointerventional Surg*. 2018;10(5):461-462.
- Gao BL, et al. Recurrence risk factors in detachable balloon embolization of traumatic direct carotid cavernous fistulas in 188 patients. *J Neurointerventional Surg*. 2018; 10(7):704-707.
- Samaniego EA, Martínez-Galdámez M, Abdo G. Treatment of direct carotid-cavernous fistulas with a double lumen balloon. *J Neurointerventional Surg*. 2016; 8(5):531-535.
- Zhang X, et al. Combined use of Onyx and coils for transarterial balloon-assisted embolization of traumatic carotid-cavernous fistulas: a report of 16 cases with 17 fistulas. *J Neurointerventional Surg*. 2016;8(12):1264-1267.
- Beer-Furlan A, et al. Transvenous onyx embolization of carotid-cavernous fistulas: mid- and long-term outcomes. *J Neurol Surg B Skull Base*. 2021;82(Suppl 3): e278-e284.
- Lu X, et al. A comparison of different transarterial embolization techniques for direct carotid cavernous fistulas: a single center experience in 32 patients. *J Vasc Interv Neurol*. 2014;7(5):35-47.
- Sanal B, et al. Endovascular treatment in traumatic and spontaneous carotid cavernous fistulas: with different embolization agents and via various vascular routes. *J Vasc Interv Neurol*. 2018;10(2):18-24.
- Rhim JK, et al. Endovascular treatment of bilateral cavernous sinus dural arteriovenous fistula: therapeutic strategy and follow-up outcomes. *Korean J Radiol*. 2018;19(2):334-341.
- Briganti F, et al. Endovascular occlusion of dural cavernous fistulas through a superior ophthalmic vein approach. *NeuroRadiol J*. 2013;26(5):565-572.
- Pashapour A, et al. Long-Term endovascular treatment outcome of 46 patients with cavernous sinus dural arteriovenous fistulas presenting with ophthalmic symptoms. A non-controlled trial with clinical and angiographic follow-up. *NeuroRadiol J*. 2014; 27(4):461-470.
- Ohlsson M, Consoli A, Rodesch G. Endovascular treatment of carotico-cavernous fistulas with acrylic glue: a series of nine cases. *Neuroradiology*. 2016;58(12): 1181-1188.
- Ramalingaiah AH, et al. Transarterial treatment of direct carotico-cavernous fistulas with coils and Onyx. *Neuroradiology*. 2013;55(10):1213-1220.
- Gonzalez LF, et al. Treatment of carotid-cavernous fistulas using intraarterial balloon assistance: case series and technical note. *Neurosurg Focus*. 2012;32(5):E14.
- Alexander MD, et al. Long-Term outcomes of endovascular treatment of indirect carotid cavernous fistulae: superior efficacy, safety, and durability of transvenous coiling over other techniques. *Neurosurgery*. 2019;85(1):E94-e100.

36. Al Saiegh F, et al. Onyx embolization of carotid-cavernous fistulas and its impact on intraocular pressure and recurrence: a case series. *Oper Neurosurg (Hagerstown)*. 2021;20(2):174–182.
37. Holland LJ, et al. Endovascular treatment of carotid–cavernous sinus fistulas: ophthalmic and visual outcomes. *Orbit*. 2019;38(4):290–299.
38. Ertl L, et al. Patient reported long-term outcome after endovascular therapy of indirect dural carotid cavernous fistulas. *PLoS One*. 2020;15(4):e0231261.
39. Plasencia AR, Santillan A. Endovascular embolization of carotid-cavernous fistulas: a pioneering experience in Peru. *Surg Neurol Int*. 2012;3:5.
40. Liu LX, et al. Application of the willis covered stent in the treatment of carotid-cavernous fistula: a single-center experience. *World Neurosurg*. 2019;122:e390–e398.
41. Ellis JA, et al. Carotid-cavernous fistulas. *Neurosurg Focus*. 2012;32(5):E9.
42. Zanaty M, et al. Endovascular treatment of carotid-cavernous fistulas. *Neurosurg Clin*. 2014;25(3):551–563.
43. Barber SM, et al. Mid-and long-term outcomes of carotid-cavernous fistula endovascular management with Onyx and n-BCA: experience of a single tertiary center. *J Neurointerventional Surg*. 2015;7(10):762–769.
44. Sanal B, et al. Endovascular treatment in traumatic and spontaneous carotid cavernous fistulas: with different embolization agents and via various vascular routes. *Journal of vascular and interventional neurology*. 2018;10(2):18.
45. Morton RP, et al. Radiographic and clinical outcomes in cavernous carotid fistula with special focus on alternative transvenous access techniques. *J Clin Neurosci*. 2015;22(5):859–864.
46. Ringer AJ, Salud L, Tomsick TA. Carotid cavernous fistulas: anatomy, classification, and treatment. *Neurosurgery Clinics*. 2005;16(2):279–295.
47. Tjounmakaris SI, Jabbour PM, Rosenwasser RH. Neuroendovascular management of carotid cavernous fistulae. *Neurosurg Clin*. 2009;20(4):447–452.
48. Halbach VV, et al. Carotid cavernous fistulae: indications for urgent treatment. *Am J Roentgenol*. 1987;149(3):587–593.
49. Helmke K, Krüger O, Laas R. The direct carotid cavernous fistula: a clinical, pathoanatomical, and physical study. *Acta Neurochir*. 1994;127(1–2):1–5.
50. Wanke I, et al. Carotid cavernous fistula due to a ruptured intracavernous aneurysm of the internal carotid artery: treatment with selective endovascular occlusion of the aneurysm. *J Neurol Neurosurg Psychiatr*. 2001;71(6):784–787.
51. Germain D, HerreraGuzman Y. Vascular ehlers-danlos syndrome. *J. Ann. gen.* 2004; 47:1.
52. Debrun GM. Angiographic workup of a carotid cavernous sinus fistula (CCF) or what information does the interventionalist need for treatment? *Surg Neurol*. 1995;44(1): 75–79.
53. Pang D, et al. External carotid-cavernous fistula in infancy: case report and review of the literature. *Neurosurgery*. 1981;8(2):212–218.
54. Zeineddine HA, et al. Embolization of carotid-cavernous fistulas: a technical note on simultaneous balloon protection of the internal carotid artery. *J Clin Neurosci*. 2020; 78:389–392.
55. Halbach VV, et al. Transarterial platinum coil embolization of carotid-cavernous fistulas. *Am J Neuroradiol*. 1991;12(3):429–433.
56. De Renzis A, et al. Balloon-Assisted coiling of the cavernous sinus to treat direct carotid cavernous fistula: a single center experience of 13 consecutive patients. *Intervent Neuroradiol*. 2013;19(3):344–352.
57. Wang Y, et al. Treatment of post-traumatic carotid-cavernous fistulas with the Willis covered stent: a preliminary prospective study. *Intervent Neuroradiol*. 2012;18(2): 172–177.
58. Cho K-C, et al. Cerebral hemorrhage after endovascular treatment of bilateral traumatic carotid cavernous fistulae with covered stents. *Journal of Korean Neurosurgical Society*. 2011;50(2):126.
59. Phan K, et al. Orbital approaches for treatment of carotid cavernous fistulas: a systematic review. *World Neurosurg*. 2016;96:243–251.
60. Yu SCH, et al. Transvenous embolization of dural carotid-cavernous fistulae with transfacial catheterization through the superior ophthalmic vein. *Neurosurgery*. 2007; 60(6):1032–1038.
61. Zanaty M, et al. Endovascular treatment of carotid-cavernous fistulas. *Neurosurg Clin*. 2014;25(3):551–563.
62. Stéphan S, et al. Endovascular treatment of carotid-cavernous fistulae: long-term efficacy and prognostic factors. *J Fr Ophthalmol*. 2016;39(1):74–81.
63. Tao Z, Yu-feng L, Song-sheng S. Multimodal endovascular treatment for traumatic carotid-cavernous fistula. *Chin J Traumatol*. 2013;16(6):334–338.
64. Miller NR, et al. Treatment of carotid-cavernous sinus fistulas using a superior ophthalmic vein approach. *J Neurosurg*. 1995;83(5):838–842.