



# Research progress on exosomal proteins as diagnostic markers of gastric cancer (review article)

Hang Su<sup>1</sup> · Weihong Ren<sup>2</sup> · Dai Zhang<sup>2</sup>

Received: 11 September 2021 / Accepted: 4 January 2022 / Published online: 22 January 2022  
© The Author(s) 2022

## Abstract

Gastric cancer (GC) is one of the most common types of tumors and the most common cause of cancer mortality worldwide. The diagnosis of GC is critical to its prevention and treatment. Available tumor markers are the crucial step for GC diagnosis. Recent studies have shown that proteins in exosomes are potential diagnostic and prognostic markers for GC. Exosomes, secreted by cells, are cup-shaped with a diameter of 30–150 nm under the electron microscope. They are also surrounded by lipid bilayers and are widely found in various body fluids. Exosomes contain proteins, lipids and nucleic acid. The examination of exosomal proteins has the advantages of quickness, easy sampling, and low pain and cost, as compared with the routine inspection method of GC, which may lead to marked developments in GC diagnosis. This article summarized the exosomal proteins with a diagnostic and prognostic potential in GC, as well as exosomal proteins involved in GC progression.

**Keywords** Exosomal protein · Gastric cancer marker · Diagnosis

## Introduction

Gastric cancer (GC) is the 5th most frequently diagnosed cancer and 3rd leading cause of cancer mortality, accounting for 1,000,000 new cases and ~783,000 deaths in 2018 [1]. At present, the lack of appropriate screening markers for GC leads to the majority of patients missing the best treatment opportunities. Therefore, the exploration and application of diagnostic markers for GC are of great significance. Studies have shown that exosomal proteins can play a role in the diagnosis and prognosis of GC [2, 3]. Exosomes have the potential to be diagnostic and prognostic markers of cancer, since they contain microRNA (defined as a kind of tiny non-coding RNA with 23–35 nucleotide and tumor diagnostic potential) and cell-specific proteins that can mediate cell communication [4–8]. Exosomal proteins have a broad prospect in the development and application of biomarkers in

GC diagnosis and prognosis, with a low cost and less pain, as compared with conventional diagnostic methods. With further research conducted, exosomal protein markers of GC with a high specificity and sensitivity may be widely used in clinical practice. In the present study, exosomal proteins with a diagnostic and prognostic potential of GC and their research status were reviewed herein.

## Serological proteomics markers commonly used in GC diagnosis

Serological markers of GC include specific markers, such as gastrin-17 and pepsinogen (PG), and non-specific ones, such as carbohydrate antigen 199, CA724 and carcinoembryonic antigen. The changes in PG level and PGI/II ratio are considered indicators of gastric mucosa atrophy. Both PGI and PGII are known to reduce the development of atrophy and loss of specialized cells. PGI usually shows a more advanced decrease than PGII, causing a low PGI/II ratio [9]. Gastrin-17 is the main component of gastrin, which is secreted by G cells in the gastric wall. Tu et al. [10] demonstrated that a low level of gastrin-17 may be a biomarker for atrophic gastritis in the gastric antrum. Yu et al. [11] performed gastroscopy on 68 patients with chronic atrophic gastritis and 86 healthy controls, and detected serum levels of PGI,

✉ Weihong Ren  
ren\_weihong@163.com

<sup>1</sup> The First Clinical Medical Institute, Henan University of Chinese Medicine, No.156 Jinshui East Road, Zhengzhou, Henan Province, China

<sup>2</sup> Department of Laboratory Medicine, The First Affiliated Hospital of Henan University of Chinese Medicine, No.19 Renmin Road, Zhengzhou 450000, Henan Province, China

PGII and gastrin-17 by enzyme-linked immunosorbent assay (ELISA). That study subsequently reported that the sensitivity, specificity and area under curve of gastrin-17 were less than the PGI, PGII and PGI/II ratio, suggesting inferior clinical value of gastrin-17 in screening chronic atrophic gastritis than that of PG. CA724 has a high specificity and dissatisfactory clinical application [12]. Currently, the serological examination of GC uses multi-index combined detection to improve diagnostic efficiency, which remains flawed due to the inadequate sensitivity or specificity of single-detection indicators. The protective effect of exosomes makes its proteins a more favorable marker, as compared with the aforementioned serum markers.

## Brief introduction of exosomes

Exosomes are a type of membranous vesicles with a diameter of 30–150 nm which contain proteins, nucleic acid, etc. (Fig. 1) [13, 14]. They can be released by a variety of cells, including fibroblasts, intestinal epithelial cells, neurons, fat cells and tumor cells [15], and exist in multiple body fluids [16, 17]. Exosomes were first discovered and named by Johnstone et al. [18]. Early studies considered exosomes the “garbage bags” inside cells. Current study has demonstrated that exosomes function in numerous biological processes

and disease courses, such as angiogenesis, intercellular communication and antigen presentation [15].


## The formation of exosomes

The formation process of exosomes consists of the following three steps: First, the endosome is generated by the invagination of the plasma membrane. Secondly, the endosome is transformed into a multivesicular body (MVB). Finally, the MVB fuses with cell membranes and releases exosomes (Fig. 2) [15].


## The separation and detection methods of exosomes

The commonly used methods for exosome extraction include ultracentrifugation and related techniques, density gradients, extraction kit, size exclusion chromatography (SEC), affinity and precipitation. Several separation methods such as tangential flow, microfluidics and field flow fractionation have shown relatively few application [19]. Each of these methods possesses respective strong points and weaknesses. Centrifugation-based isolation methods, at the present stage, remain the gold standard and are most frequently applied to the exosomal separation [20, 21]. It can extract abundant exosome but is time-consuming and inefficient in recovery and repeatability, which hinders the clinical application

**Fig. 1** Compositions of Exosomes. The main components of exosomes include DNA, RNA (mRNA, miRNA) and proteins such as transmembrane protein (CD9, CD63), tetraspanins (CD13, PGRL), heat shock proteins (Hsp-60, Hsp-70), signaling proteins (EGFR, ARF1), MVB making Proteins (Alix, Clatherin), immunoregulatory molecules (CD80, CD86, MHC-I), cytoskeletal proteins (Cofilin, Talin, Vimentin), membrane trafficking (Rabs protein, Annexins), enzymes (ATPase, PGK, PK), immunoregulatory molecules (CD80, CD86, MHC-I), etc



**Fig. 2** *Biogenesis of Exosomes.*  
The Biogenesis of exosomes contains following steps: 1. Generation of endosome. 2. Transformation of endosome into multivesicular body (MVB). 3. Exosomes release



[21–23]. Immunomagnetic beads provide better purity but can only capture exosomes with specific targeted proteins [24]. Commercial extraction kits, which are simple-handling and without costly equipment, are also incapable of avoiding chemicals contamination [24, 25]. The acquirement of highly purified exosomes is of vital importance to its researches and the extraction method still needs to be improved.

The detection of exosome involves in the diagnosis and treatment of many diseases. Conventional detection methods of exosome include optical detection methods and non-optical ones. Conventional detection techniques are mainly composed of Dynamic light scattering (DLS) and nanoparticle tracking analysis (NTA). DLS, which is able to characterize exosome with a certain steady size, can neither handle samples with different sizes nor acquire source information [21]. Compared to DLS, NTA can characterize exosomes with discrepant sizes and fluorescence, meaning it can detect distributions of antigens on exosomes [21]. Non-optical methods include transmission electron microscopy (TEM), atomic force microscopy (AFM) and enzyme-linked immunosorbent assay (ELISA). TEM combined with cryogenic technique is capable of measuring the size of exosome and is first choice of visualizing nanoparticles and proteins [26]. AFM provides the size distribution of exosomal sample.

Besides, it can characterize the abundance, structure and other features of exosomes derived from tumor samples [21]. As a common detecting method in immunology, ELISA has gradually been applied to exosomal detection. In addition to these methods mentioned above, several emerging microfluidics-based detection technologies which are considered highly sensitive, low reagent-consuming and portable with high-throughput capacity have played a role in exosome characterization [21, 27].

### Exosome in delivery system

Exosome, with tiny volume, can easily pass through vascular wall and extracellular matrix and prevent phagocytosis of mononuclear macrophages [28]. Furthermore, exosome is capable of crossing biological barriers with weak immune response, making it nanocarrier and potential superior choice of delivering therapeutic drug [28–30]. With the aid of loading technology, required drugs can be transmitted to recipient cells and tissues. Pan et al. [31] created a new Exo-PMA/Au-BSA@Ce6 nanovehicle by combining urinary exosomes and ultra-small Au-BSA@Ce6 nanocomposites via an electroporation method, which improved photodynamic therapy on cancer and achieved real-time imaging of NIR fluorescence. Besides, Research by Liang

et al. [32] shown that engineered exosomes delivery system which transfer miR-21i and chemo-therapeutic drug 5-FU to the HCT-1165FR cancer cells are capable of enhancing the therapeutic efficiency of colon cancer and reversing the drug tolerance. Overall, exosome-involved delivery system has the potential to make a contribution to cancer diagnosis with the development.

## Exosomal proteins with a potential for GC diagnosis and prognosis

Exosomal proteins are divided into two types. The first type includes the common proteins in exosomes such as transmembrane transport and integration-related proteins, tetraspanins (e.g., CD9 and CD63) and heat shock proteins (HSP) [33]. The other type includes proteins present in certain types of exosomes. For example, exosomes from tumor cells possess a variety of tumor antigens. Figure 3 and Table 1 list the gene location, molecular weight and function of the proteins serving as GC markers.

### Tripartite motif 3 (TRIM3)

TRIM3, which is located at chromosome 11p15.5, is a member of the TRIM family that includes a RING domain, one or two B-boxes, and a coiled coil domain [34–38]. The upregulation of TRIM3 inhibits tumor cell behaviors, including proliferation and metastasis [2, 38–41]. Furthermore, the suppressed TRIM3 expression was associated with poor survival rates in breast cancer patients, suggesting that TRIM3 played a role in breast cancer inhibition [42]. Lu et al. [43] verified the upregulated expression of TRIM3 in Ewing sarcoma tissues, as compared with normal tissues. The enhanced expression of TRIM3 markedly and continually hindered autophagy in Ewing sarcoma (ES) cells. The opposite phenomenon was observed in TRIM3-silenced ES cells. In addition, TRIM3 is a potential target of miRNA (miR)-454-3p to promote the proliferation of human cervical cancer cells and inhibit their apoptosis [44]. With regards to other diseases, the experiment results by Wang et al. [45] indicated that TRIM3 was more lowly expressed in rheumatoid arthritis synovial tissues than healthy controls. Moreover, the stimulation of lipopolysaccharides on cell multiplication, the release of tumor necrosis factor- $\alpha$ , interleukin-6 (IL-6) and IL-1 $\beta$ , as well as the therapeutic effect on collagen-induced arthritis in rats were suppressed by the increased expression of TRIM3. Another study has also supported the viewpoint that TRIM3 contributes to the early diagnosis and treatment of Parkinson's disease [46].


Exosomal TRIM3 in patients' serum has the potential to serve as a diagnostic marker for GC. Fu et al. [2] extracted exosomes from the serum of GC patients and cell culture

supernatant using an exosome extraction kit, and tested the proteomic characteristics of exosomes by liquid chromatography-tandem mass spectrometry (LC-MS/MS). The expression of TRIM3 in the serum and tissue was also detected by ELISA and western blotting (WB). The results reported that expression levels of TRIM3 protein in both serum and tissue were lower than those in healthy controls. The overexpression of TRIM3 facilitated the growth and metastasis of GC which contradicted with TRIM3 knockdown. All these results showed that TRIM3 has the potential to serve as a diagnostic biomarker for GC.

### Gastrokine-1 (GKN1)

GKN1, also called antral mucosal protein (AMP)-18, is a member of the GKN family that is located at chromosome 2p13.3 [47]. It is a stomach-specific protein of 181-184 amino acids with a molecular weight of 18 kDa [48–52]. GKN1 comprises a hydrophobic signal peptide in the extreme NH<sub>2</sub>-terminal, and a BRICHOS domain containing conserved amino acid residues, aspartic acid residue, cysteine acid residues and a COOH-terminal domain [52–58]. GKN1 is secreted by gastric epithelial cells, stored in cytoplasmic granules and released into the extracellular environment through exosomes [49–51]. It is expressed in normal gastric tissue but not in GC tissue [59]. Studies have confirmed that GKN1 impacts cellular behavior. For example, it can hinder cell proliferation by inhibiting the expression of gastrin receptor [60]. In addition, GKN1 restrains cell growth and induces cell senescence by activating the p16/Rb and p21 signaling pathways [61]. It can also regulate immune response and suppress gastric epithelial cell carcinoma by affecting the expression of ILs and nuclear factor kappa B (NF- $\kappa$ B) [62]. GKN1 has been proven to be involved in the diagnosis and progression of various types of GC. Moss et al. [54] reported that a suppressed GKN1 expression was regularly observed in gastric adenocarcinomas, particularly of the diffuse subtype. The study by Dokhae et al. [47] confirmed that the GKN1 mRNA in the gastric tissue of GC patients was significantly lower than that in normal gastric tissue. Yoon et al. [63] measured the serum GKN1 of 200 healthy individuals and 1268 cancer patients (including 500 patients with GC) and analyzed its clinical value. They found the serum concentration of GKN1 of healthy individuals was much higher than that of patients with GC. The diagnostic accuracy of the serum GKN1 protein at the optimum cutoff was 0.9675. Moreover, GKN1 in serum of patients with advanced gastric cancer (AGC) were lower than those of patients with early gastric cancer (EGC). The diagnostic accuracies of GKN1 at the optimum cut-off (0.9675) were 0.8912 and 0.9589 for EGC and AGC, respectively. All these evidences have shown the specificity and prospect of serum GKN1 as a biomarker for

**Fig. 3** Exosomal proteins with potential as GC marker. TRIM3 and GKN1 can inhibit the proliferation of GC cells. CD97 has the opposite effect. TGF-β1 is able to play a dual role in GC. GKN1 and HSP-60, HSP-70 participate in the immune response. CD97, TGF-β1 could serve for GC prognosis and TRIM3, GKN1 can be applied to GC diagnosis



AGC and EGC. In addition, another study indicated that serum GKN1 concentrations discriminated healthy individuals with normal stomach and subjects with atrophy without and with intestinal metaplasia (IM) from GC patients with

AUCs of 1.0000, 1.0000, and 0.9964, respectively. Also, serum GKN1 levels in patients with GC yielded AUCs of 0.9938 and 0.9987 distinguishing patients with hepatocellular and colorectal carcinomas from GC patients. Besides,



**Table 1** Potential Exosomal Protein Biomarker in GC

| English abbreviations | Gene location | Molecular weight(kDa) | Source of exosomes | References |
|-----------------------|---------------|-----------------------|--------------------|------------|
| TRIM3                 | 11p15.5       | 80.8                  | Serum, Cell Line   | [2]        |
| GKN1                  | 2p13.3        | 18                    | Tissue             | [64]       |
| TGF- $\beta$ 1        | 19q13.1       | 25                    | Cell Line          | [65]       |
| CD97                  | 19p13.12      | 75–90                 | Cell Line          | [91]       |
| HSP-60                | 2q33.1        | 60                    | Malignant Ascites  | [90]       |
| HSP-70                | 6p21.33       | 70                    | Malignant Ascites  | [90]       |

serum GKN1 concentrations distinguished patients with EGC from normal individuals and subjects with atrophic gastritis without and with IM from GC patients with AUCs of 1.0000, 1.0000 and 0.9892, respectively. All of these data shown the diagnostic value of serum GKN1 on GC [50].

The diagnostic potential of exosomal GKN1 has been illustrated by previous studies. Yoon et al. [50] found that the GKN1 protein was expressed in exosomes derived from gastric epithelial HFE-145 cells but not in exosomes from AGS and MKN1 cells. Exosomes containing GKN1 impeded the proliferation and induced the apoptosis of AGS and MKN1 cells. In addition, GKN1-positive exosomes could inhibit tumor growth in GC mice model, indicating the potential of exosomal GKN1 in the diagnosis of GC. However, this conclusion has not been validated in a large-scale sample and evaluated in a prospective study. Two years after that study, Yoon et al. [64] conducted an investigation (containing 5 sporadic GC patients who had undergone a gastrectomy) on the specific uptake of exosomes from normal human gastric mucosa epithelial cells HFE-145 and examined the anticancer activity of exosomal GKN1 by immunofluorescence, WB and protein microarray chip. Experimental results showed that the exosomal GKN1 inhibited GC growth by binding to HRas and inhibiting its combination with b-raf and c-raf, which then decreased HRas/Raf/MEK/ERK signaling in AGS, MKN1 and xenograft tumor tissues. Moreover, exosomal GKN1 was involved in the inhibition on the migration and invasion of GC cells by suppressing epithelial-mesenchymal transition (EMT), suggesting that GKN1 can also be used for GC treatment.

### Transforming growth factor-beta 1 (TGF- $\beta$ 1)

TGF- $\beta$ 1, whose gene is located on chromosome 19q13, is a 25-kDa homodimeric polypeptide secreted by immune and tumor cells [65–71]. TGF- $\beta$ 1 signaling is involved in the regulation of many biological processes, including cell

proliferation, differentiation, migration and apoptosis [72, 73]. Several studies have proven that TGF- $\beta$ 1 plays a dual role in the tumor development process. In the early stage of tumors, TGF- $\beta$ 1 plays a suppressive role by hindering cell proliferation and inducing apoptosis [74–76]. Nevertheless, it was classified as a carcinogen, as it participates in EMT and angiogenesis in the late phase of tumors [77]. In terms of GC, TGF- $\beta$ 1 has the potential to be a predictor of GC progression and is also involved in natural killer (NK) cell injury, metastasis and recurrence of GC. A study showed that the TGF- $\beta$ 1 level was increased in patients with GC, and this increase was positively correlated with tumor stage [78]. The potential of plasma TGF- $\beta$ 1 as a predictor of GC progression was demonstrated by Zou et al. [79]. Furthermore, TGF- $\beta$ 1 secreted by cancer-associated monocytes/macrophages induced functional injury in NK cells in GC [80]. The study by Han et al. [81] identified a negative correlation between the TGF- $\beta$ 1 concentration and NK cells that expressed Nkp30, Nkp46, NKG2D and dnam-1 in peripheral blood. They also further illustrated the immunosuppressive effect of TGF- $\beta$ 1 on NK cells in patients with GC. Coban et al. [82] demonstrated that mean serum TGF- $\beta$ 1 levels were higher in patients with gastric cancer (GC) or colon cancer compared to the control group ( $p = 0.001$ ). The sensitivity of TGF- $\beta$ 1 was better in patients with GC than in patients with colon cancer. TGF- $\beta$ 1 had higher sensitivity than carcinoembryonic antigen (CEA) in GC patients. In addition, TGF- $\beta$ 1 was found to participate in GC cell EMT induction, invasion and migration [83, 84]. Wang et al. [85] suggested that TGF- $\beta$ 1 was specifically involved in the postoperative distant recurrence of gastric adenocarcinoma. The TGF- $\beta$ 1 signal was capable of upregulating the expression of vascular endothelial growth factor-c (VEGF-C) which caused lymphangiogenesis in GC [77].

The correlation between TGF- $\beta$ 1 and the prognosis of gastric adenocarcinoma patients has been confirmed. Yen et al. [65] analyzed the expression of exosomal TGF- $\beta$ 1 from gastroepiploic veins of 61 gastric adenocarcinoma patients by ELISA. Their findings were as follows: The expression of exosomal TGF- $\beta$ 1 in stages II, III and IV GC patients was higher than that in stage I patients. In addition, the expression of exosomal TGF- $\beta$ 1 in patients with lymph node metastasis (LNM) was twice as high as that in patients without LNM, suggesting that the expression of exosomal TGF- $\beta$ 1 in patients with gastric adenocarcinoma was associated with pathological stage and LNM. Moreover, exosomal TGF- $\beta$ 1 may be of greater value in the prediction of GC LNM than serum TGF- $\beta$ 1 because of the controversial association between serum TGF- $\beta$ 1 levels and the clinicopathological characteristics of gastric adenocarcinoma [86–88].

## Other proteins

In addition to the aforementioned ones, some proteins in both exosomes and other body fluid including serum and bile are associated with the proliferation, invasion and metastasis of GC. HSPs exist extensively and are highly conservative. Hoshino et al. [89] demonstrated that HSP-70 had the highest specificity and 3rd sensitivity among six tumor-associated antigens (p53, heat shock protein 70, HCC-22–5, peroxiredoxin VI, KM-HN-1, and p90) and their panel. A study by Zhong et al. [90], including 18 gastric adenocarcinoma patients, showed that the concentrations of HSP-60 and HSP-70 in exosomes from heat-treated malignant ascites in gastric adenocarcinoma patients were higher than that from untreated ones. In addition, exosomes from heat-treated malignant ascites have been shown to contribute to the maturation of dendritic cells and cause a tumor-specific cytotoxic T lymphocyte response, suggesting an association between HSP and gastric adenocarcinoma. CD97 is a member of EGF-seven transmembrane (EGF-TM7) with a molecular weight of 75–90 kDa. Li et al. [91] found that CD97 facilitates the proliferation and invasion of GC cells through the exosome-mediated mitogen-activated protein kinase (MAPK) signaling pathway by exosome extraction proliferation and Matrigel invasion assays. Liu et al. [92] isolated exosomes from SGC-L cells (a cell line derived from SGC-7901 with a high lymphatic metastasis) and SGC-L cells with CD97-knockdown (SGC-L/CD97-kd), and co-cultured them with GC cells. The results revealed that exosomes from SGC-L cells increased proliferation and invasion by 20 and 30%, respectively. Also, intrafootpad injections of SGC-L cells can significantly conduce to the two aforementioned cell lines accumulation in the draining lymph nodes and enhance the expression of CD55, CD31, CD151, CD44v6,  $\alpha 5\beta 1$  and epithelial cell adhesion molecule, which suggested that the exosome-dependent CD97 plays a pivotal role in lymphatic metastasis in SGC-L cells.

## Exosomal proteins involved in GC progression

### GKN1 and TRIM3

In addition to their diagnostic potential, exosomal GKN1 and TRIM3 also get involved in the progression of GC. Exosomal GKN1 could be internalized by the gastric epithelium and delay the multiplication of AGS and MKN1 cells as previously mentioned, which could be a significant mechanism of body's self-protection to cope with GC [50]. A study by Fu et al. [2] showed that the growth and metastasis of GC can be restrained by the regulation of stem cell factors and EMT regulators induced by exosomal TRIM3.

## TGF- $\beta$ superfamily

The TGF- $\beta$  superfamily contains a huge range of cytokines participating in different biological activities, such as polypeptide function in the regulation of tumor cells, tumor-related fibroblasts and immune-related cells that exist in the tumor microenvironment [93–95]. The TGF- $\beta$  superfamily includes TGF- $\beta$ , activin and bone morphogenetic protein (BMP), which play a crucial role in regulating the growth, adhesion, migration, differentiation and apoptosis of cells. There are three subtypes of TGF- $\beta$  found in mammals: TGF- $\beta 1$ , TGF- $\beta 2$  and TGF- $\beta 3$  [96]. TGF- $\beta 1$ , most generally expressed in human tissues, has three receptors: transforming growth factor beta receptors 1 (TGF- $\beta R1$ ), TGF- $\beta R2$  and TGF- $\beta R3$  [97]. A strong correlation between TGF- $\beta$  and GC has been confirmed. A study by Liu et al. [93] indicated that the decrease in the mRNA expression of TGF- $\beta 1$  and its receptors was associated with the improved overall survival (OS) of GC patients. In addition, their knockout restrained the migration, invasion and proliferation of gastric carcinoma cells. Yang et al. [97] found that advanced GC patients with a low TGF- $\beta 1$  expression often had a favorable OS, while the high levels of TGF- $\beta 1$  were markedly associated with poor OS, and GC patients with a high TGF- $\beta 2$  expression had a better OS than that of patients with a low TGF- $\beta 2$  expression, as determined by Cancer Genome Atlas database analysis. BMPs, which are members of the TGF- $\beta$  superfamily, are glycoproteins produced by certain cells. BMPs could fall under different subfamilies, based on amino acid residues sequence homology: (i) BMP-2 and BMP-2B (BMP-4); (ii) BMP-3 and BMP-3B; (iii) BMP-5, BMP-6, BMP-7, BMP-8, BMP-9, BMP-10, and BMP-11; (iv) BMP-12, BMP-13, BMP-14 and BMP-15. However, BMP-1 does not belong to this family [98]. BMPs participate in two signaling pathways: The Smad-dependent and Smad-independent pathways [98, 99]. Studies have suggested that BMPs are involved in the progression of GC. Deng et al. [100] demonstrated that BMP4 was overexpressed in GC cell lines and facilitated the EMT and metastasis of GC cells in vitro and in vivo, the knockdown of which markedly suppressed the EMT and metastasis of GC cells. Moreover, the expression of BMP4 was constantly upregulated in gastric carcinoma tissues and involved in the harsh prognosis of GC patients [100]. Lei et al. [101] performed a study on BMP10 in GC cell lines, 52 GC tissues and normal tissues. That study confirmed that the high BMP10 expression in AGS and SGC-7901 cell lines hindered the growth and migration of these cells, with BMP10-knockdown promoting their growth, migration, and metastasis. They also found that BMP10 was downregulated in GC tissues. These results showed the inhibitory effect of BMP10 on GC. BMP1, as the special one, was also found to be associated with GC. A study by Hsieh et al. [102] suggested that the inhibition

of BMP1 impeded GC cell migration, which had a negative impact on metastasis. In addition, BMP1-overexpression was linked with a very poor long-term survival of GC patients at an advanced stage.

Exosomal TGF- $\beta$ 1 and BMP have also been found to participate in the progression of stomach cancer. In addition to predicting the LNM of GC, exosomal TGF- $\beta$ 1 is also involved in the advancement of GC. It can assist GC cells in escaping immunological surveillance by causing regulatory T cell differentiation, making GC hard to manage. Furthermore, exosomal TGF- $\beta$  from SGC-7901 cells has been proven to lead to the activation of the Smad pathway, and the succedent differentiation of human umbilical cord mesenchymal stem cells (hucMSCs) into cancer-associated fibroblasts (CAFs) through interacting with TGF- $\beta$ R1 in hucMSCs [103]. Ning et al. [104] extracted exosomes from human gastric mucosal epithelial cells GES-1 and GC cells SGC-7901 and tested CAFs marker expression and CAFs conversion-related signaling pathways through WB, immunofluorescent staining and reverse transcription quantitative polymerase chain reaction (RT-qPCR). It was revealed that BMP2 in exosomes activated the PI3K/AKT and MEK/ERK signaling pathways and resulted in the transition of pericytes to CAFs.

### Epidermal growth factor receptor (EGFR)

EGFR, also known as human epidermal growth factor receptor 1 (HER1), is a member of the HER (ErbB) family. Other members of this family are ErbB-2 (HER2), ErbB-3 (HER3), and ErbB-4 (HER4) [105]. EGFR, and the rest of the ErbB family members, contain an extracellular domain, lipophilic transmembrane region, intracellular domain containing tyrosine kinase and carboxy-terminal region [106]. EGFR is transmembrane glycoprotein, which belongs to the receptor tyrosine kinase group and contains an extracellular domain, lipophilic transmembrane region, intracellular domain containing tyrosine kinase and a carboxy-terminal region, like the rest of the ErbB family members [106, 107]. The activation of EGFR has been shown to improve cell growth, proliferation and differentiation, as well as wound healing [108, 109]. Ample evidence has revealed a strong association between EGFR and gastric diseases. The phosphorylation of EGFR has been linked to DNA injury in gastric epithelial cells infected with *Helicobacter pylori* (*H. pylori*). Enhanced levels of activated EGFR in the epithelium were observed in gastritis and atrophic gastritis, but were not observed in intestinal metaplasia, and intestinal and diffuse type gastric carcinoma. In addition, the phosphorylated EGFR level in the gastric tissues of patients with gastric disease developing to intestinal metaplasia or dysplasia were higher than those of patients without disease progression [110]. The overexpression of EGFR, which is associated

with poor prognosis, was found in 37% of gastric/esophageal adenocarcinomas [111, 112]. Zhang et al. [113] suggested that overexpressed EGFR notably heralded an adverse outcome, and could function as an indicator of poor prognosis in GC patients. In a study by Terragni et al. [114], 8 cases with a strong EGFR expression were identified among 19 canine cases (5 gastric adenomas, 5 intestinal type gastric carcinomas and 9 diffuse-type gastric carcinomas) despite the position or biological behavior of the tumor.

Exosomal EGFR was found to be linked to GC progression. Zhang et al. [115] isolated exosomes from cell culture medium and the serum of gastric adenocarcinoma patients. Exosomes extracted from SGC7901 were co-cultured with primary mouse liver cells. Next, the effect of these exosomes on hepatocyte growth factor (HGF) in liver cells was examined, and the liver HGF's impact on promoting the invasion and metastasis of SGC7901 cells in vitro, as well as its effects on the formation and growth of liver metastases in vivo, were evaluated. That study demonstrated that EGFR was enriched in exosomes derived from patients with gastric adenocarcinoma, an effect opposite to that observed in exosomes of the healthy control. EGFR could be transported to the liver and integrated on the plasma membrane of liver stromal cells. Exosomal EGFR suppressed the expression of miR-26a/b and activated HGF, which combined with the cell mesenchymal-epithelial transition factor (c-MET) receptor on the migrated cancer cells and promoted the landing and proliferation of these cells, offering beneficial conditions for the hepatic metastasis of GC.

### Cytotoxin-associated gene A (CagA)

CagA is a protein encoded by CagA with a molecular weight of 128–145 kDa. It contains a structured N-terminal region that includes domain I, domain II and domain III and accounts for 70% of the entire CagA, and an intrinsically disordered/unstructured C-terminal tail, whose polymorphisms in structure induce the change of CagA molecular weight [116–118]. CagA, on the basis of a repetitive amino acid sequence known as glutamine acid-proline-isoleucine-tyrosine-alanine at its 3' terminal, can be segmented into two types, namely CagA of Western Asia and that of East Asia [119]. CagA-positive *H. pylori* is linked to gastric adenocarcinoma and gastric mucosa-associated lymphoid tissue lymphoma of B cell origin, and CagA serves as a main virulence factor of *H. pylori*, which makes *H. pylori* cagA-positive strains the most severe risk factor of GC. The transmission of CagA into the cytoplasm of the host cell has been shown to play a vital role in *H. pylori* pathogenesis and GC progression [120, 121]. Na et al. [122] reported that patients with poorly differentiated adenocarcinoma and signet ring cell carcinoma of individuals with early GC (EGC) and *H. pylori* CagA-positive infection exhibited an enhanced



methylation of Runt-related transcription factor 3. Simán et al. [123] also drew the conclusion that *H. pylori* CagA-positive infection is a risk factor of non-cardia gastric adenocarcinoma. Takahashi-Kanemitsu et al. [124] revealed that the non-physiological scaffolding actions of CagA granted cells numerous phenotypic tumor markers and intensified the malignant conversion, namely continuing proliferation, invasion, growth suppressors elusion, etc. In addition, under chronic inflammation, the tumorigenic activity of CagA was further reinforced [125]. Palrasu et al. [126] suggested that CagA caused the phosphorylation of X-linked inhibitor of apoptosis protein E3 ubiquitin ligase and intensified ubiquitination and proteasomal degradation of the host proapoptotic factor Siva1 which was associated with the restricting effects of *H. pylori* on apoptotic responses. Furthermore, that study proved that CagA hampered apoptosis and caused an effect that may enhance the survival of impaired epithelial cells and contribute to gastric tumorigenesis.

CagA in exosomes has also been linked to the development of gastric disease. Asako Shimoda et al. [127] detected exosomal CagA in the serum of 4 GC patients with CagA-positive *H. pylori* infection. They found CagA-containing exosomes caused morphological changes in GC and gastric epithelial cells, which suggested that functional CagA transported by exosomes into cells could be associated with the progression of extragastric disorders involved in CagA-positive *H. pylori* infection.

### Apolipoprotein E (ApoE)

ApoE is a small secretory glycoprotein with a molecular weight of 39 kDa located at chromosome 19q13.2 [128, 129]. The genetic structure of apoE is polymorphous with two single nucleotide polymorphisms in the coding region, which are divided into three different alleles ( $\epsilon 2$ ,  $\epsilon 3$  and  $\epsilon 4$ ) and six apoE genotypes [three homozygotes ( $\epsilon 4/\epsilon 4$ ,  $\epsilon 3/\epsilon 3$  and  $\epsilon 2/\epsilon 2$ , and three heterozygotes ( $\epsilon 4/\epsilon 3$ ,  $\epsilon 3/\epsilon 2$ ,  $\epsilon 4/\epsilon 2$ )] [128, 129]. ApoE combines with the low-density lipoprotein receptor and thus impacts cholesterol transport, lipid metabolism and protein synthesis; Moreover, it is involved with various signal transductions by bonding with the receptors on lipid particles [130–133]. Evidence has verified the participation of ApoE in various biological processes, including antioxidant activity, tissue repair, immune response and regulation, and cell growth, proliferation, differentiation and metastasis [134–139]. In terms of gastric carcinoma, studies have proven that ApoE influences GC progression and prognosis. Sakashita et al. [140] detected the ApoE mRNA through RT-qPCR and revealed that its expression in GC tissues was enhanced, as compared with that in normal mucosas. ApoE mRNA was overexpressed with further invasion into the muscle and serosa, and more positive LNM. Moreover, GC patients with ApoE-overexpression had a worse

survival than those with ApoE-downregulation. These findings suggested that ApoE might be a biomarker of survival and predicting gastric invasion. Kang et al. [141] conducted a study on 550 GC patients and an equal number of healthy individuals and observed that the existence of APOE  $\epsilon 2$  and reduction of total cholesterol were associated with an increased risk of GC. Furthermore, there was a connection between APOE  $\epsilon 2$  and an increased risk of intestinal and diffuse histotypes of GC; completely different conditions were obtained with regards to the association between APOE  $\epsilon 2$  and tumor node classification or stage of GC patients, with APOE polymorphic alleles found to be linked to a developmental risk of GC and irrelevant to its progression.

A correlation between exosomal ApoE and GC progression was verified. Research of Zheng et al. [142] indicated that ApoE was recognized as a protein with a sharp specificity in exosomes from M2 macrophages. In addition, functional ApoE exosomes delivered from tumor-associated macrophages (TAMs) to cancer cells could induce the activation of the PI3K/Akt signaling pathway, increasing the metastasis of GC cells.

### Ubiquitin protein ligase E3 component n-recognin 2 (UBR2)

UBR2 is a member of the E3 family with a molecular weight of 200 kDa [143, 144]. It is associated with chromatin and spermatogenesis, and regulates chromatin dynamics and gene expression in germ and somatic cells [144, 145]. Kwon et al. [146] revealed that the UBR2 gene was overexpressed in germ cells and involved in the susceptibility to infertility caused by the meiotic arrest and apoptosis of germ cells. As an ingredient of the N-end rule pathway, UBR2 induces ubiquitination and degradation [147]. It has been proven to serve as an ubiquitin ligase in regulating the activation of nucleotide-binding domain leucine-rich repeat pyrin domain-containing 1B inflammasome caused by anthrax lethal toxin via working with the ubiquitin-conjugating enzyme UBE2O [148]. In addition, UBR2 is likely to be an indispensable part on regulating the muscle protein degradation in tumor cachexia [149]. Besides, UBR2 is associated with tumors. It is currently known as the only N-end rule pathway E3 whose expression is increased by cachectic stimuli, including tumor and proinflammatory cytokines [147].

Despite the small amount of studies on exosomes, UBR2 and GC, exosomal UBR2 remains linked to stomach cancer. A previous study by Mao et al. [150] found that UBR2 existed greatly in exosomes stemming from mesenchymal stem cells. Next, that study confirmed an abundance of UBR2 in exosomes produced by p53-deficient mouse bone marrow mesenchymal stem cells (p53<sup>-/-</sup>mBMMSC), as compared with p53 wild-type mouse bone marrow mesenchymal stem cells (p53<sup>+/+</sup>mBMMSC). In addition, exosomes

rich in UBR2 from p53<sup>-/-</sup>mBMMSC were delivered into p53<sup>+/+</sup>mBMMSC and murine foregastric carcinoma cells and then enhanced UBR2 expression in these cells, amplifying growth and migration through the activation of the Wnt/ $\beta$ -catenin pathway.

### High mobility group box 1 (HMGB1)

HMGB1 is a member of HMGB which belongs to the HMG protein superfamily. HMG is considered as the 2nd most plentiful protein in cells [151]. It can enhance the fine tuning of transcription for dealing with fast environmental changes by interacting with nucleosomes, transcription factors, nucleosome remodeling complexes, and histone H1, and has a global role in establishing chromatin domains [151, 152]. The HMGB family is a family of chromosomal proteins participating in DNA replication, recombination, transcription and repair [153–155]. Previous studies have demonstrated that HMGB is associated with some tumors and their characteristics, such as tumor proliferation, metastasis, apoptosis escape and invasion of tissue [156, 157]. HMGB1, extensively expressed in the nucleated cells of mammals, is a nuclear and extracellular protein with a molecular weight of 25–30 kDa which combines with nuclear proteins [158, 159]; it has extensive biological functions. First, HMGB1 is involved in the aforementioned DNA-related activities [160]. Secondly, it can be secreted into the extracellular space via active secretion or passive release and combine with the corresponding receptors and then activate several pivotal cell-signaling pathways, thus participating in the initiation and progression of tumors and other diseases [161–163]. In terms of GC progression, Zhang et al. [164] found that HMGB1-overexpression was linked to the migration of gastric adenocarcinoma (GAC), and its knockdown hindered the growth and invasion of GAC cells via the NF- $\kappa$ B pathway *in vitro* and *in vivo*. A study by Wang et al. [165] indicated that HMGB1 partially participated in the promotion of GC cell proliferation and the potential of metastasis induced by intensive human antigen R. A study by Tian et al. [159] reported the suppression of stomach cancer cell proliferation and colony formation, and the induction of apoptosis following HMGB1-knockdown. Furthermore, its silencing markedly impeded the migration and invasion of GC cells. Moreover, HMGB1 is linked to GC angiogenesis and apoptosis. Chung et al. [166] conducted a series of analyses on 27 GC patients (10 EGC, 10 advanced GC-M0 and 7 advanced GC-M1), as well as GC cells, and reported that the enhanced expression of HMGB1 promoted angiogenesis through the mediation of IL-8. In addition, the joint targeting of HMGB1 and IL-8 was able to regulate GC angiogenesis [166]. Tao et al. [167] found that aloin reduced the expression and release of HMGB1 and then prevented the recombinant human HMGB1-induced Akt-mTOR-P70S6K

and ERK-P90RSK-cAMP regulatory element binding signaling pathways from activating, which caused apoptosis in GC HGC-27 cells. There were also correlations between HMGB1 and the diagnosis and prognosis of GC. Ghweil et al. [168] showed that HMGB1 had a relatively good diagnostic efficiency in differentiating patients with gastritis from those with GC (diffuse and intestinal GC). An increased level of serum amyloid A and HMGB1 signified a higher grade and advanced stage of GC [167]. As compared with normal tissues, the expression of HMGB1, HMGB2 and HMGB3 was promoted in GC tissues, which could portend a bad prognosis of GC patients [151].

HMGB1 in exosomes has been proven to be associated with GC metastasis. Zhang et al. [169] conducted an analysis of exosomal proteins derived from GC cells (BGC-823, MGC-803 and SGC-7901) via WB and LC-MS/MS. It was then found that HMGB1 played a crucial role in the impetus of the pro-tumor activation of neutrophils. In addition, GC exosomes were found to transport HMGB1 to interact with toll-like receptor 4, which induced the activation of the NF- $\kappa$ B pathway, an enhanced autophagic response and promotion of GC cell immigration. Previous study also revealed that extracellular vesicles (EVs) of GC cell were capable of causing pro-tumor activation of neutrophils, which had as similar effect to that of HMGB1 mentioned above [170]. Their study also demonstrated that EVs, including exosomes, stemmed from the microenvironment of GC transported HMGB1 to induce the activation of signal transducer and activator of transcription 3 (STAT3) and the high expression of the programmed death-ligand 1 (PD-L1) gene, which could be reversed via STAT3 pathway prohibition and HMGB1 silencing. These findings offered an updated insight on neutrophils' pro-tumor effect in GC and revealed multiple functions of EVs in coordinating an immunosuppressed microenvironment.

### MET

MET, also known as the c-MET or HGF receptor, is a transmembrane protein located on chromosome 7 [171, 172]. The extracellular domain of mature MET mainly includes a semaphorin (sema) domain (25–516), a plexin-sema-integrin domain (517–562), and 4 immunoglobulin-like regions in the plexins and transcription factors domain (563–932). Its intracellular region consists of a juxtamembrane sequence, a catalytic region and a carboxy-terminal multifunctional docking site [173]. As an HGF receptor, MET is involved in the activation of HGF, which is normally regulated by mechanisms such as paracrine ligand delivery. Even with these mechanisms, MET-related signaling pathways are also associated with oncogenesis and the progression of several types of cancer, including GC [174, 175]. MET can also impact the progression of cancer by activating other critical

pathways (e.g., PI3K/AKT, Ras/MAPK, etc.) [176–178]. Furthermore, there is a correlation between MET and the bleak prognosis of GC patients. Ha et al. [179] reported that GC patients with a highly expressed MET had worse OS and disease-free survival than patients without MET over-expression. A study by Drebber et al. [180] revealed that the immunoreactivity of MET and p21 were independent indicators of prognosis in gastric adenocarcinoma. A study conducted by Zhang et al. [181] confirmed the intensified expression of MET in gastric adenocarcinoma patients with a poor prognosis. In addition, the enhanced expression rate of MET was higher in gastric adenocarcinoma patients with a positive HER2, suggesting that MET could serve as a predictive biomarker for GC patients.

Exosomal MET also participates in tumor progression. Che et al. [78] investigated the communication exosomes involved between GC cells infected by *H. pylori* and macrophages. They found that the MET expression, particularly the phosphorylated active form in exosomes secreted by *H. pylori*-infected GC cells, was enhanced. The exosomes involving MET were transferred to macrophages and then internalized, after which exosomal MET seemingly induced the conversion of macrophages to a pro-tumorigenesis phenotype. All these discoveries confirmed that the expression of activated exosomal MET was intensified by *H. pylori* infection, and exosomal MET could have a pro-tumorigenic effect on macrophages.

## Challenges and prospects

In spite of the advantage of exosome on diagnosis and prognosis of GC, obstacles to the application of exosomal proteins should not be ignored. First, the lack of standardized extraction methods of exosomes makes it difficult to eliminate heteroprotein contamination, which has a negative impact on the research of exosomal proteins. Secondly, despite the proven potential of GC markers and their reported excellent stability and sensitivity, potential exosomal proteins markers mentioned above, such as TRIM3, lack deep fundamental research on specificity, sensitivity and application in GC diagnosis and prognosis, which is un conducive to their clinical application. Research on exosome proteomics is a fantastic way of exploring GC marker. Ding et al. [182] discovered 443 exosomal proteins, among which 110 proteins were differentially expressed, by quantitative proteomics analyses. Besides, 10 highly vital proteins (UBA52, PSMA1, PSMA5, PSMB6, PSMA7, PSMA4, PSMA3, PSMB1, PSMA6, and FGA) were observed from 110 differentially expressed proteins. However, researches on exosome proteomics for diagnosis are highly limited, which should raise

concern, especially when it comes to early diagnosis that is of primary importance in GC diagnosis. Thirdly, the large demand of exosomal samples and complex process of exosomal protein analysis are due to the absence of a general amplification technique [183]. The development of rapid and high throughput detection of exosomal proteins is of great significance to their clinical application. On the other hand, exosomal proteins have shown several diagnostic advantages compared to conventional GC markers. More researches on exosomal proteins are still definitely needed. Among all these barriers, extraction and purification of exosomes, which are the basis for subsequent research, need to be improved urgently.

## Conclusion

Exosomes have been proven to participate in various biological processes, such as angiogenesis, antigen presentation, apoptosis and intercellular signaling [15]. Exosomal proteins can also be involved in the progression of many diseases, including cancer. Proteins with a potential diagnostic and prognostic role in GC, as confirmed by previous studies, are listed in Table 1. Exosomal proteins have been reported to have a preferable stability and sensitivity, and provide relevant information “in real time” during tumorigenesis and tumor development, as compared with conventional diagnostic GC markers [115]. Moreover, inspiring diagnostic performance is expected to appear for the advantage of quick sampling, less pain and low cost of exosomal protein detection. Nevertheless, challenges such as the lack of appropriate extraction methods and intensive study on exosomal proteins remain obvious. In conclusion, exosomal proteins are capable of overcoming the aforementioned obstacles and being used in the diagnosis and prognosis of GC as relevant research progresses.

**Acknowledgements** This study was supported by grants from Basic research project of key scientific research project plan of colleges and universities in Henan Province (Grant Nos. 19zx009)

**Authors' contributions** HS drafted the article, WHR revised the article and provided ideas and DZ helped to improve language accuracy.

**Funding** This study was supported by grants from Basic research project of key scientific research project plan of colleges and universities in Henan Province (Grant nos. 19zx009); Science and Technology Project for Tackling Key Problems of Henan Province (Grant number 212102310639).

## Declarations

**Conflict of interest** The authors declare that they have no competing interests.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin*. 2015;65:87–108.
- Fu H, Yang H, Zhang X, et al. Exosomal TRIM3 is a novel marker and therapy target for gastric cancer. *J Exp Clin Cancer Res CR*. 2018;37:162.
- Beach A, Zhang HG, Ratajczak MZ, Kakar SS. Exosomes: an overview of biogenesis, composition and role in ovarian cancer. *J Ovarian Res*. 2014;7:14.
- Levy S. Function of the Tetraspanin molecule CD81 in B and T cells. *Immunol Res*. 2014;58:179–85.
- Jiang L, Gu Y, Du Y, Liu J. Exosomes: diagnostic biomarkers and therapeutic delivery vehicles for cancer. *Mol Pharm*. 2019;16:3333–49.
- Qian X, Xie F, Wei H, Cui D. Identification of key circulating exosomal microRNAs in gastric cancer. *Front Oncol*. 2021;11:693360.
- Han Z, Li K, Wu J, et al. Diagnostic value of RNA for hepatocellular carcinoma: a network meta-analysis. *Biomark Med*. 2021;15(18):1755–1767.
- Han W, Bu X, Liu Y, et al. Clinical value of miR-135 and miR-20a combined with multi-detector computed tomography in the diagnosis of gastric cancer. *World J Surg Oncol*. 2021;19:283.
- Tong Y, Wu Y, Song Z, Yu Y, Yu X. The potential value of serum pepsinogen for the diagnosis of atrophic gastritis among the health check-up populations in China: a diagnostic clinical research. *BMC Gastroenterol*. 2017;17:88.
- Tu H, Sun L, Dong X, et al. A serological biopsy using five stomach-specific circulating biomarkers for gastric cancer risk assessment: a multi-phase study. *Am J Gastroenterol*. 2017;112:704–15.
- Yu G, Wang GX, Wang HG, Mo FF, Tang BB. The value of detecting pepsinogen and gastrin-17 levels in serum for precancerous lesion screening in gastric cancer. *Neoplasma*. 2019;66:637–40.
- Abbas M, Habib M, Naveed M, et al. The relevance of gastric cancer biomarkers in prognosis and pre- and post-chemotherapy in clinical practice. *Biomed Pharmacother Biomed Pharmacother*. 2017;95:1082–90.
- Osaki M, Okada F. Exosomes and their role in cancer progression. *Yonago Acta Med*. 2019;62:182–90.
- Wu K, Xing F, Wu SY, Watabe K. Extracellular vesicles as emerging targets in cancer: Recent development from bench to bedside. *Biochim Biophys Acta Rev Cancer*. 2017;1868:538–63.
- Gurunathan S, Kang MH, Jeyaraj M, Qasim M, Kim JH. Review of the isolation, characterization, biological function, and multifarious therapeutic approaches of exosomes. *Cells*. 2019;8(4):307.
- Vanni I, Alama A, Grossi F, Dal Bello MG, Coco S. Exosomes: a new horizon in lung cancer. *Drug Discov Today*. 2017;22:927–36.
- Vidal M. Exosomes: revisiting their role as 'garbage bags'. *Traffic (Copenhagen, Denmark)*. 2019;20(11):815–828.
- Johnstone RM, Adam M, Hammond JR, Orr L, Turbide C. Vesicle formation during reticulocyte maturation. Association of plasma membrane activities with released vesicles (exosomes). *J Biol Chem*. 1987;262:9412–20.
- Royo F, Théry C, Falcón-Pérez JM, Nieuwland R and Witwer KW. Methods for separation and characterization of extracellular vesicles: results of a worldwide survey performed by the ISEV Rigor and Standardization Subcommittee. *Cells*. 2020;9(9):1955.
- Yu Z, Lin S, Xia F, et al. ExoSD chips for high-purity immunomagnetic separation and high-sensitivity detection of gastric cancer cell-derived exosomes. *Biosens Bioelectron*. 2021;194:113594.
- Lin S, Yu Z, Chen D, et al. Progress in microfluidics-based exosome separation and detection technologies for diagnostic applications. *Small*. 2020;16:e1903916.
- Liga A, Vliegenthart AD, Oosthuizen W, Dear JW, Kersaudy-Kerhoas M. Exosome isolation: a microfluidic road-map. *Lab Chip*. 2015;15:2388–94.
- Pietrowska M, Funk S, Gawin M, et al. Isolation of exosomes for the purpose of protein cargo analysis with the use of mass spectrometry. *Methods Mol Biol (Clifton, NJ)*. 2017;1654:291–307.
- Yang F, Liao X, Tian Y and Li G. Exosome separation using microfluidic systems: size-based, immunoaffinity-based and dynamic methodologies. *Biotechnol J*. 2017;12(4). <https://doi.org/10.1002/biot.201600699>.
- Wei H, Qian X, Xie F, Cui D. Isolation of exosomes from serum of patients with lung cancer: a comparison of the ultra-high speed centrifugation and precipitation methods. *Ann Transl Med*. 2021;9:882.
- Yuana Y, Oosterkamp TH, Bahatyrova S, et al. Atomic force microscopy: a novel approach to the detection of nanosized blood microparticles. *J Thromb Haemost JTH*. 2010;8:315–23.
- Friedrich R, Block S, Alizadehheidari M, et al. A nano flow cytometer for single lipid vesicle analysis. *Lab Chip*. 2017;17:830–41.
- Zhang Y, Bi J, Huang J, Tang Y, Du S, Li P. Exosome: a review of its classification, isolation techniques, storage, diagnostic and targeted therapy applications. *Int J Nanomed*. 2020;15:6917–34.
- Elsharkasy OM, Nordin JZ, Hagey DW, et al. Extracellular vesicles as drug delivery systems: why and how? *Adv Drug Deliv Rev*. 2020;159:332–43.
- Batrakova EV, Kim MS. Using exosomes, naturally-equipped nanocarriers, for drug delivery. *J Control Release Off J Control Release Soc*. 2015;219:396–405.
- Pan S, Pei L, Zhang A, et al. Passion fruit-like exosome-PMA/Au-BSA@Ce6 nanovehicles for real-time fluorescence imaging and enhanced targeted photodynamic therapy with deep penetration and superior retention behavior in tumor. *Biomaterials*. 2020;230:119606.
- Liang G, Zhu Y, Ali DJ, et al. Engineered exosomes for targeted co-delivery of miR-21 inhibitor and chemotherapeutics to reverse drug resistance in colon cancer. *J Nanobiotechnol*. 2020;18:10.
- Xu W, Yang Z, Lu N. From pathogenesis to clinical application: insights into exosomes as transfer vectors in cancer. *J Exp Clin Cancer Res CR*. 2016;35:156.
- El-Husseini AE, Fretier P, Vincent SR. Cloning and characterization of a gene (RNF22) encoding a novel brain expressed ring finger protein (BERP) that maps to human chromosome 11p15.5. *Genomics*. 2001;71:363–7.



35. Koi M, Johnson LA, Kalikin LM, Little PF, Nakamura Y, Feinberg AP. Tumor cell growth arrest caused by subchromosomal transferable DNA fragments from chromosome 11. *Science* (New York, NY). 1993;260:361–4.
36. Kawai T, Akira S. Regulation of innate immune signalling pathways by the tripartite motif (TRIM) family proteins. *EMBO Mol Med*. 2011;3:513–27.
37. Nisole S, Stoye JP, Saïb A. TRIM family proteins: retroviral restriction and antiviral defence. *Nat Rev Microbiol*. 2005;3:799–808.
38. Ozato K, Shin DM, Chang TH, Morse HC 3rd. TRIM family proteins and their emerging roles in innate immunity. *Nat Rev Immunol*. 2008;8:849–60.
39. Song Y, Guo Q, Gao S, Hua K. Tripartite motif-containing protein 3 plays a role of tumor inhibitor in cervical cancer. *Biochem Biophys Res Commun*. 2018;498:686–92.
40. Piao MY, Cao HL, He NN, et al. Potential role of TRIM3 as a novel tumour suppressor in colorectal cancer (CRC) development. *Scand J Gastroenterol*. 2016;51:572–82.
41. Wang M, Wu J, Guo Y, Chang X, Cheng T. The tripartite motif-containing protein 3 on the proliferation and cytokine secretion of rheumatoid arthritis fibroblast-like synoviocytes. *Mol Med Rep*. 2017;15:1607–12.
42. Li Y, Zhu H, Wang J, Qian X, Li N. miR-4513 promotes breast cancer progression through targeting TRIM3. *Am J Transl Res*. 2019;11:2431–8.
43. Lu Q, Zhang Y, Ma L, et al. TRIM3 negatively regulates autophagy through promoting degradation of Beclin1 in ewing sarcoma cells. *Onco Targets Ther*. 2019;12:11587–95.
44. Song Y, Guo Q, Gao S, Hua K. miR-454-3p promotes proliferation and induces apoptosis in human cervical cancer cells by targeting TRIM3. *Biochem Biophys Res Commun*. 2019;516:872–9.
45. Wang M, Wu J, Zhou E, Chang X, Gan J, Cheng T. Forkhead box o3a suppresses lipopolysaccharide-stimulated proliferation and inflammation in fibroblast-like synoviocytes through regulating tripartite motif-containing protein 3. *J Cell Physiol*. 2019;234:20139–48.
46. Dong W, Qiu C, Gong D, et al. Proteomics and bioinformatics approaches for the identification of plasma biomarkers to detect Parkinson's disease. *Exp Ther Med*. 2019;18:2833–42.
47. Dokhaee F, Mazhari S, Galehdari M, Bahadori Monfared A, Baghaei K. Evaluation of GKN1 and GKN2 gene expression as a biomarker of gastric cancer. *Gastroenterol Hepatol Bed Bench*. 2018;11:S140-s145.
48. Koper-Lenkiewicz OM, Kamińska J, Gawrońska B, Matowicka-Karna J. The role and diagnostic potential of gastrokine 1 in gastric cancer. *Cancer Manag Res*. 2019;11:1921–31.
49. Martin TE, Powell CT, Wang Z, et al. A novel mitogenic protein that is highly expressed in cells of the gastric antrum mucosa. *American journal of physiology. Gastrointest Liver Physiol*. 2003;285:G332-343.
50. Yoon JH, Ham IH, Kim O, et al. Gastrokine 1 protein is a potential thernostic target for gastric cancer. *Gastric Cancer Off J Int Gastric Cancer Assoc Jpn Gastric Cancer Assoc*. 2018;21:956–67.
51. Yoshikawa Y, Mukai H, Hino F, Asada K, Kato I. Isolation of two novel genes, down-regulated in gastric cancer. *Jpn J Cancer Res Gann*. 2000;91:459–63.
52. Alarcón-Millán J, Martínez-Carrillo DN, Peralta-Zaragoza O, Fernández-Tilapa G. Regulation of GKN1 expression in gastric carcinogenesis: a problem to resolve (Review). *Int J Oncol*. 2019;55:555–69.
53. Menhenniott TR, Kurklu B, Giraud AS. Gastrokines: stomach-specific proteins with putative homeostatic and tumor suppressor roles. *American journal of physiology. Gastrointest Liver Physiol*. 2013;304:G109-121.
54. Moss SF, Lee JW, Sabo E, et al. Decreased expression of gastrokine 1 and the trefoil factor interacting protein TFIZ1/GKN2 in gastric cancer: influence of tumor histology and relationship to prognosis. *Clin Cancer Res Off J Am Assoc Cancer Res*. 2008;14:4161–7.
55. Sánchez-Pulido L, Devos D, Valencia A. BRICHOS: a conserved domain in proteins associated with dementia, respiratory distress and cancer. *Trends Biochem Sci*. 2002;27:329–32.
56. Hedlund J, Johansson J, Persson B. BRICHOS - a superfamily of multidomain proteins with diverse functions. *BMC Res Notes*. 2009;2:180.
57. Pavone LM, Del Vecchio P, Mallardo P, et al. Structural characterization and biological properties of human gastrokine 1. *Mol BioSyst*. 2013;9:412–21.
58. Yoon JH, Choi YJ, Choi WS, Nam SW, Lee JY, Park WS. Functional analysis of the NH2-terminal hydrophobic region and BRICHOS domain of GKN1. *Biochem Biophys Res Commun*. 2013;440:689–95.
59. Guo XY, Dong L, Qin B, Jiang J, Shi AM. Decreased expression of gastrokine 1 in gastric mucosa of gastric cancer patients. *World J Gastroenterol*. 2014;20:16702–6.
60. Kim O, Yoon JH, Choi WS, et al. Gastrokine 1 inhibits gastrin-induced cell proliferation. *Gastric Cancer Off J Int Gastric Cancer Assoc Jpn Gastric Cancer Assoc*. 2016;19:381–91.
61. Xing R, Li W, Cui J, et al. Gastrokine 1 induces senescence through p16/Rb pathway activation in gastric cancer cells. *Gut*. 2012;61:43–52.
62. Yoon JH, Cho ML, Choi YJ, et al. Gastrokine 1 regulates NF-κB signaling pathway and cytokine expression in gastric cancers. *J Cell Biochem*. 2013;114:1800–9.
63. Yoon JH, Park YG, Nam SW, Park WS. The diagnostic value of serum gastrokine 1 (GKN1) protein in gastric cancer. *Cancer Med*. 2019;8:5507–14.
64. Yoon JH, Ashktorab H, Smoot DT, Nam SW, Hur H, Park WS. Uptake and tumor-suppressive pathways of exosome-associated GKN1 protein in gastric epithelial cells. *Gastric Cancer Off J Int Gastric Cancer Assoc Jpn Gastric Cancer Assoc*. 2020.
65. Yen EY, Miaw SC, Yu JS, Lai IR. Exosomal TGF-β1 is correlated with lymphatic metastasis of gastric cancers. *Am J Cancer Res*. 2017;7:2199–208.
66. Zhang Q, Yu N, Lee C. Mysteries of TGF-β paradox in benign and malignant cells. *Front Oncol*. 2014;4:94.
67. Zhao JW, Wu ZH, Guo JW, et al. Synthesis and anti-gastric cancer activity evaluation of novel triazole nucleobase analogues containing steroidal/coumarin/quinoline moieties. *Eur J Med Chem*. 2019;181:111520.
68. Attisano L, Wrana JL. Signal transduction by members of the transforming growth factor-beta superfamily. *Cytokine Growth Factor Rev*. 1996;7:327–39.
69. Lyons RM, Gentry LE, Purchio AF, Moses HL. Mechanism of activation of latent recombinant transforming growth factor beta 1 by plasmin. *J Cell Biol*. 1990;110:1361–7.
70. Lyons RM, Keski-Oja J, Moses HL. Proteolytic activation of latent transforming growth factor-beta from fibroblast-conditioned medium. *J Cell Biol*. 1988;106:1659–65.
71. Oklü R, Hesketh R. The latent transforming growth factor beta binding protein (LTBP) family. *Biochem J*. 2000;352(Pt 3):601–10.
72. Massagué J, Blain SW, Lo RS. TGFbeta signaling in growth control, cancer, and heritable disorders. *Cell*. 2000;103:295–309.
73. Liu XS, Lin XK, Mei Y, et al. Regulatory T cells promote overexpression of Lgr5 on gastric cancer cells via TGF-beta1 and confer poor prognosis in gastric cancer. *Front Immunol*. 2019;10:1741.

74. Ikushima H, Miyazono K. TGFbeta signalling: a complex web in cancer progression. *Nat Rev Cancer*. 2010;10:415–24.
75. Achyut BR, Yang L. Transforming growth factor- $\beta$  in the gastrointestinal and hepatic tumor microenvironment. *Gastroenterology*. 2011;141:1167–78.
76. Pak KH, Kim DH, Kim H, Lee DH, Cheong JH. Differences in TGF- $\beta$ 1 signaling and clinicopathologic characteristics of histologic subtypes of gastric cancer. *BMC Cancer*. 2016;16:60.
77. Pak KH, Park KC, Cheong JH. VEGF-C induced by TGF- $\beta$ 1 signaling in gastric cancer enhances tumor-induced lymphangiogenesis. *BMC Cancer*. 2019;19:799.
78. Che Y, Geng B, Xu Y, et al. Helicobacter pylori-induced exosomal MET educates tumour-associated macrophages to promote gastric cancer progression. *J Cell Mol Med*. 2018;22:5708–19.
79. Zou Z, Zhao L, Su S, et al. The plasma levels of 12 cytokines and growth factors in patients with gastric cancer. *Medicine*. 2018;97:e0413.
80. Peng LS, Zhang JY, Teng YS, et al. Tumor-associated monocytes/macrophages impair NK-cell function via TGF $\beta$ 1 in human gastric cancer. *Cancer Immunol Res*. 2017;5:248–56.
81. Han B, Mao FY, Zhao YL, et al. Altered NKp30, NKp46, NKG2D, and DNAM-1 expression on circulating NK cells is associated with tumor progression in human gastric cancer. *J Immunol Res*. 2018;2018:6248590.
82. Coban S, Yüksel O, Koçkar MC, et al. The significance of serum transforming growth factor beta 1 in detecting of gastric and colon cancers. *Hepatogastroenterology*. 2007;54:1472–6.
83. Wang H, Zhao Y, Cao L, Zhang J, Wang Y, Xu M. Metastasis suppressor protein 1 regulated by PTEN suppresses invasion, migration, and EMT of gastric carcinoma by inactivating PI3K/AKT signaling. *J Cell Biochem*. 2019;120:3447–54.
84. Zhu J, Wen K. Astragaloside IV inhibits TGF- $\beta$ 1-induced epithelial-mesenchymal transition through inhibition of the PI3K/Akt/NF- $\kappa$ B pathway in gastric cancer cells. *Phytother Res PTR*. 2018;32:1289–96.
85. Wang B, Chen H and Zhang Y. Involvement of GASL1 in post-operative distant recurrence of gastric adenocarcinoma after gastrectomy distal resection and the possible mechanism. *J Cell Biochem*. 2019. <https://doi.org/10.1002/jcb.28423>.
86. Ma GF, Miao Q, Zeng XQ, et al. Transforming growth factor- $\beta$ 1 and - $\beta$ 2 in gastric precancer and cancer and roles in tumor-cell interactions with peripheral blood mononuclear cells in vitro. *PLoS ONE*. 2013;8:e54249.
87. Suda A, Saito N, Seshimo A, Kameoka S, Kobayashi M. Examination of transforming growth factor beta1 expression in the serum and tumor tissue of gastric cancer. *Int Surg*. 2009;94:182–8.
88. Li X, Yue ZC, Zhang YY, et al. Elevated serum level and gene polymorphisms of TGF-beta1 in gastric cancer. *J Clin Lab Anal*. 2008;22:164–71.
89. Hoshino I, Nagata M, Takiguchi N, et al. Panel of autoantibodies against multiple tumor-associated antigens for detecting gastric cancer. *Cancer Sci*. 2017;108:308–15.
90. Zhong H, Yang Y, Ma S, et al. Induction of a tumour-specific CTL response by exosomes isolated from heat-treated malignant ascites of gastric cancer patients. *Int J Hyperth Off J Eur Soc Hyperth Oncol North Am Hyperth Group*. 2011;27:604–11.
91. Li C, Liu DR, Li GG, et al. CD97 promotes gastric cancer cell proliferation and invasion through exosome-mediated MAPK signaling pathway. *World J Gastroenterol*. 2015;21:6215–28.
92. Liu D, Li C, Trojanowicz B, et al. CD97 promotion of gastric carcinoma lymphatic metastasis is exosome dependent. *Gastric Cancer Off J Int Gastric Cancer Assoc Jpn Gastric Cancer Assoc*. 2016;19:754–66.
93. Liu F, Wang H, Zhang M. Distinct prognostic values and antitumor effects of tumor growth factor  $\beta$ 1 and its receptors in gastric cancer. *Oncol Lett*. 2020;20:2621–32.
94. Qiao C, Yang J, Shen Q, et al. Traceable nanoparticles with dual targeting and ROS response for RNAi-based immunotherapy of intracranial glioblastoma treatment. *Adv Mater (Deerfield Beach, Fla)*. 2018;30:e1705054.
95. Li QL, Ito K, Sakakura C, et al. Causal relationship between the loss of RUNX3 expression and gastric cancer. *Cell*. 2002;109:113–24.
96. Memon MA, Anway MD, Covert TR, Uzumcu M, Skinner MK. Transforming growth factor beta (TGFbeta1, TGFbeta2 and TGFbeta3) null-mutant phenotypes in embryonic gonadal development. *Mol Cell Endocrinol*. 2008;294:70–80.
97. Yang T, Huang T, Zhang D, et al. TGF- $\beta$  receptor inhibitor LY2109761 enhances the radiosensitivity of gastric cancer by inactivating the TGF- $\beta$ /SMAD4 signaling pathway. *Aging*. 2019;11:8892–910.
98. Carreira AC, Zambuzzi WF, Rossi MC, Astorino Filho R, Sogayar MC, Granjeiro JM. bone morphogenetic proteins: promising molecules for bone healing, bioengineering, and regenerative medicine. *Vitam Horm*. 2015;99:293–322.
99. Cui Y, Zhang F, Jia Y, et al. The BMP antagonist, SOSTDC1, restrains gastric cancer progression via inactivation of c-Jun signaling. *Am J Cancer Res*. 2019;9:2331–48.
100. Deng G, Chen Y, Guo C, et al. BMP4 promotes the metastasis of gastric cancer by inducing epithelial-mesenchymal transition via ID1. *J Cell Sci*. 2020;133(11):jcs237222.
101. Lei H, Wang J, Lu P, et al. BMP10 inhibited the growth and migration of gastric cancer cells. *Tumour Biol J Int Soc Oncodev Biol Med*. 2016;37:3025–31.
102. Hsieh YY, Tung SY, Pan HY, et al. Upregulation of bone morphogenetic protein 1 is associated with poor prognosis of late-stage gastric Cancer patients. *BMC Cancer*. 2018;18:508.
103. Gu J, Qian H, Shen L, et al. Gastric cancer exosomes trigger differentiation of umbilical cord derived mesenchymal stem cells to carcinoma-associated fibroblasts through TGF- $\beta$ /Smad pathway. *PLoS ONE*. 2012;7:e52465.
104. Ning X, Zhang H, Wang C, Song X. Exosomes released by gastric cancer cells induce transition of pericytes into cancer-associated fibroblasts. *Med Sci Monit Int Med J Exp Clin Res*. 2018;24:2350–9.
105. Yarden Y, Sliwkowski MX. Untangling the ErbB signalling network. *Nat Rev Mol Cell Biol*. 2001;2:127–37.
106. Arienti C, Pignatta S, Tesei A. Epidermal growth factor receptor family and its role in gastric cancer. *Front Oncol*. 2019;9:1308.
107. Cohen S, Ushiro H, Stoscheck C, Chinkers M. A native 170,000 epidermal growth factor receptor-kinase complex from shed plasma membrane vesicles. *J Biol Chem*. 1982;257:1523–31.
108. Terashima M, Kitada K, Ochiai A, et al. Impact of expression of human epidermal growth factor receptors EGFR and ERBB2 on survival in stage II/III gastric cancer. *Clin Cancer Res Off J Am Assoc Cancer Res*. 2012;18:5992–6000.
109. Matsubara J, Yamada Y, Hirashima Y, et al. Impact of insulin-like growth factor type 1 receptor, epidermal growth factor receptor, and HER2 expressions on outcomes of patients with gastric cancer. *Clin Cancer Res Off J Am Assoc Cancer Res*. 2008;14:3022–9.
110. Chaturvedi R, Asim M, Piazuolo MB, et al. Activation of EGFR and ERBB2 by Helicobacter pylori results in survival of gastric epithelial cells with DNA damage. *Gastroenterology*. 2014;146:1739-1751.e1714.
111. Dulak AM, Schumacher SE, van Lieshout J, et al. Gastrointestinal adenocarcinomas of the esophagus, stomach, and colon exhibit distinct patterns of genome instability and oncogenesis. *Can Res*. 2012;72:4383–93.

112. Kim MA, Lee HS, Lee HE, Jeon YK, Yang HK, Kim WH. EGFR in gastric carcinomas: prognostic significance of protein overexpression and high gene copy number. *Histopathology*. 2008;52:738–46.
113. Zhang Z, Tang H, Lin J, et al. Clinicopathologic and prognostic significance of human epidermal growth factor receptor in patients with gastric cancer: an updated meta-analysis. *Oncotarget*. 2017;8:17202–15.
114. Terragni R, Casadei Gardini A, Sabattini S, et al. EGFR, HER-2 and KRAS in canine gastric epithelial tumors: a potential human model? *PLoS ONE*. 2014;9:e85388.
115. Zhang H, Deng T, Liu R, et al. Exosome-delivered EGFR regulates liver microenvironment to promote gastric cancer liver metastasis. *Nat Commun*. 2017;8:15016.
116. Covacci A, Censini S, Bugnoli M, et al. Molecular characterization of the 128-kDa immunodominant antigen of *Helicobacter pylori* associated with cytotoxicity and duodenal ulcer. *Proc Natl Acad Sci USA*. 1993;90:5791–5.
117. Hayashi T, Senda M, Morohashi H, et al. Tertiary structure-function analysis reveals the pathogenic signaling potentiation mechanism of *Helicobacter pylori* oncogenic effector CagA. *Cell Host Microbe*. 2012;12:20–33.
118. Kaplan-Türköz B, Jiménez-Soto LF, Dian C, et al. Structural insights into *Helicobacter pylori* oncoprotein CagA interaction with  $\beta 1$  integrin. *Proc Natl Acad Sci USA*. 2012;109:14640–5.
119. Sukri A, Hanafiah A, Mohamad Zin N, Kosai NR. Epidemiology and role of *Helicobacter pylori* virulence factors in gastric cancer carcinogenesis. *APMIS Acta Pathol Microbiol Immunol Scand*. 2020;128:150–61.
120. Ohnishi N, Yuasa H, Tanaka S, et al. Transgenic expression of *Helicobacter pylori* CagA induces gastrointestinal and hematopoietic neoplasms in mouse. *Proc Natl Acad Sci USA*. 2008;105:1003–8.
121. Park JY, Forman D, Waskito LA, Yamaoka Y and Crabtree JE. Epidemiology of *Helicobacter pylori* and CagA-positive infections and global variations in gastric cancer. *Toxins*. 2018;10(4):163.
122. Na YJ, Shim KN, Joo YH, et al. RUNX3 methylation, loss of RUNX3 expression and clinicopathologic findings according to *Helicobacter pylori* CagA in gastric carcinoma. *Korean J Gastroenterol Taehan Sohwagi Hakhoe Chi*. 2015;66:75–84.
123. Simán JH, Engstrand L, Berglund G, Forsgren A, Florén CH. *Helicobacter pylori* and CagA seropositivity and its association with gastric and oesophageal carcinoma. *Scand J Gastroenterol*. 2007;42:933–40.
124. Takahashi-Kanemitsu A, Knight CT, Hatakeyama M. Molecular anatomy and pathogenic actions of *Helicobacter pylori* CagA that underpin gastric carcinogenesis. *Cell Mol Immunol*. 2020;17:50–63.
125. Liu B, Li X, Sun F, et al. HP-CagA+ regulates the expression of CDK4/CyclinD1 via reg3 to change cell cycle and promote cell proliferation. *Int J Mol Sci*. 2019;21(1):224.
126. Palrasu M, Zaika E, El-Rifai W, et al. Bacterial CagA protein compromises tumor suppressor mechanisms in gastric epithelial cells. *J Clin Invest*. 2020;130:2422–34.
127. Shimoda A, Ueda K, Nishiumi S, et al. Exosomes as nanocarriers for systemic delivery of the *Helicobacter pylori* virulence factor CagA. *Sci Rep*. 2016;6:18346.
128. Mahley RW, Nathan BP, Pitas RE. Apolipoprotein E Structure, function, and possible roles in Alzheimer's disease. *Ann N Y Acad Sci*. 1996;777:139–45.
129. Moore LE, Brennan P, Karami S, et al. Apolipoprotein E/C1 locus variants modify renal cell carcinoma risk. *Can Res*. 2009;69:8001–8.
130. Li Y, Cam J, Bu G. Low-density lipoprotein receptor family: endocytosis and signal transduction. *Mol Neurobiol*. 2001;23:53–67.
131. Ho YY, Deckelbaum RJ, Chen Y, Vogel T, Talmage DA. Apolipoprotein E inhibits serum-stimulated cell proliferation and enhances serum-independent cell proliferation. *J Biol Chem*. 2001;276:43455–62.
132. Mahley RW, Rall SC Jr. Apolipoprotein E: far more than a lipid transport protein. *Annu Rev Genom Hum Genet*. 2000;1:507–37.
133. Shi X, Xu J, Wang J, et al. Expression analysis of apolipoprotein E and its associated genes in gastric cancer. *Oncol Lett*. 2015;10:1309–14.
134. Vogel T, Guo NH, Guy R, et al. Apolipoprotein E: a potent inhibitor of endothelial and tumor cell proliferation. *J Cell Biochem*. 1994;54:299–308.
135. Rubinsztein DC, Hanlon CS, Irving RM, et al. Apo E genotypes in multiple sclerosis, Parkinson's disease, schwannomas and late-onset Alzheimer's disease. *Mol Cell Probes*. 1994;8:519–25.
136. Ishigami M, Swertfeger DK, Hui MS, Granholm NA, Hui DY. Apolipoprotein E inhibition of vascular smooth muscle cell proliferation but not the inhibition of migration is mediated through activation of inducible nitric oxide synthase. *Arterioscler Thromb Vasc Biol*. 2000;20:1020–6.
137. Williams H, Johnson JL, Carson KG, Jackson CL. Characteristics of intact and ruptured atherosclerotic plaques in brachiocephalic arteries of apolipoprotein E knockout mice. *Arterioscler Thromb Vasc Biol*. 2002;22:788–92.
138. Weisgraber KH, Rall SC Jr, Mahley RW. Human E apoprotein heterogeneity. Cysteine-arginine interchanges in the amino acid sequence of the apo-E isoforms. *J Biol Chem*. 1981;256:9077–83.
139. Rall SC Jr, Weisgraber KH, Innerarity TL, Mahley RW. Structural basis for receptor binding heterogeneity of apolipoprotein E from type III hyperlipoproteinemic subjects. *Proc Natl Acad Sci USA*. 1982;79:4696–700.
140. Sakashita K, Tanaka F, Zhang X, et al. Clinical significance of ApoE expression in human gastric cancer. *Oncol Rep*. 2008;20:1313–9.
141. Kang R, Li P, Wang T, et al. Apolipoprotein E epsilon 2 allele and low serum cholesterol as risk factors for gastric cancer in a Chinese Han population. *Sci Rep*. 2016;6:19930.
142. Zheng P, Luo Q, Wang W, et al. Tumor-associated macrophage-derived exosomes promote the migration of gastric cancer cells by transfer of functional Apolipoprotein E. *Cell Death Dis*. 2018;9:434.
143. Ma TJ, Zhang XJ, Ding XP, Chen HH, Zhang YW, Ding M. Association of single nucleotide polymorphisms in UBR2 gene with idiopathic aspermia or oligospermia in Sichuan, China. *Andrologia*. 2016;48:1253–60.
144. Yang F, Cheng Y, An JY, et al. The ubiquitin ligase Ubr2, a recognition E3 component of the N-end rule pathway, stabilizes Tex19.1 during spermatogenesis. *PLoS ONE*. 2010;5:e14017.
145. An JY, Kim E, Zakrzewska A, et al. UBR2 of the N-end rule pathway is required for chromosome stability via histone ubiquitylation in spermatocytes and somatic cells. *PLoS ONE*. 2012;7:e37414.
146. Kwon YT, Xia Z, An JY, et al. Female lethality and apoptosis of spermatocytes in mice lacking the UBR2 ubiquitin ligase of the N-end rule pathway. *Mol Cell Biol*. 2003;23:8255–71.
147. Kwak KS, Zhou X, Solomon V, et al. Regulation of protein catabolism by muscle-specific and cytokine-inducible ubiquitin ligase E3alpha-II during cancer cachexia. *Can Res*. 2004;64:8193–8.
148. Xu H, Shi J, Gao H, et al. The N-end rule ubiquitin ligase UBR2 mediates NLRP1B inflammasome activation by anthrax lethal toxin. *EMBO J*. 2019;38:e101996.

149. Zhang G, Lin RK, Kwon YT, Li YP. Signaling mechanism of tumor cell-induced up-regulation of E3 ubiquitin ligase UBR2. *FASEB J Off Publ Fed Am Soc Exp Biol.* 2013;27:2893–901.
150. Mao J, Liang Z, Zhang B, et al. UBR2 enriched in p53 deficient mouse bone marrow mesenchymal stem cell-exosome promoted gastric cancer progression via Wnt/ $\beta$ -catenin pathway. *Stem Cells (Dayton, Ohio).* 2017;35:2267–79.
151. Fang J, Ge X, Xu W, et al. Bioinformatics analysis of the prognosis and biological significance of HMGB1, HMGB2, and HMGB3 in gastric cancer. *J Cell Physiol.* 2020;235:3438–46.
152. Dicken BJ, Bigam DL, Cass C, Mackey JR, Joy AA, Hamilton SM. Gastric adenocarcinoma: review and considerations for future directions. *Ann Surg.* 2005;241:27–39.
153. Agresti A, Bianchi ME. HMGB proteins and gene expression. *Curr Opin Genet Dev.* 2003;13:170–8.
154. Kostova N, Zlateva S, Ugrinova I, Pasheva E. The expression of HMGB1 protein and its receptor RAGE in human malignant tumors. *Mol Cell Biochem.* 2010;337:251–8.
155. Stros M. HMGB proteins: interactions with DNA and chromatin. *Biochem Biophys Acta.* 2010;1799:101–13.
156. Wu T, Zhang W, Yang G, et al. HMGB1 overexpression as a prognostic factor for survival in cancer: a meta-analysis and systematic review. *Oncotarget.* 2016;7(31):50417–27.
157. Guo ZS, Liu Z, Bartlett DL, Tang D, Lotze MT. Life after death: targeting high mobility group box 1 in emergent cancer therapies. *Am J Cancer Res.* 2013;3:1–20.
158. Goodwin GH, Sanders C, Johns EW. A new group of chromatin-associated proteins with a high content of acidic and basic amino acids. *Eur J Biochem.* 1973;38:14–9.
159. Tian L, Wang ZY, Hao J and Zhang XY. miR-505 acts as a tumor suppressor in gastric cancer progression through targeting HMGB1. *J Cell Biochem.* 2018. <https://doi.org/10.1002/jcb.28082>.
160. Wu X, Wang W, Chen Y, et al. High mobility group box protein 1 serves as a potential prognostic marker of lung cancer and promotes its invasion and metastasis by matrix metalloproteinase-2 in a nuclear factor- $\kappa$ B-dependent manner. *Biomed Res Int.* 2018;2018:3453706.
161. Evankovich J, Cho SW, Zhang R, et al. High mobility group box 1 release from hepatocytes during ischemia and reperfusion injury is mediated by decreased histone deacetylase activity. *J Biol Chem.* 2010;285:39888–97.
162. Zeng W, Shan W, Gao L, et al. Inhibition of HMGB1 release via salvianolic acid B-mediated SIRT1 up-regulation protects rats against non-alcoholic fatty liver disease. *Sci Rep.* 2015;5:16013.
163. Song B, Song WG, Li ZJ, et al. Effect of HMGB1 silencing on cell proliferation, invasion and apoptosis of MGC-803 gastric cancer cells. *Cell Biochem Funct.* 2012;30:11–7.
164. Zhang J, Kou YB, Zhu JS, Chen WX, Li S. Knockdown of HMGB1 inhibits growth and invasion of gastric cancer cells through the NF- $\kappa$ B pathway in vitro and in vivo. *Int J Oncol.* 2014;44:1268–76.
165. Wang H, Chen Y, Guo J, et al. Dysregulation of tristetraptin and human antigen R promotes gastric cancer progressions partly by upregulation of the high-mobility group box 1. *Sci Rep.* 2018;8:7080.
166. Chung HW, Lim JB. High-mobility group box-1 contributes tumor angiogenesis under interleukin-8 mediation during gastric cancer progression. *Cancer Sci.* 2017;108:1594–601.
167. Tao H, Tang T, Wang S, et al. The molecular mechanisms of Aloin induce gastric cancer cells apoptosis by targeting High Mobility Group Box 1. *Drug Des Dev Ther.* 2019;13:1221–31.
168. Ghweil AA, Osman HA, Hassan MH, et al. Validity of serum amyloid A and HMGB1 as biomarkers for early diagnosis of gastric cancer. *Cancer Manag Res.* 2020;12:117–26.
169. Zhang X, Shi H, Yuan X, Jiang P, Qian H, Xu W. Tumor-derived exosomes induce N2 polarization of neutrophils to promote gastric cancer cell migration. *Mol Cancer.* 2018;17:146.
170. Zhu Q, Zhang X, Zhang L, et al. The IL-6-STAT3 axis mediates a reciprocal crosstalk between cancer-derived mesenchymal stem cells and neutrophils to synergistically prompt gastric cancer progression. *Cell Death Disease.* 2014;5:e1295.
171. Jørgensen JT, Nielsen KB, Møllerup J, Jepsen A, Go N. Detection of MET amplification in gastroesophageal tumor specimens using IQFISH. *Ann Transl Med.* 2017;5:458.
172. Bottaro DP, Rubin JS, Faletto DL, et al. Identification of the hepatocyte growth factor receptor as the c-met proto-oncogene product. *Science (New York, NY).* 1991;251:802–4.
173. Xia C, Wang Y, Liu C, et al. Novel Peptide CM 7 Targeted c-met with antitumor activity. *Molecules (Basel, Switzerland).* 2020;25(3):451.
174. Cecchi F, Rabe DC, Bottaro DP. Targeting the HGF/Met signaling pathway in cancer therapy. *Expert Opin Ther Targets.* 2012;16:553–72.
175. Jardim DL, Tang C, Gagliato Dde M, et al. Analysis of 1,115 patients tested for MET amplification and therapy response in the MD Anderson phase I CLINIC. *Clin Cancer Res Off J Am Assoc Cancer Res.* 2014;20:6336–45.
176. Zhang Y, Xia M, Jin K, et al. Function of the c-Met receptor tyrosine kinase in carcinogenesis and associated therapeutic opportunities. *Mol Cancer.* 2018;17:45.
177. Organ SL, Tsao MS. An overview of the c-MET signaling pathway. *Ther Adv Med Oncol.* 2011;3:S7-s19.
178. Comoglio PM, Trusolino L, Boccaccio C. Known and novel roles of the MET oncogene in cancer: a coherent approach to targeted therapy. *Nat Rev Cancer.* 2018;18:341–58.
179. Ha SY, Lee J, Kang SY, et al. MET overexpression assessed by new interpretation method predicts gene amplification and poor survival in advanced gastric carcinomas. *Jl Modern Pathol Off J US Can Acad Pathol Inc.* 2013;26:1632–41.
180. Drebber U, Baldus SE, Nolden B, et al. The overexpression of c-met as a prognostic indicator for gastric carcinoma compared to p53 and p21 nuclear accumulation. *Oncol Rep.* 2008;19:1477–83.
181. Zhang J, Guo L, Liu X, Li W, Ying J. MET overexpression, gene amplification and relevant clinicopathological features in gastric adenocarcinoma. *Oncotarget.* 2017;8:10264–73.
182. Ding XQ, Wang ZY, Xia D, Wang RX, Pan XR, Tong JH. Proteomic profiling of serum exosomes from patients with metastatic gastric cancer. *Front Oncol.* 2020;10:1113.
183. Im H, Shao H, Weissleder R, Castro CM, Lee H. Nano-plasmonic exosome diagnostics. *Expert Rev Mol Diagn.* 2015;15:725–33.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.