

Case Report

LEOPARD Syndrome with a Sporadic *PTPN11* Mutation in a Saudi Patient

Hussein M. Alshamrani ¹, Luai M. Assaedi,² Jumanah A. Bahattab,³
Abdulrahman M. Mohammad,⁴ and Magdy R. Abdulghani³

¹Department of Dermatology, King Abdulaziz University, Jeddah, Saudi Arabia

²Department of Dermatology, King Abdulaziz Hospital, Mecca, Saudi Arabia

³Department of Dermatology, King Fahd Armed Forces Hospital, Jeddah, Saudi Arabia

⁴Department of Dermatology, Althaghr Hospital, Jeddah, Saudi Arabia

Correspondence should be addressed to Hussein M. Alshamrani; hussain.m.edu@gmail.com

Received 25 October 2022; Revised 6 May 2023; Accepted 13 May 2023; Published 23 May 2023

Academic Editor: Ravi Krishnan

Copyright © 2023 Hussein M. Alshamrani et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

LEOPARD syndrome (LS) is a rare autosomal dominant inherited or sporadic genetic disorder caused commonly by missense mutations in the protein-tyrosine phosphatase-nonreceptor type 11 (*PTPN11*) gene. Due to its rarity and a high chance of misdiagnosis, the epidemiological profile of LS is poorly established. To the best of our knowledge, this is the second report with a documented *PTPN11* gene mutation in Saudi Arabia.

1. Introduction

LEOPARD syndrome (LS) is a rare autosomal dominant inherited or sporadic genetic disorder caused by missense mutations in the protein-tyrosine phosphatase-nonreceptor type 11 (*PTPN11*) gene [1]. The acronym LEOPARD stands for its multisystem clinical manifestations: lentiginos, electrocardiogram conduction abnormalities, ocular hypertelorism, pulmonary stenosis, abnormal genitalia, retarded growth, and sensorineural deafness [2]. Due to its rarity and a high chance of misdiagnosis, the epidemiological profile of LS is poorly established. Here, we report a case of LS with a sporadic mutation in the *PTPN11* gene.

2. Case Presentation

A 5-year-old girl born to second-degree consanguineous parents presented with multiple lentiginos that had begun to appear at the age of 2 years over her face, trunk, and thighs.

Notably, the mother had no follow-up with the obstetrician during the pregnancy and the patient was a full-term

baby delivered through cesarean section, with normal weight and activity. All the previous pregnancies of the mother were normal with healthy children, and the family had no history of lentiginos, congenital heart diseases, learning difficulties, speech delays, or syndromic disorders. A physical examination revealed the height and weight of the child to be 101 cm (3rd centile) and 13.9 kg (below the 3rd centile), respectively, and the child had an occipitofrontal circumference of 52 cm. She showed facial dysmorphism, including a triangular face, slightly protruded mandible, broad nasal bridge, ocular hypertelorism, palpebral ptosis, low set ears, low posterior hairline, pectus carinatum, and scapula alata. No other skeletal anomalies were noted.

After birth, a pan-systolic murmur was detected. Three days after birth, an echocardiogram revealed multiple ventricular septal defects (VSDs). The patient has been continuously followed up with echocardiogram till date. At the age of 4 months, her mother started to notice that the baby was quiet and did not react to sound stimuli. An audiogram was performed at the age of 7 months and showed sensorineural hearing loss for which she underwent



FIGURE 1: Multiple lentigines distributed on the face, trunk, and extremities of the pectus carinatum. Note the *café noir* spot and *café-au-lait* spot near the knee.



FIGURE 2: Multiple lentigines found distributed on the trunk of the patient on further physical examination.

a cochlear implant at the age of 2 years. She also had delayed speech and motor development; she started crawling at 13 months and started to walk at 24 months.

At our hospital, the clinical examination of her lentigines showed multiple oval light brown macules, 2–4 mm in size, over both the sun-exposed and sun-protected body sites including trunk and upper and lower extremities (Figures 1 and 2). There was one darker lentigo, 5 mm in size (*café noir* spot) over the right knee, and a *café-au-lait* spots (brown patch), 1 cm in size, over the left thigh. Oral and genital mucosa were spared.

LS was suspected and the child underwent genetic testing, which showed a heterozygous mutation in *PTPN11*, c.836A > G, a missense variant resulting in the substitution of the tyrosine codon at amino acid position 279 with a cysteine codon (p.Tyr279Cys). LS was thus confirmed, and multidisciplinary follow-up was performed.

3. Discussion

To the best of our knowledge, this is the second report with a documented *PTPN11* gene mutation in Saudi Arabia, after the first one being reported by Alfurayh et al. [3] A comparison of the clinical features of the other cases reported in Saudi Arabia is shown in Table 1.

The negative parental and family history in our case indicates a possible sporadic form of LS that resulted from a new mutation, as shown in a recently published case report with a novel *PTPN11* mutation and negative genetic tests in the parents [4]. However, there still exists the possibility of nonpenetrance (one or two parents may have had the mutated gene but never developed the disease).

Since prenatal diagnosis is possible to be performed for LS-associated conditions as for other RASopathies [5], the age

of diagnosis in our patient could also have been more precocious. Cardiac defects such as VSDs can be detected as early as at 16–18 weeks of gestation and genetic syndromes are usually linked with features such as increased nuchal transparency through ultrasound examination [6], which was not performed in our case. However, even with those parameters combined, the diagnosis of LS would have remained difficult to confirm without lentigines, the hallmark lesions of this syndrome.

To eliminate the possibility of nonpenetrance and detect causal mutations in the fetus, genetic testing is the best screening method for LS-associated mutations. To date, 12 missense *PTPN11* gene mutations associated with LS have been detected: Tyr279Cys/Ser, Ala461Thr/Ser, Gly464Ala, Thr468Met/Pro, Arg498Leu/Trp, Gln506Pro, and Gln510-Glu/Pro [1, 7]. It is, therefore, recommended to perform the sequence analysis of *PTPN11* gene for the coding exons 7, 12, and 13, which are known to be associated with LS. If this test is negative, sequence analysis should then involve LS-linked coding exons from *RAF1* and *BRAF* genes (6-13-16 and 6-11-17, respectively). In case no mutation is identified, the remaining coding exons of *PTPN11* as well as exons of other genes associated with LS, such as *RAF1* and *BRAF*, should be analyzed as well [1].

Besides the known characteristics of LS, our patient also had atypical cardiac findings like those of multiple VSDs without hypertrophic cardiomyopathy (HCM) or pulmonary stenosis. The absence of HCM may signify a favorable prognosis as the overall survival in LS is mainly conditioned by the severity of cardiac defects and this may support the hypothesis of the possible correlation between missense mutations and less serious forms of the disease [8]. However, since the possibility of developing HCM at a later stage still exists, the patient will have to be kept on long-term and

TABLE 1: Comparison of the clinical features of LS in the reported cases in Saudi Arabia.

	Our case	Case 1 of Alfurayh et al. [3]	Case 2 of Alfurayh et al. [3]	Case 3 of Alfurayh et al. [3]
Age/sex	5 years/female	37 years/male	4 years/male	18 months/male
Age of first lentiginos	2 years	Unknown	6 months	2 months
Height (in cm) (percentile)	101 (3rd)	163	103 (50th)	88 (>95th)
Cardiac defects	VSD	ASD	-	ASD and VSD
Pulmonary stenosis	-	-	-	-
Deafness	+	-	-	-
Ocular hypertelorism	+	+	+	+
Pectus excavatum/carinatum	Pectus carinatum	Pectus excavatum	Pectus excavatum	-
Motor delay	+	+	+	+
<i>PTPN11</i> mutation	Exon 7, c.836A>G, p.Tyr279Cys	Exon 7, c.836A>G, p.Tyr279Cys	Exon 7, c.836A>G, p.Tyr279Cys	Exon 7, c.836A>G, p.Tyr279Cys

VSD, ventricular septal defects; ASD, atrial septal defects

stringent clinical and echocardiographic monitoring [9]. Furthermore, an oncology follow-up is considered due to the increased risk of tumor development similarly to those who are affected by dysregulation of the molecular mechanisms responsible for the control of cellular replication [10].

In conclusion, we presented a possible sporadic case of LS caused by a *PTPN11* gene mutation with typical clinical presentation except for the absence of HCM. Dermatological findings were crucial to suspect the diagnosis in our case. Therefore, we recommend physicians to be cautious when dealing with particular skin lesions (lentiginous and *café-au-lait* spots), specifically when associated with pediatric age, and with facial dysmorphism as in other differential diagnosis [11], as this might be the first diagnostic indicator to uncover a serious potentially deadly condition. Due to advances in genetics and molecular biology such as microarray analysis [12], significant progress has been accomplished in the last decades in understanding many complex genetic disorders such as LS and other RASopathies. However, this progress has been substantially limited by challenges regarding the development of effective therapies. Further specific guidelines on the management of LS and the collection of more epidemiologic data are required. Success in the ongoing preclinical studies would allow current efforts to headway into the human trials phase, especially those of RAS pathway inhibitors and genome-modifying therapies.

Data Availability

All data related to the article are available from corresponding authors upon request.

Consent

Consent for the publication of the photographs and medical information of the patient was taken.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Acknowledgments

The authors are thankful to the Dermatology Department in King Fahd Armed Forces Hospital for their support.

References

- [1] E. Martínez-Quintana and F. Rodríguez-González, "Leopard syndrome: clinical features and gene mutations," *Molecular Syndromology*, vol. 3, pp. 145–157, 2012.
- [2] R. J. Gorlin, R. C. Anderson, and M. Blaw, "Multiple lentiginous syndrome," *American Journal of Diseases of Children*, vol. 117, pp. 652–662, 1969.
- [3] N. A. Alfurayh, F. Alsaif, N. Alballa et al., "Leopard syndrome with *PTPN11* gene mutation in three family members presenting with different phenotypes," *Journal of Pediatric Genetics*, vol. 9, pp. 246–251, 2020.
- [4] N. Rahal, A. Sadi, E. Cohen-Barak, M. Ziv, J. Krausz, and R. P. Dodiuk-Gad, "LEOPARD syndrome: a report of a case with a novel *PTPN11* mutation," *JAAD Case Report*, vol. 11, pp. 57–59, 2021.
- [5] F. Mercadante, E. Piro, and M. Busè, "Cutis verticis gyrata and Noonan syndrome: report of two cases with pathogenetic variant in *SOS1* gene," *Italian Journal of Pediatrics*, vol. 48, p. 152, 2022.
- [6] M. C. Digilio, A. Sarkozy, A. de Zorzi et al., "LEOPARD syndrome: clinical diagnosis in the first year of life," *American Journal of Medical Genetics*, vol. 140, pp. 740–746, 2006.
- [7] J. Zhang, J. Shen, R. Cheng et al., "Identification of a *PTPN11* hot spot mutation in a child with atypical LEOPARD syndrome," *Molecular Medicine Reports*, vol. 14, pp. 2639–2643, 2016.
- [8] E. Piro, G. Serra, and V. Antona, "Novel LRPPRC compound heterozygous mutation in a child with early-onset Leigh syndrome French-Canadian type: case report of an Italian patient," *Italian Journal of Pediatrics*, vol. 46, p. 140, 2020.
- [9] J. Lauriol, K. Keith, B. Zhang, R. Bronson, K.-H. Lee, and M. I. Kontaridis, "Abstract 196: adult-onset cardiac hypertrophy in Leopard syndrome is caused by both cell autonomous and non-autonomous effects that occur during development," *Circulation Research*, vol. 113, p. A196, 2013.
- [10] G. Serra, V. Antona, M. Schierz, D. Vecchio, E. Piro, and G. Corsello, "Esophageal atresia and Beckwith–Wiedemann syndrome in one of the naturally conceived discordant newborn twins: first report," *Clinical Case Reports*, vol. 6, no. 2, p. 399, 2018.
- [11] G. Serra, V. Antona, M. Giuffrè et al., "Novel missense mutation of the *TP63* gene in a newborn with Hay-Wells/Ankyloblepharon-Ectodermal Defects-Cleft Lip/Palate (AEC) syndrome: clinical report and follow-up," *Italian Journal of Pediatrics*, vol. 47, pp. 1–6, 2021.
- [12] G. Serra, V. Antona, and M. Giuffrè, "Interstitial deletions of chromosome 1p: novel 1p31.3p22.2 microdeletion in a newborn with craniosynostosis, coloboma and cleft palate, and review of the genomic and phenotypic profiles," *Italian Journal of Pediatrics*, vol. 48, p. 38, 2022.