

# Carotid-Cavernous Fistulas: The Utility of Ocular Echography in Their Differentiation [Response to Letter]

Pawasoot Supasai, Kanwasee Kanjana, Yosanan Yospaiboon 

KKU Eye Center, Department of Ophthalmology, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand

Correspondence: Yosanan Yospaiboon, KKU Eye Center, Department of Ophthalmology, Faculty of Medicine, Khon Kaen University, 123 Mitraparb Highway, Khon Kaen, 40002, Thailand, Tel +66-4336-3010, Fax +66-4334-8383, Email [yosanan@gmail.com](mailto:yosanan@gmail.com)

## Dear editor

We appreciate the discussion of Graziano et al on the use of ocular ultrasonography in the differentiation between direct and dural carotid-cavernous fistulas (CCFs). We agree that it can be a useful tool in the evaluation of CCFs. Although it is not the primary diagnostic modality, it can provide valuable information that aids in the diagnosis of these conditions. Ocular ultrasonography can demonstrate ocular findings associated with CCFs, such as proptosis, globe enlargement, thickened extraocular muscles, dilation of the superior ophthalmic vein, and abnormal pulsations. Moreover, intraocular tumors, scleritis, and endocrine myopathy can be excluded.<sup>1</sup> Ophthalmologists may use ocular ultrasonography as part of the diagnostic evaluation for CCFs, but it is not typically the primary or sole diagnostic modality.

Differentiating between direct CCFs (high-flow) and dural CCFs (low-flow) is crucial as it impacts the management approach. Additional imaging modalities such as computed tomography angiography (CTA) or magnetic resonance angiography (MRA) are typically used to provide more comprehensive assessment of the arterial anatomy, blood flow patterns, and potential venous drainage abnormalities. They can indicate the location of the shunts and confirm the type of CCFs.<sup>1,2</sup> If an intervention, such as endovascular embolization, is planned for CCFs treatment, more precise imaging modalities such as digital subtraction angiography (DSA) are often necessary to guide the procedure and ensure accurate embolization.<sup>3</sup>

The purpose of our retrospective study<sup>4</sup> is to compare the clinical characteristics of patients with direct and dural CCFs. The results show that patients with direct CCFs were younger, associated with trauma, and had more visual impairment at presentation. Chemosis, proptosis, bruit, and dilated retinal vessels were detected more in the direct CCF than in the dural CCF. Since ocular ultrasonography is not a routine diagnostic modality of CCFs in our setting, therefore the data is incomplete and not included in the analysis. However, information on the difference in clinical characteristics may help clinicians to initially discriminate the urgent direct CCFs and promptly refer the patients for further investigation and definite treatment.

## Disclosure

The authors report no conflicts of interest in this communication.

## References

1. de Keizer R. Carotid-cavernous and orbital arteriovenous fistulas: ocular features, diagnostic and hemodynamic considerations in relation to visual impairment and morbidity. *Orbit*. 2003;22(2):121–142. doi:10.1076/orbi.22.2.121.14315
2. Chaudhry IA, Eikhamry SM, Al-Rashed W, Bosley TM. Carotid cavernous fistula: ophthalmological implications. *Middle East Afr J Ophthalmol NBS*. 2009;16(2):57–63. doi:10.4103/0974-9233.53862
3. Soulez G, Gilbert P, Giroux MF, Racicot JN, Dubois J. Interventional management of arteriovenous malformations. *Tech Vasc Interv Radiol*. 2019;22(4):100633. doi:10.1016/j.tvir.2019.100633
4. Supasai P, Kanjana K, Yospaiboon Y. Direct and dural carotid cavernous sinus fistulas: comparison of clinical characteristics. *Clin Ophthalmol*. 2023;17:1207–1214. doi:10.2147/OPHT.S410930

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Ophthalmology 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Ophthalmology editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

## Clinical Ophthalmology

Dovepress

### Publish your work in this journal

Clinical Ophthalmology is an international, peer-reviewed journal covering all subspecialties within ophthalmology. Key topics include: Optometry; Visual science; Pharmacology and drug therapy in eye diseases; Basic Sciences; Primary and Secondary eye care; Patient Safety and Quality of Care Improvements. This journal is indexed on PubMed Central and CAS, and is the official journal of The Society of Clinical Ophthalmology (SCO). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-ophthalmology-journal>

<https://doi.org/10.2147/OPTH.S422888>