



# Development and simulated validation of a dish composition database for estimating food group and nutrient intakes in Japan

Nana Shinozaki<sup>1</sup>, Kentaro Murakami<sup>2</sup>, Shizuko Masayasu<sup>3</sup> and Satoshi Sasaki<sup>1,2,\*</sup>

<sup>1</sup>Department of Social and Preventive Epidemiology, Graduate School of Medicine, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan: <sup>2</sup>Department of Social and Preventive Epidemiology, School of Public Health, The University of Tokyo, Tokyo, Japan: <sup>3</sup>Ikurien-naka, Ibaraki, Japan

Submitted 20 August 2018: Final revision received 20 December 2018: Accepted 1 February 2019: First published online 16 May 2019

## Abstract

**Objective:** To develop a dish composition database (DCD) and assess its ability to estimate dietary intake.

**Design:** The DCD was developed based on 16 d dietary records (DR). We aggregated all reported dishes into 128 dish codes and calculated mean food group and nutrient contents for each code. These data were used to calculate dietary intake in a different population that completed a 4 d DR. The estimated values were compared with those estimated using the standard food composition database (FCD) of Japan.

**Setting:** Japan.

**Participants:** A total 252 adults aged 31–81 years for the 16 d DR (3941 d in total) and 392 adults aged 20–69 years for the 4 d DR (1568 d in total) participated.

**Results:** There were significant differences in median intakes between the DCD and the FCD for eighteen and twenty (of twenty-six) food groups and for twenty-nine and twenty-two (of forty-three) nutrients (including energy) in men and women, respectively. For food group intakes, Spearman correlation coefficients between the DCD and FCD ranged from 0.19 (animal fats) to 0.90 (fruits and alcoholic beverages) in men (median: 0.61) and from 0.25 (oils) to 0.89 (noodles) in women (median: 0.58). For nutrient intakes, the corresponding values ranged from 0.25 (retinol) to 0.90 (alcohol) in men (median: 0.60) and from 0.15 (retinol) to 0.74 (alcohol) in women (median: 0.53).

**Conclusions:** Whereas it is difficult to accurately estimate absolute dietary intake values using the present DCD, it has acceptable ability to rank the intakes of many food groups and nutrients.

## Keywords

Dietary record  
Dish composition database  
Validity  
Nutrient intake  
Food group intake  
Japan

The dietary record (DR), which widely used in dietary surveys, requires participants to record all foods and beverages they have consumed in a specified period<sup>(1,2)</sup>. Although the DR has many advantages, recording all food items is time-consuming and requires a high level of motivation on the part of participants<sup>(1–3)</sup>. Moreover, foods are not typically eaten as single foods but as composite dishes prepared or cooked together with other foods<sup>(4–7)</sup>. Hence, it is difficult to accurately describe the types and amounts of all single food ingredients in mixed dishes, especially for people who are not involved in cooking<sup>(5,8)</sup>. Consequently, albeit dish names are provided in a DR, the type or amount of each ingredient

is not always captured, particularly for cooked foods or seasonings<sup>(8)</sup>.

To estimate dietary intake from mixed dishes, standard recipe databases are often used in national dietary surveys<sup>(8–15)</sup>. Several countries have also made continuous efforts to develop and harmonize food composition databases, including nutrient composition of mixed dishes, for epidemiological studies<sup>(4,16–25)</sup>. These databases of recipes or dish composition are essential to estimate dietary intake from mixed dishes in each country, and they have been used to support food-based dietary assessment methods, such as DR or dietary recall. In addition,

\*Corresponding author: Email stssasak@m.u-tokyo.ac.jp

dish-based FFQ have been developed recently in countries such as South Korea<sup>(26–29)</sup>, Bangladesh<sup>(30)</sup>, Iran<sup>(31)</sup> and Sweden<sup>(6,32)</sup>. Validation studies of dish-based FFQ against DR<sup>(28–30,32)</sup> and doubly labelled water<sup>(6)</sup> have shown adequate ranking abilities for intakes of most foods and nutrients. However, to our knowledge, there have been a limited number of studies assessing the validity of dietary intake estimated based on information collected about individual dishes<sup>(6,28–30,32)</sup>.

Because cooking methods and the composition of dishes are different among countries, a dish composition database (DCD) should be developed in each country. In particular, Japanese dishes are prepared from various ingredients using different approaches and these are mixed with many seasonings<sup>(1,3,33–35)</sup>, thus making it complicated to estimate ingredients prior to cooking<sup>(8,33,34)</sup>. Although there is some information on dish composition in the Standard Table of Food Composition of Japan<sup>(36)</sup>, the number of dishes is limited. To date, several previous studies have developed a DCD<sup>(34,35,37,38)</sup> or have assessed the validity of a DCD<sup>(3,8,34,35,37,39,40)</sup>. Most of these studies tried to establish dish-based dietary assessment methods based on the idea that use of a DCD can reduce the burden on participants and staff in a dietary survey by omitting or standardizing the process of disaggregation of dishes into food items<sup>(3,34,35,37,39,40)</sup>. However, the validity of such DCD has not been assessed<sup>(38)</sup>, or this has been assessed only in groups consisting of mostly women<sup>(3,34,35,37,39)</sup> or for only a limited number of dish items among elderly adults<sup>(8)</sup>; also, the sample sizes of all of these studies were small. Moreover, the validity was assessed by comparing a DR and a list of dish names recorded by participants on the same day as recording in the DR<sup>(3,35,37,39)</sup>. Because it is possible that participants recorded more accurate information about dishes that they consumed, as compared with the case that they did not maintain the DR, the validity of the DCD might be overestimated in these previous studies. Furthermore, the validated databases were developed

based on a 1 d DR or a 3 d DR conducted during one season among participants in certain prefectures<sup>(34,35,37)</sup>. Given that dietary intake varies depending on the season<sup>(41–43)</sup> and region<sup>(7,40,44)</sup>, these databases may not contain comprehensive information on dishes in Japan.

Hence, we developed a DCD based on data from 16 d weighed DR obtained from Japanese men and women. To assess the ability of the DCD to estimate dietary intake based on dish names, food group and nutrient intakes were calculated from a 4 d DR conducted in a different population using the DCD and the estimated values were compared with those estimated using the standard food composition database (FCD) of Japan<sup>(36)</sup>. Moreover, to determine whether information on the portion size (PS) of dishes is necessary for dietary assessment using the DCD, we assessed the consequences of using reported PS of dishes instead of standard PS in the DCD on estimated intakes. We expected that a DCD would be helpful not only in supporting food-based DR, but also in developing dish-based dietary assessment methods in future.


## Methods

The study consisted of the two phases. First, we developed the DCD using dietary data obtained from a 16 d DR. Second, the ability of the DCD to estimate dietary intake was assessed using a 4 d DR obtained from a different population, by comparing food group and nutrient intakes estimated using the DCD with those estimated using the FCD (Fig. 1).

### Development of the dish composition database

#### Data source

The DCD was developed based on dietary information derived from a 16 d DR survey administered between November 2002 and September 2003 in four areas of Japan, namely Osaka (urban), Okinawa (urban island), Nagano (rural inland) and Tottori (rural coastal). Details of the survey have been described elsewhere<sup>(45–47)</sup>. Briefly,



**Fig. 1** Study framework (DCD, dish composition database; DR, dietary record; FCD, food composition database)



apparently healthy women ( $n$  126) aged 31–69 years and their cohabitating spouses ( $n$  126) completed a four non-consecutive-day weighed DR (three weekdays and one weekend day) during each of four seasons. Women were recruited such that each 10-year age class (30–39, 40–49, 50–59 and 60–69 years) contained eight women (without consideration of age of men). None of them was a dietitian or had received dietary therapy by a doctor or dietitian, and there was no case of history of educational admission for diabetes mellitus. The participants were asked to record and weigh all beverages and foods consumed on each recording day. Registered dietitians explained to participants how to complete the DR using both written and verbal instructions, and they provided recording sheets and a digital scale. Recording sheets were checked by trained registered dietitians in the respective local centre and then again in the study centre. The records were coded by trained dietitians at the study centre in accordance with uniform procedures using the FCD<sup>(36)</sup>.

#### Generation process of the dish composition database

A total of 3941 d of DR were obtained from the participants. In the present study, 'dishes' were defined as all those recorded in the column 'dish name' in the DR by participants, even if the dish consisted of only one food item (i.e. milk or apple consumed alone was regarded as a dish). As a result, a total of 71 213 dishes appeared in the 16 d DR (Fig. 2). Dishes with identical names were combined, resulting in 2409 dish names. The mean food group and nutrient contents of dishes were calculated for each dish

name using the FCD<sup>(36)</sup>. Food groups were based mainly on the FCD and the similarity of nutrient composition or culinary use of foods (see online supplementary material, Supplemental Table 1). Dishes were aggregated into dish groups based on degree of similarity of the dish name<sup>(8,26,34)</sup>, main ingredient (food group with the largest proportion in all ingredients in a dish)<sup>(7,26,27,34,37,48)</sup>, cooking method (raw, mixed, boiled, steamed, stewed, grilled, stir-fried and deep-fried)<sup>(5,26,34,37,44,49)</sup>, and energy and nutrient contents per portion (protein, fat and carbohydrate)<sup>(5,26,48)</sup>. Dishes with low frequencies of consumption were integrated with those having high occurrence frequencies ( $\geq 20$  times) when possible. Dishes that could not be classified into a specific dish group due to a lack of the information necessary for classification were categorized as dish groups about which details are unknown. For example, grilled fish dishes were categorized based on the type of fish; however, if the type of fish was not written in the dish name, it was categorized as 'grilled fish (details unknown)'. This process yielded 383 dish groups.


We then reviewed the classification of dishes, as follows. First, the mean food group and nutrient contents of each dish group were calculated. Next, food groups accounting for 5% or more of the total weight of each dish group were identified. If there were any unusual ingredients or excessively high or low nutrient contents for a dish group, we reviewed dishes classified in that dish group and reclassified them if necessary. Furthermore, some dish groups that could be further combined based on their similarity with other groups were aggregated. After this process, all dishes were categorized into a total of 371 dish groups.

Each dish group was given a dish name and dish code (i.e. minor code). To assess the ability of the DCD to estimate dietary intake according to the aggregation level of dishes, the 371 types of dishes were further aggregated based on their similarities, and each dish was assigned another dish code (major code). For instance, the minor codes 'potato croquette' and 'cream croquette' were aggregated into the major code 'croquette' (see online supplementary material, Supplemental Table 2). This process yielded 128 major codes. We calculated the mean of weight, food group and nutrient content of individual dishes assigned to each coded dish for the minor and major codes in the DCD (Fig. 3). In the DCD, the mean of weight of dishes categorized into the same dish code was regarded as a standard PS of that dish. For example, for a major code 'hamburger', we obtained fifty-one recipes from the DR. Therefore, the standard PS of a major code 'hamburger' in the DCD was calculated as the mean of weight of these fifty-one hamburgers.

#### Simulated validation of the dish composition database

##### Data source

The simulated validation of the DCD was assessed using dietary intake information derived from a 4 d DR conducted



**Fig. 2** Flowchart for the aggregation process of dishes from the 16 d dietary record conducted among 252 Japanese adults between 2002 and 2003

The food group and nutrient content of fifty-one hamburgers reported by each participant						The food group and nutrient content of a 'hamburger' in the dish composition database	
	No. 1	No. 2	No. 3	...	No. 51		
<b>Food group (g)</b>						<b>Food group (g)</b>	
Rice	0.0	0.0	0.0	...	0.0	Rice	0.0
Noodles	0.0	0.0	0.0	...	0.0	Noodles	0.0
Bread	30.6	60.0	60.0	...	30.8	Bread	55.7
Other grain products	2.5	7.8	5.3	...	2.6	Other grain products	4.9
Nuts	0.0	0.0	0.0	...	0.0	Nuts	0.0
Pulses	0.0	0.0	0.0	...	0.0	Pulses	0.0
Potatoes	0.0	0.0	0.0	...	0.0	Potatoes	1.4
Sugar	0.0	0.0	0.0	...	0.0	Sugar	0.1
Confectioneries	0.0	0.0	0.0	...	0.0	Confectioneries	2.9
Fats	0.0	0.0	0.0	...	0.0	Fats	0.0
Oils	2.5	12.6	15.6	...	2.6	Oils	6.9
Fruits	0.0	0.0	0.0	...	0.0	Fruits	0.0
Green and yellow vegetables	12.7	0.0	0.0	...	12.8	Green and yellow vegetables	17.3
Other vegetables	2.5	0.0	0.0	...	2.6	Other vegetables	3.9
Pickled vegetables	15.3	10.0	10.0	...	15.4	Pickled vegetables	24.4
Mushrooms	0.0	0.0	0.0	...	0.0	Mushrooms	0.0
Seaweeds	0.0	0.0	0.0	...	0.0	Seaweeds	0.0
Fruit and vegetable juice	0.0	0.0	0.0	...	0.0	Fruit and vegetable juice	0.0
Seasonings and spices	5.4	12.4	6.9	...	5.4	Seasonings and spices	9.3
Alcoholic beverages	0.0	0.0	0.0	...	0.0	Alcoholic beverages	0.0
Tea and coffee	0.0	0.0	0.0	...	0.0	Tea and coffee	0.0
Soft drinks	0.0	0.0	0.0	...	0.0	Soft drinks	0.0
Fish and shellfish	0.0	0.0	0.0	...	0.0	Fish and shellfish	3.3
Meats	35.7	70.3	56.6	...	36.0	Meats	56.9
Eggs	4.1	3.9	2.7	...	4.1	Eggs	7.7
Dairy products	7.6	0.0	0.0	...	7.7	Dairy products	7.0
Total amount (portion size)	119.0	177.0	157.0	...	120.0	Total amount (standard portion size)	201.7
<b>Nutrient</b>						<b>Nutrient</b>	
Protein (g)	12.3	20.2	17.3	...	12.4	Protein (g)	20.8
Fat (g)	11.7	28.4	22.9	...	11.8	Fat (g)	19.1
Carbohydrate (g)	20.7	39.1	35.4	...	20.9	Carbohydrate (g)	40.5
Total dietary fibre (g)	1.2	1.7	1.5	...	1.2	Total dietary fibre (g)	2.3
Na (mg)	456	791	724	...	459	Na (mg)	729
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮

The amount of each food group and nutrient of all hamburgers was averaged

**Fig. 3** Example of calculation process of the standard portion size and food group and nutrient contents in the dish composition database, with 'hamburger' as an example of a major code

in February and March 2013 in twenty study areas consisting of twenty-three prefectures (including Osaka and Okinawa but not Nagano or Tottori). Details of the survey have been described elsewhere<sup>(50)</sup>. Briefly, the study targeted apparently healthy men and women aged 20–69 years working in welfare facilities in each area. Each of the twenty areas included four apparently healthy adults (two men and two women) from each of five 10-year age groups. One individual per household was permitted to participate in the survey. None of the participants was a dietitian, had received dietary therapy by a doctor or dietitian, or had history of educational admission for diabetes mellitus. In total, 196 men and 196 women completed a weighed DR for four non-consecutive days (three working days and one non-working day). The participants were asked to record all foods and drinks they consumed on each recording day. Research dietitians explained how to record foods and drinks and asked participants to weigh them with a provided digital scale or measuring spoon and cup. The main recorded items on the DR sheets were dish names, names of foods (including beverages and any ingredients in dishes), and approximate amounts or

measured weights of foods and dishes consumed. Recording sheets were checked by research dietitians and two other research dietitians at the central office of the study.

#### *Assignment of food codes to foods in the 4 d dietary records using the food composition database*

A total of 1568 d of the DR were obtained from participants. The coding of records using the FCD<sup>(36)</sup> was performed by the two research dietitians at the study centre in accordance with uniform procedures. Each item recorded in the column of names of foods (91 045 items) was assigned a food code in the FCD. The FCD has a total of 1878 food codes, whereas only sixteen items (0.9%) of those are prepared foods such as frozen croquettes. Hence, almost all recorded items were coded by a food code consisting of single food items.

#### *Assignment of dish codes to dishes in the 4 d dietary records using the dish composition database*

The coding of the DR using the DCD was performed by a registered dietitian not involved in the coding using the FCD. Each item appearing in the column of dish names

**Table 1** Basic characteristics of the participants in each study

Variable	Development of the DCD				Simulated validation of the DCD			
	Men ( <i>n</i> 126)		Women ( <i>n</i> 126)		Men ( <i>n</i> 196)		Women ( <i>n</i> 196)	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
Age (years)	52.4	12.3	49.5	11.5	44.7*	13.3	44.4*	13.5
Body height (cm)†	167.4	6.5	154.8	6.2	170.3*	5.4	157.6*	5.7
Body weight (kg)‡	66.3	10.4	53.5	7.1	69.6*	11.3	56.1*	10.0
BMI (kg/m <sup>2</sup> )§	23.6	2.9	22.3	2.8	24.0	3.5	22.6	3.7
Energy intake (kJ/d)	9874	1774	7703	1222	9870	2021	7904	1510

DCD, dish composition database.

\**P* < 0.05 compared with the corresponding value for each sex of participants completing the dietary records (DR) used for development of the DCD (two-sample *t* test).

†Measured without shoes to the nearest 0.1 cm. *n* 123 for each sex due to missing values.

‡Measured in light clothing to the nearest 0.1 kg. *n* 123 for men and *n* 122 for women due to missing values.

§Calculated by dividing body weight (kilograms) by the square of body height (metres). *n* 123 for men and *n* 122 for women due to missing values.

|| Calculated from 16 d DR for development of the DCD and 4 d DR for simulated validation of the DCD, both using the Standard Table of Food Composition in Japan<sup>(96)</sup>.

in the 4 d DR (26 642 dishes consisting of 9727 dish names) was categorized using a minor code in the DCD, based on its name<sup>(35)</sup>. If there was any special supplementary information on a dish (e.g. name of the brand or manufacturer), it was also used for categorizing the dish. If there was no corresponding dish name in the DCD, we assigned a code of a dish that was similar in its ingredients, type of dish or cooking method included in the dish name. A major code in the DCD was also assigned to each dish based on its corresponding minor code.

### Statistical analysis

Age, BMI and energy intake were compared between participants completing the 16 d and 4 d DR, using a two-sample *t* test. Daily intakes of food groups, energy and nutrients were calculated using the FCD as well as using both the major and minor codes of the DCD. The food groups used in the analysis were the same as those used in development of the DCD. In the calculation of dietary intake using the DCD, the amount consumed of a dish was replaced by the standard PS according to major or minor code in the DCD. To investigate the effect of using the PS of dishes reported by participants instead of the standard PS in estimating dietary intake, dietary intakes from the DCD adjusted by the reported PS of each dish in the DR were also computed. The values were calculated as follows: estimated food group or nutrient intake from a dish adjusted by reported PS = food group or nutrient content in the dish in the DCD (g) × reported PS of the dish in the DR (g)/standard PS of the dish in the DCD (g).

We performed simulated validation of food group and nutrient intakes estimated by the DCD, in comparison with those estimated by the FCD. The ability of the DCD to estimate median intake was evaluated using the Wilcoxon signed-rank test and ranking ability was evaluated using Spearman correlation coefficients. For food groups, energy and macronutrients, a Bland–Altman plot was used to

assess the agreement for estimated values between the DCD and the FCD<sup>(51)</sup>. Additionally, we examined differences in the ability to estimate and rank dietary intakes between the methods using major codes and minor codes, and standard PS and reported PS. The ability to estimate median dietary intake was assessed by comparing the median percentage differences in median intake for each food group or nutrient estimated by the DCD from that estimated by the FCD, using the Wilcoxon signed-rank test. The median correlation coefficients were also compared using the Wilcoxon signed-rank test. Although the DCD was developed without regard for sex, statistical analyses were conducted for men and women separately, using the statistical software package SAS version 9.4. We analysed dietary intakes using both crude values (amount per day) and energy-adjusted values based on the density method (percentage of energy for energy-providing nutrients and amount per 4184 kJ energy for food groups and other nutrients). Because similar associations with the FCD were observed for both calculations, only results for crude values are presented. Two-sided *P* < 0.05 was considered statistically significant.

### Results

Table 1 shows basic characteristics of the participants completing the 16 d DR used for developing the DCD and of participants completing the 4 d DR used for assessing the simulated validation. Mean age was higher in participants completing the 16 d DR than in those completing the 4 d DR in both sexes. Mean BMI and energy intake were not different in either sex.

Table 2 shows food group intakes estimated based on major codes of the DCD and those estimated based on the FCD. For median intakes of twenty-six food groups, the number of food groups that differed significantly

**Table 2** Comparison of food group intakes estimated based on the food composition database (FCD) and those estimated based on the dish composition database (DCD), with use of standard portion size data or reported portion size data

Food group (g/d)	Men (n 196)											Women (n 196)										
	FCD‡			DCD†								FCD‡			DCD†							
	Median	P <sub>25</sub>	P <sub>75</sub>	Standard portion size§				Reported portion size				Median	P <sub>25</sub>	P <sub>75</sub>	Standard portion size§				Reported portion size			
				Median	P <sub>25</sub>	P <sub>75</sub>	P¶	Median	P <sub>25</sub>	P <sub>75</sub>	P¶				Median	P <sub>25</sub>	P <sub>75</sub>	P¶	Median	P <sub>25</sub>	P <sub>75</sub>	P¶
Rice	390.6	290.0	480.6	320.0	253.7	380.1	<0.0001	393.6	294.8	480.1	0.002	262.3	196.6	327.5	298.9	242.3	368.7	<0.0001	284.9	202.6	338.6	<0.0001
Noodles	81.6	41.6	125.9	80.9	47.7	115.6	0.007	95.4	46.0	162.9	<0.0001	50.0	12.5	87.8	55.3	28.6	97.8	0.0001	66.7	14.0	111.3	<0.0001
Bread	30.0	12.1	62.0	31.3	14.2	55.6	0.69	32.7	11.7	58.0	0.49	36.9	16.5	57.8	36.9	16.0	57.6	0.66	33.5	13.1	54.6	0.0002
Other grain products	9.9	5.4	22.8	10.8	6.8	18.1	0.76	11.6	7.4	19.0	0.41	8.3	3.0	17.7	9.6	6.2	17.1	0.06	8.6	5.5	14.4	0.60
Nuts	1.0	0.1	2.6	1.9	1.3	2.8	0.0005	2.2	1.5	3.1	<0.0001	1.0	0.3	3.9	2.3	1.6	3.1	0.005	2.3	1.7	3.2	0.002
Pulses	56.8	28.6	89.4	52.9	35.9	79.1	0.85	60.7	36.1	83.6	0.002	51.4	28.2	84.7	60.5	37.6	80.6	0.03	59.2	35.6	84.9	0.005
Potatoes	39.9	17.6	62.9	43.3	28.5	61.7	0.02	47.0	29.4	64.5	<0.0001	29.1	16.3	50.3	39.7	26.4	55.0	<0.0003	36.0	25.1	50.3	<0.0001
Sugar	11.8	5.9	17.5	9.0	6.6	11.6	<0.0001	10.4	7.6	13.3	0.003	12.8	7.1	19.3	10.4	8.2	12.7	<0.0001	10.8	8.2	13.7	0.0001
Confectioneries	27.9	4.4	55.3	32.6	13.1	55.2	0.01	37.5	11.0	58.5	<0.0001	39.7	18.6	64.4	55.4	34.2	74.1	<0.0001	54.0	31.0	78.0	<0.0001
Animal fats	1.0	0.0	2.6	1.3	0.8	1.9	0.79	1.4	0.9	2.1	0.51	1.1	0.0	3.1	1.3	0.9	2.0	0.15	1.3	0.8	1.9	0.13
Oils	19.6	12.3	27.8	17.5	13.5	23.1	0.01	20.2	15.2	25.1	0.68	15.8	10.2	21.8	15.5	12.3	19.1	0.87	15.8	12.2	20.1	0.88
Fruits	27.6	4.6	73.3	42.2	9.8	97.1	<0.0001	33.0	9.3	75.0	<0.0001	46.0	16.3	101.4	57.8	28.2	106.5	<0.0001	46.9	19.9	99.5	0.007
Green and yellow vegetables	69.1	39.8	100.6	83.6	66.9	109.1	<0.0001	92.0	72.0	116.3	<0.0001	72.6	40.5	104.5	89.4	69.2	112.0	<0.0001	90.1	68.9	110.9	<0.0001
Other vegetables	147.1	100.5	188.4	137.8	109.6	174.1	0.12	154.0	119.9	190.7	0.02	137.8	99.3	185.6	136.5	109.3	166.0	0.46	139.4	110.9	179.8	0.96
Pickled vegetables	18.0	0.9	46.1	7.0	2.4	13.9	<0.0001	5.5	2.3	12.2	<0.0001	15.0	0.0	32.5	5.8	2.2	11.9	<0.0001	4.6	2.0	10.6	<0.0001
Mushrooms	10.0	3.4	20.9	10.9	7.8	14.0	0.18	11.7	8.8	15.7	0.90	10.8	4.9	22.6	10.2	7.7	13.6	0.002	9.7	7.5	14.6	0.002
Seaweeds	4.5	1.5	11.3	9.9	7.4	14.4	<0.0001	10.3	7.6	14.9	<0.0001	4.4	1.5	13.0	10.6	7.7	15.0	<0.0001	10.3	7.4	14.0	<0.0001
Fruit and vegetable juice	0.0	0.0	12.5	0.8	0.4	32.6	<0.0001	0.8	0.4	20.7	0.054	0.0	0.0	28.4	1.4	0.4	35.3	<0.0001	1.4	0.4	35.7	0.002
Seasonings and spices	90.9	54.3	153.4	149.4	103.6	181.8	<0.0001	162.4	111.5	211.5	<0.0001	77.2	39.2	128.9	138.1	105.4	172.9	<0.0001	138.5	112.5	174.2	<0.0001
Alcoholic beverages	43.7	4.8	357.2	54.8	2.7	241.2	<0.0001	36.9	3.1	312.0	<0.0001	6.4	2.6	36.9	3.5	2.2	30.8	0.002	3.8	2.3	23.2	0.0002
Tea and coffee	723.7	450.0	994.0	534.1	407.4	689.2	<0.0001	730.2	492.4	922.8	0.44	754.4	511.1	1040.9	653.2	502.7	784.6	<0.0001	759.1	558.4	999.1	0.54
Soft drinks	0.0	0.0	31.3	12.4	1.9	52.6	<0.0001	10.9	2.6	73.4	<0.0001	0.0	0.0	30.1	14.7	2.2	47.0	<0.0001	10.4	2.9	38.8	<0.0001
Fish and shellfish	68.9	36.3	106.1	71.8	48.9	100.3	0.10	86.3	56.8	120.8	<0.0001	54.7	29.5	82.3	63.5	42.9	88.9	<0.0001	69.9	44.8	101.3	<0.0001
Meats	98.0	67.6	140.2	91.7	68.5	120.5	0.007	101.0	75.0	133.4	0.70	70.1	45.1	102.3	78.5	59.6	99.5	0.003	78.1	55.2	104.6	0.002
Eggs	41.6	26.6	58.8	34.6	23.7	48.3	0.0001	39.7	26.4	53.2	0.10	34.9	18.9	47.0	30.4	21.5	41.6	0.29	32.2	22.6	44.2	0.67
Dairy products	60.6	18.2	145.0	65.1	27.9	144.2	0.06	72.9	31.3	144.5	0.0002	85.5	41.5	160.3	102.2	54.8	154.3	0.0005	87.1	54.8	158.0	0.004

P<sub>25</sub>, 25th percentile; P<sub>75</sub>, 75th percentile.

†Food group intake estimated using the DCD for 128 kinds of dishes developed from 16 d dietary records in 126 men and 126 women.

‡Food group intake estimated from each food using the Standard Table of Food Composition in Japan<sup>(36)</sup>.

§Food group intake estimated using the weight of each dish in the DCD.

|| Estimated food group intake calculated by adjusting weight of a food group of a dish in the DCD by reported portion size of the dish in the dietary records. The calculation method was as follows: estimated food group intake from a dish adjusted by reported portion size = weight of a food group in the dish in the DCD (g) × reported portion size of the dish in the dietary records (g)/standard portion size of the dish in the DCD (g).

¶The difference compared with values derived from the FCD was tested using the Wilcoxon signed-rank test.



between the DCD and the FCD was similar between standard PS and reported PS in both sexes, ranging from seventeen to twenty (65–77%). More than half of these values were considered overestimation rather than underestimation. In men, food group intakes were underestimated more often using standard PS than using reported PS. The median percentage differences in food group intakes did not differ between standard PS and reported PS in either sex (range: 15.3–17.2%).

Energy and nutrient intakes estimated by the FCD and major codes of the DCD are shown in Table 3. Energy intake was overestimated by the DCD but was underestimated only when the standard PS was used for men. Regarding median intakes of forty-two nutrients estimated using standard PS of the DCD, the number of nutrients that differed significantly between the FCD and the DCD was twenty-eight (67%) and twenty-one (50%) in men and women, respectively. For intakes estimated from reported PS, the respective numbers were thirty-seven (88%) and thirty-five (83%). More than 90% of the differed values were considered overestimation when standard PS was used in women or reported PS was used in men or women, whereas underestimation by DCD was observed more often (79%) when standard PS was used in men. The median percentage difference in nutrient intakes (including energy) was larger in reported PS (8.0%) than in standard PS (6.5%) in men ( $P=0.02$ ).

Spearman correlation coefficients for estimates of food group intakes in the FCD *v.* the major codes of the DCD are shown in Table 4. In both sexes, correlation coefficients for standard PS were  $\geq 0.7$  for rice, noodles, bread, pulses, fruits, pickled vegetables, alcoholic beverages, fish and shellfish, and dairy products, and  $< 0.4$  for animal fats, oils, and seasoning and spices. The median correlation coefficients between the two methods were 0.61 (range: 0.19–0.90) and 0.58 (range: 0.25–0.89) in men and women, respectively. The use of reported PS provided higher median correlation coefficients (0.73, range: 0.23–0.97 in men; 0.72, range: 0.35–0.96 in women) than use of standard PS.

Spearman correlation coefficients for estimates of energy and nutrient intakes in the FCD *v.* the major codes of the DCD are shown in Table 5. For standard PS, the median correlation coefficients were 0.60 (range: 0.25–0.90) and 0.53 (range: 0.15–0.74) in men and women, respectively. For reported PS, the respective values were 0.75 (range: 0.26–0.93) and 0.74 (range: 0.19–0.90), which were higher than the median correlation coefficients for standard PS ( $P < 0.0001$  for both).

Bland–Altman plots were used to assess the agreement between the FCD and DCD with reported PS for intakes of selected foods, energy and macronutrients. In both men (Fig. 4) and women (online supplementary material, Supplemental Fig. 1), there was moderate agreement at the group level whereas agreement at the individual level was somewhat poor.

The results for minor codes of the DCD are presented in the online supplementary material, Supplemental

Tables 3–6. The numbers of food groups and nutrients (including energy) that differed significantly from the FCD were similar to those for major codes. The median percentage differences for food group intakes did not differ between minor codes and major codes, except when standard PS was used in women (17.2% for minor codes, 17.1% for major codes,  $P=0.0497$ ). The median percentage differences for energy and nutrient intakes were, when reported PS was used, larger in major codes than minor codes in men (8.0 *v.* 7.6%,  $P=0.03$ ), while larger in minor codes than major codes in women (7.8 *v.* 8.3%,  $P=0.02$ ). The median correlation coefficients for food group intakes and those for energy and nutrient intakes were higher in minor codes than major codes, for both standard PS and reported PS, and in both sexes. For standard PS, the median correlation coefficients for food group intakes and energy and nutrient intakes were 0.65 (range: 0.26–0.90) and 0.65 (range: 0.46–0.90) in men, respectively, and 0.67 (range: 0.28–0.90) and 0.55 (range: 0.27–0.82) in women, respectively. The respective values for reported PS were 0.77 (range: 0.32–0.97) and 0.77 (range: 0.53–0.94) in men and 0.76 (range: 0.33–0.96) and 0.76 (range: 0.43–0.90) in women, each of which was higher than the corresponding values for standard PS.

## Discussion

To our knowledge, the present study is the first to develop a DCD and assess its ability to estimate food group and nutrient intakes using different dietary data in a reasonably large sample of Japanese men and women. Although median intakes of many food groups and nutrients were significantly different between the DCD and the FCD, the median correlation coefficients were moderate for both food groups and nutrients. These results show that the DCD developed in the present study had acceptable ranking ability for intakes of many food groups and nutrients.

Our results are consistent with those of previous studies showing that representative values (mean or median) of the intakes of many food groups and nutrients were different between the DCD and FCD<sup>(3,34,37,39,40)</sup>. The application of the DCD to a DR can lead to measurement error by diluting detailed information on specific foods. This error becomes larger if the PS or composition of each dish differs markedly between dietary data used for the development of a DCD and those used for assessment of applicability of the DCD. It has been suggested that PS and dish composition vary according to between-individual factors such as age, sex and geographic region and also vary within the same dish consumed by the same individual<sup>(7,40,52)</sup>. In the present study, the mean age of participants for the 16 d DR was higher than those for the 4 d DR. Furthermore, two surveys were conducted in different areas, seasons and years. These differences could cause measurement error of the DCD. We calculated food group and nutrient intakes using

**Table 3** Comparison of energy and nutrient intakes estimated based on the food composition database (FCD) and those estimated based on the dish composition database (DCD), with use of standard portion size data or reported portion size data

Nutrient	Unit	Men (n 196)												Women (n 196)											
		DCD†												DCD†											
		FCD‡			Standard portion size§			Reported portion size			FCD‡			Standard portion size§			Reported portion size								
		Median	P <sub>25</sub>	P <sub>75</sub>	Median	P <sub>25</sub>	P <sub>75</sub>	P¶	Median	P <sub>25</sub>	P <sub>75</sub>	P¶	Median	P <sub>25</sub>	P <sub>75</sub>	P¶	Median	P <sub>25</sub>	P <sub>75</sub>	P¶	Median	P <sub>25</sub>	P <sub>75</sub>	P¶	
Total energy	kJ/d	9830	8563	11 068	8688	7421	9760	<0.0001	10 018	8632	11 521	<0.0001	7717	6921	8772	8165	7016	9229	0.01	8105	7095	9067	<0.0001		
Protein	g/d	81.7	69.2	93.3	74.3	63.2	85.3	<0.0001	87.0	72.8	98.3	<0.0001	65.6	57.0	76.0	70.8	59.1	81.7	0.0003	71.8	61.6	82.6	<0.0001		
Fat	g/d	67.7	57.0	83.6	62.9	53.0	73.4	<0.0001	72.2	60.2	83.3	0.03	60.3	51.6	72.0	60.8	52.2	71.8	0.96	62.7	53.6	72.6	0.11		
SFA	g/d	18.5	15.4	23.0	17.5	14.4	20.8	0.009	19.6	16.4	24.2	0.0008	17.9	13.8	21.2	17.9	14.7	21.1	0.59	18.1	15.1	21.3	0.02		
MUFA	g/d	24.56	20.0	31.23	23.26	19.00	27.77	0.002	26.18	21.70	30.63	0.002	21.59	17.49	25.79	21.86	18.63	25.76	0.60	22.29	19.18	26.18	0.03		
PUFA	g/d	14.71	12.0	17.77	13.85	11.27	16.24	0.005	15.82	13.29	18.14	<0.0001	12.53	9.98	14.80	13.09	10.87	15.14	0.17	13.21	11.52	15.37	0.003		
n-6 PUFA	g/d	12.25	9.90	14.74	11.22	9.26	13.40	0.002	12.82	10.88	14.83	0.002	10.32	8.12	12.65	10.81	9.02	12.43	0.24	10.90	9.38	12.45	0.02		
n-3 PUFA	g/d	2.36	1.75	3.07	2.43	1.94	3.03	0.89	2.79	2.27	3.50	<0.0001	1.90	1.44	2.46	2.18	1.81	2.77	<0.0001	2.35	1.85	2.97	<0.0001		
EPA	g/d	0.22	0.08	0.41	0.24	0.15	0.36	0.41	0.31	0.18	0.46	<0.0001	0.16	0.06	0.29	0.21	0.13	0.33	<0.0001	0.24	0.13	0.39	<0.0001		
DHA	g/d	0.40	0.20	0.72	0.43	0.28	0.63	0.63	0.55	0.33	0.77	<0.0001	0.31	0.15	0.51	0.39	0.26	0.56	<0.0001	0.43	0.25	0.66	<0.0001		
α-Linolenic acid	g/d	1.47	1.16	1.90	1.50	1.24	1.86	0.43	1.79	1.42	2.04	<0.0001	1.30	1.02	1.60	1.43	1.19	1.71	0.0001	1.48	1.23	1.70	<0.0001		
Cholesterol	mg/d	366	270	441	346	281	424	0.41	393	314	478	<0.0001	305	236	383	319	253	383	0.002	336	272	401	<0.0001		
Carbohydrate	g/d	300	263	350	269	225	301	<0.0001	319	268	358	<0.0001	251	220	285	265	229	301	<0.0001	259	226	292	<0.0001		
Soluble dietary fibre	g/d	3.1	2.4	3.8	3.1	2.5	3.6	0.73	3.4	2.9	4.0	<0.0001	3.0	2.5	3.8	3.2	2.6	3.7	0.10	3.1	2.7	3.7	<0.0001		
Insoluble dietary fibre	g/d	10.0	7.8	12.5	9.8	7.9	11.4	0.04	11.0	8.9	12.7	<0.0001	9.5	7.8	11.8	10.0	8.2	11.7	0.23	9.6	8.2	11.6	0.007		
Total dietary fibre	g/d	14.0	10.8	16.8	13.6	11.0	15.9	0.06	15.3	12.4	17.5	<0.0001	13.3	10.8	16.1	14.0	11.4	16.2	0.25	13.5	11.5	16.2	0.0007		
Na	mg/d	4399	3572	4920	4916	4121	5919	<0.0001	5628	4517	6842	<0.0001	3502	2928	4233	4672	3750	5604	<0.0001	5002	3810	5855	<0.0001		
K	mg/d	2650	2233	3216	2508	2106	2984	0.0003	2858	2445	3356	<0.0001	2512	2140	2877	2564	2117	2992	0.36	2633	2223	3013	<0.0001		
Ca	mg/d	487	379	637	475	387	609	0.18	538	437	681	<0.0001	500	390	599	520	427	609	0.052	531	440	617	<0.0001		
Mg	mg/d	276	231	332	267	220	311	0.001	306	258	349	<0.0001	250	216	289	267	221	307	0.01	269	228	312	<0.0001		
P	mg/d	1170	941	1352	1053	900	1250	<0.0001	1264	1033	1428	<0.0001	984	851	1138	1051	877	1213	0.0002	1067	914	1222	<0.0001		
Fe	mg/d	8.8	7.1	10.3	8.1	6.9	9.5	0.0005	9.4	8.3	10.8	<0.0001	7.6	6.4	9.3	8.3	6.7	9.6	0.16	8.3	7.0	9.7	<0.0001		
Zn	mg/d	9.2	7.8	10.9	8.9	7.7	10.3	0.03	10.3	8.7	11.8	<0.0001	7.4	6.4	8.6	8.6	6.9	9.8	<0.0001	8.3	7.4	9.6	<0.0001		
Cu	mg/d	1.28	1.05	1.54	1.20	1.02	1.39	0.0002	1.39	1.18	1.61	<0.0001	1.08	0.90	1.27	1.22	0.99	1.40	<0.0001	1.17	1.00	1.34	<0.0001		
Mn	mg/d	3.77	3.06	4.92	3.60	2.98	4.12	<0.0001	4.21	3.54	5.16	<0.0001	3.44	2.81	4.44	3.77	3.12	4.24	0.54	3.77	3.14	4.58	0.003		
Retinol	µg/d	158	116	216	366	249	553	<0.0001	398	269	592	<0.0001	157	116	203	319	215	509	<0.0001	311	217	532	<0.0001		
α-Carotene	µg/d	484	280	802	442	355	622	0.04	503	394	663	0.68	440	267	668	472	367	588	0.60	461	367	547	0.78		
β-Carotene	µg/d	2986	1786	4481	2891	2300	3687	0.77	3324	2505	3889	0.06	2906	1917	4123	3009	2393	3633	0.73	2979	2389	3711	0.61		
Cryptoxanthin	µg/d	61	33	204	134	43	294	0.02	119	47	240	0.10	76	36	345	198	72	322	0.07	151	71	305	0.39		




**Table 3 Continued**

Nutrient	Unit	Men (n 196)											Women (n 196)										
		DCD†											DCD†										
		FCD‡			Standard portion size§				Reported portion size				FCD†			Standard portion size§				Reported portion size			
		Median	P <sub>25</sub>	P <sub>75</sub>	Median	P <sub>25</sub>	P <sub>75</sub>	P¶	Median	P <sub>25</sub>	P <sub>75</sub>	P¶	Median	P <sub>25</sub>	P <sub>75</sub>	Median	P <sub>25</sub>	P <sub>75</sub>	P¶	Median	P <sub>25</sub>	P <sub>75</sub>	P¶
β-Carotene equivalent	μg/d	3525	2093	4987	3295	2633	4178	0.56	3759	2867	4407	0.14	3334	2204	4536	3423	2713	4115	0.75	3414	2719	4103	0.84
Vitamin A (retinol equivalent)	μg/d	472	352	649	678	513	854	<0.0001	713	564	961	<0.0001	448	358	565	625	493	835	<0.0001	622	478	850	<0.0001
Vitamin D	μg/d	6.6	3.5	12.0	6.8	4.8	9.6	0.52	8.8	5.8	11.4	0.0004	5.5	3.0	9.2	6.5	4.6	8.7	0.06	6.9	4.7	9.9	0.0001
α-Tocopherol	mg/d	7.5	6.1	9.1	7.6	6.3	9.0	0.75	8.4	7.2	10.1	<0.0001	7.0	5.4	8.6	7.5	6.4	8.5	0.01	7.5	6.5	8.8	<0.0001
Vitamin K	μg/d	224	152	301	216	170	265	0.19	235	186	296	0.004	214	160	311	208	160	283	0.053	205	165	292	0.44
Thiamin	mg/d	1.09	0.88	1.31	0.99	0.84	1.13	<0.0001	1.08	0.96	1.33	0.03	0.86	0.72	1.07	0.93	0.80	1.10	0.04	0.96	0.79	1.11	<0.0001
Riboflavin	mg/d	1.39	1.10	1.69	1.27	1.06	1.49	<0.0001	1.47	1.23	1.76	<0.0001	1.25	1.03	1.54	1.29	1.07	1.50	0.99	1.35	1.11	1.59	<0.0001
Niacin	mg/d	20.6	16.8	25.0	18.4	15.4	22.2	<0.0001	22.2	18.3	25.8	<0.0001	17.0	14.0	19.9	17.0	14.1	20.3	0.10	18.1	15.5	20.7	<0.0001
Vitamin B <sub>6</sub>	mg/d	1.39	1.06	1.72	1.29	1.10	1.55	0.052	1.48	1.28	1.74	<0.0001	1.15	0.96	1.37	1.23	1.02	1.46	0.006	1.24	1.05	1.45	<0.0001
Vitamin B <sub>12</sub>	μg/d	6.1	3.7	9.3	7.7	5.6	9.6	<0.0001	9.3	6.6	11.0	<0.0001	4.2	2.8	7.7	6.6	5.1	8.8	<0.0001	7.2	5.6	9.4	<0.0001
Folate	μg/d	352	256	471	337	276	399	0.007	381	320	446	0.0001	346	271	429	339	282	396	0.14	352	286	414	0.33
Pantothenic acid	mg/d	6.52	5.46	7.94	6.12	5.16	7.17	<0.0001	6.98	5.94	8.09	<0.0001	5.69	4.65	6.70	6.17	4.86	6.99	0.002	5.97	5.05	7.01	<0.0001
Vitamin C	mg/d	102	72	144	94	72	118	<0.0001	102	80	129	0.43	109	80	143	99	79	116	<0.0001	97	81	123	<0.0001
Alcohol	g/d	4.1	0.6	28.2	5.0	0.3	19.4	<0.0001	3.8	0.4	27.1	<0.0001	0.9	0.3	3.6	0.4	0.3	4.9	<0.0001	0.5	0.3	2.5	0.0004

P<sub>25</sub>, 25th percentile; P<sub>75</sub>, 75th percentile.

†Nutrient intake estimated using the DCD for 128 kinds of dishes developed from 16 d dietary records in 126 men and 126 women.

‡Nutrient intake estimated using the Standard Table of Food Composition in Japan<sup>(36)</sup>.

§Nutrient intake estimated using the weight of each dish in the DCD.

|| Estimated nutrient intake calculated by adjusting nutrient content of a dish in the DCD by reported portion size of the dish in the dietary records. The calculation method was as follows: estimated nutrient intake from a dish adjusted by reported portion size = nutrient content of the dish in the DCD (g) × reported portion size of the dish in the dietary records (g)/standard portion size of the dish in the DCD (g).

¶The difference compared with values derived from the FCD was tested using the Wilcoxon signed-rank test.

**Table 4** Spearman correlation coefficients between food group intakes estimated based on the food composition database and those estimated based on the dish composition database (DCD)†, with use of standard portion size data or reported portion size data

Food group (g/d)	Men (n 196)		Women (n 196)	
	Standard portion size‡	Reported portion size§	Standard portion size‡	Reported portion size§
Rice	0.74	0.97	0.76	0.95
Noodles	0.89	0.90	0.89	0.92
Bread	0.86	0.94	0.86	0.92
Other grain products	0.40	0.44	0.42	0.46
Nuts	0.46	0.41	0.27	0.40
Pulses	0.76	0.80	0.72	0.74
Potatoes	0.63	0.71	0.63	0.70
Sugar	0.36	0.39	0.44	0.49
Confectioneries	0.81	0.87	0.68	0.76
Animal fats	0.19	0.23	0.37	0.35
Oils	0.36	0.51	0.25	0.42
Fruits	0.90	0.91	0.89	0.96
Green and yellow vegetables	0.51	0.70	0.49	0.71
Other vegetables	0.48	0.76	0.38	0.74
Pickled vegetables	0.74	0.77	0.72	0.73
Mushrooms	0.38	0.48	0.49	0.56
Seaweeds	0.44	0.43	0.49	0.50
Fruit and vegetable juice	0.62	0.63	0.56	0.62
Seasonings and spices	0.30	0.33	0.27	0.35
Alcoholic beverages	0.90	0.92	0.71	0.74
Tea and coffee	0.58	0.76	0.60	0.76
Soft drinks	0.56	0.54	0.57	0.53
Fish and shellfish	0.75	0.83	0.72	0.82
Meats	0.61	0.77	0.53	0.70
Eggs	0.61	0.67	0.69	0.73
Dairy products	0.83	0.87	0.75	0.84

†DCD for 128 kinds of dishes developed from 16 d dietary records in 126 men and 126 women.

‡Food group intake estimated using the weight of each dish in the DCD.

§Estimated food group intake calculated by adjusting weight of a food group of a dish in the DCD by reported portion size of the dish in the dietary records. The calculation method was as follows: estimated food group intake from a dish adjusted by reported portion size = weight of a food group in the dish in the DCD (g) × reported portion size of the dish in the dietary records (g)/standard portion size of the dish in the DCD (g).

reported PS of the 4 d DR to examine whether the use of reported PS improves the ability of the DCD to estimate dietary intakes. However, this did not reduce either the number of the food groups and nutrients in which median intakes were different between the DCD and the FCD or the median percentage difference. This suggests that there were differences in dish composition between the 16 d and 4 d DR, which would be a major factor affecting the performance of the DCD. In addition, the process of applying dish codes of the DCD to each dish can be problematic. Dish codes were assigned on the basis of dish names recorded in the DR. Hence, if an inaccurate dish name was written in the DR, an incorrect dish code was assigned and dietary intake from the dish was incorrectly estimated.

Food group and nutrient intakes were often underestimated when standard PS was used in men. Because the DCD was developed for men and women combined,

**Table 5** Spearman correlation coefficients between energy and nutrient intakes estimated based on the food composition database and those estimated based on the dish composition database (DCD)†, with use of standard portion size data or reported portion size data


	Men (n 196)		Women (n 196)	
	Standard portion size‡	Reported portion size§	Standard portion size‡	Reported portion size§
Total energy	0.63	0.93	0.52	0.88
Protein	0.65	0.87	0.53	0.79
Fat	0.57	0.77	0.36	0.69
SFA	0.55	0.72	0.43	0.70
MUFA	0.54	0.73	0.35	0.65
PUFA	0.49	0.68	0.30	0.63
n-6 PUFA	0.48	0.68	0.32	0.63
n-3 PUFA	0.57	0.71	0.57	0.70
EPA	0.69	0.77	0.72	0.74
DHA	0.69	0.76	0.69	0.74
α-Linolenic acid	0.48	0.61	0.38	0.60
Cholesterol	0.61	0.75	0.56	0.68
Carbohydrate	0.66	0.93	0.60	0.89
Soluble dietary fibre	0.66	0.87	0.62	0.85
Insoluble dietary fibre	0.65	0.90	0.59	0.88
Total dietary fibre	0.67	0.89	0.61	0.90
Na	0.46	0.53	0.46	0.63
K	0.69	0.89	0.55	0.88
Ca	0.69	0.84	0.61	0.82
Mg	0.68	0.89	0.57	0.86
P	0.68	0.89	0.58	0.83
Fe	0.59	0.80	0.59	0.80
Zn	0.63	0.85	0.50	0.74
Cu	0.66	0.91	0.62	0.86
Mn	0.65	0.72	0.57	0.68
Retinol	0.25	0.26	0.15	0.19
α-Carotene	0.45	0.56	0.40	0.59
β-Carotene	0.59	0.74	0.44	0.68
Cryptoxanthin	0.63	0.68	0.67	0.74
β-Carotene equivalent	0.59	0.75	0.44	0.69
Vitamin A (retinol equivalent)	0.37	0.48	0.29	0.42
Vitamin D	0.67	0.68	0.64	0.68
α-Tocopherol	0.59	0.77	0.43	0.71
Vitamin K	0.67	0.81	0.63	0.79
Thiamin	0.41	0.67	0.39	0.69
Riboflavin	0.58	0.73	0.51	0.66
Niacin	0.57	0.73	0.42	0.68
Vitamin B <sub>6</sub>	0.55	0.75	0.43	0.77
Vitamin B <sub>12</sub>	0.60	0.70	0.53	0.59
Folate	0.55	0.72	0.53	0.77
Pantothenic acid	0.60	0.84	0.57	0.81
Vitamin C	0.64	0.78	0.59	0.74
Alcohol	0.90	0.93	0.74	0.77

†DCD for 128 kinds of dishes developed from 16 d dietary records in 126 men and 126 women.

‡Nutrient intake estimated using the weight of each dish in the DCD.

§Estimated nutrient intake calculated by adjusting nutrient content of a dish in the DCD by reported portion size in the dietary records. The calculation method was as follows: estimated nutrient intake from a dish adjusted by reported portion size = nutrient content of the dish in the DCD (g) × reported portion size of the dish in the dietary records (g)/standard portion size of the dish in the DCD (g).

standard PS reflects the PS for both men and women. This may result in underestimation of food group and nutrient intakes in men because women generally eat smaller amounts of food than men. Nevertheless, we did not separate men and women during development of



**Fig. 4** Bland–Altman plot assessing the agreement between the food composition database (FCD) and the dish composition database (DCD) with reported portion size for intakes of (a) rice, (b) oils, (c) fruits, (d) fish, (e) energy, (f) protein, (g) fat and (h) carbohydrate in Japanese men (*n* 196). — represents the mean difference and - - - represent the lower and upper 95 % limits of agreement



the DCD, based on the fact that the reliability of dietary intake estimation using the DCD would be higher with a greater number of different types of dishes constituting the DCD<sup>(34)</sup>. Additionally, to our knowledge, there has been no DCD developed for men and women separately. However, sex differences should be taken into consideration when interpreting food group and nutrient intakes estimated using the DCD.

Despite poor agreement at individual level, the DCD showed acceptable ranking ability for food groups, energy and nutrients. The median correlation coefficients with the FCD for food groups, energy and nutrients were within the ranges reported in previous studies (0.54–0.92 for food groups<sup>(34,35,37,39,40)</sup> and 0.50–0.81 for energy and nutrients<sup>(3,34,35,37,39,40)</sup>). The correlation coefficient was relatively high ( $\geq 0.7$ ) for food groups generally eaten in large portions such as staple foods (rice, noodles, bread), those eaten as main ingredients in Japan (pulses, fish and shellfish), or those eaten as a single food (fruits, pickled vegetables, alcoholic beverages and dairy products). Meanwhile, the correlation coefficient was low ( $< 0.4$ ) for food groups consumed in small portions or that are 'integrated' within a dish such as animal fats, oils, and seasoning and spices. These food groups with high correlation or with low correlation are similar to those in previous studies<sup>(34,35,37,39,40)</sup>. It has also been reported that foods used in relatively large amounts in a dish were likely to be reflected in the dish name and had high correlation<sup>(34)</sup>. In addition, it has also been suggested that between-individual variation in intake is large for foods or beverages such as yoghurt, fruits, natto (fermented soybeans), and tea and coffee<sup>(7)</sup>. On the other hand, between-individual variation has been reported to be small for animal fats, oils, and seasoning and spices<sup>(7)</sup>, which can lead to low correlation coefficients. Nutrients with correlation coefficients of about 0.7 or above for both sexes in the present study had high correlation coefficients for those food sources; for example, EPA and DHA for fish, and alcohol for alcoholic beverages.

We compared the results on reported PS and standard PS, and major codes and minor codes. As mentioned above, the use of reported PS did not improve the estimation ability of median intake of the DCD; however, the correlation coefficients were improved between the DCD and FCD for both food groups and nutrients. This indicates that if the PS of dishes is reported by participants in addition to a dish name, the DCD can rank individuals more accurately with respect to food group and nutrient intakes. When comparing major codes and minor codes, there was no great difference in estimation of median intakes of food groups, energy and nutrients, indicating that the level of aggregation of dishes in the present study did not profoundly affect the ability of the DCD to estimate median intakes. However, compared with major codes, minor codes ranked food group and nutrient intakes well, suggesting that detailed classification of dishes in the

DCD may be effective to improve the ranking ability of the DCD. Nevertheless, the difference in correlation coefficients between minor codes and major codes was small. Given that the benefits of dish-based dietary assessment methods include reducing the burden on participants and staff involved in dietary surveys<sup>(8,34,35,40)</sup>, the use of minor codes may not always be required.

Several limitations of the present study should be acknowledged. First, the comparison of the DCD and the FCD is not exactly an appropriate evaluation of the validity of the DCD because estimation using the DCD and the FCD were conducted based on the same DR. Although this method would minimize errors regarding inaccurate recording and selection of dishes, the concordance between the intakes estimated by the DCD and by the FCD is likely overestimated. Ideally, the validity of the DCD should be assessed by comparison of results of a dish-based dietary assessment with values estimated from objective measures of dietary intake (e.g. biomarkers or duplicated methods). However, since these methods are expensive and few validation studies of DCD have been previously conducted, we conducted the present study as the initial step towards development of a dish-based dietary assessment method in Japanese people. Despite the limitation of study design, the estimated intakes of the DCD were moderately correlated with those of the FCD, thus the DCD may be useful in the practical assessment of food group and nutrient intakes. Nevertheless, a stricter evaluation of validity is required to examine the ability of the DCD to estimate dietary intake in the future. Second, the dietary survey for developing DCD was conducted in four seasons, whereas that for simulated validation was conducted in one season. Although the DCD developed reflecting seasonal variation in dietary habits<sup>(41–43)</sup> is a strength of the current study, the difference in seasons between two dietary surveys might affect the results of validation. Hence, the validation of the DCD in other seasons should be confirmed in future studies. Third, the 16 d DR was obtained from cohabiting couples (with the percentage of single foods eaten at home: 77%), which might reduce between-individual variation of dishes. However, the number of dishes used for development of the DCD in the present study (71 213) was larger than those in previous studies (4814<sup>(34)</sup>, 10 533<sup>(35)</sup>, 42 508<sup>(38)</sup> and 67 532<sup>(37)</sup>). Fourth, the participants were not randomly selected and thus may not be representative of the general Japanese population. Participants were volunteers who were considered to be more health conscious than the general population. Nevertheless, the weight and height of our participants were similar to those of the general population in Japan<sup>(53)</sup>. Finally, dish classification can be subjective, as no standardized protocol has been available. Subjectivity also exists in how participants named each dish. Furthermore, beyond the scope of the study, we did not examine whether the performance of the DCD is dependent on individual characteristics such



as education, age and obesity (because of sample size limitations). Increasing sample size and evaluating the effect of participant characteristics on dish composition should be considered in future.

## Conclusion

In conclusion, we developed a DCD and assessed its ability to estimate food group and nutrient intakes. Whereas it is difficult to accurately estimate dietary intake using the DCD developed in the present study, it has acceptable ranking ability for intakes of many food groups and nutrients commonly consumed in Japan. The DCD may be useful for future dietary surveys not only to rank individuals, but also to characterize dietary patterns of populations, because mixed dishes represent combination of foods and cooking methods. However, consideration should be given to issues concerning study design and further investigations are needed to establish a dish-based dietary assessment method.

## Acknowledgements

**Acknowledgements:** The authors are grateful to all participants and local staff for their participation in this study. They also thank N. Hirota, A. Notsu, A. Miura, M. Fukui, H. Todoriki and C. Date for data collection. **Financial support:** This study was funded by a Health and Labour Sciences Research Grant (number H23-jyunkankitou (seishuu)-ippan-001) and H13 Health Sciences Research Grant (Kenkou-kagakusougoukenkyujigyou) from the Ministry of Health, Labour and Welfare, Japan. The Ministry of Health, Labour and Welfare had no role in the design, analysis or writing of this article. **Conflict of interest:** None of the authors has any conflict of interest to declare. **Authorship:** N.S. contributed to conceptualization of the research, developed the dish composition database, performed the statistical analysis, wrote the first draft of the manuscript and prepared the revised version of the manuscript. K.M. contributed to conceptualization of the research, provided critical input to the final draft of the manuscript and contributed to the preparation of the revised version of the manuscript. S.M. contributed to data collection. S.S. directed the survey and contributed to data collection. All authors read and approved the final manuscript. **Ethics of human subject participation:** The study purpose and protocol were explained before the study and written informed consent was obtained from each participant. Use of data on the 16 d dietary record survey and the study protocol of the 4 d dietary record were approved by the Ethics Committee at the University of Tokyo, Faculty of Medicine (numbers 3421 and 10005, respectively).

## Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1368980019000600>

**Author ORCID.** Nana Shinozaki, 0000-0002-8998-5066.

## References

1. Shim J-S, Oh K & Kim HC (2014) Dietary assessment methods in epidemiologic studies. *Epidemiol Health* **36**, e2014009.
2. Thompson FE, Subar AF, Loria CM *et al.* (2010) Need for technological innovation in dietary assessment. *J Am Diet Assoc* **110**, 48–51.
3. Matsuzaki E, Michie M & Kawabata T (2017) Validity of nutrient intakes derived from an internet website dish-based dietary record for self-management of weight among Japanese women. *Nutrients* **9**, 1058.
4. Bognár A & Piekarski J (2000) Guidelines for recipe information and calculation of nutrient composition of prepared foods (dishes). *J Food Compos Anal* **13**, 391–410.
5. Date C, Yamaguchi M & Tanaka H (1996) Development of a food frequency questionnaire in Japan. *J Epidemiol* **6**, 131–136.
6. Christensen SE, Möller E, Bonn SE *et al.* (2013) Two new meal- and web-based interactive food frequency questionnaires: validation of energy and macronutrient intake. *J Med Internet Res* **15**, e109.
7. Ishihara J, Takachi R, Hosoi S *et al.* (2009) Application of digital photographic images of meals to assess dietary intake in epidemiological studies. *Jpn J Nutr Diet* **67**, 252–259 (in Japanese).
8. Imaeda N, Goto C, Fujiwara N *et al.* (2016) Comparison of nutrient contents between the standard recipes used in the National Health and Nutrition Survey and the dishes actually consumed by community-dwelling elderly individuals. *J Jpn Soc Nutr Food Sci* **69**, 237–248 (in Japanese).
9. Bates B, Lennox A, Prentice A *et al.* (2014) National Diet and Nutrition Survey: results from years 1, 2, 3 and 4 (combined) of the rolling programme (2008/2009–2011/2012). [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/594361/NDNS\\_Y1\\_to\\_4\\_UK\\_report\\_full\\_text\\_revised\\_February\\_2017.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/594361/NDNS_Y1_to_4_UK_report_full_text_revised_February_2017.pdf) (accessed June 2017).
10. Fitt E, Mak T, Stephen A *et al.* (2010) Disaggregating composite food codes in the UK National Diet and Nutrition Survey food composition databank. *Eur J Clin Nutr* **64**, 32–36.
11. Fitt E, Cole D, Ziauddeen N *et al.* (2015) DINO (Diet In Nutrients Out) – an integrated dietary assessment system. *Public Health Nutr* **18**, 234–241.
12. Stumbo PJ (2001) Structure and uses of USDA food composition databases. *J Food Compos Anal* **14**, 323–328.
13. MRC Human Nutrition Research (2017) Food Standards Agency Standard Recipes Database, 1992–2012 [data collection]. UK Data Service SN 8159. <http://doi.org/10.5255/UKDA-SN-8159-1> (accessed May 2018).
14. Fitt E, Prynne CJ, Teucher B *et al.* (2009) National Diet and Nutrition Survey: assigning mixed dishes to food groups in the nutrient databank. *J Food Compos Anal* **22**, S52–S56.
15. Bodner JE & Perloff BP (2003) Databases for analyzing dietary data – the latest word from What We Eat in America. *J Food Compos Anal* **16**, 347–358.
16. Merchant AT & Dehghan M (2006) Food composition database development for between country comparisons. *Nutr J* **5**, 2.
17. Reinivuo H, Bell S & Ovaskainen M-L (2009) Harmonisation of recipe calculation procedures in European food composition databases. *J Food Compos Anal* **22**, 410–413.
18. Joslowski G, Yang J, Andrén Aronsson C *et al.* (2017) Development of a harmonized food grouping system for

- between-country comparisons in the TEDDY Study. *J Food Compos Anal* **63**, 79–88.
19. Slimani N, Deharveng G, Unwin I *et al.* (2007) The EPIC nutrient database project (ENDB): a first attempt to standardize nutrient databases across the 10 European countries participating in the EPIC study. *Eur J Clin Nutr* **61**, 1037–1056.
  20. Durazzo A, Lisciani S, Camilli E *et al.* (2017) Nutritional composition and antioxidant properties of traditional Italian dishes. *Food Chem* **218**, 70–77.
  21. Khokhar S, Marletta L, Shahar DR *et al.* (2010) New food composition data on selected ethnic foods consumed in Europe. *Eur J Clin Nutr* **64**, Suppl. 3, S82–S87.
  22. Lander RL, Hambidge KM, Krebs NF *et al.* (2017) Repeat 24-hour recalls and locally developed food composition databases: a feasible method to estimate dietary adequacy in a multi-site preconception maternal nutrition RCT. *Food Nutr Res* **61**, 1311185.
  23. Shai I, Vardi H, Shahar DR *et al.* (2003) Adaptation of international nutrition databases and data-entry system tools to a specific population. *Public Health Nutr* **6**, 401–406.
  24. Shin S, Park E, Sun DH *et al.* (2014) Development and evaluation of a web-based computer-assisted personal interview system (CAPIS) for open-ended dietary assessments among Koreans. *Clin Nutr Res* **3**, 115–125.
  25. Gurinović M, Milešević J, Kadvan A *et al.* (2016) Establishment and advances in the online Serbian food and recipe data base harmonized with EuroFIR™ standards. *Food Chem* **193**, 30–38.
  26. Kim YO, Kim MK, Lee SA *et al.* (2009) A study testing the usefulness of a dish-based food-frequency questionnaire developed for epidemiological studies in Korea. *Br J Nutr* **101**, 1218–1227.
  27. Park MK, Kim DW, Kim J *et al.* (2011) Development of a dish-based, semi-quantitative FFQ for the Korean diet and cancer research using a database approach. *Br J Nutr* **105**, 1065–1072.
  28. Yum J & Lee S (2016) Development and evaluation of a dish-based semiquantitative food frequency questionnaire for Korean adolescents. *Nutr Res Pract* **10**, 433–441.
  29. Park MK, Noh HY, Song NY *et al.* (2012) Validity and reliability of a dish-based, semi-quantitative food frequency questionnaire for Korean diet and cancer research. *Asian Pac J Cancer Prev* **13**, 545–552.
  30. Lin P-ID, Bromage S, Mostofa MG *et al.* (2017) Validation of a dish-based semiquantitative food questionnaire in rural Bangladesh. *Nutrients* **9**, 49.
  31. Keshтели A, Esmailzadeh A, Rajaie S *et al.* (2014) A dish-based semi-quantitative food frequency questionnaire for assessment of dietary intakes in epidemiologic studies in Iran: design and development. *Int J Prev Med* **5**, 29–36.
  32. Christensen SE, Möller E, Bonn SE *et al.* (2014) Relative validity of micronutrient and fiber intake assessed with two new interactive meal- and web-based food frequency questionnaires. *J Med Internet Res* **16**, e59.
  33. Kobayashi T, Tanaka S, Toji C *et al.* (2010) Development of a food frequency questionnaire to estimate habitual dietary intake in Japanese children. *Nutr J* **9**, 17.
  34. Takachi R, Kudo Y, Watanabe S *et al.* (2006) Validity of smaller sample sizes for the dish-based component table to use with the self-reported 'dietary record by cooked dishes'. *Jpn J Nutr Diet* **64**, 97–105 (in Japanese).
  35. Kimira M, Takachi R, Kudo Y *et al.* (2004) Simplified nutritional survey by 'dietary record by cooked dishes' (DRcd). *Jpn B Heal Fit Nutr* **10**, 3–13 (in Japanese).
  36. Science and Technology Agency (2010) *Standard Tables of Food Composition in Japan, 2010*. Tokyo: Official Gazette Co-operation of Japan (in Japanese).
  37. Imai T, Otsuka R, Katou Y *et al.* (2009) Validity of nutrient intake assessed by the food balance questionnaire using a foods and dishes database with serving size information. *Jpn J Nutr Diet* **67**, 301–309 (in Japanese).
  38. Hayabuchi H, Hisano M, Matsunaga Y *et al.* (2007) A systematic analytical approach for the classification of Japanese dishes using dietary intake data based on the food weighing method. *J Jpn Soc Food Nutr* **60**, 189–198 (in Japanese).
  39. Imai T, Kato T, Otsuka R *et al.* (2013) The utility of food balance questionnaire in college students. *J Jpn Miyou Syst Assoc* **19**, 93–97 (in Japanese).
  40. Kito K, Ishihara J, Kimira M *et al.* (2012) Applicability of the dietary record by cooked dishes method for estimating dietary intake of populations in the areas other than where the database was developed. *Jpn J Public Health* **59**, 700–711 (in Japanese).
  41. Tokudome Y, Imaeda N, Nagaya T *et al.* (2002) Daily, weekly, seasonal, within- and between-individual variation in nutrient intake according to four season consecutive 7 day weighed diet records in Japanese female dietitians. *J Epidemiol* **12**, 85–92.
  42. Owaki A, Takatsuka N, Kawakami N *et al.* (1996) Seasonal variations of nutrient intake assessed by 24 hour recall method. *Jpn J Nutr Diet* **54**, 11–18.
  43. Suga H, Asakura K, Sasaki S *et al.* (2014) Effect of seasonality on the estimated mean value of nutrients and ranking ability of a self-administered diet history questionnaire. *Nutr J* **13**, 51.
  44. Tsugane S, Sasaki S, Kobayashi M *et al.* (2001) Dietary habits among the JPHC study participants at baseline survey. Japan Public Health Center-based Prospective Study on Cancer and Cardiovascular Diseases. *J Epidemiol* **11**, 6 Suppl., S30–S43.
  45. Murakami K, Sasaki S, Takahashi Y *et al.* (2008) Reproducibility and relative validity of dietary glycaemic index and load assessed with a self-administered diet-history questionnaire in Japanese adults. *Br J Nutr* **99**, 639–648.
  46. Kobayashi S, Murakami K, Sasaki S *et al.* (2011) Comparison of relative validity of food group intakes estimated by comprehensive and brief-type self-administered diet history questionnaires against 16 d dietary records in Japanese adults. *Public Health Nutr* **14**, 1200–1211.
  47. Tani Y, Asakura K, Sasaki S *et al.* (2015) The influence of season and air temperature on water intake by food groups in a sample of free-living Japanese adults. *Eur J Clin Nutr* **69**, 907–913.
  48. Hisano M, Hayabuchi H, Matsunaga Y *et al.* (2008) Classifications of dish groups: comparison of the 8-dish group classification with the dish pattern from a cluster analysis. *Jpn J Nutr Diet* **66**, 15–23.
  49. Gabriel AS, Ninomiya K & Uneyama H (2018) The role of the Japanese traditional diet in healthy and sustainable dietary patterns around the world. *Nutrients* **10**, 173.
  50. Asakura K, Uechi K, Sasaki Y *et al.* (2014) Estimation of sodium and potassium intakes assessed by two 24 h urine collections in healthy Japanese adults: a nationwide study. *Br J Nutr* **112**, 1195–1205.
  51. Bland JM & Altman DG (1995) Comparing methods of measurement: why plotting difference against standard method is misleading. *Lancet* **346**, 1085–1087.
  52. Wakimoto P & Block G (2001) Dietary intake, dietary patterns, and changes with age: an epidemiological perspective. *J Gerontol A Biol Sci Med Sci* **56**, 65–80.
  53. Ministry of Health, Labour and Welfare (2017) The National Health and Nutrition Survey in Japan, 2016 (in Japanese). <http://www.mhlw.go.jp/bunya/kenkou/eiyoudl/h28-houkuoku.pdf> (accessed June 2018).