

Infant and young child feeding practices, sociodemographic factors and their association with nutritional status of children aged <3 years in India: findings of the National Nutrition Monitoring Bureau survey, 2011–2012

Indrapal Ishwarji Meshram*, K Mallikharjun Rao, Nagalla Balakrishna, R Harikumar, N Arlappa, Kakani Sreeramakrishna and Avula Laxmaiah

Division of Community Studies, National Institute of Nutrition, Indian Council of Medical Research, Jamai-Osmania (PO), Hyderabad – 500007, Tarnaka, India

Submitted 16 May 2018; Final revision received 7 September 2018; Accepted 25 September 2018; First published online 6 November 2018

Abstract

Objective: To study infant and young child feeding (IYCF) practices and their association with nutritional status among young children.

Design: A community-based, cross-sectional study was carried out in ten states of India, using a multistage random sampling method. Anthropometric measurements such as length/height and weight were conducted and nutritional assessment was done using the WHO child growth standards.

Setting: National Nutrition Monitoring Bureau survey, 2011–2012.

Participants: Children aged <3 years and their mothers.

Results: Only 36% of infants received breast-feeding within an hour of birth and 50% were exclusively breast-fed up to 6 months. Prevalence of underweight, stunting and wasting was 38, 41 and 22%, respectively. The chance of undernutrition among <3-year-old children was significantly higher among those from scheduled caste/scheduled tribe communities, the lowest-income group, with illiterate mothers and lack of sanitary latrine. Among infants, the chance of undernutrition was significantly higher among low-birth-weight babies, and among children whose mother had not consumed iron-folic acid tablets during pregnancy. Immunization practices and minimum dietary diversity were observed to be associated with undernutrition among 12–23-month-old children.

Conclusions: Undernutrition is still an important public health problem in India and observed to be associated with low socio-economic status, illiteracy of mother, low birth weight and dietary diversity. Improving socio-economic and literacy status of mothers can help in improving maternal nutrition during pregnancy and thus low birth weight. Also, improving knowledge of mothers about IYCF practices will help in improving children's nutritional status.

Keywords
Undernutrition
Infant and young child
feeding practices
Conceptual framework
Nutritional assessment
Minimum dietary diversity

Undernutrition continues to be an important public health problem in India, despite several nutrition intervention programmes in operation over the last four decades. Undernutrition is responsible directly or indirectly for about 45% of deaths among children under 5 years of age (under-5s) globally, with these children at higher risk of death from common childhood illness such as diarrhoea, pneumonia and malaria⁽¹⁾. Of this, about two-thirds are attributable to suboptimal infant and young child feeding (IYCF) practices and occur during first year of life⁽²⁾. In developing countries, optimal breast-feeding – that is, breast-feeding within an hour of birth, exclusive breast-

feeding for 6 months and continued breast-feeding until age 2 years or longer – has the potential to prevent 12% of all deaths in under-5s⁽¹⁾.

Poor feeding practices during infancy and early childhood, resulting in malnutrition, contribute to impairment of cognitive and social development, poor school performance and reduced productivity in later life⁽³⁾.

Exclusively breast-fed children are less susceptible to diarrhoea and pneumonia and are fourteen times more likely to survive than non-breast-fed children⁽⁴⁾. IYCF is a key area to improve child survival and promote healthy growth and development. The first 2 years of a child's life

are particularly important, as optimal nutrition during this period lowers morbidity and mortality, reduces the risk of chronic disease and fosters better development overall.

Many studies have shown beneficial effects of breast-feeding on infant mortality, respiratory infections, diarrhoea⁽⁴⁾ and neonatal sepsis⁽⁵⁻⁹⁾ and thus on nutritional status. Suboptimum breast-feeding was estimated to be responsible for 1.4 million child deaths and 44 million disability-adjusted life years (10% of disability-adjusted life years in under-5s) for the year 2004⁽²⁾. WHO has recommended breast-feeding to be initiated early after birth, preferably within an hour, avoidance of prelacteal feeds and exclusive breast-feeding up to first 6 months.

UNICEF has set seventeen developmental goals under the Sustainable Development Goals 2015⁽¹⁰⁾, officially known as *Transforming Our World: The 2030 Agenda for Sustainable Development*, which include poverty alleviation, ending hunger, achieving food security and improved nutrition, and promoting sustainable agriculture. Improving nutrition includes: ending all forms of malnutrition, such as reducing stunting and wasting in under-5s by 2025 as agreed internationally; addressing the nutritional needs of adolescent girls, pregnant and lactating women and older persons; and reducing the global maternal mortality rate to less than 70/100 000 live births, the neonatal mortality rate to as low as 12/1000 live births and the under-5 mortality rate to 25/1000 live births⁽¹⁰⁾.

These targets can only be achieved by improving maternal and adolescent nutrition, proper health-care services, control of childhood diseases and reducing undernutrition among under-5s.

The present study was carried out by the National Nutrition Monitoring Bureau (NNMB) on 'diet and nutritional status of the rural population in India' during 2011–2012. Data pertaining to IYCF feeding practices and their influence on nutritional status are presented in the current paper.

Methods

Study design and setting

A community-based, cross-sectional study was carried out in ten states in India (Kerala, Tamil Nadu, Karnataka, Andhra Pradesh, Maharashtra, Gujarat, Madhya Pradesh, Orissa, West Bengal and Uttar Pradesh) adopting a multistage random sampling procedure. A total of 120 villages were covered in each NNMB state.

Selection of villages and households

From each state, the villages were selected based on probability proportional to size of the population. Ninety villages were selected from those covered in previous surveys and the remaining thirty villages were randomly selected afresh from the list of villages obtained from the

Census of India⁽¹¹⁾. From each selected village, twenty households (HH) were covered after dividing the village into five geographical areas based on streets/*moballas*/areas. It was ensured that at least one of the five areas was inhabited by scheduled caste (SC)/scheduled tribe (ST) communities, which are officially designated groups of historically disadvantaged people as per the Constitution of India, and the groups in the other areas were designated in one or other of the population categories wherever possible to give equal representation to all communities. From each area, four contiguous HH were covered by randomly selecting the first HH. Thus, a total of twenty HH were covered in each village and 2400 HH in each state. All children <3 years of age present at the time of the survey were included in the study.

Data collection

Data were collected by a team comprising a medical officer, a nutritionist and a social worker in each state, who were trained and standardized in survey methodologies by scientists from the National Institute of Nutrition, Hyderabad. All survey schedules were pre-tested before being used in the field. Information on household socio-economic and sociodemographic characteristics, such as community, religion, education, occupation and income, were collected using a pre-tested proforma. IYCF practices such as initiation of breast-feeding, age at initiating complementary feeding, colostrum feeding, etc. were also collected from mothers of children aged <2 years. Anthropometric measurements of the children were collected. Length/height (up to nearest 1 mm) was measured with an infantometer/anthropometer rod and weight (up to nearest 100 g) with a SECA weighing scale (SECA Deutschland, Hamburg, Germany) using standard anthropometric procedures⁽¹²⁾. History of morbidity such as fever, respiratory infection, diarrhoea, etc., if any, during the 15 d preceding the visit was also collected.

Definitions

'Household' is defined as those living together under one roof and sharing a common kitchen.

'Pucca house' means walls made of cement and bricks or stones with a reinforced cement concrete roof; a 'semi-pucca house' is one that has brick or stone walls and a tiled or asbestos roof; while a '*kutch*a house' has mud or thatched walls and a thatched or tiled/asbestos roof.

'Minimum dietary diversity' is the number of different foods or food groups consumed over a given reference period and is said to be met by the consumption of four or more food groups in a diet.

'Prelacteal feeds' are those foods given to a newborn before breast-feeding is established or before breast milk 'comes in', usually on the first day of life, and include honey, *jaggery* (brown sugar from sugarcane), *ghee* (clarified butter) and *ghutti* (herbal paste).

'Exclusive breast-feeding' means giving only breast milk, not even water, up to 6 months.

Ethical approval

The study was approved by the Institutional Ethical Committee of the National Institute of Nutrition as well as the Scientific Advisory Committee. Written informed consent was obtained from the mothers involved in the study.

Data analysis

The data were scrutinized and entered on computer at the National Institute of Nutrition, Hyderabad. Data cleaning was done by carrying out range and consistency checks, then data analysis was conducted using the statistical software package SPSS Statistics for Windows version 17.0. Tests of proportions (χ^2 test) and bivariate analyses (logistic regression) were carried out, and multivariate logistic regression analyses were done according to the conceptual hierarchical framework developed by Victora *et al.*⁽¹³⁾. Figure 1 shows the factors controlled for in hierarchical order for underweight, stunting and wasting, which was the basis for adjusted logistic regression analyses. Age and gender were considered inherent factors and were controlled for in all models and at each stage irrespective of significance level. The distal factors (such as community, per capita income (tertile), type of house and landholding) constituted the first stage. Age and

gender remained in the model together with significant distal factors, while intermediate factors (such as age and parity of mother, education of mother, place of delivery, birth weight, sanitary latrine and hand-washing practices of mother) were added, which constituted the second stage. The proximal variables (including breast-feeding practices, age at initiation of complementary feeding and morbidities during the preceding fortnight) were added in the third stage.

The nutritional status of children was assessed according to SD classification⁽¹⁴⁾ using the WHO Child Growth Standards⁽¹⁵⁾. Children who were below 2 SD from the reference median (median < -2 SD) on the basis of weight-for-age, height-for-age and weight-for-height indices were classified as underweight, stunted and wasted, respectively; while children who were below 3 SD from the reference median (median < -3 SD) were classified as severely underweight, severely stunted and severely wasted, respectively.

Results

Coverage

A total of 4038 (2095 boys, 51.9%) children aged 0–35 months were covered for IYCF practices and anthropometric measurements. The mean age of the children was 16.3 (SD 9.5) months.




Fig. 1 (colour online) Factors controlled for in hierarchical order in the adjusted logistic regression analyses for underweight, stunting and wasting (BF, breast-feeding; CF, complementary feeding)

Sociodemographic characteristics

The majority (88.6%) of children belonged to the Hindu religion, while 7% were Muslims; 14% belonged to ST communities, 26% to SC and 35% were from the other backward caste category. About 44% were living in a *kutchha* house, 34% in a semi-*pucca* house, while only 23% had a *pucca* house. About 43% belonged to a nuclear family while 38% were from joint families, 78% of houses had electricity, 39% of HH were using tap water for drinking purposes and 29% of HH had a sanitary latrine. Monthly per capita income was Rs 1083 only.

Feeding practices of infants

Only 36% of the mothers initiated breast-feeding within 1 h of birth, while 15% did so after 24 h. About 50% of children were exclusively breast-fed up to 6 months. Among children aged 6–11 months, 54% had complementary feeding initiated at 6–7 months old. Minimum dietary diversity was observed among 31% of 6–11-month-old children while it was 85% among 12–23-month-old children.

Prevalence of undernutrition among children

The prevalence of underweight, stunting and wasting was 38, 41 and 22%, respectively, and all prevalences except wasting were lower among girls (36% underweight, 39% stunted) compared with boys (40 and 44%, respectively).

Association between sociodemographic characteristics and undernutrition using bivariate analyses

The odds of being underweight were 1.91 times higher among children aged 12–35 months compared with 0–11-month-olds. The odds of being underweight were 1.28 times higher among boys than girls. The odds of underweight were 2.89 and 1.60 times higher among children belonging to ST and SC compared with others; 1.57 times higher among children belonging to nuclear families compared with joint families; 2.17 times higher among children of illiterate mothers compared with children of the most highly educated mothers (9th grade or above); 2.56 times higher among children from HH with per capita income in the first tertile compared with children from HH with per capita income in the third tertile; and 2.38 times higher among children belonging to HH not having a sanitary latrine. Similarly, the odds of stunting and wasting were higher among 12–35-month-olds, among boys, among children belonging to ST and SC communities, among children from nuclear families, of illiterate mothers, from HH with per capita income in the first tertile and from HH not having the facility of a sanitary latrine (Table 1).

Association of antenatal care, delivery and feeding practices with undernutrition among 0–11-month-old children using bivariate analyses

The odds of being underweight were 2.90 times higher among children whose mothers did not attend antenatal check-ups compared with those whose mothers had four or more antenatal visits; and about 2.5 times higher among children whose mothers had not consumed or consumed fewer than ninety iron-folic acid (IFA) tablets during pregnancy, compared with those whose others consumed ninety IFA tablets or more. The odds of underweight were 1.61 times higher among children delivered at home than among those delivered at hospital and 2.22 times higher among children with low birth weight than among those with normal weight at birth. Similarly, the odds of stunting were higher among children delivered at home and among low-birth-weight children. The odds of wasting were higher among children whose mothers did not attend antenatal check-ups and among children whose mothers did not consume or consumed fewer than ninety IFA tablets during pregnancy. Only stunting was observed to be significantly associated with minimum dietary diversity (child's consumption of four or more food groups; OR = 1.56; 95% CI 1.10, 2.21; Table 2).

Undernutrition in relation to feeding practices among 0–11-month-old children

Among 0–5-month-old children, it was observed that undernutrition prevalence was low among exclusively breast-fed infants up to 4 months of age; at age 5 months, the prevalence was higher among exclusively breast-fed children compared with those receiving complementary feeding in addition to breast milk. Among 6–11-month-old children, the prevalence of underweight and stunting was lower among children who received complementary feeding in addition to breast milk at age 6–8 months compared with those who were exclusively breast-fed or those who received complementary feeding after 8 months.

Association of complementary feeding practices and immunization with undernutrition among 12–23-month-old children

The prevalence of underweight and wasting was observed to be high among children who were exclusively breast-fed (78.6 and 50.0%, respectively) compared with children who received complementary feeding in addition to breast milk (43.7 and 22.0%, respectively). Underweight and stunting prevalences were observed to be high among children who received complementary feeding after age 8 months (45.2 and 58.0%, respectively) compared with children who received complementary feeding at 6–8 months of age (40.8 and 52.9%, respectively). It was also observed that the prevalence of underweight and stunting was higher among partially immunized or not

Table 1 Bivariate analyses of undernutrition according to age, gender and socio-economic status among Indian children aged <3 years, National Nutrition Monitoring Bureau survey, 2011–2012

Characteristic	n	Underweight		Stunting		Wasting	
		OR	95% CI	OR	95% CI	OR	95% CI
Age group (months)							
0–11	1436	1.00	Ref.	1.00	Ref.	1.00	Ref.
12–35	2602	1.91	1.66, 2.19	3.32	2.88, 3.83	0.64	0.55, 0.74
Pooled	4038						
Gender							
Boys	2095	1.28	1.13, 1.46	1.28	1.13, 1.45	1.20	1.03, 1.39
Girls	1943	1.00	Ref.	1.00	Ref.	1.00	Ref.
Community							
Schedule tribe	582	2.89	2.34, 3.58	2.06	1.67, 2.54	2.18	1.70, 2.78
Scheduled caste	1050	1.60	1.33, 1.92	1.48	1.24, 1.77	1.69	1.36, 2.11
Other backward caste	1396	1.36	1.14, 1.62	1.14	0.97, 1.35	1.32	1.07, 1.64
Others	1010	1.00	Ref.	1.00	Ref.	1.00	Ref.
Type of family							
Nuclear	1743	1.57	1.36, 1.82	1.35	1.17, 1.55	1.40	1.19, 1.66
Extended nuclear	750	1.20	1.00, 1.44	1.05	0.88, 1.25	1.04	0.83, 1.29
Joint	1545	1.00	Ref.	1.00	Ref.	1.00	Ref.
Mother's education							
Illiterate	1937	2.17	1.80, 2.55	2.10	1.78, 2.48	1.55	1.26, 1.90
1st–8th grade	1194	1.45	1.20, 1.75	1.37	1.14, 1.65	1.37	1.10, 1.71
9th grade or above	907	1.00	Ref.	1.00	Ref.	1.00	Ref.
Per capita income							
First tertile (< Rs 500)	1278	2.56	2.18, 3.01	2.15	1.84, 2.51	1.89	1.57, 2.28
Second tertile (Rs 500–1014)	1392	1.60	1.36, 1.88	1.28	1.09, 1.49	1.41	1.17, 1.71
Third tertile (> Rs 1014)	1368	1.00	Ref.	1.00	Ref.	1.00	Ref.
Source of drinking-water							
Tap and tube well	3365	1.00	Ref.	1.00	Ref.	1.00	Ref.
Others	673	0.86	0.72, 1.03	0.86	0.72, 1.02	0.84	0.68, 1.03
Sanitary latrine							
Present	1189	1.00	Ref.	1.00	Ref.	1.00	Ref.
Absent	2849	2.38	2.04, 2.77	2.14	1.85, 2.47	1.76	1.47, 2.10
Morbidity							
Present	541	0.92	0.76, 1.11	0.85	0.71, 1.03	1.12	0.90, 1.39
Absent	3497	1.00	Ref.	1.00	Ref.	1.00	Ref.

Ref., reference category.

immunized children (53.0 and 63.0%, respectively) compared with fully immunized children (39.5 and 50.8%, respectively). The underweight and wasting prevalences were significantly higher ($P < 0.001$) among 12–35-month-old children who did not meet minimum dietary diversity (fewer than four food groups in diet; 54.0 and 31.3%, respectively) than among children having minimum dietary diversity (39.7 and 19.8%, respectively). The odds of underweight and stunting were 1.7 and 1.6 times higher among partially/not immunized children compared with children fully immunized against six vaccine-preventable diseases, while underweight (OR = 1.65; 95% CI 1.45, 2.05) and wasting (OR = 1.66; 95% CI 1.29, 2.12) were higher among children not meeting minimum dietary diversity than among children with minimum dietary diversity (Table 2).

Stepwise logistic regression analysis for undernutrition using the conceptual framework

It was observed that the odds of underweight at entry level were 2.29 (95% CI 1.83, 2.86) and 1.40 (95% CI 1.16, 1.70) times higher among children belonging to ST and SC

communities, respectively, compared with children from the others category. The odds of underweight were 2.04 (95% CI 1.71, 2.43) times higher among children belonging to the lowest socio-economic group compared with children from the highest socio-economic group. Children from nuclear families had 1.27 (95% CI 1.09, 1.47) times more risk and children living in *kutcha* houses had 1.63 (95% CI 1.32, 2.02) times more risk of underweight than children from joint families and children living in *pucca* houses, respectively. In the second stage, in addition to the above variables, mother's education, sanitary latrine, hand-washing practices of mothers, IFA consumption during pregnancy, place of birth and birth weight were added. The risk of underweight was 1.48 (95% CI 1.02, 2.14) times higher among children of illiterate mothers compared with children whose mothers were educated to 9th grade or above. The risk of underweight was 1.37 (95% CI 1.00, 1.89) times higher among children from HH not having the facility of a sanitary latrine. Low-birth-weight children had 2.13 times higher risk of underweight (95% CI 1.43, 3.16) compared with normal-birth-weight children. Furthermore, compared with children whose mothers had consumed ninety or more, underweight risk

Table 2 Bivariate analyses of undernutrition among Indian children aged <3 years, National Nutrition Monitoring Bureau survey, 2011–2012

Characteristic	n	Underweight		Stunting		Wasting	
		OR	95% CI	OR	95% CI	OR	95% CI
ANC, delivery and IYCF practices							
For <6-month-old children							
ANC visits							
≥4	321	1.00	Ref.	1.00	Ref.	1.00	Ref.
<4	246	1.56	1.03, 2.35	1.33	0.87, 2.02	1.29	0.87, 1.86
No ANC visits	72	2.90	1.66, 5.07	0.96	0.47, 1.87	3.46	2.02, 5.88
Consumption of IFA tablets							
≥90	220	1.00	Ref.	1.00	Ref.	1.00	Ref.
<90	255	2.44	1.51, 3.94	1.40	0.87, 2.25	1.81	1.17, 2.80
Not consumed	162	2.45	1.45, 4.13	1.50	0.89, 2.54	2.92	1.84, 4.64
Type of feeding							
BF	492	1.00	Ref.	1.00	Ref.	1.00	Ref.
BF + CF	145	0.67	0.41, 1.07	0.80	0.44, 1.45	0.65	0.38, 1.11
For <12-month-old children							
Place of delivery							
Institution	1095	1.00	Ref.	1.00	Ref.	1.00	Ref.
Home	341	1.61	1.24, 2.09	1.41	1.07, 1.85	1.26	0.96, 1.65
Prelacteal feeds given							
Yes	367	1.05	0.81, 1.37	1.04	0.79, 1.37	1.25	0.96, 1.62
No	1069	1.00	Ref.	1.00	Ref.	1.00	Ref.
Time of initiation of BF							
<1 h	523	1.00	Ref.	1.00	Ref.	1.00	Ref.
1–3 h	507	1.24	0.94, 1.63	1.27	0.95, 1.70	0.92	0.70, 1.22
>3 h	406	1.25	0.94, 1.68	1.34	0.99, 1.82	1.01	0.75, 1.35
Colostrum discarded							
Yes	216	1.34	0.98, 1.83	1.15	0.83, 1.60	1.17	0.85, 1.61
No	1220	1.00	Ref.	1.00	Ref.	1.00	Ref.
Birth weight (kg)							
≥2.5	813	1.00	Ref.	1.00	Ref.	1.00	Ref.
<2.5	143	2.22	1.52, 3.26	1.74	1.17, 2.58	1.38	0.92, 2.06
Not recorded	480	2.44	1.90, 3.14	1.66	1.28, 2.16	1.92	1.49, 2.46
At 6–11 months							
Type of feeding							
BF only	136	1.45	1.00, 2.18	1.26	0.85, 1.87	1.69	1.13, 2.51
BF + CF	664	1.00	Ref.	1.00	Ref.	1.00	Ref.
Age at CF initiation (months)							
6–8	491	1.00	Ref.	1.00	Ref.	1.00	Ref.
<6	147	0.78	0.52, 1.18	0.66	0.43, 1.02	0.76	0.48, 1.20
>8	25	1.18	0.51, 2.74	1.32	0.58, 3.10	0.74	0.27, 2.03
Not yet started	136	1.38	0.94, 2.05	1.18	0.78, 1.77	1.58	1.05, 2.37
Minimum dietary diversity							
Yes	245	1.00	Ref.	1.00	Ref.	1.00	Ref.
No	555	1.20	0.87, 1.67	1.56	1.10, 2.21	1.09	0.77, 1.54
Immunization and feeding practices among 12–35-month-old children							
For 12–23-month-old children							
Immunization							
Fully immunized	1338	1.00	Ref.	1.00	Ref.	1.00	Ref.
Partially/not immunized	143	1.74	1.23, 2.46	1.59	1.11, 2.27	1.20	0.80, 1.81
Age at CF initiation (months)							
6–8	1001	1.00	Ref.	1.00	Ref.	1.00	Ref.
<6	295	0.90	0.69, 1.17	0.75	0.57, 0.97	0.86	0.62, 1.20
>8	219	1.19	0.89, 1.61	1.22	0.91, 1.64	1.15	0.81, 1.63
Not yet started	14	5.32	1.47, 19.22	0.88	0.30, 2.55	3.78	1.31, 10.92
Minimum dietary diversity (at 12–35 months)							
Yes	2200	1.00	Ref.	1.00	Ref.	1.00	Ref.
No	401	1.65	1.45, 2.05	1.28	1.03, 1.58	1.66	1.29, 2.12

ANC, antenatal care; IYCF, infant and young child; IFA, iron-folic acid; BF, breast-feeding; CF, complementary feeding; ref., reference category.

was 2.14 and 2.09 times higher among those children whose mothers had consumed fewer than ninety or not consumed IFA tablets during pregnancy, respectively. In the third stage, in addition to the above variables, time of initiation of breast-feeding, age at complementary feeding

initiation and morbidity were added, but they were not observed to be associated with underweight (Table 3).

Similarly, age and gender were associated with stunting. The risk of stunting was 1.78 (95% CI 1.42, 2.23) and 1.44 (95% CI 1.19, 1.74) times higher among children

Table 3 Adjusted logistic regression analysis of factors associated with undernutrition among Indian children aged <3 years, National Nutrition Monitoring Bureau survey, 2011–2012

	Step 1		Step 2		Step 3	
	OR	95 % CI	OR	95 % CI	OR	95 % CI
Underweight using conceptual framework						
Gender (ref. = girls)						
Boys	1.28	1.13, 1.46	1.31	1.14, 1.49	1.42	1.11, 1.82
Age (ref. = 0–11 months)						
12–35 months	1.91	1.66, 2.19	1.96	1.70, 2.26	2.00	1.4, 2.85
Community (ref. = others)						
Scheduled tribe	2.29	1.83, 2.86	1.76	1.16, 2.65	1.88	1.24, 2.84
Scheduled caste	1.40	1.16, 1.70	1.27	0.90, 1.83	1.35	0.94, 1.94
Other backward caste	1.35	1.12, 1.61	1.22	0.86, 1.71	1.22	0.86, 1.73
Per capita income (ref. = third tertile)						
First tertile	2.04	1.71, 2.43	1.48	1.07, 2.08	1.60	1.15, 2.25
Second tertile	1.40	1.18, 1.66	1.05	0.76, 1.43	1.10	0.80, 1.51
Type of family (ref. = joint)						
Nuclear	1.27	1.09, 1.47				
Extended nuclear	1.09	0.90, 1.32				
Type of house (ref. = <i>pucca</i>)						
<i>Kutcha</i>	1.63	1.32, 2.02	1.41	0.94, 2.14		
Semi- <i>pucca</i>	1.53	1.27, 1.83	1.39	1.00, 1.94		
Sanitary latrine (ref. = present)						
Absent			1.37	1.00, 1.89	1.46	1.07, 2.00
Mother's education (ref. = 9th grade or above)						
Illiterate			1.48	1.02, 2.14	1.63	1.14, 2.34
1st–8th grade			1.13	0.76, 1.65	1.22	0.83, 1.79
Birth weight (ref. = ≥ 2.5 kg)						
< 2.5 kg			2.13	1.43, 3.16	2.18	1.46, 3.24
Not measured			1.65	1.22, 2.23	1.69	1.28, 2.25
IFA tablets consumed (ref. = ≥ 90)						
< 90			2.14	1.31, 3.51	2.27	1.38, 3.73
Not consumed			2.09	1.88, 4.46	2.94	1.90, 4.56
Stunting using conceptual framework						
Gender (ref. = girls)						
Boys	1.31	1.15, 1.75	1.40	1.10, 1.80	1.44	1.12, 1.85
Age (ref. = 0–11 months)						
12–35 months	3.48	3.10, 4.02				
Community (ref. = others)						
Scheduled tribe	1.78	1.42, 2.23	1.57	1.02, 2.42	1.54	1.01, 2.36
Scheduled caste	1.44	1.19, 1.74	1.46	1.01, 2.12	1.39	0.96, 2.01
Other backward caste	1.17	0.98, 1.40	1.23	0.86, 1.76	1.20	0.84, 1.72
Per capita income (ref. = third tertile)						
First tertile	1.91	1.61, 2.28	1.48	1.04, 2.10	1.42	1.01, 2.01
Second tertile	1.21	1.03, 1.42	1.29	0.92, 1.79	1.28	0.92, 1.77
Type of house (ref. = <i>pucca</i>)						
<i>Kutcha</i>	1.31	1.08, 1.66				
Semi- <i>pucca</i>	1.44	1.21, 1.71				
Mother's education (ref. = 9th grade or above)						
Illiterate			1.69	1.17, 2.44	1.59	1.11, 2.30
1st–8th grade			1.17	0.79, 1.73	1.13	0.77, 1.67
Sanitary latrine (ref. = present)						
Absent			1.55	1.17, 2.15	1.46	1.06, 2.01
Birth weight (ref. = ≥ 2.5 kg)						
< 2.5 kg			1.64	1.09, 2.47	1.61	1.07, 2.43
Not measured			1.17	0.85, 1.61	1.08	0.81, 1.45
IFA tablets consumed (ref. = ≥ 90)						
< 90			1.28	0.79, 2.09	1.26	0.77, 2.06
Not consumed			2.08	1.38, 3.13	2.13	1.41, 3.22
Wasting using conceptual framework						
Gender (ref. = girls)						
Boys	1.22	1.05, 1.45	1.12	0.88, 1.42	1.09	0.86, 1.38
Age (ref. = 0–11 months)						
12–35 months	0.62	0.53, 0.72				
Community (ref. = others)						
Scheduled tribe	1.79	1.39, 2.31				
Scheduled caste	1.47	1.18, 1.85				
Other backward caste	1.26	1.01, 1.56				

Table 3 *Continued*

	Step 1		Step 2		Step 3	
	OR	95% CI	OR	95% CI	OR	95% CI
Per capita income (ref. = third tertile)						
First tertile	1.52	1.24, 1.85				
Second tertile	1.20	0.98, 1.46				
Type of house (ref. = <i>pucca</i>)						
<i>Kutcha</i>	1.50	1.17, 1.92	1.73	1.16, 2.57	1.90	1.32, 2.75
Semi- <i>pucca</i>	1.29	1.04, 1.59	1.29	0.93, 1.33	1.31	0.97, 1.83
Type of family (ref. = joint)						
Nuclear	1.26	1.06, 1.51				
Extended nuclear	1.00	0.79, 1.24				
Birth weight (ref. = ≥ 2.5 kg)						
< 2.5 kg			1.33	0.88, 2.08	1.33	0.88, 2.04
Not measured			1.58	1.16, 2.14	1.75	1.35, 2.27
IFA tablets consumed (ref. = ≥ 90)						
< 90			1.60	1.03, 2.50	1.75	1.12, 2.71
Not consumed			1.51	1.03, 2.21	1.64	1.11, 2.40

Ref., reference category; IFA, iron–folic acid tablets.

belonging to ST and SC communities, respectively, compared with children from the others category. The odds of stunting were 1.91 (95% CI 1.61, 2.28) times higher among children belonging to the lowest socio-economic group than among those from the highest socio-economic group. Children living in *kutcha* houses had 1.31 (95% CI 1.08, 1.66) more risk of stunting compared with children living in *pucca* houses. In the second stage, community and per capita income, mother's education, sanitary latrine, birth weight and IFA consumption were significantly associated with stunting. The risk of stunting was 1.69 (95% CI 1.17, 2.44) times higher among children of illiterate mothers compared with children whose mothers had studied to 9th grade or above. The risk of stunting was 1.55 (95% CI 1.17, 2.15) times higher among children from HH without a sanitary latrine. Low-birth-weight children had 1.64 times higher risk of stunting (95% CI 1.09, 2.47) than normal-birth-weight children. Mothers not consuming IFA tablets increased the children's risk of stunting by 2.08 (95% CI 1.38, 3.13) times compared with consumption of ninety or more IFA tablets during pregnancy. In the third stage, in addition to the above variables, infant feeding and morbidity were added, but were not observed to be associated with stunting (Table 3).

Gender was observed to be associated with wasting at entry level. In the first stage, the risk of wasting was 1.79 (95% CI 1.39, 2.31) and 1.47 (95% CI 1.18, 1.85) times higher among children belonging to ST and SC communities than among children from the others category. The odds of wasting were 1.52 (95% CI 1.24, 1.85) times higher among children belonging to the lowest socio-economic group compared with those from the highest. Children living in *kutcha* houses and children from nuclear families had 1.50 and 1.26 times more risk of wasting compared with their counterparts in *pucca* houses and joint families. In the second stage, children whose weight was not measured and children whose mothers had not consumed

or consumed fewer than ninety IFA tablets had 1.58, 1.51 and 1.60 times more risk of wasting, respectively. In the third stage, feeding practices and morbidity was added, but none of them was significant (Table 3).

Regression analysis carried out among 12–23-month-old children using variables such as education of mother, immunization, hand-washing practices before feeding, minimum dietary diversity and morbidity during the previous fortnight, keeping age and sex constant, showed that the risk of undernutrition was significantly higher among children of illiterate mothers, those not receiving any immunization or partially immunized and those not meeting minimum dietary diversity (Table 4).

Discussion

During 2011–2012, the NNMB carried out for first the time a study on IYCF practices in ten states of India representing 80% of the population. Optimal nutrition in the first 2 years of life – that is, early and exclusive breast-feeding and continued breast-feeding for 2 years or more, together with nutritionally adequate, safe, age-appropriate and responsive complementary feeding starting at 6 months – is critical to prevent stunting in infancy and early childhood and break the intergenerational cycle of undernutrition. The present study observed suboptimum IYCF practices among the 0–23-month-old children. It was observed that only 36% of infants received breast-feeding within an hour of birth, similar to UNICEF data on this IYCF indicator of 41% for India, 43% for Bangladesh, 50% for Thailand and 38% in Nigeria. About 50% of 6–11-month-old children were exclusively breast-fed up to 6 months of age, similar to the UNICEF findings which showed 57% of children aged 6–9 months in India were exclusively breast-fed up to 6 months, but lower than the findings of 74% for Bangladesh and 75% for Indonesia and Nigeria⁽¹⁶⁾. The National

Table 4 Logistic regression analysis of factors associated with undernutrition among Indian children aged 12–23 months, National Nutrition Monitoring Bureau) survey, 2011–2012

	Underweight		Stunting		Wasting	
	OR	95 % CI	OR	95 % CI	OR	95 % CI
Mother's education (ref. = 9th grade or above)						
Illiterate	2.16	1.61, 2.89	2.22	1.69, 2.91	1.65	1.15, 2.36
1st–8th grade	1.63	1.18, 2.24	1.43	1.06, 1.92	1.49	1.01, 2.20
Minimum dietary diversity (ref. = yes)						
No	1.39	1.07, 1.81	–	–	1.63	1.21, 2.19
Immunization (ref. = fully immunized)						
Partially/not immunized	1.54	1.07, 2.21	1.51	1.05, 2.18	–	–
Hand-washing practices of mother (ref. = washing before feeding)						
Not washing					1.43	1.04, 1.97
Mother is feeding					1.04	0.71, 1.53

Ref., reference category.

Family Health Survey-4 (NFHS-4) also reported similar findings for India, which is 55%⁽¹⁷⁾. Exclusively breast-fed infants are at a lower risk of diseases like diarrhoea and pneumonia. In 2011, UNICEF highlighted that breast-feeding is a preventive intervention and the most important element in reducing child mortality⁽¹⁸⁾.

WHO recommends that infants should start receiving complementary foods at 6 months of age in addition to breast milk. However, in the present study, only 54% of children aged 6–11 months received complementary feeding at 6–7 months of age, which is similar to a previous study⁽¹⁹⁾. NFHS-4 reported only 43% of children aged 6–8 months receiving complementary foods⁽¹⁷⁾.

Undernutrition was observed to be associated with antenatal and perinatal care and infant feeding practices. A study by Pokhrel *et al.*⁽²⁰⁾ in Nepal observed a significant association between antenatal care practices and undernutrition among children. Mother's education was observed to be associated with availing of antenatal care services as observed by Tayie and Lartey⁽²¹⁾ and Meshram *et al.*⁽²²⁾. No significant association was observed between undernutrition and time of breast-feeding initiation, similar to a previous study⁽¹⁹⁾; however, Bahl *et al.* observed significant associations between these practices and increase in morbidity and mortality among them⁽²³⁾. A significant association between low birth weight and undernutrition was observed in the present study, which concurs with our previous studies^(19,22). Low-birth-weight babies are more prone to infections and thus undernutrition. Breast-feeding is the best method of infant feeding to meet the nutritional, metabolic and psychological needs of the baby. Exclusive breast-feeding is a feasible strategy especially in low-income countries as it reduces the risk of infant mortality, morbidity and especially infection. Ogbo *et al.*⁽²⁴⁾ observed that early initiation of breast-feeding was associated with higher maternal education, frequent antenatal care visits and birth interval.

The prevalence of undernutrition observed in the current study is higher except wasting than in the NFHS-4 carried out in India, which reported 32, 32 and 25% for

underweight, stunting and wasting, respectively⁽¹⁷⁾. UNICEF data showed that 35% of under-5s in South Asia were stunted, while this prevalence was 34% in Africa, whereas wasting prevalence was 16% in South Asia and 6–9% in Africa⁽²⁵⁾.

Undernutrition was observed to be significantly higher among children of illiterate mothers, children from SC and ST communities, and children from HH of low socio-economic status. Educated mothers have more efficiency in the management of limited HH resources, greater utilization of health-care services, better health-promoting practices, low fertility and more child-centred caring practices⁽²⁶⁾. SC and ST communities are socio-economically deprived groups; children from these communities have low socio-economic status and thus high rates of undernutrition.

High rates of undernutrition may be due to poor knowledge of mothers about feeding practices, low education and poor socio-economic status, similar to other studies⁽²⁷⁾. There is a significant relationship between improvement in nutritional status of children and optimum infant feeding practices by their mothers. Delayed initiation of breast-feeding, deprivation from colostrum and improper complementary feeding were significant risk factors for undernutrition among under-5s as observed by Kumar *et al.*⁽²⁸⁾. Severe undernutrition was significantly higher in children when weaning was delayed⁽²⁹⁾. In the present study, the prevalence of underweight and stunting was observed to be high among children who received complementary feeding after age 8 months compared with children who received complementary feeding at 6–8 months of age. It was also observed that the prevalence of underweight and stunting was higher among partially immunized or not immunized children than among fully immunized children.

The present study found a significant association between minimum dietary diversity and undernutrition. Bentley *et al.*'s study in Mumbai observed that only 13% of children aged 6–23 months were meeting minimum dietary diversity, while 43% had minimum meal frequency, but no association was observed with nutritional status⁽³⁰⁾;

while Arimond and Ruel observed a significant association of dietary diversity with stunting⁽³¹⁾. In Ghana, Sakaa *et al.* demonstrated that the high percentage of malnutrition in the children may be attributed to the faulty consistency of foods which are traditionally fed to children as well as to less emphasis on iron- and vitamin-rich foods⁽³²⁾. Hence emphasis on providing a variety of food groups to the child, along with optimal timely complementary feeding at 6 months of age, is desirable for achieving the child's growth potential and health.

The main limitation of the present study is that most of the risk factor details were obtained from the mother because there was no other means of obtaining the information; thus there could have been recall bias. However, the outcome was measured prospectively by trained investigators and so maternal recall bias was less likely to have affected the observed associations. The major strengths of the study are its national representation covering more than 80% of the population and data being collected by trained investigators of NNMB who were working in the same field.

Conclusions

The prevalence of undernutrition among Indian children aged <3 years was observed to be high and associated with low socio-economic status, illiteracy of the mother, low birth weight and lack of dietary diversity. Improving mothers' socio-economic and literacy status can help in improving maternal nutrition during pregnancy and thus low birth weight. Also improving mothers' knowledge about IYCF practices will help in improving the nutritional status of children.

Acknowledgements

Acknowledgements: The authors are thankful to the Director of the National Institute of Nutrition, Hyderabad for support during the study. They would like to thank all technical staff of the field division for their technical support. They are grateful to Mr Bhaskar and Dr Naveenkumar for their help in analysing data. The authors extend sincere thanks to the entire field staff from different states for their efforts in the data collection. *Financial support:* This work was supported by the Indian Council of Medical Research, New Delhi. The Indian Council of Medical Research had no role in the design, analysis or writing of this article. *Conflict of interest:* None declared. *Authorship:* All authors were involved in study design, sampling and protocol development; I.I.M. drafted the manuscript and all other authors reviewed it; N.B.K. was involved in data analysis. *Ethics of human subject participation:* This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were

approved by the Institutional Ethical Committee of the National Institute of Nutrition as well as the Scientific Advisory Committee. Written informed consent was obtained from the mothers involved in the study.

References

1. Black RE, Victora CG, Walker SP *et al.* (2013) Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* **382**, 427–451.
2. Black RE, Allen LH, Bhutta ZA *et al.* (2008) Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet* **371**, 243–260.
3. Victora CG, Adair L, Fall C *et al.* (2008) Maternal and child undernutrition: consequences for adult health and human capital. *Lancet* **371**, 340–357.
4. Arifeen S, Black RE, Antelman G *et al.* (2001) Exclusive breastfeeding reduces acute respiratory infection and diarrhea deaths among infants in Dhaka slums. *Pediatrics* **108**, E67.
5. Bahl R, Frost C, Kirkwood BR *et al.* (2005) Infant feeding patterns and risks of death and hospitalization in the first half of infancy: multicentre cohort study. *Bull World Health Organ* **83**, 418–426.
6. Bhutta ZA & Yusuf K (1997) Early-onset neonatal sepsis in Pakistan: a case control study of risk factors in a birth cohort. *Am J Perinatol* **14**, 577–581.
7. Ashraf RN, Jalil F, Zaman S *et al.* (1991) Breastfeeding and protection against neonatal sepsis in a high risk population. *Arch Dis Child* **66**, 488–490.
8. Victora CG, Vaughan JP, Lombardi C *et al.* (1987) Evidence for protection by breast-feeding against infant deaths from infectious diseases in Brazil. *Lancet* **2**, 319–321.
9. Edmond KM, Zandoh C, Quigley MA *et al.* (2006) Delayed breastfeeding initiation increases risk of neonatal mortality. *Pediatrics* **117**, 380–386.
10. United Nations (2015) Transforming Our World: The 2030 Agenda for Sustainable Development. Sustainable Development Knowledge platform. <https://sustainabledevelopment.un.org/post2015/transformingourworld> (accessed August 2015).
11. Office of the Registrar General & Census Commissioner, India, Ministry of Home Affairs, Government of India (2018) 2011 Census data. <http://censusindia.gov.in> (accessed October 2018).
12. Jelliffe DB & Jelliffe EP (1989) *Community Nutritional Assessment*. Oxford: Oxford University Press.
13. Victora CG, Huttly SR, Fuchs SC *et al.* (1997) The role of conceptual framework in epidemiological analysis: a hierarchical approach. *Int J Epidemiol* **26**, 224–227.
14. World Health Organization (1983) *Measuring Changes in Nutritional Status*. Geneva: WHO.
15. World Health Organization (2006) *Child Growth Standards. Length/Height for Age, Weight for Age, Weight for Length, Weight for Height and Body Mass Index for Age. Methods and Development*. Geneva: WHO; available at <http://www.who.int/childgrowth/en>
16. UNICEF (2010) *Progress for Children: Achieving the MDGs with Equity. Number 9, September 2010*. New York: UNICEF.
17. International Institute for Population Sciences & ICF (2017) *National Family Health Survey (NFHS-4), 2015–16: India*. Mumbai: IIPS.
18. UNICEF (2011) *Infant and Young Child Feeding: Programming Guide. Nutrition Section Programmes*. New York: UNICEF.
19. Meshram II, Mallikharjun Rao K, Reddy Gal *et al.* (2015) Influence of feeding practices and associated factors on nutritional status of infants in rural areas of Madhya Pradesh State, India. *Asia Pac J Public Health* **27**, NP1345–NP1361.

20. Pokhrel K, Nanishi K, Poudel KC *et al.* (2016) Under-nutrition among infants and children in Nepal: maternal health services and their roles to prevent it. *Matern Child Health J* **20**, 2037–2049.
21. Tayie F & Lartey A (2008) Antenatal care and pregnancy outcome in Ghana, the importance of women's education. *Afr J Food Agric Nutr Dev* **8**, 291–303.
22. Meshram II, Laxmaiah A, Venkaiah K *et al.* (2012) Impact of feeding and breastfeeding practices on nutritional status of infants in the district of Andhra Pradesh, India. *Natl Med J India* **25**, 201–206.
23. Bahl R, Frost C, Kirkwood BR *et al.* (2005) Infant feeding patterns and risks of death and hospitalization in the first half of infancy: multicentre cohort study. *Bull World Health Organ* **83**, 418–426.
24. Ogbo FA, Agho KE & Page A (2015) Determinants of sub-optimal breastfeeding practices in Nigeria: evidence from the 2008 demographic and health survey. *BMC Public Health* **15**, 259.
25. UNICEF (2018) UNICEF Data: Monitoring the situation of children and women. Malnutrition. <http://data.unicef.org/topic/nutrition/malnutrition/> (accessed September 2018).
26. Bharati S, Pal M, Chakrabarty S *et al.* (2011) Trends in socioeconomic and nutritional status of children younger than 6 years in India. *Asia Pac J Public Health* **23**, 324–340.
27. Mananga MJ, Kana-Sop MM, Nolla NP *et al.* (2014) Feeding practices, food and nutrition insecurity of infants and their mothers in Bangang rural community, Cameroon. *J Nutr Food Sci* **4**, 264.
28. Kumar D, Goel NK, Mittal PC *et al.* (2006) Influence of infant-feeding practices on nutritional status of under-five children. *Indian J Pediatr* **73**, 417–421.
29. Rasanias SK & Sachdev TR (2001) Nutritional status and feeding practices of children attending MCH centre. *Indian J Community Med* **26**, 1–7.
30. Bentley A, Das S, Alcock G *et al.* (2015) *Malnutrition and infant and young child feeding in informal settlements in Mumbai, India: findings from a census.* *Food Sci Nutr* **3**, 257–271.
31. Arimond M & Ruel MT (2004) Dietary diversity is associated with child nutritional status: evidence from 11 demographic and health surveys. *J Nutr* **134**, 2579–2585.
32. Sakaa M, Wemakor A, Abizary AR *et al.* (2015) How well do WHO complementary feeding indicators relate to nutritional status of children aged 6–23 months in rural Northern Ghana? *BMC Public Health* **15**, 1157.