

Food in rural northern Norway in relation to Sami ethnicity: the SAMINOR 2 Clinical Survey

Natalia Petrenya^{1,*}, Guri Skeie¹, Marita Melhus² and Magritt Brustad¹

¹Department of Community Medicine, Faculty of Health Sciences, UiT The Arctic University of Norway, Postboks 6050 Langnes 9037, Tromsø, Norway; ²Centre for Sami Health Research, Department of Community Medicine, UiT The Arctic University of Norway, Tromsø, Norway

Submitted 2 January 2018; Final revision received 25 March 2018; Accepted 24 April 2018; First published online 29 May 2018

Abstract

Objective: To estimate current food intake in the population of northern Norway and to investigate the impact of self-perceived Sami ethnicity and region of residence on food intake.

Design: The data are part of the second cross-sectional survey of the Population-based Study on Health and Living Conditions in Regions with Sami and Norwegian Populations (the SAMINOR 2 Clinical Survey, 2012–2014). Food intake was assessed by an FFQ. Ethnic and regional differences in food intake were studied by sex-specific, multivariable-adjusted quantile regression models.

Setting: Ten municipalities (rural northern Norway).

Subjects: Males (*n* 2054) and females (*n* 2450) aged 40–69 years (2743 non-Sami, 622 multi-ethnic Sami, 1139 Sami).

Results: The diet of Sami participants contained more reindeer meat, moose meat, food made with animal blood and freshwater fish; and contained less lean fish and vegetables. In the inland region, the consumption of reindeer meat was greatest in Sami participants, followed by multi-ethnic Sami participants and non-Sami participants, who had the lowest consumption (median 25, 12 and 8 g/d, respectively). Compared with the inland region, fish roe/liver intake was higher in the coastal region and lean fish intake was twice as high (41 and 32 g/d in males and females, respectively).

Conclusions: When compared with non-Sami participants, those with solely self-perceived Sami ethnicity reported a significantly different intake of several foods, especially reindeer meat in the inland region. Multi-ethnic Sami tended to have similar diets to non-Sami. Residence in the coastal region predicted higher fish and roe/liver intake.

Keywords
Sami
Food intake
Diet
Ethnicity
Northern Norway
Indigenous

Many traditional foods in the North (e.g. reindeer meat and fish) have long been recognized to be rich in nutrients and are thus favourable for health^(1,2). Indeed, traditional local food systems are strongly related to health and well-being in Indigenous Peoples^(3,4). Lifestyle behaviours such as cigarette smoking, lack of physical activity, excessive alcohol intake and dietary changes like reduced intake of traditional local foods and increased intake of processed foods appear to contribute to rapidly growing obesity rates and incidences of CVD, type 2 diabetes, lung and colorectal cancers in certain Indigenous populations in the Arctic^(5–8).

The Sami are an Indigenous People in Norway and represent an ethnic minority in the country⁽⁹⁾. Although the majority of the Sami population can be found in Norway, they also live in Sweden, Finland and Russia.

Historically, the largest part of the Sami population in Norway has resided north of the Arctic Circle, with a smaller population in mid-Norway. In 1970, the number of Sami living in Norway was estimated at about 40 000 individuals, but there is no current official statistic on this number and it could vary according to the definition of ‘Sami’ applied⁽¹⁰⁾.

In the past, and similarly to other Indigenous inhabitants of the circumpolar geographical area, the Sami economy was based on herding and breeding reindeer, fishing, hunting, gathering and some agriculture⁽¹¹⁾. Therefore, primary traditional foods included reindeer (all parts, including meat, blood and organs) and fish, especially oily fish. In coastal communities, fish liver/roe with fresh fish-liver oil were traditionally consumed. A variety of harvested foods (wild fowl and mammals, plants and berries)

have also been important components of the Sami diet, and crucial determinants of the Sami diet included region of residence (coastal *v.* inland (mountain) regions), availability of foods in their natural harvest seasons, ability to purchase foods (e.g. flour, butter/margarine and sugar) and involvement in farming and agriculture^(12–15). Dairy products, cereals, fruit and vegetables were consumed in small amounts, whereas high consumption of boiled, unfiltered coffee has long been recognized as an important component of the Sami culture and diet^(16,17).

Existing studies on the diet of the Sami population in Norway were published between the 1960s⁽¹²⁾ and the 2000s^(17,18). However, these studies were based on small sample sizes and focused mainly on reindeer herders from inland Finnmark County^(17,18). Limited dietary data were collected in the first survey of the Population-based Study on Health and Living Conditions in Regions with Sami and Norwegian Populations (the SAMINOR 1 Survey, 2003–2004)^(19–21) and these data have been used to identify dietary patterns in relation to Sami ethnicity⁽²⁰⁾, including dietary patterns in childhood⁽²¹⁾. An interesting finding was that the differences in dietary patterns were stronger by geographical region than by Sami/Norwegian ethnicity. A childhood diet high in fish was associated with residence in a coastal region, whereas a childhood diet high in reindeer meat and other parts of the reindeer was more common in the inland regions⁽²¹⁾. It was also observed that the associations between ethnicity and dietary patterns were more prominent in inhabitants of the inland than the coastal region⁽²¹⁾.

Interesting, relevant, population-based data from Sweden regarding incidence of and mortality from CVD and

cancer in relation to aspects of the Sami diet and lifestyle have shown that: (i) oily fish was a very important dietary component for the reindeer-herding Sami of southern Lapland in the 1930s to 1950s, and it is still consumed more frequently among people of Sami ethnicity; (ii) historical Sami and present-day reindeer-herding Sami populations have higher intakes of fat, blood and boiled coffee, and lower intakes of cultivated vegetables, bread and fibre, than present-day non-Sami populations; (iii) there was no clear evidence that the studied aspects of the ‘traditional Sami’ diet have beneficial effects on health outcomes in the general northern Swedish population; and (iv) more detailed, updated information on dietary intake and lifestyle among the Sami population is required^(22–26). Therefore, we aimed to estimate current food intake in the Sami compared with the non-Sami population of selected municipalities within northern Norway and to investigate the impact of self-perceived Sami ethnicity and region of residence on food intake.

Methods

Study design and population

The SAMINOR 2 Clinical Survey is a cross-sectional study conducted by the Centre for Sami Health Research, UiT The Arctic University of Norway in 2012–2014. Data collection for the SAMINOR 2 Clinical Survey was carried out in ten municipalities (Fig. 1). All inhabitants aged 40–79 years and residing in these municipalities were invited to participate in the study by personal letter. Our sample considers only the 10 399 invitees aged 40–69 years, of




Fig. 1 Map of study sites

whom 4876 attended the survey (participation rate 47%). In this age group, data were collected through an eight-page self-administered questionnaire, which contained a four-page FFQ (see www.saminor.no for an English translation of the questionnaire), a short clinical examination and analyses of blood samples.

We excluded participants who did not provide information on ethnicity (n 115), as well as immigrants from non-Western European, Asian and African countries (n 69), because the questionnaire may not cover their diet. We further excluded twenty-nine males and sixty-two females because of incomplete FFQ responses (>50% blanks on standardized food frequency intake per week (≥ 57 food items)); two males and five females with missing height and weight measurements; and forty males and fifty females within the top and bottom 1% of the ratio of energy intake (EI) to BMR (EI:BMR)⁽²⁷⁾. Totally, 372 participants were excluded (7.6%), resulting in a final analytical sample of 4504 individuals (2054 males and 2450 females).

Questionnaire data

Ethnicity was based on self-reported information from the SAMINOR 2 Clinical Survey questionnaire, which included the question: 'What do you consider yourself to be?' Response options were 'Norwegian', 'Sami', 'Kven' and 'Other (please describe)'. Participants who chose solely 'Other' (n 142) were further divided into: immigrants from Western European countries and immigrants from non-Western European, Asian and African countries. Ethnicity was then categorized as: (i) non-Sami, including participants who considered themselves as something other than Sami (i.e. Norwegian, Kven or immigrant from Western European countries, n 2856); (ii) multi-ethnic Sami, including participants who defined themselves as Sami in combination with any kind of other ethnic background (n 643); and (iii) Sami, which included participants who defined themselves as Sami only (n 1193).

The SAMINOR 2 Clinical Survey FFQ was a slightly modified version of the FFQ from the Norwegian Women and Cancer (NOWAC) Study^(28,29), which has previously been validated for the general female population of Norway^(29–31). Thus the FFQ covered a wide range of food items commonly consumed in Norway, as well as known traditional food items. Participants reported the frequency with which listed foods and beverages were consumed over the past 12 months. Information on the usual amount consumed was also collected for some food items. We used the NOWAC Study nutrient calculation program to estimate the amounts eaten. Herein we provide consumption information for the following twenty-three broad food groups and individual food items (g/d): vegetables, fruit/berries, potatoes, dairy products, total fish, lean fish, oily fish, fish products, red meat (pork, beef and mutton)/meat products, sauces, chicken, cereal products, breads/crispbreads, fat as spread on bread, total coffee, baked

goods, salty snacks, sweets, freshwater fish (e.g. perch, grayling, pike, arctic char, common whitefish, trout), fish roe/liver, reindeer meat, moose meat and food made with animal blood. Two traditional foods, seagull eggs and tundra grouse (ptarmigan; *Lagopus* spp.), were rarely consumed and thus were not included in the analysis. The food items included in the broad food groups are shown in the online supplementary material, Table S1.

Data on education, physical activity and smoking status were taken from questionnaires. Education was categorized as <13 years and ≥ 13 years; physical activity was reported on a scale from 1 (very low) to 10 (very high) and categorized as low (1–3), moderate (4–7) or high (8–10); and smoking status was categorized as current, former or never.

Clinical examination

Height and weight were measured using an electronic Height, Weight & Fatness Measuring System device (DS-103; Dongsahn Jenix, Seoul, South Korea), with the participants wearing light clothing and no shoes. Height was measured to the nearest 0.1 cm and weight to the nearest 100 g. BMI was then calculated and categorized into two groups: normal/overweight ($< 30 \text{ kg/m}^2$) and obese ($\geq 30 \text{ kg/m}^2$).

Register-based variables

Sex, year of birth and municipality of residence were obtained from the National Registry (Folkeregisteret). Age was defined as that of participants at the end of the year in which clinical examination took place, and was categorized into 40–49, 50–59 and 60–69 years. Geographical region of residence was categorized as the inland region (including the municipalities of Karasjok and Kautokeino) and the coastal region (including the other eight municipalities), based on whether the municipalities include coastal areas or not (Fig. 1).

Ethics approval

The Regional Committee for Medical and Health Research Ethics of Northern Norway (REK-Nord) approved the SAMINOR 2 Clinical Survey. The storing of personal data for the SAMINOR 2 Clinical Survey was approved by the National Data Inspectorate. The application for the present research project was approved by the Regional Committee for Medical and Health Research Ethics of Northern Norway (REK-Nord). All participants signed an informed consent form.

Statistical analysis

We hypothesized that differences in food intake between Sami and non-Sami populations may still exist despite lifestyle changes, and that persons who regard themselves as multi-ethnic Sami (both Sami and Norwegian or other ethnic group) may have a food intake that is more similar to non-Sami than to those who regard themselves as being solely of Sami ethnicity.

All data were analysed separately in males and females, and sample characteristics are presented by ethnic group. Ethnic comparisons were made using Pearson's χ^2 test or the Mann–Whitney test. The intake of twenty-three defined food groups/foods items (g/d) is shown by sex and ethnic group. The distribution of food intake did not meet the assumption of normality. Medians and 25th–75th percentiles are given for descriptive purposes.

Non-parametric multiple quantile regression⁽³²⁾ was used to detect the influence of Sami and multi-ethnic Sami *v.* non-Sami ethnicity (reference group) on food intake, with adjustment for age only and with adjustment for age, education, physical activity level, BMI, smoking and EI (tertiles; MJ). Median regression estimates the median of the dependent variable, conditional on the values of the independent variables, similarly to least-squares regression. Standard errors (95% CI) of quantile regression coefficients were obtained using the bootstrap method (500 bootstrap replications were used). Age-adjusted and fully adjusted β coefficients and *P* values are presented.

The sample was stratified by region (inland and coastal) due to the well-known effect of region of residence on the intake of traditional Sami foods⁽²⁰⁾. Differences in the consumption of lean fish, oily fish, freshwater fish, fish roe/liver and reindeer meat were estimated between ethnic groups in each geographical region. In addition, differences in the intake of total fish, lean fish, oily fish, freshwater fish, fish roe/liver and reindeer meat were estimated for the male and female sub-samples between the inland (reference group) and the coastal population groups. The same regression models were used. Data were analysed using the statistical software package Stata version 14. All tests were two-sided with a 5% significance level.

Results

Characteristics of the study sample

The distribution of ethnic groups (non-Sami 61%, multi-ethnic Sami 14%, Sami 25%) did not differ by sex. Multi-ethnic Sami females were slightly younger (54 years) than their non-Sami and Sami counterparts (57 years; *P* = 0.0002). In males, median age was similar across ethnic groups (57–58 years).

Approximately 40% of males had more than 13 years of education in all three ethnic groups. The proportion of non-Sami *v.* Sami females with more than 13 years of education was 50 and 53%, respectively. However, a higher proportion of multi-ethnic Sami females reported more than 13 years of education (62%). A similar proportion of current smokers was observed in all ethnic groups and for both sexes. Obesity rates (BMI ≥ 30 kg/m²) were somewhat higher among Sami than among multi-ethnic Sami and non-Sami participants (34, 28 and 28% in males and 35, 29 and 25% in females, respectively). The

proportion of Sami males and females with a low level of physical activity was higher when compared with multi-ethnic Sami and non-Sami participants. Moreover, fewer multi-ethnic Sami (16%) and Sami (15%) females reported a high level of physical activity when compared with non-Sami females (21%; Table 1).

The minimum–maximum values of total EI were 1810–29 812 kJ in males (*n* 2096) and 1460–32 951 kJ in females (*n* 2505) before exclusion based on EI:BMR⁽²⁷⁾. The minimum–maximum values of total EI in the final analytical sample were 3067–22 096 kJ (3.1–22.1 MJ) in males and 2615–15 131 kJ (2.6–15.1 MJ) in females. The distribution of ethnic groups differed by tertile of EI in males (*P* < 0.0001) and females (*P* = 0.008), with a higher proportion (41%) of Sami males in the lowest tertile and a higher proportion of multi-ethnic Sami females (40%) in the highest tertile.

The three ethnic groups were not equally distributed across the two geographical regions (Pearson's χ^2 test, *P* < 0.0001). Thus, stratified analyses were done by region. In the inland region, 12% considered themselves non-Sami, 12% multi-ethnic Sami and 76% Sami. Corresponding values for the coastal region were 75, 14 and 11% in non-Sami, multi-ethnic Sami and Sami, respectively.

Percentages of non-consumers of selected foods

We assumed that some of the local food items in the FFQ were consumed to a limited degree. Thus, we recorded the percentage of non-consumers of freshwater fish, fish roe/liver, reindeer meat, moose meat, food made with animal blood, seagull eggs and tundra grouse by ethnic group (Table 2). Ethnic differences in non-consumption were detected in both males and females. We observed a strong gradient for all these local foods, with the fewest non-consumers among Sami, followed by multi-ethnic Sami and non-Sami, except for fish roe/liver, seagull eggs and tundra grouse. The difference in the proportion of non-consumers was particularly clear for reindeer meat. Intakes of seagull eggs and tundra grouse were very low in all ethnic groups; thus these food items were not selected for subsequent regression analyses. Only 7% of Sami males and 5% of Sami females did not consume reindeer meat. However, approximately one-third of the Sami participants did not consume freshwater fish, and half did not consume moose meat or food made with animal blood. Nevertheless, the proportions of non-consumers of these items were considerably higher in the non-Sami group.

Impact of ethnicity on food intake

Sami males consumed less vegetables, potatoes, total fish, lean fish, fish products, chicken, baked goods and salty snacks than their non-Sami counterparts. Conversely, intakes of fat as spread on bread, total coffee, freshwater fish, reindeer meat, moose meat and food made with animal blood were higher in Sami males than non-Sami

males (Table 3). A similar pattern was observed in Sami females (Table 4).

The diet of multi-ethnic Sami males and females was more similar to the diet of participants who considered themselves non-Sami. However, some significant differences were seen: multi-ethnic Sami males ate less dairy products and baked goods, and more oily fish, freshwater fish and reindeer meat, than non-Sami males. Multi-ethnic Sami females ate less lean fish and fish products, and more reindeer meat, than non-Sami females.

Impact of ethnicity on food intake in different geographical regions

Stratified by inland/coastal region and sex, intakes of lean fish, oily fish, freshwater fish, fish roe/liver and reindeer meat were estimated and compared by ethnic group (Table 5). Sami ethnicity had a greater influence on the intake of reindeer meat in the inland region. Participants of Sami ethnicity from the inland region had the highest median intake of reindeer meat (25 g/d in both males and females). The median intake of reindeer meat was lower for multi-ethnic Sami from the inland region (12 g/d in both males and females) and non-Sami (8 g/d in both males and females). In the coastal region, the intake of reindeer meat was 7 g/d in Sami males, 8 g/d in Sami females, 5 g/d in multi-ethnic Sami males and females, 4 g/d in non-Sami males and 2 g/d in non-Sami females.

No differences were found for lean fish and oily fish intakes between non-Sami and multi-ethnic Sami males and females, neither in the inland nor in the coastal region. However, in the coastal region, Sami males consumed more oily fish and more freshwater fish than their non-Sami counterparts, although this was not the case in females. The highest oily fish intake was observed among Sami males in the coastal region (27 g/d) and the highest freshwater fish intake was observed among Sami males from the inland region (6 g/d). Sami males in the inland region and Sami males and females in the coastal region consumed less lean fish when compared with non-Sami from the same regions.

Impact of geographical region on food intake

The consumption of reindeer meat was considerably lower, and the consumption of lean fish was considerably higher, in the coastal region than in the inland region when all ethnic groups were combined (Table 6). Indeed, when the lean fish consumption of all ethnic groups in the coastal and inland regions were compared, males consumed 41 and 17 g/d, and females consumed 32 and 14 g/d, respectively. Oily fish intake was significantly higher in females living in the coastal region, but not in males. Fish roe/liver intake was higher in the coastal region and a higher freshwater fish intake was found in the inland region.

Table 1 Characteristics of the study sample by ethnicity and sex (*n* 4504)* in the SAMINOR 2 Clinical Survey, 2012–2014

Characteristic	Males (<i>n</i> 2054)						P†	Females (<i>n</i> 2450)						P†
	Non-Sami (<i>n</i> 1248)		Multi-ethnic Sami (<i>n</i> 286)		Sami (<i>n</i> 520)			Non-Sami (<i>n</i> 1495)		Multi-ethnic Sami (<i>n</i> 336)		Sami (<i>n</i> 619)		
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%		<i>n</i>	%	<i>n</i>	%	<i>n</i>	%	
Age groups (years)														
40–49	316	25.3	82	28.7	118	22.7	0.367	417	27.9	123	36.6	156	25.2	0.001
50–59	386	30.9	90	31.5	174	33.5		485	32.4	115	34.2	217	35.1	
60–69	546	43.8	114	39.9	228	43.8		593	39.7	98	29.2	246	39.7	
Education (years)														
< 13	708	58.2	157	56.7	300	60.1	0.616	729	50.4	123	37.7	274	46.7	<0.0001
≥ 13	509	41.8	120	43.3	199	39.9		718	49.6	203	62.3	313	53.3	
Geographical region														
Inland	53	4.2	45	15.7	333	64.0	<0.0001	71	4.7	74	22.0	422	68.2	<0.0001
Coastal	1195	95.8	241	84.3	187	36.0		1424	95.3	262	78.0	197	31.8	
Physical activity level														
Low (1–3)	245	19.9	56	19.9	129	25.6	0.018	209	14.5	60	18.1	151	25.3	<0.0001
Moderate (4–7)	821	66.5	177	63.0	295	58.6		935	65.0	217	65.6	356	59.6	
High (8–10)	168	13.6	48	17.1	79	15.7		295	20.5	54	16.3	90	15.1	
BMI category														
Normal/overweight (<30 kg/m ²)	901	72.2	206	72.0	343	66.0	0.027	1117	74.7	238	70.8	405	65.4	<0.0001
Obese (≥30 kg/m ²)	347	27.8	80	28.0	177	34.0		378	25.3	98	29.2	214	34.6	
Smoking status														
Never	436	35.1	109	38.5	174	33.7	0.390	550	37.2	122	36.5	217	35.6	0.918
Former	589	47.4	118	41.7	242	46.8		600	40.6	132	39.5	254	41.7	
Current	217	17.5	56	19.8	101	19.5		328	22.2	80	24.0	138	22.7	
Energy intake (MJ)														
Tertile 1	379	30.4	96	33.6	210	40.4	<0.0001	502	33.6	88	26.2	227	36.7	0.008
Tertile 2	451	36.1	81	28.3	153	29.4		504	33.7	112	33.3	201	32.5	
Tertile 3	418	33.5	109	38.1	157	30.2		489	32.7	136	40.5	191	30.9	

*Subgroups may not total 4504 due to missing values.

†Differences in proportions between non-Sami, multi-ethnic Sami and Sami were tested by Pearson's χ^2 test.

Table 2 Percentages of non-consumers of selected traditional food items by ethnicity and sex (n 4504) in the SAMINOR 2 Clinical Survey, 2012–2014

	Percentage of non-consumers among males (n 2054)							Percentage of non-consumers among females (n 2450)						
	Freshwater fish	Fish roe/liver	Reindeer meat	Moose meat	Food made with animal blood	Seagull eggs	Tundra grouse	Freshwater fish	Fish roe/liver	Reindeer meat	Moose meat	Food made with animal blood	Seagull eggs	Tundra grouse
Non-Sami	60.0	14.3	41.0	66.9	82.0	76.4	94.0	66.8	16.3	46.5	67.8	86.2	89.6	96.7
Multi-ethnic Sami	40.2	12.6	18.2	54.5	66.4	71.3	91.6	48.8	18.8	17.0	55.7	71.7	81.8	96.4
Sami	27.7	15.4	6.5	44.4	47.7	83.5	91.7	35.4	22.1	5.2	49.6	46.0	93.4	96.3
P*	<0.0001	0.55	<0.0001	<0.0001	<0.0001	<0.0001	0.13	<0.0001	0.007	<0.0001	<0.0001	<0.0001	<0.0001	0.91

*Differences in proportions between non-Sami, multi-ethnic Sami and Sami were tested by Pearson's χ^2 test.

Discussion

To the best of our knowledge, the present study is the first large-scale, mixed-sex, population-based study on food intake which covers the extensive geographical area of northern Norway and focuses on ethnicity and the Indigenous Sami population. In the present paper we provide a comprehensive description and comparison of the intakes of main food groups/food items among males and females with non-Sami, multi-ethnic Sami and solely Sami self-perceived ethnicity. The present study shows that food intake among males and females with Sami ethnicity is different from that of non-Sami males and females. In addition, we observed that the diet of those who regarded themselves solely as Sami differed more from that of non-Sami than did the diet of those who perceived themselves as multi-ethnic Sami, especially with respect to intake of reindeer meat in the inland region. Additionally, geographical area of residence was a strong determinant of fish, fish roe/liver and reindeer meat consumption.

Our reference non-Sami population resided in the same rural geographical areas as the Sami and multi-ethnic Sami populations. However, this non-Sami population may differ from those in urban or other rural areas in terms of diet. The studied non-Sami population may have better access to traditional Sami food items and may have a greater acceptance of the dietary habits of Sami culture. Also, individuals who regard themselves as non-Sami, but have Sami ancestors, may have retained parts of that culture, such as certain dietary habits. Nevertheless, we found that the average consumption of reindeer meat, moose meat and food made with animal blood was higher in Sami males and females. As expected, the consumption of reindeer meat was the highest among Sami males (25.0 g/d) and females (25.0 g/d) who live in the inland region (Finnmark County), as reindeer herding takes place primarily in this geographical area. The latest study on daily reindeer consumption among reindeer-herding Sami in Finnmark County was carried out by the Norwegian Radiation Protection Authority (year 2002), which aimed to investigate ^{137}Cs and ^{90}Sr deposition⁽¹⁸⁾. According to that report, the median consumption of reindeer meat (not including blood and organs) was 74 g/d. The difference with the intake of reindeer meat in our present study may be attributable to differences in methodology, sample size and the studied population group. Indeed, both the FFQ and the calculation of reindeer meat intake were different in the present and the aforementioned study. Additionally, our population was not restricted to reindeer-herding Sami. It is likely that Sami participants who are not involved in reindeer herding eat less reindeer meat. It is also possible that the consumption of reindeer meat is decreasing in this Indigenous population.

We found that geographical area of residence was an important factor associated with the intake of reindeer meat. In fact, in the coastal region the median consumption of

Table 3 Food intake (g/d) by non-Sami, multi-ethnic Sami and Sami males (n 2054) in the SAMINOR 2 Clinical Survey, 2012–2014

Food	Entire study sample		Non-Sami males (n 1248)		Multi-ethnic Sami males (n 286)		Age-adjusted: multi-ethnic Sami v. non-Sami males (ref.)		Fully adjusted: multi-ethnic Sami v. non-Sami males (ref.)		Sami males (n 520)		Age-adjusted: Sami v. non-Sami males (ref.)		Fully adjusted: Sami v. non-Sami males (ref.)	
	Median	P25–P75	Median	P25–P75	Median	P25–P75	β^*	<i>P</i> *	β^*,\dagger	<i>P^*,\dagger</i>	Median	P25–P75	β^*	<i>P</i> *	β^*,\dagger	<i>P^*,\dagger</i>
Vegetables	115.4	63.9–184.3	120.5	69.2–192.8	120.3	68.3–190.9	-0.1	0.99	-6.2	0.37	100.8	51.8–158.5	-21.1	0.001	-13.9	0.01
Fruit and berries	162.2	86.5–250.8	159.3	85.2–253.9	166.4	95.9–243.9	8.6	0.30	-3.6	0.69	162.6	88.2–249.7	0.9	0.91	9.1	0.21
Potatoes	110.2	74.3–209.9	112.9	74.3–209.9	110.2	56.2–207.5	-4.5	0.55	-3.3	0.65	94.5	56.2–173.3	-20.2	<0.0001	-12.9	0.006
Dairy products	272.7	128.1–580.3	281.8	133.8–580.0	251.5	100.0–574.9	-29.0	0.17	-57.5	0.03	272.0	126.6–585.8	-15.3	0.47	-3.9	0.84
Total fish	67.2	42.1–104.7	70.9	44.2–108.5	71.4	47.2–111.6	3.5	0.36	3.7	0.38	55.5	33.4–91.5	-15.2	<0.0001	-12.2	<0.0001
Lean fish	33.1	15.8–63.1	40.8	19.5–70.1	37.8	21.2–66.9	-4.6 × 10 ⁻¹⁴	1.0	0.1	0.98	19.0	8.0–39.6	-21.9	<0.0001	-20.0	<0.0001
Oily fish	23.4	12.4–37.6	23.1	11.5–36.6	24.3	15.7–40.1	0.8	0.64	2.8	0.02	23.3	12.2–37.7	0.1	0.94	1.1	0.39
Fish products	29.8	16.5–51.5	33.0	18.1–55.0	29.9	16.6–52.9	-3.1	0.16	0.7	0.76	23.0	11.6–44.7	-10.2	<0.0001	-6.6	<0.0001
Red meat/meat products	72.7	47.7–101.5	74.4	50.1–102.8	77.1	48.1–105.6	0.1	0.99	1.1	0.72	66.0	43.1–93.8	-10.0	<0.0001	-4.1	0.10
Sauces	23.0	12.3–38.3	23.1	12.4–37.9	24.3	12.9–40.0	1.1	0.53	0.9	0.50	21.3	11.1–38.3	-1.4	0.33	0.2	0.82
Chicken	11.1	4.6–18.1	11.1	4.6–20.0	11.1	4.6–20.8	5.8 × 10 ⁻¹⁵	1.0	-1.3	0.22	6.4	0.0–16.2	5.3 × 10 ⁻¹⁵	1.0	-1.9	0.004
Cereal, excl. bread	59.2	32.1–97.2	62.6	33.7–97.7	59.0	30.2–94.4	-0.9	0.84	-2.0	0.61	54.3	27.4–93.5	-8.2	0.019	-0.04	1.0
Bread and crispbread	180.0	102.6–195.1	180.0	104.4–193.3	180.0	100.0–210.6	-7.8 × 10 ⁻¹⁶	1.0	0.5	0.92	175.0	100.0–195.1	-5.0	0.62	0.7	0.80
Fat as spread on bread	18.6	8.6–29.6	18.0	7.9–26.9	20.0	8.6–30.0	2.1	0.19	0.04	0.98	20.0	9.3–30.0	2.3	0.15	3.2	0.001
Total coffee	945.0	630.0–1383.5	945.0	580.5–1365.0	945.0	630.0–1365.0	9.6 × 10 ⁻¹⁴	1.0	1.1 × 10 ⁻¹⁴	1.0	1365.0	945.0–1680.0	315	0.004	210	<0.0001
Baked goods	35.7	21.0–51.1	36.3	21.4–51.6	34.4	20.0–51.5	-3.3	0.017	-4.3	0.01	33.0	17.3–45.8	-3.3	0.010	-3.6	0.01
Salty snacks	5.0	2.0–10.0	6.7	2.0–10.7	5.0	2.0–10.7	4.5 × 10 ⁻¹⁵	1.0	-0.3	0.62	4.0	0.0–9.1	-2.7	<0.0001	-1.9	<0.0001
Sweets	30.1	15.3–54.6	32.0	15.8–55.8	27.7	15.4–53.2	-4.3	0.014	-3.3	0.11	27.2	13.7–51.3	-5.4	0.001	-2.3	0.15
Traditional food items																
Freshwater fish	1.1	0.0–6.3	0.0	0.0–4.0	2.3	0.0–7.8	2.3	<0.0001	1.5	0.01	4.7	0.0–13.5	4.7	<0.0001	4.7	<0.0001
Fish roe/liver	1.0	0.8–2.4	1.0	0.8–2.4	1.0	0.8–2.8	-3.0 × 10 ⁻¹⁶	1.0	0.07	0.28	0.9	0.6–2.4	-0.1	0.06	-0.05	0.39
Reindeer meat	5.3	0.0–12.3	3.5	0.0–7.0	7.0	3.5–12.3	3.5	<0.0001	3.5	<0.0001	16.4	7.0–37.5	11.1	<0.0001	11.1	<0.0001
Moose meat	0.0	0.0–5.3	0.0	0.0–5.3	0.0	0.0–7.0	4.0 × 10 ⁻¹⁵	1.0	5.4 × 10 ⁻¹⁵	1.0	3.5	0.0–8.8	3.5	<0.0001	3.5	<0.0001
Food made with animal blood	0.0	0.0–5.0	0.0	0.0–0.0	0.0	0.0–5.0	2.2 × 10 ⁻¹⁴	0.035	-2.3 × 10 ⁻¹⁵	0.75	5.0	0.0–5.0	5.0	<0.0001	5.0	<0.0001

P25, 25th percentile; P75, 75th percentile; ref., reference category.

* β and *P* value from quantile regression (at the median), adjusted for age (40–49, 50–59, 60–69 years), education (<13, ≥13 years), physical activity level (three groups), BMI (<30, ≥30 kg/m²), smoking status (current, former, never) and energy intake (tertiles; MJ).

†In bold font if statistically significant in the fully adjusted model.

reindeer meat was approximately 20 g/d lower when compared with the inland region.

In the small study by Nilsen *et al.* from 1999, dietary history was obtained during interviews with Sami living in Finnmark County (municipalities of Kautokeino, Karasjok and Tana) and with Norwegians living in the small city of Alta⁽¹⁷⁾. When compared with Norwegians, Sami (mixed-sex sample, adjusted for age and sex) received a higher proportion of their energy from total red meat (reindeer meat was included in calculation) and reindeer meat as estimated separately. Intakes of oily fish, fat added to food (mainly margarine on bread), coffee, sugar with coffee and dairy butter were also higher among Sami. The Norwegians received more energy from lean fish, fish products, dairy products, pork, beef, vegetables and fruits, and total fish intake was not different between Sami and Norwegians⁽¹⁷⁾. To some extent, our findings were consistent with these previous results. We found that Sami males and females consumed less vegetables, lean fish and fish products; and more fat as spread on bread, coffee, freshwater fish, reindeer meat, moose meat and food made with animal blood. We also found that Sami males and females consumed less total fish, chicken, baked goods and salty snacks than their non-Sami counterparts. Additionally, Sami females ate less oily fish and red meat (pork, beef and mutton)/meat products, and Sami males consumed fewer potatoes than their non-Sami counterparts. Interestingly, red meat estimated as the consumption of pork, beef, mutton and meat products did not differ between Sami and non-Sami males. Compared with Nilsen *et al.*'s study, we did not find differences in the consumption of dairy and fruits for both sexes. It is possible that the consumption of these products has increased among Sami.

Food and food traditions are often described as important carriers, markers and tools for cultural identity and belonging. Food provides nutrients, but what we eat and our food traditions also nourish cultural and ethnic identity and belonging. Our results indicate that having self-perceived Sami ethnicity implies stronger ties to Sami reindeer-herding food traditions in the inland region, which is consistent with the previous results from dietary studies among Sami adolescents⁽³³⁾ and Sami adults⁽²⁰⁾ living in northern Norway. Reindeer meat, moose meat, food made with animal blood and freshwater fish are traditional foods that remain important in Sami culture. Higher intakes of traditional meats and food made with animal blood may contribute to a higher concentration of nutrients like Fe^(19,33) and vitamin B₁₂ among the Sami in northern Norway.

In the present paper, we applied region-stratified statistical analysis, because previous studies have shown that diet may have stronger relationship with geographical area than with ethnicity⁽²⁰⁾. As expected, our results showed that the coastal population ate more total fish, lean fish and fish liver/roe; and that the inland population

ate more reindeer meat and local freshwater fish. Comparison of oily fish intake between the coastal region and the inland region did not provide a clear result. When we considered the coastal regions in Finnmark County alone (municipalities of Porsanger, Tana and Nesseby), the oily fish intake was higher by approximately 4 g/d among males and females (data not shown). However, when the coastal region included Finnmark, Troms and Nordland counties, the oily fish intake was significantly higher in females, but not in males. We found that the inland population eats more freshwater fish. However, the consumption of freshwater fish was generally low and did not contribute much to the total fish intake of Sami living in the inland region. In our study, the average freshwater fish consumption among Sami males and females living in the inland regions was considerably lower (6.0 g/d in males and 3.9 g/d in females) than that previously reported among reindeer-herding Sami males and females (18 g/d)⁽¹⁸⁾.

Lifestyle changes within the last century have brought about a rapid transition in nutrition, characterized by a decreasing consumption of traditional foods and an associated increase in the consumption of processed, shop-bought foods. These changes have also been observed in non-Indigenous populations, but the negative effects are suggested to affect the Indigenous populations to a larger degree^(8,34). In fact, a high prevalence of obesity and metabolic syndrome was found in the SAMINOR 1 Survey in both the Sami and the non-Sami populations of rural northern Norway^(35,36). The prevalence of obesity was higher in Sami females⁽³⁵⁾ and the prevalence of metabolic syndrome was higher in Sami females younger than 50 years than their non-Sami counterparts⁽³⁶⁾. However, there was an overall lower prevalence of metabolic syndrome in Sami males compared with non-Sami males⁽³⁶⁾. Even though Sami women were more obese, no differences in the incidence of diabetes mellitus have been found between Sami and Norwegian populations based on 14 years of follow-up (in the period 1974–1989)⁽³⁷⁾. Similarly, no ethnic differences in the prevalence of diabetes between Sami and non-Sami populations have been reported in the SAMINOR 1 Survey (2003–2004)⁽³⁶⁾. Overall, age-standardized prevalence of diabetes mellitus in Sami males and females was 5.5 and 4.8%, respectively⁽³⁸⁾. When data from the SAMINOR 1 Survey were stratified by region, Sami males living in the inland region and Sami females living in Porsanger, Tana and Nesseby municipalities had lower prevalence of diabetes mellitus (2.8 and 2.4%, respectively) compared with their non-Sami counterparts (8.6 and 4.7%, respectively). However, the opposite result was seen for both Sami males (9.3%) and females (7.7%) living in the southern-most region, where Sami participants constituted the minority⁽³⁸⁾. Indeed, dietary patterns were associated with insulin resistance and cardiometabolic risk in other Indigenous populations^(39,40). The diet and lifestyle factors in relation

Table 4 Food intake (g/d) in non-Sami, multi-ethnic Sami and Sami females (n 2450) in the SAMINOR 2 Clinical Survey, 2012–2014

Food	Entire study sample		Non-Sami females (n 1495)		Multi-ethnic Sami females (n 336)		Age-adjusted: multi-ethnic Sami v. non-Sami females (ref.)		Fully adjusted: multi-ethnic Sami v. non-Sami females (ref.)		Sami females (n 619)		Age-adjusted: Sami v. non-Sami females (ref.)		Fully adjusted: Sami v. non-Sami females (ref.)	
	Median	P25–P75	Median	P25–P75	Median	P25–P75	β^*	<i>P</i> *	β^*, \dagger	<i>P^*, \dagger</i>	Median	P25–P75	β^*	<i>P</i> *	β^*, \dagger	<i>P^*, \dagger</i>
Vegetables	166.4	104.2–249.0	174.7	112.2–258.0	173.5	113.4–250.0	–1.4	0.87	–6.8	0.40	147.1	81.6–229.1	–27.4	<0.0001	–34.3	<0.0001
Fruit and berries	190.3	117.0–288.4	191.3	122.8–291.7	191.8	113.5–283.9	–0.5	0.96	0.7	0.94	186.3	101.8–282.7	–2.5	0.78	1.4	0.85
Potatoes	82.1	56.2–102.3	88.9	56.2–102.3	74.3	56.2–94.5	–7.6 × 10 ^{–15}	1.0	–5.6	0.09	74.3	56.2–94.5	4.1 × 10 ^{–15}	1.0	–2.6	0.33
Dairy products	206.9	96.4–340.3	210.7	96.3–331.6	221.6	98.5–420.3	15.0	0.27	9.5	0.48	187.0	96.5–329.1	–20.0	0.08	0.8	0.93
Total fish	55.8	32.0–84.8	58.1	34.1–87.2	58.4	34.3–83.0	–0.6	0.83	–0.1	0.97	47.2	25.6–79.5	–9.8	<0.0001	–8.4	0.001
Lean fish	27.2	11.8–48.9	31.5	15.2–54.3	27.6	11.9–47.3	–3.8	0.03	–4.7	0.003	17.0	6.3–37.8	–13.8	<0.0001	–11.8	<0.0001
Oily fish	19.9	9.7–32.8	20.3	9.5–32.9	22.0	12.7–35.8	2.8	0.07	2.1	0.11	16.7	8.4–29.4	–2.5	0.003	–2.3	0.03
Fish products	29.0	15.3–47.4	32.1	17.4–49.2	28.3	15.8–47.4	–4.2	0.04	–5.4	0.001	22.4	9.9–41.5	–9.2	<0.0001	–8.5	<0.0001
Red meat/meat products	50.4	33.2–71.0	51.1	34.8–72.1	54.0	35.8–74.5	1.4	0.52	0.4	0.82	45.7	28.7–67.2	–6.2	<0.0001	–5.5	0.001
Sauces	17.8	9.9–30.2	17.4	9.9–30.2	21.1	12.3–32.5	3.01	0.03	1.5	0.13	17.2	8.8–29.3	–0.1	0.89	–0.7	0.39
Chicken	11.6	4.6–20.0	11.6	6.4–20.0	11.6	6.4–26.5	4.5 × 10 ^{–15}	1.0	–0.04	0.96	11.1	4.6–18.1	–1.9	0.22	–1.9	0.004
Cereal, excl. bread	70.8	41.1–108.6	70.0	40.2–106.4	74.6	41.7–122.7	3.1	0.42	3.3	0.42	68.6	41.7–105.7	2.1	0.48	3.4	0.21
Bread and crispbread	110.5	69.4–180.0	110.5	64.9–180.0	121.9	77.0–184.4	12.1	0.07	11.9	0.05	110.5	65.8–180.0	–7.8 × 10 ^{–15}	1.0	5.8	0.25
Fat as spread on bread	14.3	4.3–24.3	13.5	3.8–22.5	14.6	4.3–24.3	1.8	0.002	1.0	0.32	14.6	6.8–25.0	1.8	0.002	2.2	0.01
Total coffee	945.0	525.0–1365.0	945.0	525.0–1155.0	945.0	525.0–1365.0	0	1.0	81.7	0.12	945.0	525.0–1365.0	0	1.0	163.3	<0.0001
Baked goods	32.9	18.2–44.8	34.8	20.6–47.4	32.1	14.9–44.1	–2.8	0.12	–3.3	0.05	30.1	14.7–41.9	–3.2	0.01	–3.4	0.004
Salty snacks	5.0	2.0–10.7	6.7	2.0–12.0	7.0	2.7–12.4	–1.4 × 10 ^{–15}	1.0	–0.3	0.57	4.3	0.0–9.0	–2.7	<0.0001	–2.4	<0.0001
Sweets	24.0	11.4–42.1	23.4	11.2–41.8	24.5	12.9–41.7	1.2	0.56	–0.2	0.88	24.4	11.1–44.0	0.9	0.50	0.6	0.68
Traditional food items																
Freshwater fish	0.0	0.0–4.0	0.0	0.0–2.0	0.9	0.0–4.7	1.0	0.07	0.95	0.08	3.0	0.0–9.0	3.0	<0.0001	3.0	<0.0001
Fish roe/liver	0.9	0.6–1.7	0.9	0.6–1.7	0.9	0.6–1.8	–2.2 × 10 ^{–16}	1.0	0.012	0.65	0.9	0.2–1.1	–1.4 × 10 ^{–16}	1.0	–0.02	0.32
Reindeer meat	4.1	0.0–12.3	1.8	0.0–5.3	5.3	3.5–12.3	3.5	<0.0001	3.5	<0.0001	16.4	8.2–37.5	12.9	0.001	12.2	<0.0001
Moose meat	0.0	0.0–5.3	0.0	0.0–3.5	0.0	0.0–5.3	–1.7 × 10 ^{–16}	1.0	1.0 × 10 ^{–17}	1.0	1.8	0.0–7.0	1.8	<0.0001	1.8	<0.0001
Food made with animal blood	0.0	0.0–5.0	0.0	0.0–0.0	0.0	0.0–5.0	–1.8 × 10 ^{–14}	0.24	1.95 × 10 ^{–15}	0.86	5.0	0.0–5.0	5.0	<0.0001	5.0	<0.0001

P25, 25th percentile; P75, 75th percentile; ref., reference category.

* β and *P* value from quantile regression (at the median), adjusted for age (40–49, 50–59, 60–69 years), education (<13, ≥13 years), physical activity level (three groups), BMI (<30, ≥30 kg/m²), smoking status (current, former, never) and energy intake (tertiles; MJ).

†In bold font if statistically significant in the fully adjusted model.

Table 5 Selected food intake (g/d) in non-Sami, multi-ethnic Sami and Sami participants, stratified by inland/coastal region and sex (n 4504), in the SAMINOR 2 Clinical Survey, 2012–2014

	Males (n 2054)										Females (n 2450)									
	Lean fish		Oily fish		Freshwater fish		Fish roe/liver		Reindeer meat		Lean fish		Oily fish		Freshwater fish		Fish roe/liver		Reindeer meat	
	Median	P25–P75	Median	P25–P75	Median	P25–P75	Median	P25–P75	Median	P25–P75	Median	P25–P75	Median	P25–P75	Median	P25–P75	Median	P25–P75	Median	P25–P75
Inland region																				
Non-Sami	26.2	11.8–41.4	20.3	14.9–38.4	4.0	0.0–9.4	0.9	0.6–1.1	8.2	5.3–12.3	18.8	7.6–34.5	23.9	12.4–35.9	1.1	0.0–6.0	0.8	0.2–1.3	8.2	3.5–12.5
Multi-ethnic Sami	21.7	11.0–37.8	20.9	14.7–33.8	4.0	0.0–12.0	0.9	0.0–1.7	12.5	8.2–50.0	14.1	6.3–30.4	21.2	10.7–32.2	1.5	0.0–7.9	0.7	0.0–0.9	12.3	5.3–35.0
Sami	15.2	6.3–31.5	20.8	10.7–34.0	6.0	1.1–15.8	0.9	0.3–1.9	25.0	12.3–50.0	13.9	4.8–31.5	16.2	7.4–27.3	3.9	0.0–10.2	0.8	0.0–1.0	25.0	12.3–37.5
Multi-ethnic Sami v. non-Sami (ref.)																				
β^*, \dagger	0.4		6.0		1.4		0.04		9.8		0.5		0.2		0.5		–0.2		1.3	
P^*, \dagger	0.93		0.12		0.57		0.82		0.08		0.90		0.97		0.67		0.20		0.67	
Sami v. non-Sami (ref.)																				
β^*, \dagger	–9.2		3.9		1.6		0.04		15.2		–2.8		–4.6		1.9		2.9×10^{-16}		16.8	
P^*, \dagger	0.01		0.13		0.26		0.67		< 0.0001		0.41		0.26		0.05		1.0		< 0.0001	
Coastal region																				
Non-Sami	41.9	20.5–70.1	23.3	11.5–36.6	0.0	0.0–4.0	1.0	0.8–2.4	3.5	0.0–7.0	32.0	15.8–54.3	20.1	9.3–32.9	0.0	0.0–1.9	0.9	0.6–1.7	1.8	0.0–5.3
Multi-ethnic Sami	42.8	23.6–70.1	24.6	16.0–41.3	1.5	0.0–6.3	1.0	0.9–2.8	5.3	3.5–12.3	31.5	15.8–49.6	22.2	13.9–36.5	0.0	0.0–4.0	0.9	0.7–2.1	5.3	1.8–8.2
Sami	31.5	15.2–60.6	26.6	17.4–40.5	3.9	0.0–11.8	1.0	0.8–2.4	7.0	3.5–16.4	30.3	13.0–53.5	20.4	11.5–35.7	1.1	0.0–6.0	0.9	0.7–2.0	8.2	3.5–12.5
Multi-ethnic Sami v. non-Sami (ref.)																				
β^*, \dagger	1.7		2.4		1.5		0.1		3.0		–2.4		1.4		-2.3×10^{-16}		0.1		3.5	
P^*, \dagger	0.63		0.07		0.004		0.22		< 0.0001		0.23		0.39		1.0		0.12		< 0.0001	
Sami v. non-Sami (ref.)																				
β^*, \dagger	–12.3		4.8		3.9		0.1		4.7		–6.2		1.3		1.1		0.02		5.3	
P^*, \dagger	< 0.0001		0.02		< 0.0001		0.44		< 0.0001		0.03		0.42		0.09		0.57		< 0.0001	

P25, 25th percentile; P75, 75th percentile; ref., reference category.

* β and P value from quantile regression (at the median), adjusted for age (40–49, 50–59, 60–69 years), education (<13, \geq 13 years), physical activity level (three groups), BMI (<30, \geq 30 kg/m²), smoking status (current, former, never) and energy intake (tertiles; MJ).

†In bold font if statistically significant in the fully adjusted model.

Table 6 Selected food intake (g/d) in the inland v. coastal region, stratified by sex (n 4504), in the SAMINOR 2 Clinical Survey, 2012–2014

	Males (n 2054)						Females (n 2450)					
	Inland region (n 431)		Coastal region (n 1623)		Coastal v. inland region (ref.)		Inland region (n 567)		Coastal region (n 1883)		Coastal v. inland region (ref.)	
	Median	P25–P75	Median	P25–P75	β^*, \dagger	P^*, \dagger	Median	P25–P75	Median	P25–P75	β^*, \dagger	P^*, \dagger
Total fish	51.5	31.6–82.8	71.9	45.8–110.1	15.4	< 0.0001	42.9	24.2–70.3	59.5	35.8–88.5	13.6	< 0.0001
Lean fish	16.7	6.3–33.1	40.8	20.3–70.1	20.8	< 0.0001	14.1	5.1–31.5	31.5	15.8–54.3	12.9	< 0.0001
Oily fish	20.8	11.3–34.4	24.1	12.5–38.0	1.9	0.07	16.7	8.4–29.3	20.4	10.3–33.5	3.1	0.002
Fresh water fish	5.1	1.1–13.5	0.0	0.0–5.0	-5.0	< 0.0001	3.1	0.0–10.1	0.0	0.0–3.0	-3.1	< 0.0001
Fish roe/liver	0.9	0.3–1.9	1.0	0.8–2.4	0.2	0.004	0.8	0.0–1.0	0.9	0.6–1.8	0.1	0.04
Reindeer meat	25.0	8.2–37.5	4.1	0.0–7.0	-18.5	< 0.0001	25.0	8.2–37.5	3.5	0.0–5.3	-20.3	< 0.0001

P25, 25th percentile; P75, 75th percentile; ref., reference category.

* β and P value from quantile regression (at the median), adjusted for age (40–49, 50–59, 60–69 years), education (<13, \geq 13 years), physical activity level (three groups), BMI (<30, \geq 30 kg/m²), smoking status (current, former, never) and energy intake (tertiles; MJ).

†In bold font if statistically significant.

to prevalence of dietary-related chronic diseases in the Sami population need to be investigated in more detail to explain the observed heterogeneity across sexes and geographical regions in this population.

Limitations

We detected lower EI in males with Sami ethnicity than in males with non-Sami and multi-ethnic Sami ethnicity. In females, the highest EI was observed in multi-ethnic Sami. A limitation of the present study was that the questionnaire has not been specifically validated in males or in the Sami population. Thus, it is possible that the EI in Sami males was underestimated, because the FFQ may not be well-suited for Sami males. A more detailed FFQ in terms of traditional Sami foods and dishes may be needed to better estimate the current food intake in Sami people. For instance, the traditional Sami diet may contain dried reindeer meat that is not necessarily consumed at regular meals but eaten in between meals. There are many different combinations of a traditional, reindeer-based Sami meal. Indeed, in addition to meat, meals may consist of reindeer liver, bone marrow, tongue, tallow and other reindeer parts. Traditional Sami cuisine includes reindeer broth and soup/stew. New recipes, which are a combination of the old traditional diet and the modern diet, are emerging. However, it was not feasible to include related questions, as it would have resulted in an FFQ that was too long and detailed for the present, large population-based study. Thus, adjustments were made to the NOWAC FFQ, mainly through the addition of some known traditional Sami food items, including freshwater fish, reindeer meat, moose meat and food made with animal blood. These food items are most often consumed by the Sami population and contributed considerably to nutrient intakes.

The identification of ethnicity in multi-ethnic population-based studies is challenging⁽⁴¹⁾. A harsh assimilation policy made many Sami abandon their culture and language, and in certain regions, people of Sami origin are likely to say that they regard themselves as Norwegian instead of Sami. There are no standardized and validated methods to determine ethnicity, but there are two main approaches. The first one includes objective measures, based on ancestry or a connection to Sami language, and the second one is based on self-perception. Self-perception as a determinant of ethnicity is widely used in international research⁽⁴²⁾ and was used in the current project. We hypothesized that self-perceived ethnicity may be strongly associated with cultural practices like dietary habits. Therefore, our results may not be applicable to Sami populations whose ethnicity is determined using a different approach.

One more limitation can be attributed to the reduced generalizability and external validity of the results because the response rate was only 47%, the study sample was limited to individuals aged 40–69 years and only ten municipalities were included. The response rate among

younger males in the SAMINOR 2 Clinical Survey was lower. In addition, the response rate among some categories of participants, namely individuals with fewer health concerns and lower education level, may be lower.

Self-reported data on dietary intake and lifestyle factors are known to be subject to recall biases in population-based studies. We do not have information on the tendency to report healthier lifestyle among the solely Sami, multi-ethnic Sami and non-Sami ethnic groups, and it might be interesting to develop this knowledge.

Strengths

The present study is a follow-up of the SAMINOR 1 Survey, which included only a limited number of dietary questions. The SAMINOR 2 Clinical Survey questionnaire is much more comprehensive and gives a better assessment of total diet. Other strengths of the present study include its large sample size, detailed information on self-perceived ethnicity, and recruitment of participants from a large geographical area that covered both the inland and coastal regions of northern Norway. We adjusted for important covariates related to food intake that were different between the ethnic groups studied.

Conclusion

Food intake in rural areas of northern Norway is related to ethnicity and geographical region. Individuals with solely self-perceived Sami ethnicity differ more from non-Sami participants than participants with a self-perceived multi-ethnic Sami ethnicity, especially with respect to the intake of reindeer meat in the inland region. Our results contribute to the knowledge on current food intake in the rural population of northern Norway according to self-perceived ethnicity.

Acknowledgements

Acknowledgements: The authors gratefully acknowledge the participants and staff of the SAMINOR 2 Clinical Survey, and the Centre for Sami Health Research, UiT The Arctic University of Norway, for access to the data. *Financial support:* The SAMINOR 2 Clinical Survey was financed by the Norwegian Ministry of Health and Care Services; the Northern Norway Regional Health Authority; the Regional Research Fund of Northern Norway; the Sami Parliament; the Sami Norwegian National Advisory Unit on Mental Health and Substance Use; and the Finnmark, Troms and Nordland County Councils. The present post-doctoral research project was financially supported by the Department of Community Medicine, UiT The Arctic University of Norway. The funders had no role in the design, analysis or writing of this study. *Conflict of interest:* The authors declare that they have no conflict of interest. *Authorship:* N.P. analysed the data and wrote the manuscript.

G.S. calculated food intake and revised the paper. M.M. was responsible for the database, helped with statistical analyses, designed the map of the study and revised the paper. M.B. designed the study and revised the paper. All authors approved the final version of manuscript. *Ethics of human subject participation:* The Regional Committee for Medical and Health Research Ethics of Northern Norway (REK-Nord) approved the SAMINOR 2 Clinical Survey. The storing of personal data for the SAMINOR 2 Clinical Survey was approved by the National Data Inspectorate. The application for this research project was approved by the Regional Committee for Medical and Health Research Ethics of Northern Norway (REK-Nord). All participants signed an informed consent form.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1368980018001374>

References

1. Kuhnlein HV, Chan HM, Leggee D *et al.* (2002) Macro-nutrient, mineral and fatty acid composition of Canadian arctic traditional food. *J Food Compos Anal* **15**, 545–566.
2. Hassan AA, Sandanger TM & Brustad M (2012) Selected vitamins and essential elements in meat from semi-domesticated reindeer (*Rangifer tarandus tarandus* L.) in mid- and northern Norway: geographical variations and effect of animal population density. *Nutrients* **4**, 724–739.
3. Kuhnlein H, Erasmus B, Creed-Kanashiro H *et al.* (2006) Indigenous peoples' food systems for health: finding interventions that work. *Public Health Nutr* **9**, 1013–1019.
4. Kuhnlein HV (2015) Food system sustainability for health and well-being of Indigenous Peoples. *Public Health Nutr* **18**, 2415–2424.
5. McLaughlin JB, Middaugh JP, Utermohle CJ *et al.* (2004) Changing patterns of risk factors and mortality for coronary heart disease among Alaska Natives, 1979–2002. *JAMA* **291**, 2545–2546.
6. Friborg JT & Melbye M (2008) Cancer patterns in Inuit populations. *Lancet Oncol* **9**, 892–900.
7. Jernigan VB, Duran B, Ahn D *et al.* (2010) Changing patterns in health behaviors and risk factors related to cardiovascular disease among American Indians and Alaska Natives. *Am J Public Health* **100**, 677–683.
8. Kuhnlein HV, Receveur O, Soueida R *et al.* (2004) Arctic indigenous peoples experience the nutrition transition with changing dietary patterns and obesity. *J Nutr* **134**, 1447–1453.
9. International Labour Office (2003) ILO Convention on Indigenous and Tribal Peoples (1989) (No. 169): A Manual. http://www.ilo.org/wcmsp5/groups/public/@ed_norm/@normes/documents/instructionalmaterial/wcms_088485.pdf (accessed February 2016).
10. Aubert V (1978) *Den samiske befolkningen i Nord-Norge*. vol. 107: *Såmi ål'bmüt Davvi-Norgas (The Lappish Population in Northern Norway)*. vol. 107: *Såmi ål'bmüt Davvi-Norgas*. Oslo: Statistisk sentralbyrå.
11. Solbakk JT & Biti BH (2006) *Samene: en handbok (The Sami People: A Handbook)*. Karasjok: Davvi Girji AS.
12. Solvang A & Øgrim ME (1967) *Kostholdsundersøkelser i Kautokeimo og Karasjok (Dietary Surveys in Kautokeimo*

- and Karasjok). Oslo: Landsforeningen for kosthold og helse.
13. Bardølsgaard K & Øgrim ME (1963) *Kostholdsundersøkelser i Havøysund (Dietary Surveys in Havøysund)*. Oslo: Landsforeningen for kosthold og helse.
 14. Solvang A & Øgrim ME (1963) *Kostholdsundersøkelser i Gamvik herred (Dietary Surveys in Gamvik)*. Oslo: Landsforeningen for kosthold og helse.
 15. Sørum HM & Øgrim ME (1962) *Kostholdsundersøkelser i Polmak (Dietary Surveys in Polmak)*. Oslo: Landsforeningen for kosthold og helse.
 16. Nystad T, Melhus M, Brustad M *et al.* (2010) The effect of coffee consumption on serum total cholesterol in the Sami and Norwegian populations. *Public Health Nutr* **13**, 1818–1825.
 17. Nilssen H, Utsi E & Bønaa KH (1999) Dietary and nutrient intake of a Sami population living in traditional reindeer herding areas in north Norway: comparisons with a group of Norwegians. *Int J Circumpolar Health* **58**, 120–133.
 18. Thørring H, Hosseini A & Skuterud L (2004) *Kostholdsundersøkelser 1999 og 2002: reindrifstøvere i Kautokeino (Dietary Surveys in 1999 and 2002: Reindeer Herders in Kautokeino)*. Østerås: Statens strålevern.
 19. Broderstad AR, Melhus M, Brustad M *et al.* (2011) Iron stores in relation to dietary patterns in a multiethnic population: the SAMINOR study. *Public Health Nutr* **14**, 1039–1046.
 20. Brustad M, Parr CL, Melhus M *et al.* (2008) Dietary patterns in the population living in the Sami core areas of Norway – the SAMINOR study. *Int J Circumpolar Health* **67**, 82–96.
 21. Brustad M, Parr CL, Melhus M *et al.* (2008) Childhood diet in relation to Sami and Norwegian ethnicity in northern and mid-Norway – the SAMINOR study. *Public Health Nutr* **11**, 168–175.
 22. Nilsson LM, Dahlgren L, Johansson I *et al.* (2011) Diet and lifestyle of the Sami of southern Lapland in the 1930s–1950s and today. *Int J Circumpolar Health* **70**, 301–318.
 23. Nilsson LM, Winkvist A, Brustad M *et al.* (2012) A traditional Sami diet score as a determinant of mortality in a general northern Swedish population. *Int J Circumpolar Health* **71**, 1–12.
 24. Nilsson LM, Winkvist A, Eliasson M *et al.* (2012) Low-carbohydrate, high-protein score and mortality in a northern Swedish population-based cohort. *Eur J Clin Nutr* **66**, 694–700.
 25. Nilsson LM, Johansson I, Lenner P *et al.* (2010) Consumption of filtered and boiled coffee and the risk of incident cancer: a prospective cohort study. *Cancer Causes Control* **21**, 1533–1544.
 26. Nilsson LM, Wennberg M, Lindahl B *et al.* (2010) Consumption of filtered and boiled coffee and the risk of first acute myocardial infarction; a nested case/referent study. *Nutr Metab Cardiovasc Dis* **20**, 527–535.
 27. World Health Organization (1985) *Energy and Protein Requirements. Report of a Joint FAO/WHO/UNU Expert Consultation. WHO Technical Report Series* no. 724. Geneva: WHO.
 28. Lund E, Melhus M, Hansen KL *et al.* (2007) Population based study of health and living conditions in areas with both Sami and Norwegian populations – the SAMINOR study. *Int J Circumpolar Health* **66**, 113–128.
 29. Parr CL, Veierød MB, Laake P *et al.* (2006) Test–retest reproducibility of a food frequency questionnaire (FFQ) and estimated effects on disease risk in the Norwegian Women and Cancer Study (NOWAC). *Nutr J* **5**, 4.
 30. Hjartåker A, Lund E & Bjerve KS (1997) Serum phospholipid fatty acid composition and habitual intake of marine foods registered by a semi-quantitative food frequency questionnaire. *Eur J Clin Nutr* **51**, 736–742.
 31. Hjartåker A, Andersen LF & Lund E (2007) Comparison of diet measures from a food-frequency questionnaire with measures from repeated 24-hour dietary recalls. The Norwegian Women and Cancer Study. *Public Health Nutr* **10**, 1094–1103.
 32. Davino C, Furno M & Vistocco D (2014) *Quantile Regression*. Oxford: John Wiley & Sons Ltd.
 33. Brox J, Bjørnstad E & Olaussen K (2003) Hemoglobin, iron, nutrition and life-style among adolescents in a coastal and an inland community in northern Norway. *Int J Circumpolar Health* **62**, 130–141.
 34. Andersen S, Rex KF, Noahsen P *et al.* (2014) Forty-five year trends in overweight and obesity in an indigenous arctic Inuit society in transition and spatiotemporal trends. *Am J Hum Biol* **26**, 511–517.
 35. Nystad T, Melhus M, Brustad M *et al.* (2010) Ethnic differences in the prevalence of general and central obesity among the Sami and Norwegian populations: the SAMINOR study. *Scand J Public Health* **38**, 17–24.
 36. Broderstad AR & Melhus M (2016) Prevalence of metabolic syndrome and diabetes mellitus in Sami and Norwegian populations. The SAMINOR – a cross-sectional study. *BMJ Open* **6**, e009474.
 37. Njølstad I, Arnesen E & Lund-Larsen PG (1998) Cardiovascular diseases and diabetes mellitus in different ethnic groups: the Finnmark study. *Epidemiology* **9**, 550–556.
 38. Naseribafrouei A, Eliassen BM, Melhus M *et al.* (2016) Ethnic difference in the prevalence of pre-diabetes and diabetes mellitus in regions with Sami and non-Sami populations in Norway – the SAMINOR1 study. *Int J Circumpolar Health* **75**, 31697.
 39. Johnson-Down L, Labonte ME, Martin ID *et al.* (2015) Quality of diet is associated with insulin resistance in the Cree (Eeyouch) indigenous population of northern Quebec. *Nutr Metab Cardiovasc Dis* **25**, 85–92.
 40. Ryman TK, Boyer BB, Hopkins S *et al.* (2015) Associations between diet and cardiometabolic risk among Yup'ik Alaska Native people using food frequency questionnaire dietary patterns. *Nutr Metab Cardiovasc Dis* **25**, 1140–1145.
 41. Siri SRA (2015) *Use of Sami Ethnicity in Health Research and Registries*. Tromsø: Centre for Sami Health Research, UiT The Arctic University of Norway.
 42. Pettersen T (2015) Sami ethnicity as a variable. PhD Thesis, UiT The Arctic University of Norway.