

Maternal depression symptoms are highly prevalent among food-insecure households in Ethiopia

Seifu Hagos Gebreyesus^{1,2,*}, Bilal Shikur Endris¹, Charlotte Hanlon^{2,3} and Bernt Lindtjørn⁴

¹Department of Reproductive Health and Health Service Management, School of Public Health, College of Health Sciences, Addis Ababa University, PO Box 2140, Addis Ababa, Ethiopia: ²Department of Psychiatry, School of Medicine, College of Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia: ³Centre for Global Mental Health, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK: ⁴Centre for International Health, University of Bergen, Bergen, Norway

Submitted 13 April 2017: Final revision received 17 August 2017: Accepted 6 September 2017: First published online 20 November 2017

Abstract

Objective: We aimed to evaluate the association between household food insecurity and maternal depression in Ethiopia.

Design/Setting/Subjects: In 2014, we conducted a cross-sectional study in southern Ethiopia, including 591 food-secure and 2500 food-insecure households. We measured depression status of women using the Patient Health Questionnaire-9 validated for Ethiopia, with a cut-off of ≥ 5 . We evaluated household-level food insecurity using a validated Household Food Insecurity Access Scale. We applied Bayesian modelling to evaluate the relationship between food insecurity and maternal depression accounting for other observed characteristics.

Results: Among the analytic sample, 80.8 (95% CI 79.4, 82.2) % of women were living in food-insecure households. The overall prevalence of probable depression (mild and moderate forms) was 4.7 (95% CI 4.1, 5.6) %. All individual depressive symptoms had a significantly higher prevalence in the food-insecure group, except for suicidal ideation (but small numbers; $P < 0.001$). In the Bayesian model adjusting for paternal characteristics, there was a significant dose–response linear relationship (trend) between household food insecurity and maternal depression ($P < 0.01$). The adjusted OR (95% Bayesian credible interval) for depression for differing levels of food insecurity were: mild food insecurity, 3.29 (1.63, 6.18); moderate, 3.82 (1.91, 7.45); severe, 12.50 (3.38, 32.70).

Conclusions: The study documented a high burden of depression among women who lived in food-insecure households. Given this finding, we recommend integrating mental health in the livelihood programmes in areas suffering from food insecurity.

Keywords
Food insecurity
Depression
Ethiopia
Bayesian
Maternal nutrition

Depressive disorders are one of the major public health problems globally. According to the 2015 Global Burden of Disease study, depressive disorders are the third leading cause of disability⁽¹⁾. A systematic review in low- and lower-middle-income countries showed that about one in six pregnant women and one in five women who had recently given birth experienced common perinatal mental disorders⁽²⁾. That review revealed different determinants of common perinatal mental disorders, such as financial difficulties, unemployment, difficulties in an intimate partner relationship, insufficient social support and adverse reproductive outcomes. Furthermore, in developing countries a high prevalence of maternal common mental disorders (CMD), ranging from 21% in Vietnam to 33% in Ethiopia, is reported⁽³⁾.

In Ethiopia, although there are no nationally representative data, maternal depression ranging from 5 to 37% was reported from different pocket studies^(4–8). In a systematic review of twelve studies conducted from 1990 to 2012 in Ethiopia, the population prevalence of depression was estimated to be 11%⁽⁹⁾. Female gender, marital loss, poor social support, low wealth index, tobacco smoking, hazardous use of alcohol, older age, having a co-morbid illness, living alone, stressful life events and increasing age were associated with CMD in Ethiopia^(10,11).

In 2001, the FAO defined food security as: 'a situation that exists when all people, at all times, have physical, social and economic access to sufficient, safe and nutritious food that meets their dietary needs and food preferences for an active and healthy life'. According to FAO

estimates, 795 million people remained undernourished in 2015⁽¹²⁾. The estimate showed that East African countries had a bigger share the problem. In Ethiopia the prevalence of household food insecurity was estimated to be as high as 70%⁽¹³⁾. Although Ethiopia has made significant progress against undernutrition, 40% of children in the country are stunted⁽¹⁴⁾.

The health consequences of food insecurity go beyond undernutrition. There is accumulating evidence that shows a positive association between CMD and food insecurity^(8,15–18). The association between depression and food insecurity is complex. A bidirectional causal relationship between household food insecurity and depression is documented⁽¹⁷⁾. Food insecurity is indicated as a risk factor for depression. Different mechanisms such as psychological distress and micronutrient deficiency are reported^(18–22). On the other hand, depression can be a risk factor for food insecurity through unemployment, lost productivity and increased health expenditure^(21–23).

However, there is limited evidence on the association between food insecurity and depression among women in Ethiopia, where the problem of food insecurity is rampant⁽¹³⁾. In addition, the exiting literature has limitations in terms of study size, representativeness and the use of locally validated tools^(24,25). In the present study, we hypothesized that maternal depression is associated with food insecurity in a highly food-insecure setting and tested this using a large representative sample and robust analytic method. Understanding maternal depression is very important since studies have reported that maternal depression could be one of the pathways through which household food insecurity could impact child undernutrition^(3,26–28).

Materials and methods

Study setting

The study was conducted in Meskan and Mareko districts, in the Gurage Zone of the Southern Nations and Nationalities regional state. Butajira, located 135 km south of Addis Ababa (the capital city of Ethiopia), is the districts' capital and has an estimated population of 199 771.

The agroecology of the district is characterized by an elevation ranging from approximately 1500 to 3500 m above sea level, with an average temperature ranging from 10 to 24°C and an average annual rainfall ranging from 900 to 1400 mm. The main means of livelihood in the district is rain-fed agriculture, which is characterized by production of subsistence crops (primarily enset, cereals, legumes, vegetables, fruits) and some cash crops such as *khat* (*Catha edulis*). Most of the households are affected by chronic food insecurity and are prone to drought.

Butajira Health and Demographic Surveillance Site, located in Butajira, is one of the oldest surveillance sites in Africa, being established in 1986. It consists of nine rural and one urban *kebeles* (the smallest administrative unit in

Ethiopia) from different ecological zones. Each *kebele* has one or two permanent enumerators who are residents of the respective *kebeles*. Seven health centres, forty health posts and two hospitals serve the residents in the district.

Study design and period

The present study employed a cross-sectional design. The data were collected during the postharvest season, i.e. between December 2013 and April 2014.

Sample size and sampling

The analytical sample for the study was based on a sample size calculated for another objective that aimed to examine spatial variations in the distribution of undernutrition in southern Ethiopia⁽²⁹⁾. The sample size available for the present analysis was 3091 women with children between the ages of 0 and 59 months. We recalculated the power of the analytic sample for its adequacy to answer the present research question. The power based on normal approximation with continuity correction was 98.8%.

The health extension workers provided us with a list of houses, names of the household head and the family profile. This was obtained from the family registry forms and used simple random sampling to select eligible houses that have one or more children under the age of 5 years. We visited all houses with support from the health extension workers, local guides and study supervisors.

Outcome measures

The main outcome of the study was maternal depression. Maternal depression was measured using the Patient Health Questionnaire-9 (PHQ-9)⁽³⁰⁾. The PHQ-9 is a self-report measure used to screen major depressive disorders. The PHQ-9 has been found to be valid in two studies from Ethiopia in different treatment settings: tertiary hospital medical outpatients⁽³¹⁾ and primary-care outpatients in the study area⁽³²⁾. The PHQ-9 is composed of nine items asked with a recall period of two weeks. The items inquire how often respondents have been bothered by depressive symptoms, with response categories of 0 = 'not at all', 1 = 'several days', 2 = 'more than half the days' and 3 = 'nearly every day'.

In the present study, the internal consistency of the PHQ-9 was moderate (Cronbach's α value of 0.65). Validation work in another study has shown evidence of construct validity: factor analysis indicated a unidimensional scale. Core depression items (feeling down, little interest) were among the items most strongly related to the underlying construct⁽³¹⁾.

Other measures

The main explanatory (exposure) variable considered was the level of household food insecurity. Secondary variables measured in the analytic sample were woman's age, marital status, religion, ethnicity, education, husband's occupation, household head's gender, household wealth and dietary diversity. Socio-economic status was measured

using a customized questionnaire asking about ownership and size of land; type of house and construction materials; availability of radio, television, house telephone, mobile phone, bed, chair, refrigerator, electric stove, clock/watch and electricity; possession of domestic animals; and sanitation facilities and source of water.

Household food insecurity and dietary diversity

We used the Household Food Insecurity Access Scale (HFIAS) to measure the magnitude of food insecurity at household level⁽³³⁾. The HFIAS tool was validated and has been shown to perform well with minor adaptations in the same setting⁽³⁴⁾. The HFIAS is a nine-item scale with self-reported items using a recall period of four weeks and response categories relating to the frequency of occurrence.

We collected data on food groups eaten by the respective household during the day and night preceding the survey. A dietary diversity score for each household was generated using the sum of all food groups consumed. The total number of food groups consumed by households could range from 0 to 12. Based on the reported number of food groups consumed, we classified households into lowest, medium and high dietary diversity⁽³⁵⁾.

Data collection

We recruited twenty research assistants and two supervisors who had experience in data collection and supervision. We conducted a comprehensive training on the administration of the questionnaire and interviewing skills. One of the authors (S.H.G.) translated the questionnaire into the Amharic language (official language) and reviewed it together with the research assistants who were residents in the study district. We conducted a pre-test in a nearby district with socio-economic characteristics similar to those in the study district.

Data analysis

We used the software EpiData version 3.1 for data entry and the statistical software package Stata version 13.0 for data cleaning and initial analysis. Data were cleaned for inconsistencies and missing values. All variables with missing data were reported. Descriptive analysis was done on explanatory variables using frequency distributions and summary statistics. We evaluated the comparability between the women in the food-secure and food-insecure groups using χ^2 and *t* tests as appropriate. Maternal depression was indicated by categorization of total PHQ-9 score as follows: 5–9, mild depression; 10–14, moderate depression; 15–19, moderately severe depression; and ≥ 20 , severe depression.

We used the HFIAS tool to classify households into four levels of food insecurity: food secure, mildly food insecure, moderately food insecure and severely food insecure. The nine items in the HFIAS are ordered in

such a way to reflect an increasing severity of food insecurity. A food-secure household experiences none of the nine items in the HFIAS or experienced the first item only (worry about food), but rarely. A severely food-insecure household experiences one of the last three items (items 7–9: running out of food, going to bed hungry, or going a whole day and night without eating). The complete procedure used to categorize a household to one of the four food insecurity levels is described elsewhere⁽²⁹⁾.

We used principal component analysis to construct a household wealth index. The variables used in the principal component analysis included ownership and size of land, type of house and construction materials, availability of fixed assets such as radio, television, phone, bed, chair and other household items, possession of domestic animals, type of water source for drinking and cooking, and availability and type of latrine. The resulting scores were divided into quintiles that indicated poorest, poor, medium, rich and richest.

A logistic regression model was fitted using the statistical software package Stata version 11.0 to identify variables for Bayesian modelling. Explanatory variables such as the level of household food insecurity, maternal age, ethnicity, religion, husband's occupation, household wealth and gender of the household head were statistically significant at the 20% significance level (at $P < 0.2$) and considered as potential explanatory variables for the Bayesian modelling.

Bayesian logistic regression modelling was undertaken using WinBUGS version 1.4.3 (MRC Biostatistics Unit, Cambridge and Imperial College London, UK; see online supplementary material). We started with non-informative prior and 10 000 iterations. We checked for convergence of parameters visually using history plot and kernel density. Convergence was successfully achieved after 10 000 iterations. After convergence, a further 10 000 iterations were run, and values were thinned by 10 and stored. The stored samples were used to calculate summary statistics (mean, sd and 95% Bayesian credible interval (BCI)) of the parameters.

Dose-response analysis

We ran a dose-response analysis to evaluate if the prevalence of depression increased with increasing severity of household food insecurity. A test was performed using the extended Mantel-Haenszel χ^2 for linear trend using OpenEpi software.

Results

Participant characteristics

The analytic sample included a total of 3091 mothers with one or more children under the age of 5 years. Only fifteen (0.35%) households (i.e. mothers) were not included in the study because of unavailability after repeated visits.

Table 1 Sociodemographic characteristics of women (*n* 3091) according to household food security status in south central Ethiopia, 2014

Characteristic	Food secure		Food insecure		<i>P</i> value
	Mean or <i>n</i>	SD or %	Mean or <i>n</i>	SD or %	
Mother's age (years), mean and sd	33.9	6.9	33.7	6.4	0.618
Mother's age group, <i>n</i> and %					0.031
<20 years	5	0.9	11	0.4	
20–34 years	274	46.4	1198	47.9	
35–49 years	282	47.7	1219	48.8	
>49 years	30	5.1	72	2.9	
Mother's education, <i>n</i> and %					<0.001
Illiterate	396	67.0	1905	76.2	
Able to read and write	45	7.6	73	1.9	
Primary school and above	143	24.2	503	20.1	
Marital status, <i>n</i> and %					0.484
Currently married	552	93.4	2354	94.2	
Currently unmarried	39	6.6	146	5.8	
Maternal occupation, <i>n</i> and %					0.001
Housewife	374	63.3	1781	71.2	
Farmer and housewife	49	8.3	208	8.3	
Merchant	135	22.8	398	15.9	
Other (daily labourer, student, etc.)	17	2.7	48	1.9	
Religion, <i>n</i> and %					<0.001
Orthodox Christian	161	27.2	546	21.8	
Muslim	363	61.4	1781	71.2	
Other (Protestant, Catholic)	62	10.5	159	6.4	
Husband's occupation, <i>n</i> and %					<0.001
Farmer	501	84.8	2262	90.4	
Daily labourer	11	1.9	81	3.2	
Employee/private/merchant	79	13.4	158	6.3	
Household head's gender, <i>n</i> and %					<0.001
Female	48	8.1	181	7.2	
Male	529	89.5	2303	92.1	
Wealth quintile, <i>n</i> and %					<0.001
Poorest	105	17.7	640	25.6	
Poor	90	15.2	476	19.0	
Middle	97	16.4	467	18.7	
Rich	137	23.2	548	21.9	
Richest	162	27.4	369	14.8	
Ethnicity, <i>n</i> and %					0.015
Guraghe	476	80.5	1921	76.8	
Silte	65	11.0	402	16.1	
Mareko	28	4.7	93	3.7	
Other	22	3.7	84	3.4	
Dietary diversity score, mean and sd	4.3	0.9	4.1	0.8	<0.001
Dietary diversity, <i>n</i> and %					<0.001
Lowest	79	13.4	890	15.6	
Medium	451	76.3	1994	79.8	
High	61	10.3	116	4.6	
Depression status					<0.001
PHQ-9 score, mean and sd	0.4	1.15	1.5	1.9	
Depressed, <i>n</i> and %	8	1.3	139	5.6	<0.01

PHQ-9, Patient Health Questionnaire-9.

The mean age of the mothers at the time of the interview was 33 (SD 6.08) years. Most of the respondents were illiterate (74.4%) and housewives (69.7%). The great majority (94.0%) were married. Table 1 shows the differences in the background characteristics of the respondents who lived in food-secure and food-insecure households. Women who lived in the food-insecure households were illiterate ($\chi^2 = 37.8$, $P < 0.001$), poor ($\chi^2 = 62.2$, $P < 0.001$) and had a lower dietary diversity score ($\chi^2 = 30.3$, $P < 0.001$) compared with women who lived in food-secure households.

Magnitude of household food insecurity and maternal depression

The overall prevalence of food insecurity was 80.8 (95% CI 79.4, 82.2)%. Of the women, 1087 (35.2%) lived in mildly food-insecure households, 1352 (43.7%) lived in moderately food-insecure households and thirty-eight (1.2%) lived in severely food-insecure households.

The overall prevalence of depression (PHQ-9 score of ≥ 5) was 4.7 (95% CI 4.1, 5.6)%. The overall prevalence of depression at the mild, moderate and severe threshold

Table 2 Patient Health Questionnaire-9 (PHQ-9) item responses of women living in food-secure and food-insecure households, south central Ethiopia, 2014

Depressive symptoms (PHQ-9 scale item)	Food-secure women (n 591)		Food-insecure women (n 2500)		P value
	None (%)	Several days per week or more (%)	None (%)	Several days per week or more (%)	
Q1. Little interest or pleasure in doing things	97.6	2.4	85.3	14.7	<0.001
Q2. Feeling down, depressed or hopeless	95.4	4.6	86.7	13.3	<0.001
Q3. Trouble falling or staying asleep	95.9	4.1	89.4	10.6	<0.001
Q4. Feeling tired or having little energy	88.0	12.0	54.9	45.1	<0.001
Q5. Poor appetite or overeating	96.1	3.9	87.7	12.3	<0.001
Q6. Feeling bad about self	98.6	1.4	94.6	5.4	0.001
Q7. Trouble concentrating	98.3	1.7	85.3	14.7	<0.001
Q8. Moving or speaking slowly	99.3	0.7	97.5	2.5	<0.05
Q9. Thoughts of being better off dead	99.7	0.3	99.0	1.0	0.297

levels was 3.9 (95% CI 3.3, 4.6) %, 0.9 (95% CI 0.6, 1.2) % and 0 %, respectively.

The overall mean PHQ-9 score was 1.27 (SD 1.87; 95% CI 1.20, 1.34), ranging from 0 to 14 points. The mean PHQ-9 score was 1.48 (95% CI 1.40, 1.56) in women who lived in food-insecure households and 0.38 (95% CI 0.29, 0.48) in women who lived in food-secure households (i.e. 1.08 points lower (95% CI -1.26, -0.93)).

The distribution of responses to the PHQ-9 items by household food security status is presented in Table 2. The responses to the problems were categorized into two groups: (i) those without any problems (none); and (ii) those with the problems occurring in any frequency (several days, more than half the days or nearly every day) over the past two weeks. PHQ-9 item frequency ranged from 0.3 to 12.0 % and from 1.0 to 45.1 % among women in the food-secure and food-insecure groups, respectively. 'Thoughts of being better off dead' received the lowest affirmative responses in both groups of women.

All individual depressive symptoms had a significantly higher prevalence in the food-insecure group, except for suicidal ideation (but small numbers; $P < 0.001$). For example, women in food-insecure households, compared with the food-secure group, reported seven times more on item 1 ('little interest or pleasure in doing things') and eight times more on item 7 ('trouble concentrating'; Table 2).

Household food insecurity and depression

In the Bayesian logistic regression model, women who live in food-insecure households had higher odds of depression than women who lived in food-secure households (Table 3). The model adjusted for mother's age, household wealth, husband's occupation, household head's gender, religion and ethnicity.

The adjusted OR (AOR) for depression for differing levels of food insecurity were: mild food insecurity, AOR = 3.29 (95% BCI 1.63, 6.18); moderate, AOR = 3.82 (95% BCI 1.91, 7.45); and severe, AOR = 12.50 (95% BCI 3.38, 32.70).

Other variables were also associated with depression: female-headed households (AOR = 1.85; 95% BCI 1.02,

3.01) and women who lived in the poorest households (AOR = 1.85; 95% BCI: 1.00, 3.24) were more likely to be depressed. Women's age was also associated with depression; those women who were older were more likely to be depressed than women who were younger (AOR = 1.03; 95% BCI 1.00, 1.06).

Dose-response relationship

The magnitude of depression increased as the level of household food insecurity increased (Fig. 1). We found a significant dose-response linear relationship between household food insecurity and women's depression (dose-response extended Mantel-Haenszel χ^2 for linear trend = 20.54, $P < 0.01$).

Discussion

The present analysis employed a large cross-sectional study to evaluate the association between household food insecurity and maternal depression in southern Ethiopia. We used validated measures for both depression and food insecurity. After adjustment for a range of confounders, women who lived in food-insecure households had significantly higher odds of depression compared with women who lived in food-secure households.

The relationship between food insecurity and depression may be explained through different mechanisms. One of the mechanisms could be related to increased psychological stress associated with food insecurity. Household food insecurity is likely to be associated with worries about not having enough food for the family in general and for children in particular. Furthermore, shame because of eating a socially unacceptable diet may lead to stress. Due to these and other psychological stressors, women will be at higher risk of developing CMD⁽¹⁸⁾. Another possible mechanism that could explain why food-insecure women are at higher risk of CMD is related to nutritional deficiencies. There is an emerging evidence base indicating that deficiencies of specific nutrients, such as folate, vitamin D, vitamin B₁₂, Zn and *n*-3 fatty acids, are associated with CMD^(19,20).

Table 3 Bayesian logistic model for the relationship between food security and maternal depression, south central Ethiopia, 2014

Background characteristic	Depression			Bayesian logistic regression model	
	Posterior mean	SD	Median	AOR	95% BCI
Household food security					
Food secure	–	–	–	1.00	Ref.
Mild food insecurity	3.29	1.2	3.1	3.29	1.63, 6.18
Moderate food insecurity	3.82	1.4	3.6	3.82	1.91, 7.45
Severe food insecurity	12.5	7.6	10.7	12.50	3.38, 32.70
Household head's gender					
Male	–	–	–	1.00	Ref.
Female	1.85	0.49	1.80	1.85	1.02, 3.01
Household wealth					
Richest	–	–	–	1.00	Ref.
Rich	1.00	0.36	0.94	1.00	0.48, 1.82
Middle	1.35	0.47	1.28	1.35	0.69, 2.47
Poor	1.87	0.60	1.76	1.87	0.95, 3.63
Poorest	1.85	0.59	1.74	1.85	1.00, 3.24
Mother's age (years)*	1.03	0.01	1.03	1.03	1.00, 1.06
Ethnicity					
Guraghe	–	–	–	1.00	Ref.
Silte	0.98	0.24	0.96	0.98	0.57, 1.53
Mareko	1.15	0.52	1.06	1.15	0.41, 2.45
Others	1.97	0.78	1.85	1.97	0.82, 3.81
Husband's occupation					
Farmer	2.45	1.54	2.03	2.45	0.87, 6.70
Daily labourer	1.79	1.59	1.36	1.79	0.29, 5.72
Employee/private/merchant	–	–	–	1.00	Ref.
Religion					
Orthodox Christian	1.54	0.60	1.42	1.54	0.72, 3.43
Muslim	1.00	0.38	0.92	1.00	0.48, 1.99
Other	–	–	–	1.00	Ref.

AOR, adjusted odds ratio; BCI, Bayesian credible interval; Ref., reference category.

Significant results are indicated in bold font.

*Maternal age is treated as a continuous variable.




Fig. 1 Percentage of women ($n = 3091$) with depression by household food security status, south central Ethiopia, 2014 (dose–response extended Mantel–Haenszel χ^2 for linear trend = 20.54, $P < 0.01$)

The finding of higher odds of depression in women who lived in food-insecure households is consistent with a number of previous studies^(8,15–17,25,36,37). Furthermore, the present study's finding is in agreement with a systematic review of eleven qualitative and sixteen

quantitative studies conducted in low- and middle-income countries which confirmed a positive association between food insecurity and CMD⁽¹⁸⁾. We also found that the odds of depression increased as the level of household food insecurity increased^(37,38).

Our analysis found that women who lived in the poorest households were more likely to be depressed than those who lived in the richest households. Although a similar finding is documented elsewhere^(22,39,40), the relationship between depression and poverty is complex⁽⁴¹⁾. According to the social causation hypothesis, poverty raises the risk of mental illness through different pathways including increased stress, social segregation, violence and trauma. On the other hand, according to the social selection hypothesis, people with mental illness are at increased risk of poverty through increased health expenditure, reduced productivity, stigma, and loss of employment and associated earnings. The social causation pathway might be more relevant to CMD such as depression, whereas the social selection hypothesis might be more appropriate to disorders such as schizophrenia and intellectual disabilities⁽⁴²⁾.

The findings of the present study should be interpreted within the context of the following limitations. Due to the cross-sectional nature of the study, we cannot rule out the possibility of reverse causality, i.e. depression may cause household food insecurity. Second, the study did not account for different determinants of depression including poor social support, recent stressful life events, previous depressive illness, violence and physical illness, which might not be evenly distributed among households of different levels of food insecurity. Moreover, respondent bias with the anticipation of some form of support might overestimate the prevalence of food insecurity, although this may not affect the association between depression and food insecurity as the anticipation of support might not differ between depressed and non-depressed women. In addition to these limitations, we admit that the present analysis is subjected to recall bias. It has been documented that depressed women have negative cognitive distortion and may be more likely to give affirmative responses to the household food insecurity questions.

The use of locally validated tools to measure depression (PHQ-9) and food insecurity (HFIAS) are important strengths of the present study. In addition, we used large sample size with adequate power and this helped us to analyse the dose–response trend among different levels of food insecurity and depression. Moreover, we applied a robust statistical analysis in a Bayesian framework which enabled us to get robust estimates.

Conclusion

In conclusion, household food insecurity was found to be an important determinant of maternal depression in rural Ethiopia. However, further research employing robust research designs is recommended to answer the relationship between food security and depression. For this, we recommend a prospective study design with a cohort of food-secure and food-insecure households that are screened for depression at a baseline. Furthermore, we

recommend a study that aims to evaluate the mechanism(s) explaining the relationship between food insecurity and depression. This might involve testing the hypothesis that some micronutrient deficiencies could explain the relationship between food insecurity and depression. In addition, intervention studies are required to test whether interventions to reduce food insecurity lead to decreased maternal depression.

Acknowledgements

Acknowledgements: The authors would like to thank the Centre for International Health at the University of Bergen for funding the study and all study participants for their willingness to participate in the study. *Financial support:* This work was supported by the Centre for International Health at the University of Bergen. The funding body had no role in designing the study, in the collection, analysis and interpretation of data, in the writing of this manuscript and in the decision to submit for publication. *Conflict of interest:* The authors declare that they have no competing interests. *Authorship:* S.H.G. and B.L. designed the study. S.H.G., B.L., B.S.E. and C.H. participated in the data analysis and drafting of the manuscript. All authors read and approved the final manuscript. *Ethics of human subject participation:* The study protocol was approved by institutional review boards from Addis Ababa University, College of Health Sciences. The study was also approved by the Regional Committee for Medical and Research Ethics, Western Norway (REK Vest). Information on the research objective was read to the participants and verbal informed consent was received. Privacy and confidentiality of respondents were also maintained.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1368980017003056>

References

1. Vos T, Allen C, Arora M *et al.* (2016) Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* **388**, 1545–1602.
2. Fisher J, Cabral de Mello M, Patel V *et al.* (2012) Prevalence and determinants of common perinatal mental disorders in women in low- and lower-middle-income countries: a systematic review. *Bull World Health Organ* **90**, 139–149.
3. Harpham T, Huttly S, De Silva MJ *et al.* (2005) Maternal mental health and child nutritional status in four developing countries. *J Epidemiol Community Health* **59**, 1060–1064.
4. Deyessa N, Berhane Y, Alem A *et al.* (2009) Intimate partner violence and depression among women in rural Ethiopia: a cross-sectional study. *Clin Pract Epidemiol Mental Health* **5**, 8.

5. Hanlon C, Medhin G, Alem A *et al.* (2009) Impact of antenatal common mental disorders upon perinatal outcomes in Ethiopia: the P-MaMiE population-based cohort study. *Trop Med Int Health* **14**, 156–166.
6. Bitew T, Hanlon C, Kebede E *et al.* (2017) Antenatal depressive symptoms and utilisation of delivery and postnatal care: a prospective study in rural Ethiopia. *BMC Pregnancy Childbirth* **17**, 206.
7. Dibaba Y, Fantahun M & Hindin MJ (2013) The association of unwanted pregnancy and social support with depressive symptoms in pregnancy: evidence from rural Southwestern Ethiopia. *BMC Pregnancy Childbirth* **13**, 135.
8. Hadley C, Tegegn A, Tessema F *et al.* (2008) Food insecurity, stressful life events and symptoms of anxiety and depression in east Africa: evidence from the Gilgel Gibe growth and development study. *J Epidemiol Community Health* **62**, 980–986.
9. Bitew T (2014) Prevalence and risk factors of depression in Ethiopia: a review. *Ethiop J Health Sci* **24**, 161–169.
10. Fekadu A, Medhin G, Selamu M *et al.* (2014) Population level mental distress in rural Ethiopia. *BMC Psychiatry* **14**, 194.
11. Molla GL, Sebhat HM, Hussen ZN *et al.* (2016) Depression among Ethiopian adults: cross-sectional study. *Psychiatry J* **2016**, 1468120.
12. Food and Agriculture Organization of the United Nations, International Fund for Agricultural Development & World Food Programme (2015) *The State of Food Insecurity in the World 2015 – Meeting the 2015 International Hunger Targets: Taking Stock of Uneven Progress*. Rome: FAO.
13. Endale W, Mengesha ZB, Atinafu A *et al.* (2014) Food Insecurity in Farta District, Northwest Ethiopia: a community based cross-sectional study. *BMC Res Notes* **7**, 130.
14. Headey D (2014) *An Analysis of Trends and Determinants of Child Undernutrition in Ethiopia, 2000–2011*. Washington, DC: International Food Policy Research Institute.
15. Hadley C & Patil CL (2006) Food insecurity in rural Tanzania is associated with maternal anxiety and depression. *Am J Hum Biol* **18**, 359–368.
16. Noonan K, Corman H & Reichman NE (2016) Effects of maternal depression on family food insecurity. *Econ Hum Biol* **22**, 201–215.
17. Huddleston-Casas C, Charnigo R & Simmons LA (2009) Food insecurity and maternal depression in rural, low-income families: a longitudinal investigation. *Public Health Nutr* **12**, 1133–1140.
18. Weaver LJ & Hadley C (2009) Moving beyond hunger and nutrition: a systematic review of the evidence linking food insecurity and mental health in developing countries. *Ecol Food Nutr* **48**, 263–284.
19. Jacka FN, Maes M, Pasco JA *et al.* (2012) Nutrient intakes and the common mental disorders in women. *J Affect Disord* **141**, 79–85.
20. Rao TS, Asha M, Ramesh B *et al.* (2008) Understanding nutrition, depression and mental illnesses. *Indian J Psychiatry* **50**, 77–82.
21. Beck A, Crain AL, Solberg LI *et al.* (2011) Severity of depression and magnitude of productivity loss. *Ann Fam Med* **9**, 305–311.
22. Patel V & Kleinman A (2003) Poverty and common mental disorders in developing countries. *Bull World Health Organ* **81**, 609–615.
23. Stewart WF, Ricci JA, Chee E *et al.* (2003) Cost of lost productive work time among US workers with depression. *JAMA* **289**, 3135–3144.
24. Maes KC, Hadley C, Tesfaye F *et al.* (2010) Food insecurity and mental health: surprising trends among community health volunteers in Addis Ababa, Ethiopia during the 2008 food crisis. *Soc Sci Med* **70**, 1450–1457.
25. Jebena MG, Taha M, Nakajima M *et al.* (2015) Household food insecurity and mental distress among pregnant women in Southwestern Ethiopia: a cross sectional study design. *BMC Pregnancy Childbirth* **15**, 250.
26. Hadley C, Tessema F & Muluneh AT (2012) Household food insecurity and caregiver distress: equal threats to child nutritional status? *Am J Hum Biol* **24**, 149–157.
27. Jyoti DF, Frongillo EA & Jones SJ (2005) Food insecurity affects school children's academic performance, weight gain, and social skills. *J Nutr* **135**, 2831–2839.
28. Wemakor A & Mensah KA (2016) Association between maternal depression and child stunting in Northern Ghana: a cross-sectional study. *BMC Public Health* **16**, 869.
29. Seid O, Tadesse K & Gebremedhin A (2015) Iron deficiency anemia is moderate public health problem among school going adolescent girls in Berahle district, Afar, northeast Ethiopia. *J Food Nutr Sci* **3**, 10–16.
30. Kroenke K & Spitzer RL (2002) The PHQ-9: a new depression diagnostic and severity measure. *Psychiatr Ann* **32**, 509–515.
31. Gelaye B, Williams MA, Lemma S *et al.* (2013) Validity of the Patient Health Questionnaire-9 for depression screening and diagnosis in East Africa. *Psychiatry Res* **210**, 653–661.
32. Hanlon C, Medhin G, Selamu M *et al.* (2015) Validity of brief screening questionnaires to detect depression in primary care in Ethiopia. *J Affect Disord* **186**, 32–39.
33. Swindale A & Bilinsky P (2007) *Household Food Insecurity Access Scale (HFIAS) for Measurement of Household Food Access: Indicator Guide (v. 3)*. Washington, DC: Food and Nutrition Technical Assistance Project, Academy for Educational Development.
34. Gebreyesus SH, Lunde T, Mariam DH *et al.* (2015) Is the adapted Household Food Insecurity Access Scale (HFIAS) developed internationally to measure food insecurity valid in urban and rural households of Ethiopia? *BMC Nutr* **1**, 2.
35. Kennedy G, Ballard T & Dop MC (2011) *Guidelines for Measuring Household and Individual Dietary Diversity*. Rome: FAO.
36. Sorsdahl K, Slopen N, Siefert K *et al.* (2011) Household food insufficiency and mental health in South Africa. *J Epidemiol Community Health* **65**, 426–431.
37. Whitaker RC, Phillips SM & Orzol SM (2006) Food insecurity and the risks of depression and anxiety in mothers and behavior problems in their preschool-aged children. *Pediatrics* **118**, e859–e868.
38. Dewing S, Tomlinson M, le Roux IM *et al.* (2013) Food insecurity and its association with co-occurring postnatal depression, hazardous drinking, and suicidality among women in peri-urban South Africa. *J Affect Disord* **150**, 460–465.
39. Brown GW & Moran PM (1997) Single mothers, poverty and depression. *Psychol Med* **27**, 21–33.
40. Martikainen P, Adda J, Ferrie J *et al.* (2003) Effects of income and wealth on GHQ depression and poor self rated health in white collar women and men in the Whitehall II study. *J Epidemiol Community Health* **57**, 718–723.
41. Lund C, Breen A, Flisher AJ *et al.* (2010) Poverty and common mental disorders in low and middle income countries: a systematic review. *Soc Sci Med* **71**, 517–528.
42. Hudson CG (2005) Socioeconomic status and mental illness: tests of the social causation and selection hypotheses. *Am J Orthopsychiatry* **75**, 3–18.