

# Subclinical inflammation affects iron and vitamin A but not zinc status assessment in Senegalese children and Cambodian children and women‡

Marion Fiorentino<sup>1,\*</sup>, Marlène Perignon<sup>1</sup>, Khov Kuong<sup>2</sup>, Chhoun Chamnan<sup>2</sup>, Jacques Berger<sup>1</sup> and Frank T Wieringa<sup>1</sup>

<sup>1</sup>Institut de Recherche pour le Développement (IRD), UMR 204 Nutripass, IRD-UM2-UM1, Montpellier, France:

<sup>2</sup>Department of Fisheries Post-Harvest Technologies and Quality Control, Ministry of Agriculture, Forestry and Fisheries, Phnom Penh, Cambodia

Submitted 31 May 2017: Final revision received 3 November 2017: Accepted 20 November 2017: First published online 18 January 2018

## Abstract

**Objective:** To assess the impact of the acute-phase response (APR) during inflammation on Fe, Zn and vitamin A biomarkers to allow accurate evaluation of micronutrient status in populations.

**Design:** Ferritin (FER), soluble transferrin receptor (TfR), retinol-binding protein (RBP), Zn,  $\alpha_1$ -acid glycoprotein and C-reactive protein concentrations were measured. Correction factors (CF) for each biomarker were calculated as the ratio for groups at different stages of inflammation *v.* the reference group without inflammation.

**Setting/Subjects:** Senegalese (*n* 594) and Cambodian schoolchildren (*n* 2471); Cambodian women of reproductive age (*n* 2117).

**Results:** TfR was higher during the incubation phase (CF = 1.17) and lower during early and late convalescence (CF = 0.87 and 0.78). FER was higher during all phases (CF = 0.83, 0.48 and 0.65, respectively). RBP was higher during incubation (CF = 0.88) and lower during early convalescence (CF = 1.21). No effect of inflammation on Zn status was found.

**Conclusions:** Inflammation led to overestimation of Fe status and underestimation of vitamin A status. The response of the biomarker for vitamin A status to inflammation depended on the vitamin A status of the populations. Surprisingly, the assessment of Zn status was hardly affected by inflammation. Different phases of the APR had opposite effects on the assessment of Fe status using TfR. More research is needed to define the correct methods to adjust for inflammation in nutritional studies.

**Keywords**  
Inflammation  
Ferritin  
Soluble transferrin receptor  
Zinc  
Retinol-binding protein

Micronutrient deficiencies are a major public health problem, especially in low-income countries, mostly affecting children and women of reproductive age<sup>(1)</sup>. Micronutrient deficiencies are associated with increased prevalence of infectious diseases and hampered cognitive development in childhood. By impairing growth and school achievement in children and affecting reproductive functions and fetal development in women<sup>(1)</sup>, they accentuate the intergenerational cycle of malnutrition<sup>(2)</sup>.

The accuracy of micronutrient status assessment can be hampered by factors affecting biomarkers independently from status. Inflammation, for example, is known to affect biomarkers for Fe status (plasma ferritin (FER)

concentration) and vitamin A status (plasma retinol and retinol-binding protein (RBP) concentrations) even in apparently healthy individuals<sup>(3)</sup>. A complex chain reaction involving immunoregulatory cytokines follows injury or infection and contributes to a redistribution of nutrients to different body compartments. This redistribution results in a withholding of essential nutrients to infectious agents ('nutritional immunity') as well as benefits repair processes<sup>(4)</sup>. However, this redistribution also affects the concentrations of biomarkers for micronutrient status in the circulation, while overall micronutrient status remains the same, unless an infection is prolonged or severe. At the same time, it appears that micronutrient deficiency can affect the acute-phase response (APR)<sup>(4)</sup>. Therefore, more precision is needed about the biological response to inflammation of biomarkers for micronutrient status. For biomarkers of vitamin A status (plasma retinol and RBP

‡ The original version of this article was published with incorrect author affiliation information. A notice detailing this has been published and the error rectified in the online PDF and HTML copies.

concentrations)<sup>(5-7)</sup>, correction factors based on a meta-analysis have been proposed to adjust values in subjects during different stages of inflammation<sup>(5)</sup>. A similar approach has been used for FER concentrations<sup>(8)</sup>. Fe deficiency (ID) is indicated by depleted Fe stores, corresponding to low plasma FER concentrations, and/or tissue Fe deficiency, indicated by elevated soluble transferrin receptor (TfR) concentrations<sup>(9)</sup>. The impact of inflammation on TfR concentrations is less clear, with some studies suggesting that TfR is not or less sensitive to inflammation than FER<sup>(10)</sup>. Other studies, however, showed a significant impact of the APR on TfR values<sup>(11,12)</sup>. The effect of the APR on plasma Zn concentration remains unclear too<sup>(13,14)</sup>.

The objective of the present study was therefore to investigate the impact of subclinical inflammation on the assessment of micronutrient deficiencies in three different populations and settings (Senegalese schoolchildren, Cambodian schoolchildren, Cambodian women of reproductive age) by investigating the impact of having elevated acute-phase proteins (APP), namely C-reactive protein (CRP) and  $\alpha_1$ -acid glycoprotein (AGP), on biomarkers for Fe, vitamin A and Zn status.

## Participants and methods

### **Study area, design, population surveyed and ethics**

Three data sets were available for the current study as follows.

#### *Schoolchildren from Senegal*

The study was a representative cross-sectional survey conducted in 2010 in children from primary state schools of Dakar, Senegal<sup>(15)</sup>. A two-stage cluster sampling method was chosen, with schools considered primary sampling units. Within thirty randomly selected schools, and without criteria for age and gender, twenty children were randomized as final sampling units in each school, resulting in a sample of 594 apparently healthy schoolchildren.

#### *Schoolchildren from Cambodia*

Data were collected as part of a randomized placebo-controlled trial investigating the impact of multi-micronutrient-fortified rice on the health and development of Cambodian schoolchildren: the FORISCA (FOrtified Rice for School meals in Cambodia) UltraRice<sup>®</sup>+NutriRice<sup>®</sup> study. Baseline data collection was conducted in November 2012 in twenty primary schools from Kampong Speu Province in Cambodia<sup>(16)</sup>. The schools were randomly selected from primary schools participating in school meal or take-home ration programmes of the UN World Food Programme. Children attending the selected schools were eligible to be part of the study if they were between 6 and 16 years of age, had written informed consent from parents/caregivers and did not have any mental or severe physical

handicap. In each school, 132 children were randomly selected after stratification by sex and grade; hence 2640 children in total. One hundred and sixty-nine children were not recruited because they were absent on the day of data collection or refused to participate. Thus, a total of 2471 apparently healthy schoolchildren, aged 6–16 years, participated in the blood sample collection.

#### *Women of reproductive age from Cambodia*

The last data set was provided by a serological survey for antibodies for tetanus, rubella and measles conducted in women of reproductive age in Cambodia in 2012. Six hundred and eleven enumeration areas were selected from the 28 764 enumeration areas in the 2008 Cambodia General Population Census by probability proportional to size<sup>(17)</sup>. In each enumeration area, twenty-two households were randomly selected. All eligible women in selected households were invited to participate after providing written information about the survey and obtaining consent from women. In total, 2117 apparently healthy women participated in the blood sample collection. We used residual serum from this survey to determine FER, TfR, RBP, CRP and AGP concentrations.

### **Blood collection and laboratory analyses**

Venous blood samples (4–5 ml) were collected by experienced phlebotomists using sterile single-use material: into a vacutainer with heparin (Heparin Venosafe<sup>®</sup>; Terumo, Japan) in Senegal and in Cambodian women, and into a vacutainer with no anticoagulant (Vacuette<sup>®</sup>; Greiner Bio One, Austria) in Cambodian children, all trace-element-free. Blood samples were stored immediately in the field in an icebox containing ice-packs and transported to the laboratory within a maximum of 5 h after the first sample withdrawal. Plasma/serum samples were separated by centrifugation, aliquoted and stored at  $-20^{\circ}\text{C}$ . Centrifugation resulted in serum samples in Cambodian children and in plasma samples in Cambodian women and Senegalese children. However, all samples are referred to as 'plasma' herein for simplicity.

For plasma Zn measurement, samples collected from Senegalese schoolchildren were sent on dry ice to the Nutripass laboratory of the Institut de Recherche pour le Développement (IRD, Montpellier, France); and samples collected from Cambodian schoolchildren to the National Institute of Nutrition (NIN, Hanoi, Vietnam). Zn status was not measured in Cambodian women of reproductive age. In both laboratories, plasma Zn was measured by flame atomic absorption spectrophotometry, using trace-element-free procedures and certified controls.

Samples from the three studies were sent on dry ice to the VitMin Laboratory (Willstaett, Germany) for determination of RBP, CRP, FER, TfR and AGP concentrations. RBP, FER, TfR, CRP and AGP were measured by a sandwich ELISA technique<sup>(18)</sup>.

Inflammation was defined as elevated CRP (>5 mg/l) and/or elevated AGP (>1 g/l), allowing differentiation between incubation phase (high CRP and normal AGP), early convalescence phase (both high AGP and CRP) and late convalescence phase (high AGP and normal CRP)<sup>(19)</sup>.

Tissue Fe deficiency was defined as TfR > 8.3 mg/l<sup>(18)</sup> and depleted Fe stores was defined as FER < 15 µg/l<sup>(20)</sup>. ID was defined as depleted Fe stores (low FER) and/or tissue Fe deficiency (high TfR).

Body Fe was calculated according to the formula of Cook *et al.*<sup>(9)</sup>:  $\text{body Fe (mg/kg)} = -[\log(\text{TfR}/\text{FER}) - 2.8229]/0.1207$ . Body Fe deficiency (BID) was defined as body Fe < 0 mg/kg<sup>(8)</sup>.

A third way to evaluate ID as 'high TfR/log FER index', called ID-TFI herein, was defined using a cut-off of 7.06 ( $= 8.3/\log(15)$ )<sup>(9)</sup>.

Vitamin A status was determined by RBP concentration, which reflects plasma retinol concentration. RBP occurs in a 1:1:1 complex with retinol and transthyretin, which is assumed to be less affected by inflammation than retinol concentrations<sup>(21)</sup>. Vitamin A deficiency (VAD) was defined as RBP < 0.7 µmol/l<sup>(22)</sup>. Marginal VAD was defined as RBP value  $\geq 0.7$  µmol/l and < 1.05 µmol/l, as it was initially defined in adults<sup>(23,24)</sup>. Zn deficiency was defined as plasma Zn < 9.9 µmol/l, 10.1 µmol/l or 10.7 µmol/l for children aged < 10 years, girls aged > 10 years and boys aged > 10 years, respectively<sup>(25)</sup>. Children were considered severely Zn deficient when plasma Zn was < 7.7 µmol/l<sup>(14)</sup>.

### Statistical analysis

Data entry, including quality checks and validation by double entry of questionnaires, was performed with EpiData version 3.1 (EpiData, Odense, Denmark). Data management and analyses were performed using the statistical software package IBM SPSS Statistics version 20.0. Significance was defined as  $P < 0.05$ . The distributions of biomarker concentrations were checked for normality using normality plots and Kolmogorov–Smirnov tests. Because distributions of AGP, CRP, TfR and FER were skewed, they were log-transformed before statistical analysis. Spearman's rank correlation coefficients were determined to assess relationships between APP and micronutrient status biomarkers. For biomarkers that were correlated to APP concentrations, we calculated geometric mean values in each subgroup with inflammation (incubation, early convalescence, late convalescence, high AGP or high CRP) and in each reference group (no inflammation, normal AGP or normal CRP). We then calculated the ratio of the geometric mean value in the subgroup with inflammation to the mean value in the corresponding reference group without inflammation<sup>(5,19,26,27)</sup>. The correction factor (CF) was calculated as 1/ratio. Prevalence of poor micronutrient status was calculated: (i) without correction; (ii) only in the subjects with no inflammation;

(iii) using biomarker concentrations adjusted with CF calculated in the present study; (iv) using FER and RBP concentrations adjusted with CF recommended by Thurnham and co-workers<sup>(5,19)</sup>; and (v) using 30 µg/l as low FER cut-off in children with inflammation<sup>(8)</sup>. Corrected prevalences were compared with the uncorrected prevalences using McNemar's  $\chi^2$  test.

### Results

Biochemical characteristics of the participants from the three different samples are presented in Table 1. Although similar in gender proportions (half of children were female) and age (approximately 10 years), the populations of schoolchildren from Cambodia and Senegal differed in inflammatory and micronutrient status. Prevalence of inflammation was less than 15% in Senegalese children and Cambodian women but was 40% in Cambodian children, most of them being in the late convalescence phase (high AGP and normal CRP). ID was prevalent, being present in one in three children in the Senegalese study population, half of the children and in one in seven women in the Cambodian study populations. In Cambodian children, ID was mostly related to a high TfR (only 1% had low FER); while in Senegalese children, respectively one-third and one-fifth had a high TfR and a low FER. In Cambodian women, the proportion of high TfR was similar to the proportion of low FER (8–10%). Prevalence of VAD was less than 5% in all samples. Marginal vitamin A status was present in more than 40% of Senegalese children, while less than 15% of the Cambodian subjects had a marginal vitamin A status. Zn deficiency affected one-quarter of the children in Senegal and more than 90% of the Cambodian children.

Correlations between APP and biomarkers of Fe, vitamin A and Zn status are presented in Table 2. CRP and AGP were highly positively correlated in all samples. FER was highly positively correlated with both CRP and AGP in all samples. TfR was positively correlated with AGP in all samples, and with CRP in the Cambodian schoolchildren. Contradictory results were found between indicators based on the TfR/FER ratio (body Fe and TfR/log FER index) and APP among the different groups: TfR/log FER index was negatively correlated with CRP in both Senegalese children and Cambodian women, while this index was positively correlated with AGP in Cambodian children. Similarly, correlations between body Fe and AGP were positive in Senegalese children and Cambodian women, but negative in Cambodian children.

In most subjects, RBP was negatively correlated with CRP and AGP. In Cambodian schoolchildren, however, RBP was negatively associated with CRP but positively associated with AGP. No correlation between Zn and CRP or AGP was found in subgroups but in all subjects combined, Zn was negatively associated with CRP and AGP. For biomarkers that were significantly correlated with CRP

**Table 1** Demographic and biochemical characteristics of participants: schoolchildren from Senegal, schoolchildren from Cambodia and women of reproductive age (WRA) from Cambodia

	Senegalese schoolchildren (n 594)		Cambodian schoolchildren (n 2471)		Cambodian WRA (n 2117)	
	Mean	SD	Mean	SD	Mean	SD
Age (years)	10.2	2.4	9.6	2.4	26.3	7.0
Female (%)	52.5		50.3		100.0	
CRP (mg/l)	1.2	2.6	1.4	5.0	1.7	4.7
High CRP (>5 mg/l; %)	5.7		5.6		6.8	
AGP (g/l)	0.8	0.2	1.0	0.4	0.7	0.2
High AGP (>1 g/l; %)	10.6		39.9		10.3	
Inflammation status						
Inflammation (high CRP or high AGP; %)	12.1		40.3		14.1	
No inflammation (normal CRP and normal AGP; %)	87.9		59.7		85.9	
Incubation (high CRP and normal AGP; %)	1.5		0.4		3.8	
Early convalescence (high CRP and high AGP; %)	4.2		5.3		3.0	
Late convalescence (normal CRP and high AGP; %)	6.4		34.6		7.3	
FER (µg/l)	29.7	17.3	88.4	46.4	79.8	58.1
Low FER (<15 µg/l; %)	20.4		1.3		7.6	
TfR (mg/l)	8.1	3.1	8.8	2.7	6.1	2.6
High TfR (>8.3 mg/l; %)	33.3		51.4		10.1	
ID (low FER and/or high TfR; %)	38.7		51.6		13.5	
Body Fe (mg/kg)	2.6	2.9	6.1	2.2	6.8	3.7
BID (body Fe <0 mg/kg; %)	16.7		1.7		5.1	
TfR/log FER index	6.5	6.0	4.8	2.2	3.9	3.3
ID-TFI (TfR/log FER index > 7.06; %)	23.9		7.0		7.2	
RBP (µmol/l)	1.1	0.3	1.5	0.4	1.8	0.7
VAD (RBP <0.7 µmol/l; %)	3.7		1.0		0.5	
Marginal VAD (RBP < 1.05 µmol/l; %)	42.1		11.4		6.0	
Zn (µmol/l)	11.4	2.1	7.5	1.9	6.8	3.7
Zn deficiency† (%)	27.2		92.8		–	
Severe Zn deficiency‡ (%)	4.5		54.2		–	

CRP, C-reactive protein; AGP,  $\alpha_1$ -acid glycoprotein; FER, ferritin; TfR, soluble transferrin receptor; ID, Fe deficiency; BID, body Fe deficiency; ID-TFI, Fe deficiency defined by high TfR/log FER index; RBP, retinol-binding protein; VAD, vitamin A deficiency.

Data are presented as mean and standard deviation or as percentage.

†Zn < 9.9 µmol/l, 10.1 µmol/l or 10.7 µmol/l, respectively, in children aged <10 years, girls aged >10 years and boys aged >10 years.

‡Zn < 7.7 µmol/l.

and/or AGP (FER, TfR, RBP and Zn), CF were calculated for groups according to inflammatory status (Tables 3–6).

For indicators of Fe status, the effect of inflammation was stronger on FER than on TfR with all CF being lower than 0.8. Mean FER was significantly higher in participants in incubation, early convalescence and late convalescence phase, or with elevated AGP or CRP in all subjects (Table 3). Elevated CRP had no influence on TfR concentrations (Table 4), whereas elevated AGP was associated with higher TfR concentrations. TfR concentrations were significantly lower in all subjects combined during the early incubation phase, and higher in later stages of inflammation. The maximum inflammation effect on TfR concentrations in the later stages of inflammation was less than 25% (lowest CF being 0.78).

In children, but not in women, RBP concentrations were significantly lower when CRP was elevated (Table 5). During incubation, mean RBP concentrations were lower in children but higher in women. Concentrations were lower to the same extent in all populations during early convalescence.

In all children combined, Zn concentrations tended to be lower in early convalescence and were significantly lower during late convalescence and with elevated AGP (Table 6). These effects were not significant in the two subgroups of Senegalese children and Cambodian children.

The effect of correcting micronutrient biomarkers for inflammation on the estimated prevalence of deficiency is shown in Table 7. Uncorrected prevalence of high TfR was higher than the corrected prevalence in all samples combined, as well as in subgroups of Cambodian children and women (Table 7). Uncorrected prevalence of Fe stores deficiency was lower than the corrected prevalence, especially in Senegalese children and Cambodian women. The difference between the corrected and uncorrected prevalence of ID was lower than 2% in Senegalese children and Cambodian women, but about 8% in Cambodian children. BID and ID-TFI were less sensitive to inflammation. Prevalence of VAD was below 4% in all samples and difference between adjusted and unadjusted prevalence was below 1%.

**Table 2** Spearman correlation coefficients ( $\rho$ ) between inflammatory biomarkers and micronutrient status variables in Senegalese schoolchildren, Cambodian schoolchildren and Cambodian women of reproductive age (WRA)

	Overall (n 5182)	Senegalese schoolchildren (n 594)	Cambodian schoolchildren (n 2471)	Cambodian WRA (n 2117)
<b>CRP</b>				
AGP	0.44**	0.58**	0.62**	0.34**
FER	0.26**	0.27**	0.26**	0.24
TfR	-0.02	0.02	0.13**	-0.04
BI	0.16**	0.16**	0.03	0.16**
TFI	-0.12**	-0.11*	0.02	-0.13**
RBP	0.03*	-0.23**	-0.15**	0.14**
Zn	-0.08**	0.00	-0.01	-
<b>AGP</b>				
FER	0.30**	0.26**	0.35**	0.20
TfR	0.40**	0.09*	0.34**	0.16**
BI	-0.01	0.11*	-0.05*	0.08**
TFI	0.20**	-0.06	0.17**	0.01
RBP	-0.01	-0.11**	0.11**	-0.05*
Zn	-0.16**	-0.04	0.01	-

CRP, C-reactive protein; AGP,  $\alpha_1$ -acid glycoprotein; FER, ferritin; TfR, soluble transferrin receptor; BI, body Fe; TFI, TfR/fer index; RBP, retinol-binding protein.

\* $P < 0.05$ ; \*\* $P < 0.001$ .

## Discussion

Inflammation is known to affect indicators of Fe status (FER concentration) and vitamin A status (retinol and RBP concentrations), an effect we confirmed in the present study. Other indicators such as plasma Zn and TfR concentrations are thought to be affected as well, but these indicators have been less studied. In animal models, the fall in plasma Zn concentration occurs before clinical symptoms arise (hence just after infection) and returns to normal quickly<sup>(13)</sup>. A rapid fall in plasma Zn following infection was also observed in human subjects<sup>(28)</sup>. However, in our cohorts of schoolchildren in Cambodia and Senegal, inflammation affected plasma concentrations of Zn only slightly. The prevalence of Zn deficiency after adjustment was similar to the prevalence among children without inflammation. Duncan *et al.*<sup>(29)</sup> reported Zn concentrations of Scottish hospital patients to be significantly negatively correlated to CRP ( $\rho = -0.16$ ). However, in these hospital patients, the degree of inflammation was much more severe than in our normal populations with the causes of inflammation differing, and therefore the subjects in the Scottish study cannot be classified as having 'subclinical inflammation'. Indeed, in the study of Duncan *et al.*, Zn concentrations were not significantly lower in patients with CRP concentrations  $< 20$  g/l, a situation similar to ours. Therefore, our findings suggest that for determining Zn status in populations with a low prevalence of subclinical infection, taking the APR into account will not lead to a significant improvement in the estimate of the prevalence of Zn deficiency. This is consistent with the conclusions of an earlier review indicating no effect of inflammation on Zn status in children<sup>(3)</sup>.

To our knowledge, no correction factors for adjusting TfR concentration for inflammation have been suggested. TfR was positively correlated with AGP, as well as with CRP in Cambodian schoolchildren. However, TfR concentrations were higher only in groups in early and late convalescence phases. This is consistent with studies suggesting that FER reacts faster to the APR than TfR<sup>(15,30)</sup>. It is possible that TfR responds only indirectly to the APR, as a regulating response during convalescence. During the first phase of inflammation FER responds directly, regulated through higher hepcidin concentrations reducing circulating Fe and Fe absorption<sup>(31)</sup>. Then when the convalescence starts, the shortage of tissue Fe indicated by elevated TfR may result from the compensation of reduced circulating Fe. However, significantly lower TfR concentrations were observed in all subjects combined during the incubation phase, similar to reports by Righetti *et al.*<sup>(26)</sup>. Although TfR has been long considered more reliable than FER to indicate poor Fe status of populations when inflammation is prevalent<sup>(32)</sup>, TfR was significantly higher during inflammation phases in Cambodian women and children, as also found in other studies<sup>(33)</sup>. In our cohorts, TfR concentrations were higher during inflammation leading to an overestimation of ID when using TfR, which is again in line with results from Righetti *et al.*<sup>(26)</sup>. In the cohort of Cambodian children, who had a high prevalence of inflammation, the prevalence of high TfR (indicative of tissue Fe deficiency) was almost 10% higher, as compared with children without inflammation (51 *v.* 42%). Similar effects on TfR were reported for Ivorian infants who also had a high prevalence of inflammation<sup>(5-7)</sup>. However, when inflammation was not very prevalent, such as in the Cambodian women and Senegalese children, adjusting the prevalence of tissue Fe deficiency for inflammation did not modify the estimated prevalence of tissue Fe deficiency in these populations.

Use of TfR/log FER rather than TfR/fer<sup>(4)</sup> to predict ID has been recommended, especially in areas with endemic infection<sup>(7)</sup>. Instead of calculating CF to apply to TfR/log FER index as has been done by Righetti *et al.*<sup>(26)</sup>, we used corrected values of FER and TfR to calculate TfR/log FER index or body Fe, with prevalence of ID defined by TfR/log FER index  $> 7.06$  or by a negative value for body Fe. Depending on the biomarker for Fe status, that is whether ID was indicated primarily by low FER concentration (depleted Fe stores) or by elevated TfR concentration (tissue Fe deficiency), inflammation resulted in either over- or underestimation of the prevalence of ID defined by high TfR/log FER index (ID-TFI) or negative body Fe (BID). Overall, the difference of corrected prevalence between uncorrected prevalence and the reference prevalence in children without inflammation was low ( $< 2\%$ ), suggesting that indicators based on both TfR and FER were least sensitive to inflammation, compared with others. Thus, only when CRP and AGP cannot be measured do our findings suggest that indicators based on both TfR and FER, such as

**Table 3** Ferritin (FER) concentrations, ratios with 95% confidence intervals and correction factors (CF) by inflammatory status in Senegalese schoolchildren, Cambodian schoolchildren and Cambodian women of reproductive age (WRA)

	n	%	FER ( $\mu\text{g/l}$ )†		P value‡	Ratio§	95% CI	CF
			Mean	SD				
All subjects combined								
No inflammation	3817	74	53.7	2.2	—	—	—	—
Incubation	98	2	64.6	2.3	0.023	1.20	1.19, 1.21	0.83
Early convalescence	219	4	112.2	1.8	0.000	2.09	2.08, 2.09	0.48
Late convalescence	1048	20	83.2	1.9	0.000	1.55	1.55, 1.55	0.65
Senegalese schoolchildren								
No inflammation	522	88	23.5	1.9	—	—	—	—
Incubation	9	2	28.5	1.7	0.339	1.22	1.16, 1.26	0.82
Early convalescence	25	4	46.9	1.7	0.000	2.00	1.96, 2.03	0.50
Late convalescence	38	6	35.1	1.6	0.000	1.50	1.47, 1.52	0.67
Cambodian schoolchildren								
No inflammation	1476	60	68.9	1.7	—	—	—	—
Incubation	9	0	88.9	1.7	0.141	1.29	1.27, 1.31	0.77
Early convalescence	130	5	125.8	1.6	0.000	1.83	1.82, 1.83	0.55
Late convalescence	856	35	87.5	1.7	0.000	1.27	1.27, 1.27	0.79
Cambodian WRA								
No inflammation	1819	86	56.4	2.3	—	—	—	—
Incubation	80	4	67.9	2.4	0.051	1.20	1.19, 1.21	0.83
Early convalescence	64	3	120.4	1.7	0.000	2.14	2.13, 2.14	0.47
Late convalescence	154	7	73.8	2.4	0.000	1.31	1.30, 1.32	0.76
All subjects combined								
Normal AGP	3915	76	54.3	2.2	—	—	—	—
High AGP	1267	24	86.9	1.9	0.000	1.60	1.60, 1.60	0.62
Senegalese schoolchildren								
Normal AGP	531	89	23.5	1.9	—	—	—	—
High AGP	63	11	38.9	1.7	0.000	1.65	1.63, 1.68	0.60
Cambodian schoolchildren								
Normal AGP	1485	60	69.0	1.7	—	—	—	—
High AGP	986	40	91.2	1.7	0.000	1.32	1.32, 1.32	0.76
Cambodian WRA								
Normal AGP	1899	90	56.8	2.3	—	—	—	—
High AGP	218	10	85.1	2.3	0.000	1.50	1.49, 1.5	0.67
All subjects combined								
Normal CRP	4865	94	59.2	2.2	—	—	—	—
High CRP	317	6	93.7	2.0	0.000	1.58	1.58, 1.59	0.63
Senegalese schoolchildren								
Normal CRP	560	94	24.1	1.9	—	—	—	—
High CRP	34	6	41.1	1.8	0.000	1.71	1.68, 1.73	0.59
Cambodian schoolchildren								
Normal CRP	2332	94	74.1	1.0	—	—	—	—
High CRP	139	6	123.0	1.6	0.000	1.66	1.66, 1.66	0.60
Cambodian WRA								
Normal CRP	1973	93	57.6	2.3	—	—	—	—
High CRP	144	7	87.1	2.2	0.000	1.51	1.51, 1.52	0.66

AGP,  $\alpha_1$ -acid glycoprotein; CRP, C-reactive protein.

†Geometric means and SD.

‡ANOVA on log-transformed FER means of positive group v. control group.

§Ratio of back-transformed FER concentrations of positive group v. control group.

||CF = 1/ratio.

TfR/log FER index, may be used to assess Fe status and ID prevalence of populations with a high prevalence of inflammation. However, assessment of Fe store levels and tissue Fe levels would still be affected by inflammation.

The adjusted prevalence of ID using our CF for both FER and TfR was very close to that obtained using Thurnham's CF for FER in Cambodian women and Senegalese children, suggesting that in these cohorts adjusting FER concentrations for inflammation is more important than adjusting TfR for inflammation to obtain a more precise estimate of ID. However, using Thurnham's CF for FER only in the Cambodian children, without adjusting for TfR, led to

overestimation of ID defined by high TfR and/or low FER, because in this sample FER concentrations were high, making TfR the most important factor for indicating ID.

Not adjusting FER concentrations for inflammation can lead to underestimation of the prevalence of low Fe stores, especially in populations with endemic inflammation. Thurnham's CF were shown to be accurate to adjust FER status<sup>(34)</sup>. Also in the present study, the prevalence of depleted Fe stores in the three samples adjusted with either Thurnham's CF or ours was similar to the prevalence of depleted Fe stores in subgroups with no inflammation. The small effect of adjusting the prevalence

**Table 4** Soluble transferrin receptor (TfR) concentrations, ratios with 95 % confidence intervals and correction factors (CF) by inflammatory status in Senegalese schoolchildren, Cambodian schoolchildren and Cambodian women of reproductive age (WRA)

	<i>n</i>	%	TfR (mg/l)†		<i>P</i> value‡	Ratio§	95 % CI	CF
			Mean	SD				
All subjects combined								
No inflammation	3817	74	6.8	1.4	–	–	–	–
Incubation	98	2	5.8	1.3	0.000	0.85	0.81, 0.89	1.17
Early convalescence	219	4	7.8	1.4	0.000	1.16	1.12, 1.18	0.87
Late convalescence	1048	20	8.7	1.3	0.000	1.29	1.27, 1.29	0.78
Senegalese schoolchildren								
No inflammation	522	88	7.7	1.3	–	–	–	–
Incubation	9	2	7.5	7.7	0.746	0.97	0.32, 1.63	1.03
Early convalescence	25	4	8.2	1.3	0.266	1.07	1.00, 1.13	0.94
Late convalescence	38	6	7.9	1.2	0.653	1.02	0.97, 1.08	0.98
Cambodian schoolchildren								
No inflammation	1476	60	8.0	1.3	–	–	–	–
Incubation	9	0	7.2	1.3	0.206	0.90	0.79, 1.01	1.12
Early convalescence	130	5	8.8	1.3	0.000	1.10	1.07, 1.13	0.91
Late convalescence	856	35	9.3	1.3	0.000	1.15	1.15, 1.18	0.87
Cambodian WRA								
No inflammation	1819	86	5.7	1.4	–	–	–	–
Incubation	80	4	5.5	1.3	0.296	0.96	0.91, 1.02	1.04
Early convalescence	64	3	6.0	1.4	0.145	1.06	0.99, 1.11	0.94
Late convalescence	154	7	6.4	1.4	0.000	1.13	1.08, 1.16	0.89
All subjects combined								
Normal AGP	3915	76	6.8	1.4	–	–	–	–
High AGP	1267	24	8.6	1.4	0.000	1.27	1.25, 1.28	0.79
Senegalese schoolchildren								
Normal AGP	531	89	7.7	1.3	–	–	–	–
High AGP	63	11	8.0	1.3	0.301	1.04	0.99, 1.08	0.96
Cambodian schoolchildren								
Normal AGP	1485	60	8.0	1.3	–	–	–	–
High AGP	986	40	9.2	1.3	0.000	1.15	1.14, 1.16	0.87
Cambodian WRA								
Normal AGP	1899	90	5.7	1.4	–	–	–	–
High AGP	218	10	6.3	1.4	0.000	1.11	1.07, 1.14	0.90
All subjects combined								
Normal CRP	4865	94	7.2	1.4	–	–	–	–
High CRP	317	6	7.1	1.4	0.847	1.00	0.96, 1.01	1.00
Senegalese schoolchildren								
Normal CRP	560	94	7.7	1.3	–	–	–	–
High CRP	34	6	8.0	1.3	0.449	1.04	0.98, 1.10	0.96
Cambodian schoolchildren								
Normal CRP	2332	94	8.5	1.3	–	–	–	–
High CRP	139	6	8.7	1.3	0.238	1.03	1.00, 1.05	0.97
Cambodian WRA								
Normal CRP	1973	93	5.7	1.4	–	–	–	–
High CRP	144	7	5.7	1.4	0.170	1.00	0.96, 1.04	1.00

AGP,  $\alpha_1$ -acid glycoprotein; CRP, C-reactive protein.

†Geometric means and SD.

‡ANOVA on log-transformed TfR means of positive group *v.* control group.§Ratio of back-transformed TfR concentrations of positive group *v.* control group.

||CF = 1/ratio.

of depleted Fe stores for inflammation in the three populations in our study is readily explained either by the low prevalence of inflammation in Cambodian women and Senegalese children or by the low prevalence of low Fe stores in Cambodian children and women. Hence, inflammation only had a large impact on the estimate of the prevalence of low Fe stores when both the prevalence of inflammation and the prevalence of low Fe stores were high. Overall, using 30  $\mu\text{g/l}$  as a cut-off for FER in children with inflammation as suggested by WHO<sup>(35)</sup> performed less efficiently than adjustment for inflammation by use of APP for accurate prevalence of ID.

In children from Senegal and Cambodia, RBP was lower in the incubation and early convalescence phases, which is consistent with other studies<sup>(5–7)</sup>. Indeed, retinol is known to fall rapidly, within 48 h of the onset of inflammation<sup>(36)</sup>. During late convalescence, RBP was still lower in Senegalese children but higher in Cambodian children compared with participants with no inflammation; the latter has been reported earlier, in a small sample of Zambian schoolchildren<sup>(37)</sup>. A meta-analysis using the same CRP and AGP cut-offs as in the present study showed that the APR-induced decrease of retinol lasted during the late convalescence phase<sup>(5)</sup>. However, this

**Table 5** Retinol-binding protein (RBP) concentrations, ratios with 95% confidence intervals and correction factors (CF) in Senegalese schoolchildren, Cambodian schoolchildren and Cambodian women of reproductive age (WRA)

	<i>n</i>	%	RBP ( $\mu\text{mol/l}$ )†		<i>P</i> value‡	Ratio§	95% CI	CF
			Mean	SD				
All subjects combined								
No inflammation	3817	74	1.6	0.6	—	—	—	—
Incubation	98	2	1.8	0.7	0.000	0.88	0.81, 0.95	0.88
Early convalescence	219	4	1.3	0.4	0.000	0.83	0.79, 0.86	1.21
Late convalescence	1048	20	1.6	0.5	0.996	1.00	0.98, 1.02	1.00
Senegalese schoolchildren								
No inflammation	522	88	1.1	0.3	—	—	—	—
Incubation	9	2	0.9	0.2	0.002	0.76	0.64, 0.88	1.31
Early convalescence	25	4	1.0	0.3	0.003	0.86	0.75, 0.97	1.16
Late convalescence	38	6	1.1	0.3	0.092	0.94	0.85, 1.03	1.07
Cambodian schoolchildren								
No inflammation	1476	60	1.5	0.4	—	—	—	—
Incubation	9	0	1.3	0.5	0.090	0.85	0.63, 1.07	1.17
Early convalescence	130	5	1.3	0.3	0.000	0.85	0.82, 0.89	1.17
Late convalescence	856	35	1.6	0.4	0.000	1.05	1.03, 1.08	0.95
Cambodian WRA								
No inflammation	1819	86	1.8	0.7	—	—	—	—
Incubation	80	4	2.0	0.6	0.015	1.09	1.02, 1.17	0.91
Early convalescence	64	3	1.5	0.6	0.002	0.85	0.77, 0.94	1.17
Late convalescence	154	7	1.8	0.7	0.736	1.01	0.94, 1.07	0.99
All subjects combined								
Normal AGP	3915	76	1.6	0.6	—	—	—	—
High AGP	1267	24	1.5	0.5	0.003	0.97	0.93, 0.95	1.03
Senegalese schoolchildren								
Normal AGP	531	89	1.1	0.3	—	—	—	—
High AGP	63	11	1.0	0.3	0.003	0.91	0.84, 0.98	1.10
Cambodian schoolchildren								
Normal AGP	1485	60	1.5	0.4	—	—	—	—
High AGP	986	40	1.5	0.4	0.014	1.03	1.01, 1.05	0.97
Cambodian WRA								
Normal AGP	1899	90	1.8	0.7	—	—	—	—
High AGP	218	10	1.7	0.6	0.136	0.96	0.91, 1.01	1.04
All subjects combined								
Normal CRP	4865	94	1.6	0.5	—	—	—	—
High CRP	317	6	1.5	0.6	0.000	0.92	0.90, 0.98	1.08
Senegalese schoolchildren								
Normal CRP	560	94	1.1	0.3	—	—	—	—
High CRP	34	6	0.9	0.3	0.000	0.83	0.74, 0.92	1.20
Cambodian schoolchildren								
Normal CRP	2332	94	1.5	0.4	—	—	—	—
High CRP	139	6	1.3	0.4	0.000	0.85	0.80, 0.89	1.18
Cambodian WRA								
Normal CRP	1973	93	1.8	0.7	—	—	—	—
High CRP	144	7	1.8	0.7	0.804	0.99	0.92, 1.06	1.01

AGP,  $\alpha_1$ -acid glycoprotein; CRP, C-reactive protein.

†Geometric means and SD.

‡ANOVA on log-transformed RBP means of positive group *v.* control group.§Ratio of back-transformed RBP concentrations of positive group *v.* control group.

||CF = 1/ratio.

meta-analysis was conducted in populations with a poor vitamin A status (median retinol < 1.05  $\mu\text{mol/l}$ ). Retinol (and hence RBP) concentrations are expected to rebound during convalescence<sup>(13)</sup>. We believe that baseline vitamin A status might explain the difference between Senegalese and Cambodian children. Forty per cent of the Senegalese children had marginal vitamin A status, whereas vitamin A status in Cambodian children was adequate. Therefore, the reaction of vitamin A biomarkers to the different phases of inflammation may differ according to the initial vitamin A status, suggesting that baseline vitamin A status should be taken into account when correcting for

inflammation. The interaction between inflammation and initial vitamin A status and its effect on the assessment of vitamin A status should be further investigated. More data on populations with a relatively good vitamin A status are needed to confirm this. But if found correct, it could affect the way to correct vitamin A status for inflammation. In Senegalese children, the CF proposed by Thurnham (and calculated using populations with poor vitamin A status) resulted in a prevalence of VAD very close to the prevalence in children without inflammation. In Cambodian subjects, however, the CF proposed by Thurnham led to a significant but slight underestimation of the prevalence of



**Table 6** Zinc concentrations, ratios with 95% confidence intervals and correction factors (CF) in Senegalese schoolchildren, Cambodian schoolchildren and Cambodian women of reproductive age (WRA)

	<i>n</i>	%	Zn ( $\mu\text{mol/l}$ )†		<i>P</i> value‡	Ratio§	95% CI	CF
			Mean	SD				
All subjects combined								
No inflammation	1716	67	8.4	1.3	–	–	–	–
Incubation	15	1	9.2	1.4	0.219	1.09	0.98, 1.11	0.92
Early convalescence	120	5	8.0	1.3	0.053	0.95	0.85, 1.00	1.05
Late convalescence	698	27	7.7	1.3	0.000	0.91	0.95, 0.97	1.09
Senegalese schoolchildren								
No inflammation	1203	88	11.2	7.4	–	–	–	–
Incubation	7	2	11.7	1.1	0.512	1.05	0.96, 1.08	0.95
Early convalescence	95	4	10.8	1.3	0.340	0.96	0.95, 1.02	1.04
Late convalescence	662	6	11.3	1.2	0.703	1.01	0.98, 1.03	0.99
Cambodian schoolchildren								
No inflammation	513	60	7.4	1.2	–	–	–	–
Incubation	8	0	6.9	1.2	0.399	0.93	0.88, 1.05	1.07
Early convalescence	25	5	7.4	1.3	0.795	1.00	0.98, 1.02	1.00
Late convalescence	36	35	7.5	1.3	0.278	1.01	1.00, 1.02	0.99
All children combined								
Normal AGP	1731	68	8.4	1.3	–	–	–	–
High AGP	818	32	7.7	1.3	0.000	0.92	0.91, 0.93	1.09
Senegalese schoolchildren								
Normal AGP	521	90	11.2	1.2	–	–	–	–
High AGP	61	10	11.1	1.2	0.740	0.99	0.98, 1.02	1.01
Cambodian schoolchildren								
Normal AGP	1210	62	7.4	1.2	–	–	–	–
High AGP	757	38	7.5	1.3	0.339	1.01	1.00, 1.01	0.99
All children combined								
Normal CRP	135	5	8.2	1.3	–	–	–	–
High CRP	2414	95	8.1	1.3	0.712	0.99	0.96, 1.02	1.01
Senegalese schoolchildren								
Normal CRP	549	94	11.2	1.2	–	–	–	–
High CRP	33	6	11.0	1.2	0.593	0.98	0.98, 1.06	1.02
Cambodian schoolchildren								
Normal CRP	1865	95	7.5	1.2	–	–	–	–
High CRP	102	5	7.4	1.3	0.514	0.99	0.98, 1.05	1.01

AGP,  $\alpha_1$ -acid glycoprotein; CRP, C-reactive protein.

†Geometric means and sd.

‡ANOVA on log-transformed Zn means of positive group *v.* control group.§Ratio of back-transformed Zn concentrations of positive group *v.* control group.

||CF = 1/ratio.

VAD compared with adjusting by using the CF calculated in the present study, presumably because vitamin A concentrations rebounded earlier in this population.

Prevalence of inflammation is considered low under 15% and to have only a modest impact on the assessment of micronutrient deficiencies<sup>(38,39)</sup>. However, consensus about cut-offs to define what should be considered an elevated APP is needed. Indeed, the sensitivity of the <5 mg/l cut-off commonly used for CRP has recently been questioned<sup>(40)</sup>. Several studies used a cut-off different from 5 mg/l for CRP and/or 1 g/l for AGP, or a series of cut-offs to define categories of inflammation<sup>(38)</sup>. Some studies showed similar patterns of impact of the APR on micronutrient biomarkers using either 5 or 10 mg/l for the CRP cut-off<sup>(41)</sup>, while one study indicated that the best CRP cut-off could vary from 5, 10 to 20 mg/l depending on the measured biomarker<sup>(42)</sup>. Moreover, different cut-offs for different ages may be needed<sup>(5)</sup>. Finally, the definition of 'apparently healthy individuals' might have differed between researchers involved in the different studies.

Hence, an international recommendation about APP cut-offs to categorize inflammation is needed as well as a meta-analysis on the impact of inflammation on a broad range of micronutrient biomarkers.

Our study had several limitations. First, the Fe status of populations was heterogeneous with Cambodian children having adequate Fe stores but a high proportion of high TfR, while ID in Cambodian women and in Senegalese children was both due to low Fe stores and Fe tissue deficiency. However, this gave us the opportunity to examine the impact of inflammation on micronutrient status in different population profiles and to bring out original observations compared with previous research. Second, the prevalence of inflammation in Senegalese children and Cambodian women was low, reducing the power of the study to investigate the impact of inflammation. Third, the sample size of Senegalese children was small, hence more data should be gathered in African populations on the impact of inflammation on indicators of micronutrient status.

**Table 7** Effect of correcting ferritin (FER), soluble transferrin receptor (TfR) and retinol-binding protein (RBP) concentrations on the prevalence of low iron status and low vitamin A status in Senegalese schoolchildren, Cambodian schoolchildren and Cambodian women of reproductive age (WRA)

	Senegalese schoolchildren (n 594)		Cambodian schoolchildren (n 2471)		Cambodian WRA (n 2117)	
	%	P value†	%	P value†	%	P value†
Low FER‡						
Uncorrected	20.4	NA	1.3	NA	7.6	NA
In participants with no inflammation	22.6	NA	1.4	NA	8.0	NA
Corrected for three phases of inflammation§	21.5	0.016	1.4	0.250	7.9	0.008
Corrected for three phases of inflammation (Thurnham)	22.9	0.000	1.8	0.000	8.0	0.002
FER < 30 µg/l in children with inflammation	23.2	0.000	2.1	0.000	8.9	0.000
High TfR¶						
Uncorrected	33.3	NA	51.4	NA	10.1	NA
In participants with no inflammation	32.6	NA	42.2	NA	9.5	NA
Corrected for three phases of inflammation††	33.2	1.000	42.9	0.000	9.7	0.021
ID (high TfR and/or low FER)‡,¶						
Uncorrected	38.7	NA	51.6	NA	13.5	NA
In participants with no inflammation	38.5	NA	42.5	NA	13.2	NA
Corrected for three phases of inflammation§,††	37.9	0.180	44.5	0.000	13.4	0.791
Corrected for three phases of inflammation (Thurnham)	39.1	0.500	51.6	1.000	13.8	0.016
FER < 30 µg/l in children with inflammation	40.2	0.004	51.7	0.500	14.3	0.000
BID (low body Fe‡‡)						
Uncorrected	16.7	NA	1.7	NA	5.1	NA
In participants with no inflammation	18.6	NA	1.7	NA	5.3	NA
Corrected for three phases of inflammation§,††	17.5	0.002	1.8	0.002	5.2	0.002
Corrected for three phases of inflammation (Thurnham)	17.7	0.000	1.9	0.000	5.3	0.000
ID-TFI (high TfR/log FER index§§)						
Uncorrected	23.9	NA	7.0	NA	7.2	NA
In participants with no inflammation	25.7	NA	6.3	NA	7.1	NA
Corrected for three phases of inflammation§,††	24.6	0.125	6.3	0.000	7.2	1.000
Corrected for three phases of inflammation (Thurnham)	24.6	0.125	7.7	0.000	7.3	0.500
FER < 30 µg/l for children with inflammation	26.9	0.000	13.2	0.000	7.9	0.000
Vitamin A deficiency¶¶						
Uncorrected	3.7	NA	1.0	NA	0.5	NA
In participants with no inflammation	2.9	NA	0.9	NA	0.4	NA
Corrected for three phases of inflammation†††	3.4	0.500	0.9	1.000	0.4	0.500
Corrected for three phases of inflammation (Thurnham)‡‡‡	3.0	0.125	0.7	0.031	0.3	0.250
Marginal vitamin A status§§§						
Uncorrected	42.1	NA	11.4	NA	6.0	NA
In participants with no inflammation	39.7	NA	11.0	NA	5.7	NA
Corrected for three phases of inflammation†††	39.6	0.000	11.7	0.427	5.9	0.687
Corrected for three phases of inflammation (Thurnham)‡‡‡	38.9	0.000	8.7	0.000	5.6	0.008
Zn deficiency						
Uncorrected	27.2	NA	92.8	NA	NA	NA
In participants with no inflammation	27.5	NA	93.3	NA	NA	NA
Corrected for three phases of inflammation¶¶¶¶	27.3	1.0	92.8	0.5	NA	NA

ID, Fe deficiency; BID, body Fe deficiency; ID-TFI, Fe deficiency defined by high TfR/log FER index; NA, not applicable; CF, correction factor.  
 †McNemar's  $\chi^2$  test of proportion to compare uncorrected prevalence and corrected prevalence.  
 ‡FER < 15 µg/l for children and WRA, except if notified.  
 §Using CF in Table 3.  
 || Thurnham's CF for incubation, early convalescence and late convalescence: 0.64, 0.39 and 0.65 in children; 0.73, 0.58 and 0.85 in women.  
 ¶TfR > 8.3 mg/l.  
 ††Using CF in Table 4.  
 ‡‡Body Fe < 0 mg/kg.  
 §§TfR/logFER > 7.05 (corresponding to 8.3/log15).  
 ||| TfR/logFER > 5.3 (corresponding to 8.3/log30).  
 ¶¶RBP < 0.7 µmol/l.  
 †††Using CF in Table 5.  
 ‡‡‡Thurnham's CF for incubation, early convalescence and late convalescence: 0.87, 0.76 and 0.89.  
 §§§RBP < 1.05 µmol/l.  
 |||| Zn < 9.9 µmol/l, 10.1 µmol/l or 10.7 µmol/l, respectively, in children aged <10 years, girls aged >10 years and boys aged >10 years.  
 ¶¶¶¶Using CF in Table 6.

**Conclusion**

Inflammation affected the biomarkers of vitamin A and Fe status, but had little impact on the biomarker for Zn status in the populations studied. This suggests that ideally FER,

TfR and RBP should be adjusted for inflammation measured by both CRP and AGP to differentiate the different phases of inflammation, especially in areas with a high infection pressure. We also proposed a correcting factor for TfR that has to be confirmed, ideally through a

large meta-analysis or pooled analyses such as those carried out on retinol and FER<sup>(5,19)</sup>. Divergent responses of RBP to incubation and convalescence phases were observed according to the studied populations, presumably depending on the vitamin A status of the population. More research should be conducted on APP cut-offs to define the different phases of inflammation and the sensitivity of micronutrient biomarkers, especially TfR and RBP, to inflammation in relation to the underlying micronutrient status.

## Acknowledgements

**Acknowledgements:** The authors thank all the children and their families from Senegal and Cambodia, and the women from Cambodia, who participated into the study. They would like to thank Richard de Groot for his help with statistical analysis. **Financial support:** The study on Cambodian children was funded by the US Department of Agriculture/Foreign Agricultural Service (USDA/FAS), the UN World Food Programme (WFP)–DSM consortium and the Institut de Recherche pour le Développement (IRD). The USDA/FAS and the WFP–DSM consortium had no role in the design, data analysis or writing of this article. The study on Cambodian women was funded by the Centers for Disease Control and Prevention (CDC). The CDC had no role in the design, data analysis or writing of this article. The study in Senegal was funded by Danone. Danone had no role in the design, data analysis or writing of this article. **Conflict of interest:** None. **Authorship:** M.F., F.T.W. and J.B. formulated the research questions. M.F., F.T.W., M.P., K.K. and C.C. designed and carried out the study in Cambodian children, and M.F. and J.B. the study in Senegalese children. M.F. and J.B. designed and M.F. carried out the data analysis. M.F., F.T.W. and J.B. and wrote the article that was reviewed and agreed by all co-authors. **Ethics of human subject participation:** The study was conducted according to the guidelines laid down in the Declaration of Helsinki. All procedures involving human subjects were approved by the National Health Research of Senegal (for the study in Senegal); by the National Ethics Committee for Health Research (NECHR) of the Ministry of Health, Phnom Penh, Cambodia (for the study in Cambodian children and in Cambodian women); by the Ministry of Education, Youth and Sports, Phnom Penh, Cambodia and the Research Ethical Committee (REC) of PATH, Seattle, WA, USA (for the study in Cambodian children); and by the CDC, Atlanta, GA, USA (for the study in Cambodian women). Written informed consent was obtained from all women enrolled in the study and from all parents of children enrolled in the study.

## References

- Black RE, Victora CG, Walker SP *et al.* (2013) Maternal and child undernutrition and overweight in low-income and middle-income countries. *Lancet* **382**, 427–451.
- Delisle HF (2008) Poverty: the double burden of malnutrition in mothers and the intergenerational impact. *Ann N Y Acad Sci* **1136**, 172–184.
- Thurnham DI, Mburu AS, Mwaniki DL *et al.* (2005) Micronutrients in childhood and the influence of subclinical inflammation. *Proc Nutr Soc* **64**, 502–509.
- Thurnham DI (2014) Interactions between nutrition and immune function: using inflammation biomarkers to interpret micronutrient status. *Proc Nutr Soc* **73**, 1–8.
- Thurnham DI, McCabe GP, Northrop-Clewes CA *et al.* (2003) Effects of subclinical infection on plasma retinol concentrations and assessment of prevalence of vitamin A deficiency: meta-analysis. *Lancet* **362**, 2052–2058.
- Wieringa FT, Dijkhuizen MA, West CE *et al.* (2002) Estimation of the effect of the acute phase response on indicators of micronutrient status in Indonesian infants. *J Nutr* **132**, 3061–3066.
- Rosales FJ, Ritter S, Zolfaghari R *et al.* (1996) Effects of acute inflammation on plasma retinol, retinol-binding protein, and its mRNA in the liver and kidneys of vitamin A-sufficient rats. *J Lipid Res* **37**, 962–971.
- World Health Organization (2007) *Assessing the Iron Status of Populations: Including Literature Reviews. Report of a Joint World Health Organization/Centers for Disease Control and Prevention Technical Consultation on the Assessment of Iron Status at the Population Level, Geneva, Switzerland, 6–8 April 2004*. Geneva: WHO/CDC.
- Cook JD, Flowers CH & Skikne BS (2003) The quantitative assessment of body iron. *Blood* **101**, 3359–3363.
- Beguín Y (2003) Soluble transferrin receptor for the evaluation of erythropoiesis and iron status. *Clin Chim Acta* **329**, 9–22.
- Phiri K, Calis J, Siyasiya A *et al.* (2009) New cut-off values for ferritin and soluble transferrin receptor for the assessment of iron deficiency in children in a high infection pressure area. *J Clin Pathol* **62**, 1103–1106.
- Punnonen K, Irjala K & Rajamäki A (1997) Serum transferrin receptor and its ratio to serum ferritin in the diagnosis of iron deficiency. *Blood* **89**, 1052–1057.
- Brown KH (1998) Effect of infections on plasma zinc concentration and implications for zinc status assessment in low-income countries. *Am J Clin Nutr* **68**, 2 Suppl., 425S–429S.
- Brown KH, Rivera JA, Bhutta Z *et al.* (2004) International Zinc Nutrition Consultative Group (IZiNCG) technical document #1. Assessment of the risk of zinc deficiency in populations and options for its control. *Food Nutr Bull* **25**, 1 Suppl. 2, S99–S203.
- Fiorentino M, Bastard G, Sembène M *et al.* (2013) Anthropometric and micronutrient status of school-children in an urban West Africa setting: a cross-sectional study in Dakar (Senegal). *PLoS One* **8**, e84328.
- Perignon M, Fiorentino M, Kuong K *et al.* (2016) Impact of multi-micronutrient fortified rice on hemoglobin, iron and vitamin A status of Cambodian schoolchildren: a double-blind cluster-randomized controlled trial. *Nutrients* **8**, 29.
- Scobie HM, Mao B, Buth S *et al.* (2016) Tetanus immunity among women aged 15 to 39 years in Cambodia: a national population-based serosurvey, 2012. *Clin Vaccine Immunol* **23**, 546–554.
- Erhardt JG, Estes JE, Pfeiffer CM *et al.* (2004) Combined measurement of ferritin, soluble transferrin receptor, retinol binding protein, and C-reactive protein by an inexpensive, sensitive, and simple sandwich enzyme-linked immunosorbent assay technique. *J Nutr* **134**, 3127–3132.
- Thurnham DI, McCabe LD, Haldar S *et al.* (2010) Adjusting plasma ferritin concentrations to remove the effects of subclinical inflammation in the assessment of iron deficiency: a meta-analysis. *Am J Clin Nutr* **92**, 546–555.

20. World Health Organization (2011) *Serum Ferritin Concentrations for the Assessment of Iron Status and Iron Deficiency in Populations*. Geneva: WHO.
21. de Pee S & Dary O (2002) Biochemical indicators of vitamin A deficiency: serum retinol and serum retinol binding protein. *J Nutr* **132**, 9 Suppl., 2895S–2901S.
22. Engle-Stone R, Haskell MJ, Ndjebayi AO *et al.* (2011) Plasma retinol-binding protein predicts plasma retinol concentration in both infected and uninfected Cameroonian women and children. *J Nutr* **141**, 2233–2241.
23. West KP Jr (2002) Extent of vitamin A deficiency among preschool children and women of reproductive age. *J Nutr* **132**, Suppl. 9, 2857S–2866S.
24. Tanumihardjo SA (2004) Assessing vitamin A status: past, present and future. *J Nutr* **134**, issue 1, 290S–293S.
25. Gibson RS (1990) *Principles of Nutritional Assessment*. New York: Oxford University Press.
26. Righetti AA, Glinz D, Adiossan LG *et al.* (2012) Interactions and potential implications of *Plasmodium falciparum*–hookworm coinfection in different age groups in south-central Côte d'Ivoire. *PLoS Negl Trop Dis* **6**, e1889.
27. Grant FK, Suchdev PS, Flores-Ayala R *et al.* (2012) Correcting for inflammation changes estimates of iron deficiency among rural Kenyan preschool children. *J Nutr* **142**, 105–111.
28. Beisel WR (1976) Trace element in infectious processes. *Med Clin North Am* **60**, 831–849.
29. Duncan A, Talwar D, McMillan DC *et al.* (2012) Quantitative data on the magnitude of the systemic inflammatory response and its effect on micronutrient status based on plasma measurements. *Am J Clin Nutr* **95**, 64–71.
30. Perignon M, Fiorentino M, Kuong K *et al.* (2014) Stunting, poor iron status and parasite infection are significant risk factors for lower cognitive performance in Cambodian school-aged children. *PLoS One* **9**, e112605.
31. Wessling-Resnick M (2010) Iron homeostasis and the inflammatory response. *Annu Rev Nutr* **30**, 105–122.
32. Bui VQ, Stein AD, DiGirolamo AM *et al.* (2012) Associations between serum C-reactive protein and serum zinc, ferritin, and copper in Guatemalan school children. *Biol Trace Elem Res* **148**, 154–160.
33. Black R (2003) Micronutrient deficiency – an underlying cause of morbidity and mortality. *Bull World Health Organ* **81**, 79.
34. Cichon B, Ritz C, Fabiansen C *et al.* (2017) Assessment of regression models for adjustment of iron status biomarkers for inflammation in children with moderate acute malnutrition in Burkina Faso. *J Nutr* **147**, 125–132.
35. World Health Organization (2001) *Iron Deficiency Anaemia Assessment, Prevention and Control. A Guide for Programme Managers*. Geneva: WHO.
36. Goodman MN (1991) Tumor necrosis factor induces skeletal muscle protein breakdown in rats. *Am J Physiol* **260**, E727–E730.
37. Bresnahan K, Chileshe J & Tanumihardjo S (2014) Acute phase response is associated with decreased nutrient intake and altered markers of micronutrient status in Zambian children (624.16). *FASEB J* **28**, Suppl. 1, 624.616.
38. Beisel WR (1998) Infection-induced depression of serum retinol – a component of the acute phase response or a consequence? *Am J Clin Nutr* **68**, 993–994.
39. Mitra AK, Alvarez JO, Guay-Woodford L *et al.* (1998) Urinary retinol excretion and kidney function in children with shigellosis. *Am J Clin Nutr* **68**, 1095–1103.
40. Dillon D, Htet MK & Chiwile FP (2010) The application of correction factors on serum retinol of Indonesia school children. *Med J Indones* **19**, 258–263.
41. Hotz C, Chileshe J, Siamusantu W *et al.* (2012) Vitamin A intake and infection are associated with plasma retinol among pre-school children in rural Zambia. *Public Health Nutr* **15**, 1688–1696.
42. Wieringa F, Dijkhuizen M, West C *et al.* (2004) Reduced production of immunoregulatory cytokines in vitamin A- and zinc-deficient Indonesian infants. *Eur J Clin Nutr* **58**, 1498–1504.