

Hypertension and diabetes prevalence among adults with moderately increased BMI (23.0–24.9 kg/m²): findings from a nationwide survey in Bangladesh

Muntasirur Rahman*, Gail Williams and Abdullah Al Mamun

School of Public Health, The University of Queensland, Level 2, Public Health Building, Herston, QLD 4006, Australia

Submitted 29 June 2016: Final revision received 24 November 2016: Accepted 30 November 2016: First published online 23 January 2017

Abstract

Objective: BMI is a proxy for fat accumulation in the body. Increased diabetes and CVD risks have been observed for Asian populations at lower BMI than the WHO-recommended BMI cut-off points for overweight (≥ 25.0 kg/m²) and obesity (≥ 30.0 kg/m²). The current study aimed to quantify the increased hypertension (HTN) and type 2 diabetes mellitus (T2DM) prevalence in Bangladeshi adults with moderately increased BMI (23.0–24.9 kg/m²).

Design: Data from the most recent Bangladesh Demographic and Health Survey (2011) were analysed. Modified Poisson regression models with robust error variance were used to calculate prevalence ratios (PR) for HTN or T2DM by BMI category, considering BMI = 18.5–22.9 kg/m² as the reference. All analyses incorporated the complex sampling design of the survey.

Setting: BMI, blood pressure, blood sugar and related information were collected from a nationally representative sample.

Subjects: Adults ($n = 7433$) aged ≥ 35 years.

Results: About 12% of Bangladeshi adults, both male and female, were within the BMI range 23.0–24.9 kg/m² or moderately overweight. Compared with the reference BMI group (18.5–22.9 kg/m²), they had an increased PR for HTN (1.55–1.77) and T2DM (1.54–1.93). These increased PR are similar to those for the WHO-defined overweight group (BMI = 25.0–29.9 kg/m²).

Conclusions: Our findings support the recommendation that calls for setting the optimum BMI for Asian populations to 18.5–23.0 kg/m² for health promotion and for public health interventions like leisure-time physical activity. WHO cut-off points for overweight (≥ 25 kg/m²) should be used to facilitate international comparisons.

Keywords
BMI
Hypertension
Diabetes
Public health
Policy
Overweight

Bangladesh is experiencing ‘double burden’ malnutrition⁽¹⁾. Data show a steady increase in the prevalence of overweight and a steady decrease in underweight among Bangladeshi women of reproductive age^(1,2) and children^(2–4). Compared with pre-1960s, mothers in a post-1985 cohort were 22% less likely to be underweight and four times more likely to be overweight, indicating a secular trend⁽⁵⁾.

A rising secular trend in hypertension (HTN) and type 2 diabetes mellitus (T2DM) has also been observed in Bangladesh⁽⁶⁾. According to a recent national estimate, one in four adults had HTN and one in ten adults had T2DM^(7,8). Studies have identified older age, female sex, higher education and wealth, less physical activity and higher BMI as independent risk factors for both HTN and T2DM in Bangladesh^(7–9). Among these, physical inactivity

and overweight/obesity are important modifiable risk factors^(10,11).

Bangladesh is developing economically at a rapid pace and is aiming to become a middle-income country by 2021⁽¹²⁾. Wealthier people are more likely to be overweight/obese and to engage in sedentary lifestyles and less physical labour. Therefore problems of overweight and obesity, especially among the urban dwellers, are likely to increase in future.

BMI, calculated by dividing weight in kilograms by the square of height in metres (kg/m²), is widely used in public health to identify high-risk individuals, reinforce health promotion and assess the effect of public health interventions. BMI cut-off points are a proxy for body fatness or fat accumulation in the body. However, BMI cannot distinguish between body fat and lean body mass,

*Corresponding author: Email muntasirur.rahman@uq.net.au

which are affected by body structure and muscularity. Lean muscle mass may also decrease with ageing. The relationship between BMI and body fat percentage depends on age, sex and ethnicity. Consequently universal BMI cut-off points may not be appropriate⁽¹³⁾.

There is increasing evidence that Asian populations differ in body fat percentage, body fat distribution and BMI-related CVD risks, and that they have higher percentage of body fat at a lower BMI than white or European populations⁽¹⁴⁾. The WHO's recommended BMI cut-off points for overweight ($\geq 25.0 \text{ kg/m}^2$) and obesity ($\geq 30.0 \text{ kg/m}^2$) are based on increased non-communicable disease risks among European populations⁽¹⁵⁾. However, compared with Europeans, increased T2DM and CVD risks have been observed for Asian populations at lower BMI ($22.0\text{--}25.0 \text{ kg/m}^2$)^(16,17). The WHO has acknowledged that countries may need to adopt different BMI cut-off points such as 23.0 kg/m^2 or 27.5 kg/m^2 for potential public health action, but has refrained from redefining population-specific cut-off points⁽¹⁵⁾.

Most studies from Bangladesh have analysed CVD risks associated with higher BMI/overnutrition based on the WHO guidelines. Among those very few studies that explored risks for moderately overweight individuals, Chen *et al.* reported an increased risk of all-cause deaths (hazard ratio = 1.35; 95% CI 1.05, 1.73) and CVD deaths (hazard ratio = 1.34; 95% CI 0.94, 1.90) for individuals with BMI $\geq 21.5 \text{ kg/m}^2$, compared with BMI of $19.3\text{--}21.4 \text{ kg/m}^2$, but found no association between BMI and cancer deaths⁽¹⁸⁾. An increased risk for heart disease (age- and sex-adjusted hazard ratio = 1.79; 95% CI 1.08, 2.97) with BMI greater than 22.0 kg/m^2 has also been reported⁽¹¹⁾. To our knowledge, no study in Bangladesh has tried to quantify HTN and T2DM risk associated with a moderately increased BMI ($23.0\text{--}24.9 \text{ kg/m}^2$), i.e. among those who are 'at risk of overweight'. The purpose of the current study was to assess the increased prevalence of HTN and T2DM with excess weight, especially among those with moderately increased BMI ($23.0\text{--}24.9 \text{ kg/m}^2$), from cross-sectional data of a nationally representative sample of adults, compared with a reference BMI group of $18.5\text{--}22.9 \text{ kg/m}^2$.

Methods

Data source: Bangladesh Demographic and Health Survey 2011

The most recently available Bangladesh Demographic and Health Survey (6th BDHS) data set was used for the present study. Participants were identified using multistage sampling with a geographically clustered, probability-based sample of households. Data collection was carried out between July and December, 2011. The detailed methodology of the data collection procedure has been described previously⁽¹⁹⁾.

The study population was a nationally representative sample of women and men aged 35 years or older, who

provided consent for blood pressure and fasting plasma glucose measurements. Participation was 86% among men (n 4524) and 92% among women (n 4311). The final sample size with complete information on blood pressure and fasting plasma glucose was 7434 (3685 males, 3749 females) and 7286 (3606 males, 3680 female), respectively. All participants provided informed verbal consent.

Variables

Dependent variables

HTN and T2DM prevalence were the main outcomes of interest.

For analytical purposes, the blood pressure data were made dichotomous (persons with/without HTN). Blood pressure was measured using a LifeSource[®] UA-767 Plus Blood Pressure Monitor. Three measurements were taken by trained health workers, with the participant in the seated position, at approximately 10 min intervals. The average of the second and third measurements was used to record blood pressure. Persons with systolic blood pressure $\geq 140 \text{ mmHg}$ or diastolic blood pressure $\geq 90 \text{ mmHg}$ or current use of antihypertensive medication (self-reported) were considered as having HTN.

Similarly, blood glucose data were dichotomised (persons with/without T2DM). Whole blood was obtained from the middle or ring finger from adults after an overnight fast. The HemoCue 201+ blood glucose analyser was used. Each fasting whole-blood glucose value was multiplied by 1.11 to adjust to the corresponding fasting plasma glucose value. Persons with fasting plasma glucose $\geq 7.0 \text{ mmol/l}$ or taking antidiabetes medication at the time of the survey (self-reported) were considered as having T2DM.

Independent variables

Demographic variables. Demographic variables considered were age at the time of interview (35–55 years, 56 years or older) and sex (male, female).

Socio-economic variables. Place of residence (urban or rural), respondent's wealth index based on principal component analysis (poorest/poorer, middle, richer/richest) and highest educational attainment (no education/pre-school, 1–5 year/primary, secondary or higher) were the socio-economic variables. The wealth quintile index was constructed using information on housing materials, sources of drinking-water, sanitation facilities, use of soap and water for hand washing, availability of electricity, housing amenities, possession of household durable goods, and home and land ownership⁽¹⁹⁾.

BMI. Height (in centimetres) and weight (in kilograms) of the respondents were measured using a standard protocol. Height was measured with an adjustable measuring board, which can provide accurate measurements to the nearest 0.1 cm. Weight was measured using an electronic scale with a precision of 0.1 kg. BMI (kg/m^2) was calculated as the ratio of weight to height squared.

BMI (kg/m^2) data were categorised based on cut-offs for Asian populations into seven BMI categories: <16.0 (severe underweight), 16.0–16.9 (moderate underweight), 17.0–18.4 (mild underweight), 18.5–22.9 (normal weight), 23.0–24.9 (moderately overweight), 25.0–29.9 (overweight) and ≥ 30.0 (obese)⁽¹⁵⁾.

Statistical analysis

We initially provided descriptive information of the distributions of categorical variables by different BMI cut-off points. For descriptive analysis we stratified the population according to seven BMI categories. Mean blood pressure and blood glucose (systolic blood pressure, diastolic blood pressure, fasting plasma glucose) by BMI category were calculated, again separately for males and females.

HTN and T2DM prevalence by BMI category were calculated for selected categorical variables, separately for males and females. The χ^2 test or *t* test was used to explore the differences in proportions and means of dependent variables (HTN, T2DM) for selected independent variables as required. Trends in HTN and T2DM prevalence (by increasing BMI cut-offs) were assessed. The analysis compared HTN and T2DM prevalence for one BMI category with the previous highest BMI category, using the *lincom* command. A *P* value of 0.05 or lower was considered as statistically significant.

Using modified Poisson regression models with robust error variance, we calculated the prevalence ratio (PR) and 95% confidence interval for HTN or T2DM for different BMI categories, considering BMI = 18.5–22.9 kg/m^2 as the reference, separately for males and females. For this purpose, we considered four different models as follows. (i) Model I (unadjusted): HTN/T2DM and BMI; (ii) model II: model I adjusted for age; (iii) model III: model I additionally adjusted for wealth and education; (iv) model IV: combination of models II and III additionally adjusted for residence.

Statistical analyses were performed using the statistical software package Stata version 12 for Windows.

Results

According to the modified BMI cut-offs for Asian populations, about half of the population had BMI within the normal range (BMI = 18.5–22.9 kg/m^2), 2% were obese (BMI ≥ 30.0 kg/m^2) and about 6% were severely underweight (BMI < 16.0 kg/m^2). About one in ten or 12% (both males and females) were moderately overweight (BMI = 23.0–24.9 kg/m^2) and about a quarter (25.1%, 1872/7433) of the adults had BMI above 23.0 kg/m^2 (see online supplementary material, Supplemental Table 1).

According to the modified cut-off, overweight or obesity (BMI ≥ 23.0 kg/m^2) prevalence was significantly higher among women (30% *v.* male 20%; *P* < 0.001) and was significantly associated with young age (28% *v.* older age 18%), urban areas (43% *v.* rural 20%), higher wealth (richer 41% *v.* poorer 10%) and higher education (43% *v.* primary 23% and no education 17%; all *P* < 0.001). About one-third of the population was underweight (BMI < 18.5 kg/m^2). No sex difference was observed in underweight prevalence (see online supplementary material, Supplemental Table 1).

Overall, 19.3% of males and 31.6% of females had HTN. T2DM prevalence was 10.7 and 11.3% among males and females, respectively. Figure 1 presents HTN and T2DM prevalence by the seven BMI categories, for male and females separately. Further analyses by different factors are presented in the online supplementary material, Supplemental Tables 2 (HTN) and 3 (T2DM). In general, HTN prevalence of moderately overweight persons (BMI = 23.0–24.9 kg/m^2) was significantly higher than that of normal-weight persons (BMI = 18.5–22.9 kg/m^2), among both males and females. HTN prevalence of those moderately overweight (BMI = 23.0–24.9 kg/m^2) was statistically similar to that of overweight persons




Fig. 1 Hypertension (HTN) and type 2 diabetes mellitus (T2DM) prevalence (weighted estimates) by BMI category and sex (■, HTN, males; □, HTN, females; ▨, T2DM, males; ▩, T2DM, females) among Bangladeshi adults (*n* 7433) aged ≥ 35 years, Bangladesh Demographic and Health Survey, 2011

(BMI = 25.0–29.9 kg/m²). About half of obese adults, both male and female, had HTN. HTN prevalence was significantly higher among obese people than among those overweight, even though obesity prevalence was very low in the nationally representative sample (males 0.7% and females 2%).

T2DM prevalence was similar (9%) among normal-weight males and females (BMI = 18.5–22.9 kg/m²). Moderately overweight males and females (BMI = 23.0–24.9 kg/m²) had significantly higher frequency of T2DM than those with normal BMI. Like HTN, moderately overweight (BMI = 23.0–24.9 kg/m²) and overweight (BMI = 25.0–29.9 kg/m²) individuals had similar T2DM prevalence. Although obese persons (BMI ≥ 30.0 kg/m²) had much higher T2DM prevalence than those who were overweight, the difference was not statistically significant probably due to the small number of obese persons.

Table 1 presents sex-specific PR for HTN based on various models, separately for males and females. Unadjusted results (model I) showed significantly higher HTN prevalence for overnourished adults (BMI ≥ 23.0 kg/m²), both males and females, compared with normal-weight adults (BMI = 18.5–22.9 kg/m²). There was a trend for prevalence of HTN to increase from moderate-overweight BMI across higher BMI categories. In general, underweight was protective against HTN.

The multivariable analysis showed that higher BMI (23.0–24.9, 25.0–29.9, ≥30.0 kg/m²), older age (56 years or more) and upper wealth quintiles (richer/richest) were associated with higher HTN prevalence for both men and women. Urban living (PR = 1.14; 95% CI 0.99, 1.30) and education above primary (PR = 1.22; 95% CI 1.02, 1.44) were associated with higher HTN prevalence for men, but not for women. HTN prevalence was significantly lower among educated women compared with no education. The final model showed that underweight (BMI = 17.0–18.4, 16.0–16.9, <16.0 kg/m²) was protective against risk of HTN, but the effect was not statistically significant for severely undernourished (<16.0 kg/m²) males.

Table 2 shows that, similar to HTN, overweight men and women (BMI 23.0–24.9, 25.0–29.9, ≥30.0 kg/m²), in the older age group (56 years or more) and in the upper wealth quintiles (richer/richest) were more likely to develop T2DM. However, no difference was observed between urban and rural areas, whereas above primary education was associated with significantly higher T2DM prevalence for both men and women. In the final model, the protective effect of underweight (BMI 17.0–18.4, 16.0–16.9, <16.0 kg/m²) for T2DM was not statistically significant, except in males with BMI = 16.0–16.9 kg/m².

Discussion

To our knowledge, the present paper is the first one from Bangladesh that reports an association between

moderately increased BMI (23.0–24.9 kg/m²) and increased HTN/T2DM prevalence, based on information from a nationally representative sample of adults. Our analysis showed that moderately overweight individuals (BMI = 23.0–24.9 kg/m²) had similar HTN and T2DM PR as those of WHO-defined overweight persons (BMI = 25.0–29.9 kg/m²), across different factors. It should be noted that about 12% of Bangladeshi adults, both male and female, were moderately overweight. Apart from BMI, we found that increased age was the strongest predictor of increased prevalence of HTN (adjusted PR: males 2.14, females 1.89) and T2DM (adjusted PR: males 1.74, females 1.66) for both sexes.

We found about one in four (25%) Bangladeshi adults in this nationally representative sample had BMI ≥ 23.0 kg/m² and about one-third was undernourished (BMI < 18.5 kg/m²). A recent study reported underweight (BMI < 18.5 kg/m²), overweight (BMI = 23.0–24.9 kg/m²) and obesity (BMI ≥ 25.0 kg/m²) prevalence as 14, 18 and 26%, respectively, in rural Bangladeshi adults, with no significant sex difference⁽²⁰⁾. Another study in a rural area in Bangladesh found about 33% of the adults had BMI ≥ 23.0 kg/m² and an undernutrition prevalence of 21%, indicating a transition in lifestyle even among the rural population of Bangladesh⁽²¹⁾.

We found one-third of the females were overweight (BMI ≥ 23.0 kg/m²), significantly more than males (20%). One publication reported in 2011 that 17% of Bangladeshi women of reproductive age were overweight using a BMI cut-off of 25.0 kg/m² and that the increasing trend of overnutrition was similar in urban and rural areas⁽²⁾. Higher BMI among females may have been due to parity, menopausal stage, use of oral contraceptive pills, physical inactivity or high level of central adiposity.

Our findings support the recommendation that calls for setting the optimum BMI for Asian populations to 18.5–23.0 kg/m²⁽²²⁾. A report from Bangladesh on detection of T2DM also recommended a BMI cut-off of 21.2 and 21.8 kg/m², for males and females, respectively⁽²³⁾. Our findings suggested that HTN prevalence increased among males from the BMI range 18.5–22.9 kg/m². For females the corresponding BMI range was little higher, 23.0–24.9 kg/m². Our findings are similar to a previous report that found a lower threshold for CHD risk among males compared with females⁽²⁴⁾. We did not observe any sex difference in BMI threshold for T2DM prevalence, which also agrees with previous findings⁽²⁵⁾.

Studies quantifying the relationship between BMI and CHD are rare in Bangladesh. A study in a tertiary-level hospital in Bangladesh found that, compared with the lowest quartile BMI (<23.15 kg/m²), the adjusted OR for CHD was about three times higher among adults in the second BMI quartile (23.16–24.60 kg/m²) and five to nine times higher for men and women in the highest BMI quartile (26.33–30.90 kg/m²), respectively⁽²⁶⁾. A study from the USA showed that, compared with BMI = 18.5–22.9 kg/m², males with a moderately higher BMI (23.0–24.9 kg/m²) had an

Table 1 Modified Poisson regression models showing prevalence ratios (PR) and 95 % confidence intervals for hypertension (weighted estimates) by BMI category and sex among Bangladeshi adults (*n* 7433) aged ≥ 35 years, Bangladesh Demographic and Health Survey, 2011

		Males				Females			
		Model I	Model II	Model III	Model IV	Model I	Model II	Model III	Model IV
BMI category									
Normal (18.5–22.9 kg/m ²)	Ref.	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Underweight									
<16.0 kg/m ²	PR	0.90	0.72	0.98	0.79	0.78*	0.71**	0.78*	0.72**
	95 % CI	0.63, 1.29	0.50, 1.02	0.68, 1.41	0.55, 1.12	0.61, 1.01	0.64, 0.98	0.61, 0.99	0.56, 0.92
16.0–16.9 kg/m ²	PR	0.77	0.67*	0.83	0.73*	0.88	0.79*	0.88	0.81*
	95 % CI	0.56, 1.06	0.49, 0.91	0.61, 1.15	0.53, 0.99	0.70, 1.10	0.64, 0.98	0.71, 1.11	0.65, 1.01
17.0–18.4 kg/m ²	PR	0.60*	0.58***	0.63***	0.62***	0.84	0.80*	0.85	0.82*
	95 % CI	0.47, 0.76	0.46, 0.74	0.50, 0.81	0.49, 0.79	0.71, 1.01	0.68, 0.95	0.71, 1.01	0.69, 0.97
Overweight									
23.0–24.9 kg/m ²	PR	1.68***	1.81***	1.47***	1.57***	1.55***	1.61***	1.51***	1.55***
	95 % CI	1.42, 2.00	1.53, 2.14	1.23, 1.76	1.32, 1.87	1.36, 1.78	1.41, 1.83	1.32, 1.73	1.35, 1.76
25.0–29.9 kg/m ²	PR	1.96***	2.16***	1.63***	1.77***	1.67***	1.76***	1.65***	1.69***
	95 % CI	1.65, 2.32	1.83, 2.56	1.37, 1.96	1.47, 2.11	1.48, 1.89	1.57, 1.99	1.46, 1.88	1.49, 1.91
≥ 30.0 kg/m ²	PR	2.68***	2.63***	2.12***	2.04***	2.18***	2.31***	2.16***	2.16***
	95 % CI	1.88, 3.82	1.89, 3.65	1.47, 3.06	1.45, 2.89	1.87, 2.56	1.97, 2.71	1.83, 2.55	1.82, 2.55
Age category									
Younger	Ref.	–	1.00	–	1.00	–	1.00	–	1.00
Older	PR	–	2.12***	–	2.14***	–	1.95***	–	1.89***
	95 % CI	–	1.88, 2.40	–	1.89, 2.42	–	1.78, 2.12	–	1.72, 2.08
Residence									
Rural	Ref.	–	–	–	1.00	–	–	–	1.00
Urban	PR	–	–	–	1.14*	–	–	–	1.08
	95 % CI	–	–	–	0.99, 1.30	–	–	–	0.98, 1.19
Wealth category									
Poorest/poorer	Ref.	–	–	1.00	1.00	–	–	1.00	1.00
Middle	PR	–	–	1.17	1.15	–	–	1.03	0.99
	95 % CI	–	–	0.94, 1.44	0.93, 1.42	–	–	0.89, 1.20	0.86, 1.15
Richer/richest	PR	–	–	1.49***	1.34**	–	–	1.31***	1.17*
	95 % CI	–	–	1.25, 1.78	1.12, 1.61	–	–	1.16, 1.48	1.03, 1.32
Education									
No education	Ref.	–	1.00	1.00	–	–	1.00	1.00	–
Primary	PR	–	0.93	1.01	–	–	0.77***	0.88*	–
	95 % CI	–	0.78, 1.12	0.85, 1.21	–	–	0.69, 0.86	0.79, 0.99	–
Above primary	PR	–	1.10	1.22*	–	–	0.73***	0.91	–
	95 % CI	–	0.93, 1.31	1.02, 1.44	–	–	0.64, 0.83	0.80, 1.05	–

Ref., reference category.

Model I, unadjusted; model II, model I adjusted for age; model III, model I additionally adjusted for wealth and education; model IV, combination of models II and III additionally adjusted for residence.

* $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

Table 2 Modified Poisson regression models showing prevalence ratios (PR) and 95% confidence intervals for type 2 diabetes mellitus (weighted estimates) by BMI category and sex among Bangladeshi adults (*n* 7433) aged ≥ 35 years, Bangladesh Demographic and Health Survey, 2011

		Males				Females			
		Model I	Model II	Model III	Model IV	Model I	Model II	Model III	Model IV
BMI category									
Normal (18.5–22.9 kg/m ²)	Ref.	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Underweight									
<16.0 kg/m ²	PR	0.97	0.82	1.10	0.94	0.75	0.70	0.84	0.79
	95% CI	0.58, 1.60	0.49, 1.36	0.67, 1.83	0.57, 1.56	0.47, 1.21	0.44, 1.13	0.52, 1.36	0.49, 1.28
16.0–16.9 kg/m ²	PR	0.54*	0.49*	0.62	0.56*	0.62	0.58*	0.68	0.64
	95% CI	0.31, 0.94	0.28, 0.85	0.35, 1.07	0.32, 0.97	0.37, 1.02	0.35, 0.95	0.41, 1.12	0.39, 1.05
17.0–18.4 kg/m ²	PR	0.92	0.90	1.00	0.99	0.71	0.69*	0.78	0.76
	95% CI	0.68, 1.23	0.67, 1.21	0.75, 1.35	0.74, 1.33	0.50, 1.02	0.48, 0.98	0.54, 1.11	0.53, 1.08
Overweight									
23.0–24.9 kg/m ²	PR	1.78***	1.87***	1.48**	1.54**	1.75***	1.79***	1.54**	1.57**
	95% CI	1.38, 2.30	1.45, 2.41	1.14, 1.91	1.19, 1.99	1.35, 2.27	1.38, 2.32	1.18, 2.00	1.21, 2.04
25.0–29.9 kg/m ²	PR	2.36***	2.54***	1.84***	1.93***	2.18***	2.26***	1.79***	1.81***
	95% CI	1.86, 3.01	2.00, 3.23	1.43, 2.37	1.50, 2.49	1.74, 2.74	1.80, 2.83	1.41, 2.27	1.43, 2.29
≥ 30.0 kg/m ²	PR	3.32***	3.28***	2.41***	2.34**	2.85***	2.95***	2.19***	2.17***
	95% CI	1.99, 5.52	2.03, 5.30	1.43, 4.07	1.42, 3.85	2.09, 3.90	2.15, 4.06	1.58, 3.04	1.55, 3.04
Age category									
Younger	Ref.	–	1.00	–	1.00	–	1.00	–	1.00
Older	PR	–	1.72***	–	1.74***	–	1.53***	–	1.66***
	95% CI	–	1.43, 2.06	–	1.45, 2.09	–	1.28, 1.84	–	1.37, 2.00
Residence									
Rural	Ref.	–	–	–	1.00	–	–	–	1.00
Urban	PR	–	–	–	1.14	–	–	–	1.01
	95% CI	–	–	–	0.94, 1.38	–	–	–	0.91, 1.34
Wealth category									
Poorest/poorer	Ref.	–	–	1.00	1.00	–	–	1.00	1.00
Middle	PR	–	–	0.79	0.78	–	–	1.06	1.03
	95% CI	–	–	0.56, 1.10	0.56, 1.09	–	–	0.78, 1.45	0.76, 1.41
Richer/richest	PR	–	–	1.47**	1.35*	–	–	1.66***	1.50**
	95% CI	–	–	1.15, 1.86	1.05, 1.73	–	–	1.30, 2.11	1.16, 1.96
Education									
No education	Ref.	–	–	1.00	1.00	–	–	1.00	1.00
Primary	PR	–	–	1.17	1.24	–	–	1.17	1.29*
	95% CI	–	–	0.90, 1.53	0.95, 1.62	–	–	0.94, 1.44	1.04, 1.60
Above primary	PR	–	–	1.35*	1.44**	–	–	1.93	1.42**
	95% CI	–	–	1.05, 1.74	1.12, 1.85	–	–	0.95, 1.50	1.11, 1.81

Ref., reference category.

Model I, unadjusted; model II, model I adjusted for age; model III, model I additionally adjusted for wealth and education; model IV, combination of models II and III additionally adjusted for residence.

* $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$.

adjusted relative risk for CHD of 1.22 (95% 1.04, 1.43) and females had a non-significant relative risk of 1.10⁽²⁴⁾. The relative risk of CHD for the obese category (BMI \geq 30.0 kg/m²) was 1.81 for males and 2.16 for females. Although these two studies differ in ethnicity and design, the findings support that excess risk of CHD is associated with moderately high BMI.

Bangladesh is one of the few developing countries in the world to achieve Millennium Development Goals 5 and 4: considerable reduction in maternal and child mortality⁽²⁷⁾. However, the growing burden of non-communicable diseases, the increased number of older people and the lack of health-system preparedness for management of chronic diseases will be major public health challenges⁽²⁸⁾. Bangladeshi adults are particularly vulnerable to CVD at younger ages, compared with other countries. Studies have shown that the mean age of experiencing a first acute myocardial infarction for Bangladeshi adults is 51.9 (SD 11.0) years, which is the lowest in South Asia, and much lower than the global mean (58.1 (SD 12.2) years)⁽²⁹⁾. Approximately 40% of stroke deaths in Bangladesh occurred in persons below 60 years, indicating non-communicable disease risk prevention programmes must target younger adults⁽³⁰⁾.

BMI is a WHO-prescribed indicator of nutritional status measurement that is used internationally and is important for comparison across different countries. The main purpose of a BMI cut-off point is to identify people with a high risk of having an adverse health outcome. It should therefore depend on population based-data of 'increased risk'. The present study proposes using lower BMI cut-off points to identify high-risk individuals for screening, health promotion and monitoring of clinical and public health interventions in Bangladesh and similar settings. However, we suggest that WHO-recommended categories should also be used, especially for reporting purposes, to facilitate international comparisons⁽¹⁵⁾.

Several methodological issues need to be kept in mind while interpreting the observed associations. First of all, the study did not collect information on waist-to-hip ratio, which could be useful in avoiding the potential confounding by different body proportions. Some studies found waist circumference and waist-to-hip ratio to be better markers of obesity, some did not, while some argued that both BMI and waist circumference strongly predict future CHD risks⁽²⁴⁾. Bhowmik *et al.* found waist-to-hip ratio, waist circumference and waist-to-height ratio as better predictors for cardiometabolic risk factors in Bangladesh⁽³¹⁾. However, Khan *et al.* recommended using both BMI and waist circumference for assessing CHD risks, rather than a single obesity indicator, in both clinical and community settings⁽²⁶⁾.

Second, the current study did not collect information on co-morbidities. Therefore weight loss (very low BMI) resulting from existing illness may distort the relationship. However, while exploring associations between low BMI

and increased mortality in a cohort, Pierce *et al.* did not find any role of possible confounding by co-morbidity⁽³²⁾.

Third, the BDHS 2011 did not collect information on tobacco smoking habits at the individual level (only collected information on second-hand smoke at household level). Therefore, we did not include a smoking variable in the regression model, which could potentially influence our findings. It should be noted that even though Bangladeshi men were more likely to smoke tobacco compared with women (55% *v.* females 1%), smokeless tobacco use was high among women (34% *v.* males 29%) especially in poor, rural areas⁽³³⁾.

Conclusions

Policy-level guidelines should be adapted to advocate lower BMI targets (18.5–23.0 kg/m²) by public health managers and health-care professionals. Physicians and public health workers need to be trained on proper counselling and management to promote lifestyle change programmes including leisure-time physical activity.

Acknowledgements

Acknowledgements: The authors thank MEASURE DHS Data Archive, ICF International for providing access to the BDHS 2011 data. *Financial support:* This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. *Conflict of interest:* The authors declare no conflict of interest. *Authorship:* M.R. formulated the research question, designed the study, and carried out data analysis, interpretation and writing of the article. G.W. and A.A.M. supervised data analysis and participated in manuscript writing. All authors reviewed the final draft. *Ethics of human subject participation:* Ethical permission was obtained when the Demographic and Health Surveys were carried out. All data in the present study were anonymous and no additional ethical permission was required.

Supplementary material

To view supplementary material for this article, please visit <https://doi.org/10.1017/S1368980016003566>

References

1. Shafique S, Akhter N, Stallkamp G *et al.* (2007) Trends of under- and overweight among rural and urban poor women indicate the double burden of malnutrition in Bangladesh. *Int J Epidemiol* **36**, 449–457.
2. Khan SH & Talukder SH (2013) Nutrition transition in Bangladesh: is the country ready for this double burden. *Obes Rev* **14**, Suppl. 2, 126–133.

3. Hoque ME, Long KZ, Niessen LW *et al.* (2015) Rapid shift toward overweight from double burden of underweight and overweight among Bangladeshi women: a systematic review and pooled analysis. *Nutr Rev* **73**, 438–447.
4. Das SK, Chisti MJ, Malek MA *et al.* (2015) Changing childhood malnutrition in Bangladesh: trends over the last two decades in urban–rural differentials (1993–2012). *Public Health Nutr* **18**, 1718–1727.
5. Mohsena M, Goto R & Mascie-Taylor CN (2016) Maternal nutritional status (as measured by height, weight and BMI) in Bangladesh: trends and socio-economic association over the period 1996 to 2007. *Public Health Nutr* **19**, 1438–1445.
6. Saquib N, Saquib J, Ahmed T *et al.* (2012) Cardiovascular diseases and type 2 diabetes in Bangladesh: a systematic review and meta-analysis of studies between 1995 and 2010. *BMC Public Health* **12**, 434.
7. Rahman MM, Gilmour S, Akter S *et al.* (2015) Prevalence and control of hypertension in Bangladesh: a multilevel analysis of a nationwide population-based survey. *J Hypertens* **33**, 465–472.
8. Akter S, Rahman MM, Abe SK *et al.* (2014) Prevalence of diabetes and prediabetes and their risk factors among Bangladeshi adults: a nationwide survey. *Bull World Health Organ* **92**, 204–213, 213A.
9. Tareque MI, Koshio A, Tiedt AD *et al.* (2015) Are the rates of hypertension and diabetes higher in people from lower socioeconomic status in Bangladesh? Results from a nationally representative survey. *PLoS One* **10**, e0127954.
10. Rahman M, Nakamura K, Seino K *et al.* (2015) Sociodemographic factors and the risk of developing cardiovascular disease in Bangladesh. *Am J Prev Med* **48**, 456–461.
11. Chen Y, McClintock TR, Segers S *et al.* (2013) Prospective investigation of major dietary patterns and risk of cardiovascular mortality in Bangladesh. *Int J Cardiol* **167**, 1495–1501.
12. Bishwajit G (2015) Nutrition transition in South Asia: the emergence of non-communicable chronic diseases. *F1000 Res* **4**, 8.
13. Deurenberg-Yap M, Chew SK & Deurenberg P (2002) Elevated body fat percentage and cardiovascular risks at low body mass index levels among Singaporean Chinese, Malays and Indians. *Obes Rev* **3**, 209–215.
14. Deurenberg P, Deurenberg-Yap M & Guricci S (2002) Asians are different from Caucasians and from each other in their body mass index/body fat per cent relationship. *Obes Rev* **3**, 141–146.
15. WHO Expert Consultation (2004) Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* **363**, 157–163.
16. Razak F, Anand SS, Shannon H *et al.* (2007) Defining obesity cut points in a multiethnic population. *Circulation* **115**, 2111–2118.
17. Narasawat K, Podang J, Punyarathabundu P *et al.* (2007) Waist circumference, body mass index and health risk factors among middle aged Thais. *Asia Pac J Public Health* **19**, 10–15.
18. Chen Y, Ge W, Parvez F *et al.* (2014) A prospective study of arm circumference and risk of death in Bangladesh. *Int J Epidemiol* **43**, 1187–1196.
19. National Institute of Population Research and Training, Mitra and Associates, & MEASURE DHS (2013) *Bangladesh Demographic and Health Survey 2011*. Dhaka and Calverton MD: NIPORT, Mitra and Associates, and ICF International.
20. Siddiquee T, Bhowmik B, Da Vale Moreira NC *et al.* (2015) Prevalence of obesity in a rural Asian Indian (Bangladeshi) population and its determinants. *BMC Public Health* **15**, 860.
21. Fatema K, Zwar NA, Zeba Z *et al.* (2015) Clinical and biochemical characterization of high risk and not high risk for cardiovascular disease adults in a population from peripheral region of Bangladesh. *BMC Public Health* **15**, 559.
22. Choo V (2002) WHO reassesses appropriate body-mass index for Asian populations. *Lancet* **360**, 235.
23. Bhowmik B, Munir SB, Ahmed KR *et al.* (2014) Anthropometric indices of obesity and type 2 diabetes in Bangladeshi population: Chandra Rural Diabetes Study (CRDS). *Obes Res Clin Pract* **8**, e201–e298.
24. Flint AJ, Rexrode KM, Hu FB *et al.* (2010) Body mass index, waist circumference, and risk of coronary heart disease: a prospective study among men and women. *Obes Res Clin Pract* **4**, e171–e181.
25. Boffetta P, McLerran D, Chen Y *et al.* (2011) Body mass index and diabetes in Asia: a cross-sectional pooled analysis of 900,000 individuals in the Asia cohort consortium. *PLoS One* **6**, e19930.
26. Khan RJ, Harvey DJ, Leistikow BN *et al.* (2015) Relationship between obesity and coronary heart disease among urban Bangladeshi men and women. *Integr Obes Diabetes* **1**, 49–55.
27. Chowdhury AM, Bhuiya A, Chowdhury ME *et al.* (2013) The Bangladesh paradox: exceptional health achievement despite economic poverty. *Lancet* **382**, 1734–1745.
28. Streatfield PK & Karar ZA (2008) Population challenges for Bangladesh in the coming decades. *J Health Popul Nutr* **26**, 261–272.
29. Joshi P, Islam S, Pais P *et al.* (2007) Risk factors for early myocardial infarction in South Asians compared with individuals in other countries. *JAMA* **297**, 286–294.
30. Mateen FJ, Carone M, Alam N *et al.* (2012) A population-based case-control study of 1250 stroke deaths in rural Bangladesh. *Eur J Neurol* **19**, 999–1006.
31. Bhowmik B, Munir SB, Diep LM *et al.* (2013) Anthropometric indicators of obesity for identifying cardiometabolic risk factors in a rural Bangladeshi population. *J Diabetes Investig* **4**, 361–368.
32. Pierce BL, Kalra T, Argos M *et al.* (2010) A prospective study of body mass index and mortality in Bangladesh. *Int J Epidemiol* **39**, 1037–1045.
33. World Health Organization (2011) *Non-Communicable Disease Risk Factor Survey Bangladesh 2010*. New Delhi: WHO Regional Office for South-East Asia.