

Providing the most appropriate care to our *individual* patients

Andrea Cipriani,^{1,2} Anneka Tomlinson^{1,2}

With 350 million people affected in the world, depressive disorder is one of the top 20 causes of the overall global burden of disease.¹ The high costs, both direct and indirect, of major depressive disorder are largely due to the significant deficits in treatment provision² and therefore remediable with current therapies. A key international challenge is to determine how best to implement currently available and effective treatments.

There are a number of efficacious pharmacological and non-pharmacological interventions for depressive disorder.³ Antidepressant drugs are recommended and frequently used as first-line therapy for adults with moderate to severe depressive disorder, and in the UK, about 80% of people in primary care receive an antidepressant prescription in the first year of diagnosis.⁴ However, a significant proportion of these prescriptions are for less than 30 days, while an adequate trial of antidepressants is generally recommended to be 6–8 weeks before changing or stopping the medication.³ A too short duration of treatment limits both the therapeutic effect^{5,6} and increases the risk of withdrawal symptoms.

A number of factors contribute to the suboptimal treatment duration of antidepressant drugs, and the two most recognised contributing factors include the initial side effects of the medication and their perceived marginal efficacy. These factors are exacerbated by our current limited ability to predict which drug will cause the fewest undesirable side effects for a specific patient and which will work most effectively. Improved methods to tailor specific treatments to individual patients are needed. While the need for such an approach is widely recognised by the National Institute for Health and Care Excellence guidelines, this still remains an unmet clinical need, with no currently reliable ways of doing so.³

In fact, however, there are *major* opportunities to improve patient outcomes using

existing therapies by the efficient use of available clinical trial data combined with the technical advances in data synthesis and long-term outcomes from real-world datasets.⁶ Such analyses can now predict the probability of response for a specific subgroup of patients or estimate the chances that a person will have a particular side effect.⁷ By matching individual patients to individual specific antidepressant drugs, clinicians can precisely customise treatment to patients' individual needs and preferences to ultimately improve their clinical outcome.

Internationally, precision medicine is now a leading aim of health-care.⁸ Currently, the process of matching patients to treatments is too often by trial and error, delaying clinical improvement and increasing the risks and costs associated with ineffective treatment. Pooling an analysis of data from clinical trials may provide 'personalised' estimates of comparative effectiveness, stratified for specific subgroups of patients to predict 'individualised' response to treatment. Despite important progress in trying to identify depressed patients who may respond differently,⁹ psychiatry continues to lag behind other specialties like cardiology, oncology and stroke (where simple clinical variables were used to target aspirin and heparin to individual patients).

Factors influencing an individual's drug response in depressive disorder include environmental (eg, co-medications, smoking and food), clinical (eg, severity of illness, previous treatments) and personal or demographic variables (eg, age, gender and family history). However, the wealth and variety of these factors creates its own challenges.¹⁰ New methodologies and tools are needed which:

- ▶ Are based on robust evidence;
- ▶ Are acceptable to patients and clinicians;
- ▶ Guide and direct treatment personalisation, incorporating patients' views and values, and clinical judgement;
- ▶ Support probabilistic decision-making. For instance, a difference in efficacy between interventions of, say, 5% might mean more to one patient than to another and it does not precisely exclude a benefit that clinicians and patients might find meaningful. Or,

vice versa, the same result could allow some doctors and patients to choose to avoid the treatment after carefully considered tolerability, risk and uncertainty.

When several treatment options are available, standard meta-analyses provide only partial information because they can only answer questions about pairs of treatments.¹¹ This fragmented approach does not support optimal clinical decision-making,¹² and the need for a robust method to summarise evidence across several, indeed many, interventions has been increasingly recognised.¹³ Network meta-analysis has been developed to (1) synthesise evidence across a network of randomised trials, (2) allow the estimation of the relative effectiveness of several interventions and (3) produce ranked treatment options.¹⁴ A further advancement is individual patient data network meta-analysis (IPD-NMA).¹⁵ IPD-NMA allows the comparative efficacy of different treatments to be assessed, as well as providing a prediction of the clinical outcome, looking for instance at time to response or temporal trajectory of side effects. This is a novel and exciting development which will hopefully drive forward evidence-based practice.

Treatment algorithms have contributed to advances in many fields of medicine including psychiatry, but studies have consistently shown that the initial benefits of algorithm implementation are not sustained once the implementation support is withdrawn. Computerised decision systems have been developed to provide ongoing assistance to clinicians. However, all algorithms developed to date lack the ability to apply the best knowledge directly to the individual patient and selectively provide information relevant to the characteristics and circumstances of that patient in their specific situation. This requires human interpretation and clinician–patient interaction.¹⁶ So, as we recently said,¹⁷ 'is it now time to move away from a 'one size fits all approach' and move towards precision mental health, providing the most appropriate care to our patients individually?'

Contributors AC drafted the editorial and AT critically revised it.

Funding AC is supported by the National Institute for Health Research (NIHR) Oxford Cognitive Health Clinical Research Facility. This study was funded by the National Institute for Health Research (NIHR) Oxford Health Biomedical Research Centre (BRC-1215-20005). The views expressed are those of the authors and not necessarily those of the UK National Health Service, the NIHR, or the UK Department of Health.

Competing interests None declared.

¹Department of Psychiatry, University of Oxford, Oxford, UK

²Oxford Health NHS Foundation Trust, Warneford Hospital, Oxford, UK

Correspondence to Professor Andrea Cipriani, Department of Psychiatry, Warneford Hospital, University of Oxford, Oxford OX3 7JX, UK; andrea.cipriani@psych.ox.ac.uk

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Cipriani A, Tomlinson A. *Evid Based Mental Health* 2019;22.

Received 2 January 2019
 Accepted 2 January 2019
 Published Online First 21 January 2019

Evid Based Mental Health 2019;22.
 doi:10.1136/ebmental-2019-300080

REFERENCES

- 1 GBD 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. *Lancet* 2017;390:1211–59.
- 2 Docherty M, Shaw K, Goulding L, et al. Evidence-based guideline implementation in low and middle income countries: lessons for mental health care. *Int J Ment Health Syst* 2017;11:8.
- 3 National Institute for Health and Care Excellence (NICE). Depression in adults: recognition and management (NICE Guideline CG90). 2017 <https://www.nice.org.uk/guidance/cg90>.
- 4 Moore M, Yuen HM, Dunn N, et al. Explaining the rise in antidepressant prescribing: a descriptive study using the general practice research database. *BMJ* 2009;339:b3999.
- 5 Batelaan NM, Bosman RC, Muntingh A, et al. Risk of relapse after antidepressant discontinuation in anxiety disorders, obsessive-compulsive disorder, and post-traumatic stress disorder: systematic review and meta-analysis of relapse prevention trials. *BMJ* 2017;358:j3927.
- 6 Califf RM, Robb MA, Bindman AB, et al. Transforming Evidence Generation to Support Health and Health Care Decisions. *N Engl J Med* 2016;375:2395–400.
- 7 Chekroud AM, Zotti RJ, Shehzad Z, et al. Cross-trial prediction of treatment outcome in depression: a machine learning approach. *Lancet Psychiatry* 2016;3:243–50.
- 8 Manrai AK, Patel CJ, Ioannidis JPA, et al. In the Era of Precision Medicine and Big Data, Who Is Normal? *JAMA* 2018;319:1981–2.
- 9 Chekroud AM, Gueorguieva R, Krumholz HM, et al. Reevaluating the Efficacy and Predictability of Antidepressant Treatments: A Symptom Clustering Approach. *JAMA Psychiatry* 2017;74:370–8.
- 10 Koutsouleris N, Kahn RS, Chekroud AM, et al. Multisite prediction of 4-week and 52-week treatment outcomes in patients with first-episode psychosis: a machine learning approach. *Lancet Psychiatry* 2016;3:935–46.
- 11 Efthimiou O. Practical guide to the meta-analysis of rare events. *Evid Based Ment Health* 2018;21:72–6.
- 12 Fusar-Poli P, Radua J. Ten simple rules for conducting umbrella reviews. *Evid Based Ment Health* 2018;21:95–100.
- 13 Mavridis D, Giannatsi M, Cipriani A, et al. A primer on network meta-analysis with emphasis on mental health. *Evid Based Ment Health* 2015;18:40–6.
- 14 Leucht S, Chaimani A, Cipriani AS, et al. Network meta-analyses should be the highest level of evidence in treatment guidelines. *Eur Arch Psychiatry Clin Neurosci* 2016;266:477–80.
- 15 Furukawa TA, Efthimiou O, Weitz ES, et al. Cognitive-behavioral analysis system of psychotherapy, drug, or their combination for persistent depressive disorder: personalizing the treatment choice using individual participant data network meta-regression. *Psychother Psychosom* 2018;87:140–53.
- 16 Henshall C, Marzano L, Smith K, et al. A web-based clinical decision tool to support treatment decision-making in psychiatry: a pilot focus group study with clinicians, patients and carers. *BMC Psychiatry* 2017;17:265.
- 17 Tomlinson A, Cipriani A. The times they are a-changin'. *Evid Based Ment Health* 2018;21:81.