

Research article

High-susceptibility of photosynthesis to photoinhibition in the tropical plant *Ficus microcarpa* L. f. cv. Golden Leaves

Shunichi Takahashi¹, Ayumu Tamashiro¹, Yasuko Sakihama¹, Yasusi Yamamoto², Yoshinobu Kawamitsu³ and Hideo Yamasaki*^{1,4}

Address: ¹Laboratory of Cell and Functional Biology, Faculty of Science, University of the Ryukyus, Okinawa, Japan, ²Department of Biology, Faculty of Science, Okayama University, Okayama, Japan, ³Laboratory of Crop Science, Faculty of Agriculture, University of the Ryukyus, Okinawa, Japan and ⁴Division of Functional Genomics, Center of Molecular Biosciences (COMB), University of the Ryukyus, Okinawa, Japan

E-mail: Shunichi Takahashi - takahashishunichi@hotmail.com; Ayumu Tamashiro - ayumu@ii-okinawa.ne.jp; Yasuko Sakihama - yasuko@ii-okinawa.ne.jp; Yasusi Yamamoto - yasusiya@cc.okyama-u.ac.jp; Yoshinobu Kawamitsu - kawamitu@agr.u-ryukyu.ac.jp; Hideo Yamasaki* - yamasaki@comb.u-ryukyu.ac.jp

*Corresponding author

Published: 2 April 2002

Received: 12 December 2001

BMC Plant Biology 2002, 2:2

Accepted: 2 April 2002

This article is available from: <http://www.biomedcentral.com/1471-2229/2/2>

© 2002 Takahashi et al; licensee BioMed Central Ltd. Verbatim copying and redistribution of this article are permitted in any medium for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: The tropical plant *Ficus microcarpa* L. f. cv. Golden Leaves (GL) is a high-light sensitive tropical fig tree in which sun-leaves are yellow and shade-leaves are green. We compared the response of photosynthetic activities to strong light between GL and its wild-type (WT, *Ficus microcarpa* L. f.).

Results: Field measurements of maximum photosystem II (PSII) efficiency (F_v/F_m) of intact sun-leaves in GL showed that photo synthetic activity was severely photoinhibited during the daytime ($F_v/F_m = 0.46$) and subsequently recovered in the evening ($F_v/F_m = 0.76$). In contrast, WT did not show any substantial changes of F_v/F_m values throughout the day (between 0.82 and 0.78). Light dependency of the CO₂ assimilation rate in detached shade-leaves of GL showed a response similar to that in WT, suggesting no substantial difference in photosynthetic performance between them. Several indicators of photoinhibition, including declines in PSII reaction center protein (D1) content, F_v/F_m value, and O₂ evolution and CO₂ assimilation rates, all indicated that GL is much more susceptible to photoinhibition than WT. Kinetics of PAM chlorophyll *a* fluorescence revealed that nonphotochemical quenching (NPQ) capacity of GL was lower than that of WT.

Conclusion: We conclude that the photosynthetic apparatus of GL is more highly susceptible to photoinhibition than that of WT.

Background

Exposure of leaves to strong light sometimes causes the reduction of photosynthetic efficiency, a phenomenon referred to as photoinhibition [1–3]. The susceptibility of plants to photoinhibition depends on the species and growth light-environments [4]. In general, shade plants or

low-light grown plants are more susceptible to photoinhibition than sun plants or high-light grown plants [4]. Since photoinhibition has a potential to lower productivity and plant growth, avoidance of photoinhibition is critical for the fitness and survival of plants in natural habitats [2,5,6].

It is now widely accepted that harmful Reactive Oxygen Species (ROS) produced upon illumination are involved in the mechanism of photoinhibition [7]. Singlet-excited oxygen (1O_2) can be generated by the interaction of O_2 with triplet-excited chlorophyll (3Chl) formed in the PSII reaction center [8]. Superoxide radical (O_2^-) is unavoidably produced by the Mehler reaction via electron transfer to O_2 at photosystem I. Dismutation of O_2^- results in the formation of hydrogen peroxide (H_2O_2) and the reaction between H_2O_2 and transition metal ions generates hydroxyl radical ($\bullet OH$) which is the most reactive radical among ROS [7]. These ROS can oxidize molecules in chloroplasts including D1 protein in the PSII reaction center [9] and thiol enzymes in the Calvin-Benson cycle [10,11] to inhibit partial reactions of photosynthesis eventually leading to photoinhibition. Recently, ROS have been shown to also bring about photoinhibition through the inhibition of *de novo* synthesis of D1 protein of PSII that is essential to recover from photoinhibition [12].

To protect photosynthetic machinery from ROS-mediated photoinhibition, chloroplasts contain a high amount of ascorbate in stroma at 20 to 300 mM [13]. In the stroma, ascorbate contributes to suppress the accumulation of photoproduced- H_2O_2 by acting as the electron donor for ascorbate peroxidase (APX) which detoxifies H_2O_2 to H_2O [14]. Ascorbate can be also involved in the detoxification of 1O_2 , O_2^- and $\bullet OH$ via nonenzymatic reduction [13]. In these reactions, ascorbate is univalently oxidized to monodehydroascorbate (MDA). To maintain ascorbate concentration in the chloroplasts, MDA should be promptly regenerated to ascorbate during illumination. MDA can be reduced to ascorbate by NAD(P)H in a reaction catalyzed by monodehydroascorbate reductase (MDAR) [15] or directly by reduced ferredoxin [16]. MDA, an organic radical, can undergo spontaneous dismutation to produce ascorbate and dehydroascorbate (DHA) in the absence of MDA reduction. DHA can be reduced by glutathione either through nonenzymatic reaction at a slow rate or enzymatic reaction at a much higher rate [17]. Dehydroascorbate reductase (DHAR, EC 1.8.5.1) is considered to be involved in enzymatic DHA reduction in the chloroplasts [17]. Until now, however, direct *in vivo* evidence showing the physiological significance of DHAR has been yet not available.

We previously reported that leaves of *Ficus microcarpa* L. f. cv. Golden Leaves (GL), a tropical fig tree, lack heat-stable DHAR activity [18]. In GL, the canopy sun-leaves, which are always exposed to direct sunlight, show characteristic yellow whereas those in wild-type (WT, *Ficus microcarpa* L. f.) exhibit normal green even when exposed to direct sunlight. The mechanism for yellow leaves production in GL is unknown. In barley, it has been reported that leaves incubated with CO_2 -enriched air show yellow color due to




Figure 1
Photograph of *Ficus microcarpa* L. f. (WT) and *Ficus microcarpa* L. f. cv. Golden Leaves (GL). Each branch was collected from canopy of mature trees grown in the field. Characteristic yellow leaves are observed only in sun-leaves of GL.

photoinhibition of photosynthesis [19,20], a phenomenon apparently similar to that observed in GL. We hypothesized that GL possess an incomplete machinery of photosynthesis which is susceptible to photoinhibition. The aim of this study was to directly examine the hypothesis. The side-by-side comparisons shown in this study demonstrate that GL is much more susceptible to photoinhibition of photosynthesis than WT.

Results

Photoinhibition in the field

GL is a cultivar of the tropical fig tree *Ficus microcarpa* L. f. (WT) that is natively distributed in subtropical/tropical regions of Asia and Oceania. A significant characteristic of GL is the presence of yellow leaves that contain high amounts of flavonoids and negligible amounts of chlorophyll and carotenoids [21]. The reduced content of photosynthetic pigments in the yellow leaves can be increased by shading them from high-light. In shade conditions, therefore, GL exhibits normal green leaves which cannot be morphologically distinguished from WT leaves (Fig. 1). We have suggested that high susceptibility of GL to photoinhibition could explain the phenomenon [21]. However, there was no direct evidence available to show the susceptibility of GL to photoinhibition.

Figure 2 shows diurnal change of maximum PSII efficiency (F_v/F_m) in field grown attached sun-leaves collected from the canopy. Chlorophyll content of sun-leaves in GL (yellow) was 0.05 g m^{-2} and that in WT (green) was 0.44 g m^{-2} . There was no significant difference between GL and WT in F_v/F_m values measured early in the morning (6 a.m.). The value of F_v/F_m in GL decreased from 0.77 (6




Figure 2
Photoinhibition of GL observed in the field-grown conditions. Top panel, a typical diurnal change in PPFD (\bullet) on the leaves and atmospheric temperature (\square) in the field. Bottom panel, a diurnal change in maximum PSII efficiency of intact attached leaves in the field. (\bullet , blue), WT; (Δ red), GL. F_v/F_m value was measured as PSII efficiency with a portable chlorophyll *a* fluorometer (PAM-2000). Each data point is the mean of 10 separate measurements \pm SD.

a.m.) to 0.46 (2 p.m.) and subsequently recovered to 0.74 in the evening (6 p.m.). In contrast to GL, F_v/F_m values in WT were almost constant throughout a day. Shade-leaves of GL (green), collected from the inside of tree, did not show any decrease in the F_v/F_m value (data not shown). These data obtained in the field conditions suggest that GL is highly susceptible to high-light.

Photosynthetic performance

We compared photosynthetic performance between GL and WT to confirm that the fundamental photosynthetic machinery of GL is normal. Shade-leaves of GL and WT, both of which are green, were used for this purpose. Figure 3 shows the light response curves of CO_2 assimilation rate (based on surface area) in detached shade-leaves. The chlorophyll content of shade-leaves in GL (green) was $0.58 \text{ g Chl m}^{-2}$ and that in WT (green) was $0.70 \text{ g Chl m}^{-2}$. GL showed a very similar light-response curve to WT and there was no substantial difference in the maximum activity of CO_2 assimilation between them. These results clearly demonstrate that the photosynthetic capacity of GL is almost identical to that of WT, suggesting that photosynthetic machinery of GL is functionally not defective.




Figure 3
Light-response curves for the rate of photosynthetic CO_2 assimilation in WT and GL. Shade-leaves, which exhibited identical green coloration in both WT and GL, were used for measurements. Light intensities of irradiation were changed from high to low. (\bullet , blue), WT; (Δ , red), GL. Points are the means of three (\bullet) or four (Δ) separate measurements \pm SD.

Photoinhibition in shade-leaves of Golden Leaves

Figure 4 shows effects of high-light on several indicators of photoinhibition: D1 protein content, F_v/F_m value, O_2 evolution rate and CO_2 assimilation rate. Before high-light exposure, there was no substantial difference between WT and GL shade-leaves in D1 protein content on protein basis (Fig. 4A) and F_v/F_m value (Fig. 4C). GL showed a significant decrease in the D1 protein content and F_v/F_m value upon high-light illumination (Fig. 4A, 4B, 4C). Similar to the indicators specific for PSII activity, those for net photosynthetic activity (i.e. O_2 evolution and CO_2 assimilation rates) also decreased upon high-light illumination (Fig. 4D, 4E). In contrast to these responses observed in GL, WT showed only a small decrease in D1 protein content and F_v/F_m value (Fig. 4A, 4B, 4C). The O_2 evolution and CO_2 assimilation rates in WT did not change even under high-light condition (Fig. 4D, 4E).

Reduced NPQ capacity in Golden Leaves

When plants are exposed to high-light that exceeds the capacity of photosynthesis, excess absorbed light energy can be safely dissipated as heat [22]. This energy dissipation can be measured as the NPQ of PAM chlorophyll *a* fluorescence analysis. NPQ includes the quantum yield of PSII as an index of heat energy dissipation, photoinactivation of PSII and distribution of photon acceptance between PSII and PSI due to the state transition. It is known that the state transition can be controlled by the redox state of Q_A [23], which can be measured as 1-qP. Figure 5 shows that there is no significant difference between GL and WT in yield and 1-qP values. Furthermore, measurements of




Figure 4

High-light induced photoinhibition of GL assessed by several parameters. Shade-leaves were exposed to high-light ($2300 \mu\text{mol m}^{-2} \text{s}^{-1}$) at zero time after preexposure to medium-light ($1000 \mu\text{mol m}^{-2} \text{s}^{-1}$) for 30 min. A schematic illustration shows the outline of photosynthesis including the electron transport process in thylakoid membranes. Letters (A-D) in filled circles, which correspond to those of the panels A-D, represent the sites where the photo synthetic parameters can be measured. **A**, Immuno blotting of the D1 protein contained in leaf-extract. **B**, Degradation of the *in vivo* D1 protein induced by high-light. Each point was plotted using a relative density of band on the gel as shown in A. **C**, Decline of F_v/F_m induced by high-light. Note that the time indicated represents total illumination time. **D**, High-light induced inhibition of the activity of O_2 evolution from intact detached leaves. **E**, High-light induced inhibition of the activity of CO_2 assimilation in intact detached leaves. Points are taken by a separate measurement in different leaves (A, B, D) or a leaf (C, E). (•, blue), WT; (Δ , red), GL.

the electron transport rate (ETR) showed essentially no difference in the sensitivity of PSII photoinactivation between GL and WT (data not shown). However, GL showed a much lower level of NPQ compared to WT (Fig. 5C,5F). These results suggest that low NPQ in GL can be attributed to dysfunction of the heat dissipation mechanism.

Discussion

This study has demonstrated that photosynthetic activities of GL are highly susceptible to photoinhibition under high-light conditions (Figs. 2, 4). There was essentially no difference in photosynthetic efficiency between GL and WT at least up to $700 \mu\text{mol m}^{-2} \text{s}^{-1}$ for a short time (Fig. 3), but NPQ in GL was lower than that in WT at $2300 \mu\text{mol m}^{-2} \text{s}^{-1}$ (Fig. 5C,5F). It has been reported that a *npq1* mutant of *Arabidopsis thaliana* shows increased susceptibility to photoinhibition, but is unaffected in short-term photosynthetic O_2 evolution and CO_2 assimilation [24,25]. For the operation of NPQ, special xanthophyll pigments in the light-harvesting complexes of PSII are

known to have critical roles [26]. The extent of NPQ in plants is strongly correlated with the levels of zeaxanthin and antheraxanthin that are formed from violaxanthin by the enzyme violaxanthin deepoxidase (VDE) located on the luminal side of thylakoid membranes [22]. In excessive light conditions, VDE is activated and converts violaxanthin to zeaxanthin via antheraxanthin when acidification in thylakoid lumen reaches a critical threshold [27]. The activity of VDE is also regulated by the concentration of ascorbate in the lumen because VDE can utilize only ascorbate as a reductant [28]. It has been shown that decreases in ascorbate availability severely affect VDE activity *in vivo* [28]. Because thylakoid membranes do not have any active transport mechanisms for ascorbate, the concentration of ascorbate in the stroma determines that in the lumen by a passive diffusion mechanism [29]. In *Arabidopsis*, it has been clearly demonstrated that NPQ performance *in vivo* is reduced by a mutation that causes ascorbate deficiency [13]. We previously reported that the ascorbate content of GL is severely de-




Figure 5

Yield of PSII (A, D), 1-qP reflecting the reduction state of the q_a pool (B, E), NPQ (C, F) in shade-leaves of WT (\bullet , blue) and GL (Δ , red). Each measurement was made after the exposure of leaves to actinic light for 3 min at different intensities (A-C). In D, E and F, high light (2300 $\mu\text{mol m}^{-2} \text{s}^{-1}$) was used as actinic light to record time courses of Yield (D), 1-qP (E), and NPQ (F). Points are presented as the means of 5 (A-C) or 3 (D-F) separate measurements \pm SEM.

creased in daytime under field conditions [30]. Thus, it is reasonable to assume that decrease of ascorbate content would be a cause of dysfunction of NPQ in GL.

Among antioxidant enzymes, we have shown that GL lacks heat-stable DHAR activity [18]. Plant cells possess several kinds of enzymes that exhibit DHAR activity: thioredoxin [31], glutaredoxin [32] and disulfide isomerase [33]. Recently, two distinct chloroplast proteins that exhibit DHAR activity have been isolated from spinach; one is Kunitz-type trypsin inhibitor [31] and the other is specific DHAR [34]. Because both chloroplastic DHARs show higher heat-stable activity than non-chloroplastic DHAR [31,34], GL may lack chloroplastic DHAR(s) though molecular information is not available. In fact, heat-stable activity cannot be detected in nonphotosynthetic organs such as fruits, that do not include chloro-

plasts [18]. Although chloroplastic DHAR(s) have been presumed to function in the ascorbate regeneration system for maintaining the ascorbate concentration in the chloroplasts [7], the physiological significance of DHAR in chloroplasts is still controversial [35–38]. Since the activity of stromal MDAR is high, the spontaneous disproportionation from MDA to DHA and ascorbate may not occur under *in vivo* situations. Therefore, there is an argument that DHAR might be dispensable for the ascorbate-glutathione cycle to protect chloroplasts from high-light stress [36]. However, there is another location in chloroplasts where the stromal enzymes cannot directly regenerate ascorbate, namely, the lumen space.

In the lumen, oxidation of ascorbate by xanthophylls, α -tocopherol and PSII result in the production of DHA and all of these reactions could be promoted under high-light conditions [13,35]. Therefore, stromal DHAR would be essential to reduce DHA produced in the lumen side. In this context, chloroplastic DHAR must be physiologically indispensable for photoprotection mechanisms particularly when leaves are exposed to high-light stress conditions. This is consistent with previous reports that DHAR activity in leaves of *A. thaliana* is increased by high-light conditions [39]. Thus, we consider that ascorbate availability of GL is decreased by high-light due to a lack of DHAR activity which eventually results in lowered performance of NPQ (Fig. 5C,5F). Ascorbate is involved in photoprotection not only through NPQ but also primarily through ROS scavenging [7]. It should be noted that high susceptibility of GL to photoinhibition could be also explained by a low ROS scavenging capacity which may result in photodamage of target molecules in chloroplast [7] and the inhibition of D1 *de novo* synthesis [12].

The mechanism for yellow leaf production is still unknown. Sun-leaves of GL contain low amounts of chlorophyll and carotenoids but shade-leaves show comparable amount of those to WT, implying that exhibition of yellow leaves in GL is associated with a long-termed irradiation of high-light [21]. When leaves of GL were exposed to high-light, they showed significant symptoms of photoinhibition (Figs. 2, 4). It has been suggested that long-termed photoinhibition (or chronic photoinhibition) can lead photobleaching of photosynthetic pigments such as chlorophyll and carotenoids [3]. The *npq1 Arabidopsis* mutant which is partially defective in NPQ has been reported to show the photobleaching of chlorophyll in high-light conditions [40], a phenomenon apparently very similar to that observed in GL. It is plausible that photoinhibition is involved in an early stage of the yellow leaf-producing mechanism. Unlike herbaceous plants, the leaf-yellowing phenomenon does not cause senescence or cell death [21]. It has been shown that flavonoids accumulated in leaves could protect tissues from oxidative damage by

complementing the ascorbate-glutathione cycle [15,41,42]. In addition to the ROS scavenging function, flavonoids have long been known to be effective sun-screen pigments particularly against UV radiation [43]. Increased flavonoid pigments in GL may play roles including ROS scavenging function, UV-protection and light-attenuation for shadeleaves. The mechanisms of photoinhibition susceptibility and, in particular of high-light tolerance in GL leaves remain to be fully clarified. Although it is difficult to apply molecular biology techniques to GL, we consider that the tropical tree could provide a unique opportunity for examining the important determinants of survival in high-light environments.

Materials and Methods

Plant materials

Plant materials were *Ficus microcarpa* L. f. cv. Golden Leaves (GL) and *Ficus microcarpa* L. f. (WT) grown in the campus of the University of the Ryukyus on Okinawa island (26°15'N, 127°45'E) in Japan. The study period was from March to June in 1999–2000. The photo synthetic photon flux density (PPFD) on the plant canopy was c.a. 2500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ in full sunlight conditions during the period. There was no significant difference in growth conditions of PPFD between WT and GL. Green leaves, grown under light conditions of about 100 $\mu\text{mol m}^{-2} \text{s}^{-1}$ as the maximum, were harvested and used for laboratory experiments as shade-leaves. To avoid the drought of leaves, leaf petioles were continuously kept in distilled water throughout the measurements. To activate the photosynthetic activity, detached leaves were exposed to an illumination of 1000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 30 min before experiments.

Illumination of leaves

Leaves were illuminated with white light from three halogen lamps (400 W). Leaf temperature was maintained at 30°C with an electric fan or a thermostatically controlled water bath during illumination. Various light intensities were obtained by placing wire screens in front of the light source.

Measurements of chlorophyll fluorescence

Chlorophyll fluorescence was measured with a PAM-2000 chlorophyll fluorometer system under atmospheric conditions (Heinz Walz, Effeltrich, Germany). After a dark adaptation period of 15 min, minimum fluorescence (F_o) was determined by a weak red light. Maximum fluorescence of dark-adapted leaf (F_m) was measured during a subsequent saturating light pulse of white light (8000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for 0.4 s). Maximum fluorescence (F_m') and steady-state fluorescence (F_s) of illuminated leaf were measured upon a subsequent saturating light pulse of white light (8000 $\mu\text{mol m}^{-2} \text{s}^{-1}$, for 0.4 s) to determine NPQ ($(F_m - F_m')/F_m'$), yield of PSII [$(F_m' - F_s)/F_m'$] and qP

$(F_m' - F_s)/F_m' - F_o$] [44,45]. A 650 nm light-emitting-diode (LED) equipped with PAM-2000 was used for the illumination of leaf as actinic light.

Measurements of gas exchange

CO_2 assimilation was determined by the difference in CO_2 concentration between inlet and outlet of the assimilation chamber with a CO_2 gas analyzer (LI-6251; LiCor, Inc., Lincoln, Nebraska) in 0.036% CO_2 and 21% O_2 [46]. Measurements of CO_2 assimilation rates were carried out under an illumination of 2300 $\mu\text{mol m}^{-2} \text{s}^{-1}$, except for light response curve measurements. Approximately 25 cm^2 of leaf were employed for an experiment. The air in the chamber was stirred rapidly to maintain a high boundary layer conductance for CO_2 diffusion. O_2 evolution was determined using a gas-phase Walker-type oxygen electrode system (Model LD-1; Hansatech, Norfolk, U.K.) in 5% CO_2 and 21% O_2 [47]. Measurements of O_2 evolution were carried out on 4 cm^2 of leaf disc under an illumination of 600 $\mu\text{mol m}^{-2} \text{s}^{-1}$.

Measurements of D1 protein content

Leaf (0.5 g) was homogenized for 30 s at 0°C in a grinding medium containing 50 mM potassium phosphate (pH 7.0), 5 mM phenylmethylsulfonyl fluoride and 5% (w/v) polyvinylpyrrolidone. The homogenate (4 ml) was mixed with 4 ml of 125 mM Tris (pH 6.8) containing 5% (w/v) SDS, and then incubated at 80°C for 3 min to solubilize proteins. After cooling down the sample to room temperature, aggregates were removed by the centrifugation at 1,500 \times g for 30 s. The supernatant (800 μl) was mixed with the same amount of a solution that contained 0.4 M sucrose, 8 M urea, 5 mM EDTA and 5% (w/v) 2-mercaptoethanol.

SDS-PAGE was carried out according to a previously reported method [48]. Polyacrylamide gels containing 6.0 M urea were used for the stacking gel (4.5 %) and separation gel (13 %). Samples that contained 8 μg protein were loaded in each lane. Immunoblot analysis was performed using specific polyclonal antibodies raised against the D1 protein [48]. The intensities of the protein bands were determined with NIH image system version 1.61 (NIH, USA).

List of abbreviations used

APX, ascorbate peroxidase; ^1Chl , singlet excited chlorophyll; ^3Chl , triplet excited chlorophyll; DHA, dehydroascorbate; DHAR, dehydroascorbate reductase; D1, the reaction center-forming protein of photosystem II; F_v/F_m' , ratio of variable to maximum chlorophyll *a* fluorescence; GL, Golden Leaves; MDA, monodehydroascorbate; MDAR, monodehydroascorbate reductase; NPQ, non-photochemical quenching; $\bullet\text{OH}$, hydroxyl radical; $^1\text{O}_2$, singlet excited O_2 ; O_2^- , superoxide radical; PPFD, photo-

synthetic photon flux density; ROS, reactive oxygen species; WT, wild-type.

Acknowledgements

We are grateful to Dr. M. F. Cohen of University of the Ryukyus for critical reading of the manuscript. This work was supported by a Grant-in-Aid for Scientific Research (B) and (C) from Japan Society for the Promotion of Science to HY.

References

- Demming-Adams B, Adams WW: **Photoprotection and other responses of plants to high light stress.** *Annu Rev Plant Physiol Plant Mol Biol* 1992, **43**:599-626
- Long SP, Humphries S, Falkowski PG: **Photoinhibition of photosynthesis in nature.** *Annu Rev Plant Physiol Plant Mol Biol* 1994, **45**:633-662
- Powles SB: **Photoinhibition of photosynthesis induced by visible light.** *Ann Rev Plant Physiol* 1984, **35**:15-44
- Osmond CB: **What is photoinhibition? Some insights from comparisons of shade and sun plants.** In: *Photoinhibition of photosynthesis: from molecular mechanisms to the field* 1994, 1-24
- Ball MC: **The role of photoinhibition during tree seedling establishment at low temperatures.** In: *Photoinhibition of photosynthesis: from molecular mechanisms to the field* 1994, 365-376
- Kitao M, Lei TT, Koike T, Tobita H, Maruyama Y, Matsumoto Y, Ang LH: **Temperature response and photoinhibition investigated by chlorophyll fluorescence measurements for four distinct species of dipterocarp trees.** *Physiol Plant* 2000, **109**:284-290
- Asada K: **The water-water cycle in chloroplasts: Scavenging of active oxygens and dissipation of excess photons.** *Annu Rev Plant Physiol Plant Mol Biol* 1999, **50**:601-639
- Macpherson AN, Telfer A, Barber J, Truscott TG: **Direct detection of singlet oxygen from isolated photosystem II reaction centers.** *Biochim Biophys Acta* 1993, **1143**:301-309
- Yamamoto Y: **Quality control of photosystem II.** *Plant Cell Physiol* 2001, **42**:121-128
- Kaiser WM: **Reversible inhibition of the Calvin cycle and activation of oxidative pentose phosphate cycle in isolated intact chloroplasts by hydrogen peroxide.** *Planta* 1979, **145**:377-382
- Tanaka K, Otsubo T, Kondo N: **Participation of hydrogen peroxide in the inactivation of Calvin-cycle SH enzymes in SO₂-fumigated spinach leaves.** *Plant Cell Physiol* 1982, **23**:1009-1018
- Nishiyama Y, Yamamoto H, Allakhverdiev SI, Inaba M, Yokota A, Murata N: **Oxidative stress inhibits the repair of photodamage to the photosynthetic machinery.** *EMBO J* 2001, **20**:5587-5594
- Smirnoff N: **Ascorbate biosynthesis and function in photoprotection.** *Phil Trans R Soc Lond B* 2000, **355**:1455-1464
- Asada K: **Ascorbate peroxidase - a hydrogen peroxide-scavenging enzyme in plants.** *Physiol Plant* 1992, **85**:235-241
- Sakihama Y, Mano J, Sano S, Asada K, Yamasaki H: **Reduction of phenoxyl radicals mediated by monodehydroascorbate reductase.** *Biochem Biophys Res Commun* 2000, **279**:949-954
- Miyake C, Asada K: **Ferredoxin-dependent photoreduction of the monodehydroascorbate radical in spinach thylakoids.** *Plant Cell Physiol* 1994, **35**:539-549
- Hossain MA, Asada K: **Purification of dehydroascorbate reductase from spinach and its characterization as a thiol enzyme.** *Plant Cell Physiol* 1984, **25**:85-92
- Yamasaki H, Takahashi S, Heshiki R: **The tropical fig *Ficus microcarpa* L. f. cv. Golden Leaves lacks heat-stable dehydroascorbate reductase activity.** *Plant Cell Physiol* 1999, **40**:640-646
- Sicher RC: **Yellowing and photosynthetic decline of barley primary leaves in response to atmospheric CO₂ enrichment.** *Physiol Plant* 1998, **103**:193-200
- Sicher RC: **Photosystem-II activity is decreased by yellowing of barley primary leaves during growth in elevated carbon dioxide.** *Int J Plant Sci* 1999, **160**:849-854
- Yamasaki H, Heshiki R, Ikehara N: **Leaf-goldenning induced by high light in *Ficus microcarpa* L. f., a tropical fig.** *J Plant Res* 1995, **108**:171-180
- Gilmore AM: **Mechanistic aspects of xanthophyll cycle-dependent photoprotection in higher plant chloroplasts and leaves.** *Physiol Plant* 1997, **99**:197-209
- Krause GH, Weis E: **Chlorophyll fluorescence and photosynthesis: the basics.** *Annu Rev Plant Physiol Plant Mol Biol* 1991, **42**:313-349
- Niyogi KK, Grossman AR, Björkman O: **Arabidopsis mutants define a central role for the xanthophyll cycle in the regulation of photosynthetic energy conversion.** *Plant Cell* 1998, **10**:1121-1134
- Havaux M, Niyogi KK: **The violaxanthin cycle protects plants from photooxidative damage by more than one mechanism.** *Proc Natl Acad Sci USA* 1999, **96**:8762-8767
- Gilmore AM: **Xanthophyll cycle-dependent nonphotochemical quenching in Photosystem II: Mechanistic insights gained from Arabidopsis thaliana L. mutants that lack violaxanthin de-epoxidase activity and/or lutein.** *Photosynth Res* 2001, **67**:89-101
- Gilmore AM, Yamasaki H: **9-aminoacridine and dibucaine exhibit competitive interactions and complicated inhibitory effects that interfere with measurements of ΔpH and xanthophyll cycle-dependent photosystem II energy dissipation.** *Photosynth Res* 1998, **57**:159-174
- Bratt CE, Arvidsson P-O, Carlsson M, Åkerlund H-E: **Regulation of violaxanthin de-epoxidase activity by pH and ascorbate concentration.** *Photosynth Res* 1995, **45**:169-175
- Foyer CH, Leiadais M: **A comparison of the relative rates of transport of ascorbate and glucose across the thylakoid, chloroplast and plasmalemma membranes of pea leaf mesophyll cell.** *J Plant Physiol* 1996, **148**:391-398
- Yamasaki H, Heshiki R, Yamasu T, Sakihama Y, Ikehara N: **Physiological significance of the ascorbate regenerating system for the high-light tolerance of chloroplasts.** In: *Photosynthesis: from light to biosphere* 1995, 291-294
- Trümper S, Follmann H, Häberlein I: **A novel dehydroascorbate reductase from spinach chloroplasts homologous to plant trypsin inhibitor.** *FEBS Lett* 1994, **352**:159-162
- Sha S, Minakuchi K, Higaki N, Sato K, Ohtsuki K, Kurata A, Yoshikawa H, Kotaru M, Masumura T, Ichihara K, Tanaka K: **Purification and characterization of glutaredoxin (thioltransferase) from rice (*Oryza sativa* L.).** *J Biochem* 1997, **121**:842-848
- Wells WW, Xu DP: **Dehydroascorbate reduction.** *J Bioenerg Biomembr* 1994, **26**:369-77
- Shimaoka T, Yokota A, Miyake C: **Purification and characterization of chloroplast dehydroascorbate reductase from spinach leaves.** *Plant Cell Physiol* 2000, **41**:1110-1118
- Mano J, Ushimaru T, Asada K: **Ascorbate in thylakoid lumen as an endogenous electron donor to Photosystem II: Protection of thylakoids from photoinhibition and regeneration of ascorbate in stroma by dehydroascorbate reductase.** *Photosynth Res* 1997, **53**:197-204
- Morell S, Follmann H, DeTullio M, Häberlein I: **Dehydroascorbate and dehydroascorbate reductase are phantom indicators of oxidative stress in plants.** *FEBS Lett* 1997, **414**:567-570
- Morell S, Follmann H, DeTullio M, Häberlein I: **Dehydroascorbate reduction: the phantom remaining.** *FEBS Lett* 1998, **425**:530-531
- Foyer CH, Mullineaux PM: **The presence of dehydroascorbate and dehydroascorbate reductase in plant tissue.** *FEBS Lett* 1998, **425**:528-529
- Kubo A, Aono M, Nakajima N, Saji H, Tanaka K, Kondo N: **Differential responses in activity of antioxidant enzymes to different environmental stresses in Arabidopsis thaliana.** *J Plant Res* 1999, **112**:279-290
- Havaux M, Bonfils J-P, Lütz C, Niyogi KK: **Photodamage of the photosynthetic apparatus and its dependence on the leaf developmental stage in the *npq1* Arabidopsis mutant deficient in the xanthophyll cycle enzyme violaxanthin de-epoxidase.** *Plant Physiol* 2000, **124**:273-284
- Yamasaki H, Sakihama Y, Ikehara N: **Flavonoid-peroxidase reaction as a detoxification mechanism of plant cells against H₂O₂.** *Plant Physiol* 1997, **115**:1405-1412
- Yamasaki H: **A function of colour.** *Trends Plant Sci* 1997, **2**:7-8
- Cohen MF, Sakihama Y, Yamasaki H: **Roles of plant flavonoids in interactions with microbes: from protection against pathogens to the mediation of mutualism.** In: *Recent research developments in plant physiology* 2001, 157-173
- Schreiber U: **Detection of rapid induction kinetics with a new type of high-frequency modulated chlorophyll fluorometer.** *Photosynth Res* 1986, **9**:261-272

45. Genty B, Briantais J-M, Baker NR: **The relationship between the quantum yield of photosynthetic electron transport and quenching of chlorophyll fluorescence.** *Biochim Biophys Acta* 1989, **990**:87-92
46. Agata W, Kawamitsu Y, Hakoyama S, Shima S: **A system for measuring leaf gas exchange based on regulating vapour pressure difference.** *Photosynth Res* 1986, **9**:345-357
47. Kawamitsu Y, Boyer JS: **Photosynthesis and carbon storage between tides in a brown alga, *Fucus vesiculosus* L.** *Marine Biol* 1999, **133**:361-369
48. Yamamoto Y, Akasaka T: **Degradation of antenna chlorophyll-binding protein CP43 during photoinhibition of photosystem II.** *Biochemistry* 1995, **34**:9038-9045

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMedcentral will be the most significant development for disseminating the results of biomedical research in our lifetime."

Paul Nurse, Director-General, Imperial Cancer Research Fund

Publish with **BMC** and your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours - you keep the copyright

Submit your manuscript here:

<http://www.biomedcentral.com/manuscript/>



editorial@biomedcentral.com