



CD47 Activation by Thrombospondin-1 in Lymphatic Endothelial Cells Suppresses Lymphangiogenesis and Promotes Atherosclerosis

Bhupesh Singla¹, Ravi Varma Aithbathula, Naveed Pervaiz¹, Ishita Kathuria, Mallory Swanson, Frederick Adams Ekuban¹, WonMo Ahn, Frank Park, Maxwell Gyamfi, Mary Cherian-Shaw, Udai P. Singh¹, Santosh Kumar

BACKGROUND: TSP1 (thrombospondin-1)—a well-known angiogenesis inhibitor—mediates differential effects via interacting with cell surface receptors including CD36 (cluster of differentiation) and CD47. However, the role of TSP1 in regulating lymphangiogenesis is not clear. Our previous study suggested the importance of cell-specific CD47 blockade in limiting atherosclerosis. Further, our experiments revealed CD47 as a dominant TSP1 receptor in lymphatic endothelial cells (LECs). As the lymphatic vasculature is functionally linked to atherosclerosis, we aimed to investigate the effects of LEC TSP1-CD47 signaling inhibition on lymphangiogenesis and atherosclerosis.

METHODS: Murine atherosclerotic and nonatherosclerotic arteries were utilized to investigate TSP1 expression using Western blotting and immunostaining. LEC-specific knockout mice were used to determine the *in vivo* role of LEC *Cd47* in lymphangiogenesis and atherosclerosis. Various *in vitro* cell-based assays, *in vivo* Matrigel plug implantation, molecular biological techniques, and immunohistological approaches were used to evaluate the underlying signaling mechanisms.

RESULTS: Elevated TSP1 expression was observed in mouse atherosclerotic aortic tissue compared with nonatherosclerotic control tissue. TSP1 at pathological concentrations suppressed both *in vitro* and *in vivo* lymphangiogenesis. Mechanistically, TSP1 inhibited VEGF (vascular endothelial growth factor)-C-induced AKT and eNOS activation in LEC and attenuated NO (nitric oxide) production. Further, *CD47* silencing in LEC prevented the effects of TSP1 on lymphangiogenic AKT-eNOS signaling and lymphangiogenesis. Atheroprone AAV (adeno-associated virus) 8-*PCSK9*-injected LEC-specific *Cd47* knockout mice (*Cd47*^{ΔLEC}) had reduced atherosclerosis in both aorta and aortic root compared with control mice (*Cd47*^{ΔWT}). However, no differences in metabolic parameters including body weight, plasma total cholesterol levels, and fasting blood glucose were observed. Additional immunostaining experiments performed on aortic root cross-sections indicated higher lymphatic vessel density in *Cd47*^{ΔLEC} mice in comparison to controls.

CONCLUSIONS: These findings demonstrate that TSP1 inhibits lymphangiogenesis via activation of CD47 in LEC, and loss of LEC *Cd47* attenuates atherosclerotic lesion formation. Collectively, these results identify LEC CD47 as a potential therapeutic target in atherosclerosis.

GRAPHIC ABSTRACT: A [graphic abstract](#) is available for this article.

Key Words: aorta ■ atherosclerosis ■ endothelial cells ■ fasting ■ lymphangiogenesis

Atherosclerosis—a chronic vascular inflammatory disease—is characterized by the accumulation of lipid-loaded cells in the subendothelial space of

medium- and large-sized arteries and develops at certain focal points. Being an underlying cause of myocardial infarction, stable and unstable angina, and stroke, it

Correspondence to: Bhupesh Singla, PhD, Department of Pharmaceutical Sciences, College of Pharmacy, The University of Tennessee Health Science Center, Room 446, 881 Madison Ave, Memphis, TN 38163. Email bsingla@uthsc.edu

Supplemental Material is available at <https://www.ahajournals.org/doi/suppl/10.1161/ATVBAHA.122.318904>.

For Sources of Funding and Disclosures, see page 1248.

© 2023 The Authors. *Arteriosclerosis, Thrombosis, and Vascular Biology* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the [Creative Commons Attribution Non-Commercial-NoDerivs](#) License, which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited, the use is noncommercial, and no modifications or adaptations are made.

Arterioscler Thromb Vasc Biol is available at www.ahajournals.org/journal/atvb

Nonstandard Abbreviations and Acronyms

Arg1	arginase 1
Cdk	cyclin-dependent kinase
IL	interleukin
iNOS	inducible NO synthase
LDL	low-density lipoprotein
LEC	lymphatic endothelial cell
LV	lymphatic vessel
LYVE-1	lymphatic vessel endothelial hyaluronan receptor-1
ORO	oil red O
RCT	reverse cholesterol transport
TSP1	thrombospondin-1

accounts for the majority of deaths worldwide.¹ Earlier studies have demonstrated that adventitial lymphatic vasculature represents the primary route of cholesterol removal from atherosclerotic vessels, and improved lymphatic function leads to atherosclerosis regression.^{2,3} Further, inhibition of arterial lymphatic drainage via surgical and genetic approaches induces atherosclerotic lesion formation.^{2,4,5} Moreover, early treatment with lymphangiogenic factor VEGF (vascular endothelial growth factor)-C (152S) has been shown to improve lymphatic function and attenuate high-fat diet-induced atherosclerosis in *Ldlr* (low-density lipoprotein receptor) knockout mice.⁶ Therefore, arterial lipid accumulation is an outcome of the disturbed equilibrium between the influx of plasma lipoproteins and the efflux of lipids from the vessel wall via reverse cholesterol transport (RCT) mediated by lymphatic vessels (LVs). Despite this information, the molecular factors and downstream signaling mechanisms that govern arterial LV formation (lymphangiogenesis) are understudied.

TSP1 (thrombospondin-1) is a secreted matricellular glycoprotein, which mediates differential physiological or pathological effects via interacting with integrins and cell surface receptors including CD36 (cluster of differentiation) and CD47.⁷ It regulates various biological processes, including cell proliferation, migration, angiogenesis, and inflammation.^{8,9} TSP1 expression is low in healthy blood vessels; however, elevated TSP1 levels have been associated with cardiovascular disease including injury-induced neointima formation and atherosclerosis.^{10–12} Human *TSP1* gene polymorphism and its protein expression are linked with the increased risk of myocardial infarction and atherogenesis, respectively.^{12,13} Additional studies utilizing global genetic deletion of *Tsp1* in mice have reported contradictory roles of TSP1 in atherosclerosis development.^{14,15} However, the mechanisms by which TSP1 contributes to atherosclerotic plaque formation and its cell-specific role in its pathogenesis

Highlights

- Thrombospondin-1-induced lymphatic endothelial cell CD47 activation inhibits lymphangiogenesis.
- CD47 deletion in lymphatic endothelial cells augments both in vitro and in vivo lymphangiogenesis.
- Lymphatic endothelial cell-specific CD47 deletion suppresses atherosclerotic lesion formation.

are understudied. TSP1 after binding to CD47 present in vascular endothelial cells suppresses angiogenesis via blocking required NO (nitric oxide) signaling.¹⁶ NO signaling in lymphatic endothelial cells (LECs) is also pivotal for lymphangiogenesis and the maintenance of LVs.¹⁷ Kojima et al have shown that the administration of CD47-blocking antibodies in murine models of atherosclerosis inhibits plaque formation via promoting efferocytosis and reduces vascular inflammation in human patients with cancer.^{18,19} We recently reported that global deletion of *Cd47* in mice protects from atherosclerosis; however, myeloid cell-specific *Cd47* loss augments lesion formation, suggesting the importance of cell-specific CD47 blockade in atherosclerosis.²⁰ Nonetheless, the importance of TSP1-induced LEC CD47 activation in regulating arterial lymphangiogenesis and atherosclerosis remains unexplored.

In the current study, we examined the role of LEC TSP1-CD47 signaling axis in governing both in vitro and in vivo lymphangiogenesis and development of atherosclerosis. Our data demonstrate that (1) TSP1-mediated LEC CD47 activation inhibits lymphangiogenesis and (2) blockade of LEC CD47 signaling suppresses atherosclerosis. Altogether, the findings of the present study identify LEC CD47 as a potential therapeutic target to promote arterial lymphangiogenesis and reduce atherosclerosis.

MATERIALS AND METHODS

The data that support the findings of this study are available from the corresponding author upon reasonable request. Please see the extended Methods and Major Resources Table in the [Supplemental Material](#).

Animals

All animal experimental procedures were approved by the Institutional Animal Care and Use Committee of the University of Tennessee Health Science Center, Memphis, TN, or Augusta University. The procedures were conducted following the National Institutes of Health Guide for the Care and Use of Laboratory Animals. Eight- to 10-week-old male wild-type C57BL/6J (stock number 000664), *ApoE*^{-/-} (stock number 002052), and *Lyve-1 Cre*⁺ (stock number 012601) mice were procured from The Jackson Laboratory (Bar Harbor). *Cd47*-floxed (*Cd47*^{f/f}) mice were cryorecovered from frozen germplasm maintained in the KOMP Repository (*Cd47*^{tm1a(KOMP)Mbp},

Mouse Biology Program, University of California, Davis).²⁰ LEC-specific *Cd47* knockout mice (*Cd47^{fl} Lyve-1 Cre^{-/-}*, referred as *Cd47^{ΔLEC}*) were generated by crossing *Cd47^{fl}* mice with *Lyve-1 Cre⁺* mice. Littermate *Cd47^{fl} Lyve-1 Cre^{-/-}* and *Lyve-1 Cre^{+/-}* mice were used as controls (*Cd47^{WT}*). All mice were genotyped by polymerase chain reaction amplification of tail genomic DNA using primers listed in Table S1, and LEC-specific deletion was also confirmed using double immunostaining experiments.

Atherosclerotic Lesion Analysis

Male mice were used in the present study because sex hormones in females are known to affect atherosclerotic lesion formation and plasma cholesterol levels.^{21,22} Male mice were injected with a recombinant adeno-associated viral vector expressing a gain-of-function mutant form of human proprotein convertase subtilisin/kexin type 9 (AAV8-*hPCSK9*-D374Y, Vigene Biosciences, Rockville, referred as AAV8-*PCSK9*, 1×10^{11} viral genomes, IP) once²³ and fed a Western diet (TD.88137; Envigo, Indianapolis, IN) for 16 weeks to induce liver-specific *Ldlr* degradation, hypercholesterolemia, and atherosclerosis. Whole-body fat, lean and fluid mass (Bruker Minispec Live Mice Analyzer, LF90), and fasting blood glucose (ReliOn Prime Blood Glucose Monitoring System) were determined. Mice were fasted overnight (16 hours) before euthanasia to determine fasting blood glucose and plasma total cholesterol levels. Mice were anesthetized by isoflurane inhalation (3%), and blood was collected in a heparinized syringe via cardiac puncture for further analysis. In situ images of the aortic arch and aorta were captured using an Olympus SZX16 stereomicroscope. Mouse aorta and heart were harvested and fixed in 4% paraformaldehyde for 24 hours. Heart tissue was embedded in the optimum cutting temperature (23-730-571; Fisher Healthcare, Houston, TX) and stored at -80°C until use. Amplex Red cholesterol assay (A12216; Molecular Probes, Eugene, OR) was used to quantify plasma total cholesterol levels. Plasma triglyceride levels were examined using reagents and standards obtained from Wako Chemicals (Richmond, VA).

For en face analysis, whole paraformaldehyde-fixed aortas were stained with 2% oil red O (ORO) solution (Sigma-Aldrich, MO) for 30 minutes at room temperature, washed with 60% isopropanol (30 s) and H_2O (5 minutes). Then, aortas were opened longitudinally and imaged with an Olympus SZX16 stereomicroscope, and ORO-positive areas were quantified. Further, the upper halves of fixed hearts were dissected and embedded in the optimum cutting temperature compound. Serial aortic root frozen sections (9 μm) were stained with 2% ORO solution, washed with 60% isopropanol, and counterstained with hematoxylin (22-220-100; Fisher Healthcare, Pittsburgh, PA).²⁰ Images were acquired using an Olympus BX43 phase-contrast microscope to determine the lipid deposition in the aortic sinus. Image analysis was performed utilizing the Image-Pro Plus software (Media Cybernetics, Bethesda, MD).

Cell Culture and Gene Silencing

Primary human dermal LECs isolated from adult skin and human aortic endothelial cells were purchased from PromoCell GmbH (Heidelberg, Germany) and cultured in endothelial cell growth medium MV 2 (PromoCell) containing 5% heat-inactivated fetal bovine serum, 100 IU/mL penicillin, 100 $\mu\text{g}/\text{mL}$ streptomycin, and growth factors bullet kit. Cells were

maintained in a humidified incubator at 37°C and used till passage 7. Primary mouse lung LECs were isolated after digestion of lungs with collagenase type 2 (1 mg/mL) for 1 hour at 37°C as we described previously.²⁴ LYVE-1 (lymphatic vessel endothelial hyaluronan receptor-1)-positive cells were separated and cultured.

A smart pool of siRNA for human *CD47* (M-019505-01-0005) and a nontargeting control siRNA (D-001210-01-05) were purchased from Horizon Discovery (Cambridge, United Kingdom). Cells were transfected with siRNAs using the TransIT-TKO transfection reagent (Mirus Bio LLC, Madison, WI) according to the manufacturer's instructions. Cells were used for further experiments 48 hours post-transfection, and gene silencing was confirmed using quantitative real-time polymerase chain reaction. Protein knockdown was validated by Western blot experiments.

Western Blot

Cells/tissues were lysed in RIPA (radio-immunoprecipitation assay) lysis buffer supplemented with protease and phosphatase inhibitors. Equal amounts of proteins (10–15 μg per lane) were separated using SDS-PAGE gels, transferred onto nitrocellulose membranes (Li-Cor Biosciences, Lincoln, NE), and membranes blocked with Intercept blocking buffer (Li-Cor Biosciences) were probed with the following primary antibodies at indicated concentrations: TSP1 (1:400), vinculin 1 (1:1000), GAPDH (1:2000), pAKT^{Ser473} (1:1000), total AKT (1:1000), peNOS^{Ser1177} (1:1000), total eNOS (1:1000), pERK1/2^{Thr202/Tyr204} (1:1000), total ERK1/2 (1:1000), CD47 (1:500), VEGFR3 (1:500), epsin 1 (1:500), p53 (1:1000), and p21 (1:1000). Total AKT (2920S), pAKT^{Ser473} (4060S), peNOS^{Ser1177} (9571S), total eNOS (5880S), pERK1/2^{Thr202/Tyr204} (9101S), total ERK1/2 (4695S), and p21 (2947) were purchased from Cell Signaling Technology (Danvers, MA). CD47 antibody (NBP2-31106; Novus Biologicals, LLC, Centennial, CO) and ab175388 (Abcam, Cambridge, MA) were used. We obtained VEGFR3 (sc-321), epsin 1 (sc-55556), p53 (sc-126), and GAPDH (sc-365062) antibodies from Santa Cruz Biotechnology (Dallas, TX). TSP1 (18304-1-AP) antibody was acquired from Proteintech (Rosemont, IL). Antibody for vinculin 1 (V4505) was purchased from Sigma-Aldrich, Inc (St. Louis, MO). After washing, membranes were incubated with IRDye-conjugated secondary antibodies (Li-Cor Biosciences) and scanned with the Odyssey CLx/DLx Infrared Imaging System (Li-Cor Biosciences) to detect proteins of interest. The densitometry analysis was performed utilizing the ImageJ software (National Institutes of Health).

Immunohistochemistry

Paraffin sections were deparaffinized, rehydrated, and underwent antigen retrieval by boiling in citrate buffer at 98°C for 10 minutes, while frozen sections were washed with PBS twice and permeabilized with 0.1% Triton X-100 (10 minutes). Then, sections were blocked with 10% goat serum (1 hour) at room temperature and incubated with primary antibodies against TSP1 (Proteintech; 18304-1-AP, 1:80), SMA (Abcam; ab7817, 1:100), CD68 (Thermo Fisher Scientific, Rockford, IL; MA5-13324, 1:100), LYVE-1 (Abcam; ab14917, 1:100), CD11c (Thermo Fisher Scientific; 53-0114-80, 1:80), CD45 (Cell Signaling Technology; 70257S 1:70), CD34 (Abcam; ab81289, 1:70), iNOS (inducible NO synthase; Abcam; ab3523, 1:100),

Arg1 (arginase 1; Sigma-Aldrich; AV45673, 1:100), podoplanin (R&D Systems; AF3244, 1:40), CD31 (Abcam; ab28364, 1:70), CD3e (Thermo Fisher Scientific; 53-0031-80, 1:80), and CD47 (Novus Biologicals; NBP2-31106, 1:100 or R&D Systems; AF1866, 1:70) overnight at 4°C. Next, sections were incubated with Alexa Fluor 488/Alexa Fluor 594-labeled secondary antibodies (Life Technologies Corporation, Eugene, OR; A11008, 1:700; A11032, 1:500; A11037, 1:500; A11001, 1:1000; and A11055, 1:400) for 1 hour at room temperature, followed by mounting with DAPI (4',6-diamidino-2-phenylindole) containing Fluoromount-G (Thermo Fisher Scientific). Images were captured using a Zeiss 710 inverted confocal microscope.

Mouse aortic root frozen sections were stained with hematoxylin and eosin (Fisher Healthcare; 22-220-104) and Masson trichrome (Richard Allan Scientific LLC, Kalamazoo, MI; 22-110-648) using standard protocols. EpreDia UltraVision LP HRP polymer and DAB Detection System (Richard Allan Scientific LLC; TL015HD) was utilized for staining aortic root sections with CD68 antibody according to the manufacturer's instructions. Images were acquired using an Olympus BX43 phase-contrast microscope. For ORO, hematoxylin and eosin, Masson trichrome, and CD68 immunostaining, 4 sections per mouse spaced 100 µm apart were stained and analyzed, and mean area of 4 sections was shown. All image analyses were performed utilizing the Image-Pro Plus software (Media Cybernetics, Bethesda, MD).

Statistical Analysis

All statistical analyses were performed using the GraphPad Prism 9 (La Jolla, CA). The data are represented as mean±SEM. The sample number (n) for each experiment/group is mentioned in the figure legends. The normality of data was investigated using the Shapiro-Wilk normality test. Two-tailed Student *t* test (parametric data) or Mann-Whitney *U* test (nonparametric data) were used to compare 2 groups, while >2 groups were analyzed using 1- or 2-way ANOVA followed by appropriate post hoc test for multiple comparisons where appropriate. Kruskal-Wallis test was used to compare >2 groups of nonparametric data. While using parametric tests to compare groups, it was assumed that all groups have the same standard SD. A *P* value <0.05 was considered statistically significant.

RESULTS

TSP1 Protein Expression Is Elevated in Atherosclerotic Arteries

Previous human studies have associated polymorphism in the *TSP1* gene and TSP1 protein expression with the increased risk of myocardial infarction and atherosclerosis development, respectively.^{12,13} Despite these associations, the role of TSP1 in atherosclerosis development is largely unknown. To confirm whether TSP1 expression correlates with atherosclerosis development in mice, we collected murine atherosclerotic and nonatherosclerotic aortic tissue and quantified TSP1 expression using immunoblotting. The presence and absence of atherosclerotic lesions were confirmed with ORO staining. Western blot experiments showed increased TSP1

protein expression in atherosclerotic inner curvature of 12-week Western diet-fed *Apoe*^{-/-} mice compared with plaque-free descending aorta region (Figure 1A and 1B).

Next, immunostaining experiments were performed utilizing aortic root cross-sections from nonatherosclerotic wild-type mice and atherosclerotic *Apoe*^{-/-} mice (12 weeks Western diet) to determine the layers of the atherosclerotic arterial wall expressing TSP1. Immunostaining data demonstrated elevated TSP1 expression in *Apoe*^{-/-} mice compared with wild-type mice (Figure 1C and 1D). In particular, TSP1 expression was detected in some intraplaque CD68 (macrophage marker)-positive areas and adventitial layer (Figure 1G). Taken together, these results demonstrate that TSP1 expression is upregulated in atherosclerotic arteries, and it localizes in the plaque area and adventitial layer (where LVs are primarily localized). However, the role of TSP1 in regulating arterial lymphangiogenesis and atherosclerosis development is not known.

TSP1 Suppresses Both In Vitro and In Vivo Lymphangiogenesis

LVs play an important role in the removal of cholesterol from atherosclerotic blood vessels via mediating RCT, and disruption of arterial lymphatic drainage induces adventitial inflammation and aggravates atherosclerosis development in hypercholesterolemic mice.²⁻⁴ We identified elevated TSP1 expression (Figure 1C and 1D) in the adventitial layer of atherosclerotic arteries where LVs are mainly present and TSP1 is a potent inhibitor of angiogenesis⁷; yet surprisingly, its direct effect on lymphangiogenesis has not been investigated. Therefore, we sought to investigate the role of TSP1 in regulating lymphangiogenesis. First, we performed WST (water-soluble tetrazolium)-1 assay to examine the effect of treatment with different TSP1 concentrations on LEC proliferation. As shown in Figure 2A and Figure S1A, TSP1 at 11- and 22-nM concentrations remarkably reduced VEGF-C-induced (100 ng/mL) LEC proliferation. Therefore, we chose 11- and 22-nM TSP1 concentrations for further experiments investigating TSP1's effects on LEC migration and tube formation. Plasma levels of TSP1 in patients with vascular pathologies have been shown elevated (117–6500 ng/mL) compared with control individuals (13–137 ng/mL).²⁵ Therefore, TSP1 concentrations used in the present study are pathologically relevant (5000 ng/mL=11 nM). However, exact TSP1 concentrations in the atherosclerotic and nonatherosclerotic arteries are not known. Next, immunostaining experiments were executed to determine Ki67-positive nuclei as a measure of cell proliferation and demonstrated a significantly decreased number of Ki67-positive nuclei in the TSP1-treated group compared with the control group (Figure 2B). The results of scratch wound healing assay indicated that VEGF-C-stimulated LEC migration was




Figure 1. TSP1 (thrombospondin-1) protein levels are upregulated in murine atherosclerotic arteries.

A, Representative Western blot images of TSP1 and β -actin expression in plaque (P)-free descending aorta (DA) segments and atherosclerotic inner curvature (IC) of male *Apoe*^{-/-} mice fed with a Western diet for 12 weeks. **B**, Bar graph shows mean TSP1 protein levels ($n=4$). **C** and **D**, Wild-type and *Apoe*^{-/-} mice (12-wk Western diet) aortic sinus cross-sections were immunostained to investigate TSP1 (green), SMA (smooth muscle cell actin; red), and CD68 (cluster of differentiation; macrophage marker, red) expression. Yellow arrowheads indicate TSP1 expression in adventitia (A) and intraplaque areas ($n=3$). Scale bar, 50 μ m. Statistical analyses were performed using a 2-tailed unpaired Student *t* test (**B**). Data represent mean \pm SEM. L indicates lumen; and M, media.

inhibited by TSP1 treatment (Figure 2C; Figure S1B). Then, we performed the Matrigel tube formation assay to examine the effect of TSP1 on the vessel-forming ability of LEC in vitro. Consistent with the findings of proliferation and migration assays, TSP1-challenged cells had reduced ability of tube formation as suggested by

decreased tube length and lower number of branching points compared with vehicle-treated cells (Figure 2D through 2F; Figure S1C). Additional cell cycle analysis experiments showed increased percentage of cells in the G1 phase and reduced cell percentage in the S phase with TSP1 treatment, suggesting cell cycle inhibition in




Figure 2. TSP1 (thrombospondin-1) suppresses both in vitro and in vivo lymphangiogenesis.

A, Vehicle- and TSP1-pretreated (22 nM, 4 h) human lymphatic endothelial cells (LECs) were stimulated with VEGF (vascular endothelial growth factor)-C (100 ng/mL) and proliferation investigated after 48 hours using the WST (water-soluble tetrazolium)-1 assay. Data are representative of 3 independent experiments performed at least in duplicate. **B**, LECs plated on coverslips were pretreated and stimulated as in **A** for 24 hours. Cells were immunostained for Ki67 (red) and nuclei counterstained with DAPI (4',6-diamidino-2-phenylindole; blue). Images of at least 4 random microscopic fields were captured. Representative images are shown. Scale bar, 20 μ m. Bar graph represents the percentage of Ki67-positive nuclei ($n=4-5$). **C**, LEC migration in response to vehicle (VEGF-C) and TSP1 (VEGF-C+TSP1) was investigated (*Continued*)

TSP1-treated cells (Figure S2). Collectively, these data suggest that TSP1 inhibits lymphangiogenesis in vitro.

To further investigate the antilymphangiogenic potential of TSP1 in vivo—a well-established in vivo lymphangiogenesis model^{26,27}—the Matrigel plug assay was used. Matrigel solution containing VEGF-C with or without TSP1 was injected subcutaneously into wild-type mice on a C57BL/6 background. On day 10 after injection, Matrigel plugs were harvested, fixed, embedded, and processed for immunostaining with the LYVE-1 antibody to determine LV density. As expected, there were remarkably reduced LYVE-1–positive areas in plugs containing VEGF-C+TSP1 compared with VEGF-C–comprising plugs (Figure 2G and 2H). These results of in vivo and in vitro experiments demonstrate that TSP1 possesses an antilymphangiogenic potential.

CD47 Mediates TSP1-Induced Inhibition of Lymphangiogenesis

TSP1 is known to influence cellular responses via binding cell surface receptors mainly CD36 and CD47.^{28,29} Therefore, we next investigated the mRNA levels of *CD47* and *CD36* in LEC using quantitative real-time polymerase chain reaction and compared the levels with human aortic endothelial cells. The quantitative real-time polymerase chain reaction data revealed significantly higher expression of *CD47* mRNA in LEC in comparison to *CD36* (55-fold; Figure 3A). Additionally, LEC *CD47* transcript levels were significantly higher than those in vascular endothelial cells (Figure 3A). Consistently, higher expression of CD47 protein was observed in LECs compared with vascular endothelial cells (Figure S3). These results suggest CD47 as a dominant TSP1 receptor in LEC. Then, to examine the engagement of CD47 in TSP1-stimulated inhibition of lymphangiogenesis, *CD47* was silenced using a gene-specific siRNA pool (Figure 3B). As expected, *CD47*-silenced LEC had a higher proliferation rate compared with control siRNA-treated cells with or without TSP1 challenge (Figure 3C). Interestingly, *CD47* silencing abrogated the effects of TSP1 on LEC migration (Figure 3D) and tube formation (Figure 3E through 3G); however, the migration and tube formation ability of *CD47* knockdown cells were comparable to control cells after vehicle treatment. The results of the Matrigel plug assay showed increased LYVE-1 staining in plugs containing VEGF-C+TSP1+CD47–blocking antibody compared with VEGF-C+TSP1–containing

plugs (Figure 3H). However, no differences in CD45 (immune cell)- and CD34 (angiogenesis)-positive areas were observed in plugs among different groups (Figure S4). Taken together, these data demonstrate that CD47 is involved in mediating antilymphangiogenic effects of TSP1.

TSP1-Induced CD47 Activation Blocks VEGF-C–Stimulated Lymphangiogenic Signaling

Activation of VEGFR3 (vascular endothelial growth factor receptor 3) by its ligand VEGF-C is critical for LV formation and leads to downstream phosphorylation of molecules including AKT, ERK, and eNOS.³⁰ Therefore, we examined whether TSP1 regulates VEGF-C–stimulated prolymphangiogenic signaling in LEC. Briefly, we treated human LEC with TSP1 (22 nM, 16 hours), stimulated with VEGF-C (100 ng/mL, 15 minutes), and subjected cell lysates to immunoblotting. Cells were induced with VEGF-C for 15 minutes because maximum phosphorylation of AKT and ERK1/2 was observed after 15 minutes of incubation with VEGF-C (Figure S5). As expected, VEGF-C treatment induced phosphorylation of AKT (Ser-473), eNOS (Ser-1177), and ERK1/2 (Thr-202/Tyr204) in human LEC (Figure 4A through 4D). However, TSP1-treated cells had no increase in AKT and eNOS activation following VEGF-C stimulation. No differences in ERK1/2 phosphorylation were observed between VEGF-C and VEGF-C+TSP1–treated cells (Figure 4A and 4D). Further, TSP1 treatment suppressed cell proliferation and migration following stimulation with VEGFR3-specific mutant VEGF-C (C156S; Figure S6A and S6B). TSP1 also reduced VEGF-C (C156S)–stimulated AKT and eNOS phosphorylation compared with control cells (Figure S6C). In separate experiments, we determined the role of TSP1 in regulating the expression of VEGFR3. As shown in Figure S7A, TSP1 treatment significantly reduced VEGFR3 protein levels in LEC, suggesting that TSP1 may inhibit lymphangiogenesis via downregulating VEGFR3 expression. However, no differences in *FLT4* (*VEGFR3*) mRNA levels were observed between control and TSP1–treated cells (Figure S7B).

As TSP1 suppressed LEC proliferation and caused cell cycle arrest, we next evaluated the expression of proteins including p21 and p53, which inhibit the cell cycle. Elevated levels of p53 and p21 have been shown to induce cell cycle arrest via inhibiting CDK (cyclin-dependent

Figure 2 Continued. after 24 hours using Culture-Insert 2 Well 24 (ibidi USA). Representative images of wounds at 0 and 24 hours are shown. Scale bar, 1000 μ m. Bar diagram shows quantification of wound closure (n=9). **D**, Vehicle- or TSP1-pretreated LECs were seeded in wells of a Matrigel-coated plate in basal medium containing VEGF-C \pm TSP1 and tube formation determined after 6 hours. Representative images of tube formation are shown. Scale bar, 1000 μ m. Images of random fields were captured and tube length (**E**) and number of branching points (**F**) quantified (n=6). **G** and **H**, Wild-type male mice were injected SC with Matrigel solutions premixed with either VEGF-C or VEGF-C+TSP1. Plugs were isolated after 10 days, sectioned, and immunostained for LYVE-1 (lymphatic vessel endothelial hyaluronan receptor-1). **G**, Representative images of LYVE-1 staining of cross-sections of the Matrigel plugs are shown. Scale bar, 20 μ m. **H**, Quantification of LYVE-1–positive area (n=5–7). Statistical analyses were performed using a 2-tailed unpaired Student *t* test (**A–C**, **E**, and **F**) and a Mann-Whitney *U* test (**H**). Data represent mean \pm SEM.




Figure 3. CD47 (cluster of differentiation) mediates TSP1 (thrombospondin-1)-induced inhibition of lymphangiogenesis.

A, Quantitative real-time polymerase chain reaction (qRT-PCR) was performed to determine the relative mRNA levels of *CD47* and *CD36* in human lymphatic endothelial cells (HLECs) and human aortic endothelial cells (HAoEC). Bar graph represents mRNA levels in comparison to *CD36* ($n=9$). **B**, Control and *CD47* siRNA-treated lymphatic endothelial cells (LECs; 48 h) were utilized to quantify *CD47* transcript expression using qRT-PCR ($n=6$). **C**, WST (water-soluble tetrazolium)-1 assay was conducted to investigate the effects of *CD47* silencing on LEC proliferation in response to TSP1 treatment as described in Figure 2A. Data are representative of 3 independent experiments performed at least in quadruplicate. **D**, Control and *CD47*-silenced LECs were used to evaluate cell migration. Scale bar, 200 μm . (Continued)

kinase) activity.³¹ Treatment with TSP1 stimulated the expression of CDK inhibitors p53 and p21 in LEC after 24 hours (Figure S7C). Further, we investigated the effects of TSP1 treatment on the expression of epsin 1, which degrades VEGFR3 in LEC.²⁶ Immunoblotting data did not show any differences in epsin 1 levels after TSP1 treatment (Figure S7D).

Since CD47 is the highest expressing TSP1's receptor in LECs and silencing of *CD47* in LEC abrogated the effects of TSP1 on lymphangiogenesis, we next investigated the role of CD47 in TSP1-mediated inhibition of AKT and eNOS activation. *CD47* silencing prevented TSP1-induced suppression (16 hours) of AKT and eNOS phosphorylation in LECs (Figure 4E through 4I). The time course of TSP1 treatment experiments with control and *CD47*-knockdown cells showed improved AKT and ERK1/2 phosphorylation in *CD47*-siRNA-treated cells after 8 hours; however, no differences were observed after 4-hour TSP1 treatment (Figure S8). Interestingly, basal eNOS activation was significantly higher in *CD47*-silenced cells compared with control siRNA-treated cells (Figure 4F and 4H). Further, we noticed increased NO production by *CD47*-silenced cells following TSP1 treatment in comparison to TSP1-treated control cells (Figure 4J). As NO bioavailability can be decreased due to enhanced ROS (reactive oxygen species) production, additional experiments were performed to measure ROS generation by control and *CD47*-silenced LECs. The data demonstrated reduced ROS production with *CD47* silencing (Figure 4K). Altogether, these results suggest that TSP1 reduces lymphangiogenesis via CD47-mediated suppression of AKT and eNOS activation.

LEC-Specific Deletion of *Cd47* Suppresses Atherosclerosis

The lymphatic vasculature represents the primary route of cholesterol removal from atherosclerotic vessels, and blockade of arterial lymphatic drainage aggravates atherosclerosis, and knockdown of *CD47* from human LEC prevents TSP1-induced inhibition of lymphangiogenesis (Figure 3). Therefore, to investigate the involvement of LEC *Cd47* in the formation of atherosclerotic lesions, we generated LEC-specific *Cd47* knockout mice (*Cd47^{fl/fl} Lyve-1 Cre^{+/-}*, *Cd47^{ΔLEC}*) by breeding *Cd47^{fl/fl}* with *Lyve-1 Cre^{+/-}* mice (Figure S9A). Deletion of *Cd47* protein from LYVE-1⁺ LV was confirmed using immunostaining experiments

(Figure S9B and S9C). Further, *Cd47* mRNA levels were reduced >90% in LECs isolated from *Cd47^{ΔLEC}* mice compared with those from control mice (Figure S9D and S9E). Eight- to 10-week-old *Cd47^{ΔLEC}* and control *Cd47^{WT}* mice were injected with AAV8-*PCSK9* and fed a Western diet for 16 weeks. En face ORO staining of aortas collected from these mice showed substantially reduced atherosclerosis in *Cd47^{ΔLEC}* mice compared with *Cd47^{WT}* control mice (Figure 5A through 5C). However, no significant differences in plasma total cholesterol, plasma triglycerides, fasting blood glucose levels, weight gain, and body composition (fat, lean, and fluid mass) were observed between *Cd47^{WT}* and *Cd47^{ΔLEC}* mice (Figure 5D through 5G; Figure S10A). Additional hematoxylin and eosin and ORO staining performed on aortic sinus cross-sections from these mice demonstrated smaller lesion areas and reduced lipid accumulation in *Cd47^{ΔLEC}* mice compared with control mice (Figure 5H through 5J). Further histological staining conducted to characterize the aortic root atherosclerotic lesions exhibited increased collagen content and reduced CD68-positive area in *Cd47^{ΔLEC}* mice in comparison to *Cd47^{WT}* mice (Figure 5H, 5K, and 5L). Interaction of CD47 with phagocyte SIRPα (signal regulatory protein alpha) negatively regulates efferocytic clearance of apoptotic cells, which contributes to necrotic core formation in atherosclerotic arteries.¹⁸ Therefore, we quantified the necrotic area in aortic root cross-sections and found no differences between the groups, suggesting no effects of LEC *Cd47* deletion on necrotic core formation and efferocytosis (Figure 5H and 5M; Figure S10B). Next, to determine the effects of LEC-specific *Cd47* deletion on systemic inflammation in hypercholesterolemic atherosclerotic mice, 13 cytokines of commercially available mouse inflammation panel (BioLegend) were measured. Plasma levels of pro/anti-inflammatory cytokines including Il (interleukin)-12p70, Il-23, Il-1α, Ifn-γ, Tnf-α, Mcp-1, Il-6, Il-27, Il-17A, Ifn-β, Gm-csf, and Il-1β, except Il-10, were not different between *Cd47^{WT}* and *Cd47^{ΔLEC}* mice (Figure S11). Altogether, these results indicate that deletion of *Cd47* selectively in LEC attenuates the development of atherosclerosis in hypercholesterolemic mice.

LEC-Specific *Cd47* Knockout Mice Have Increased Arterial LV Density

To investigate the proinflammatory and anti-inflammatory nature of lesional macrophages, aortic root

Figure 3 Continued. Bar graph represents the percentage of migrated cells (n=3–4). **E** through **G**, Control and *CD47*-silenced cells were pretreated as in Figure 2D and seeded in wells of a Matrigel-coated plate in basal medium containing VEGF (vascular endothelial growth factor)-C±TSP1 and tube formation determined. Representative images of tube formation are shown (**E**). Scale bar, 1000 μm. Tube length (**F**) and number of branching points (**G**) quantified (n=7). **H**, Wild-type male mice were injected SC with Matrigel solutions premixed with either VEGF-C, VEGF-C+TSP1+IgG, or VEGF-C+TSP1+CD47-blocking antibody. Plugs were isolated after 10 days, sectioned, and immunostained for LYVE-1 (lymphatic vessel endothelial hyaluronan receptor-1). Representative images of LYVE-1 staining of the cross-sections of the Matrigel plugs and quantification of LYVE-1-positive area are shown (n=5–7). Scale bar, 20 μm. Statistical analyses were performed using 2-way ANOVA (**A**, **C**, **D**, **F**, and **G**) with Bonferroni (**A**), Tukey (**C**, **F**, and **G**), and Sidak (**D**) multiple comparisons test, 2-tailed unpaired Student *t* test (**B**), and Kruskal-Wallis test for multiple comparisons (**H**). Data represent mean±SEM.




Figure 4. TSP1 (thrombospondin-1)-induced CD47 (cluster of differentiation) activation blocks VEGF (vascular endothelial growth factor)-C-stimulated lymphangiogenic signaling.

A through **D**, Lymphatic endothelial cells (LECs) were pretreated with TSP1 (22 nM, 16 h) in 0.5% fetal bovine serum (FBS) containing basal media MV2, stimulated with VEGF-C (15 min), and cell lysates subjected to Western blot analysis. **A**, Representative Western blot images are shown. **B** through **D**, Bar diagrams represent mean protein levels expressed as a ratio of phospho-to-total proteins, AKT (**B**), eNOS (**C**), and ERK1/2 (**D**; n=3). **E**, LECs were treated with control or CD47-siRNA (48 h) and immunoblotting done to determine CD47 expression (ab175388). **F** through **I**, Control or CD47-siRNA-treated cells were treated as in Figure 4A and Western blot experiments (*Continued*)

cross-sections of *Cd47^{WT}* and *Cd47^{ΔLEC}* mice were immunostained for CD68 and iNOS (M1 marker) or Arg1 (M2 marker). Immunostaining analysis demonstrated elevated Arg1 expression in plaque CD68⁺ cells of *Cd47^{ΔLEC}* mice compared with *Cd47^{WT}* control mice; however, similar expression of iNOS was observed (Figure 6A and 6B). These data suggest that *Cd47* deficiency in LECs promotes M2 (proresolving) polarization of lesional macrophages. To determine whether decreased atherosclerotic lesion formation in LEC-specific *Cd47* knockout mice associates with increased adventitial lymphangiogenesis, immunostaining was executed on aortic sinus cross-sections of hypercholesterolemic *Cd47^{WT}* and *Cd47^{ΔLEC}* mice (16 weeks Western diet). Consistent with increased lymphangiogenesis in in vitro and in vivo Matrigel plug assays with inhibition of CD47-mediated signaling, quantitation of LYVE-1-positive vessel-like structures revealed increased adventitial LV density in *Cd47^{ΔLEC}* mice compared with *Cd47^{WT}* mice (Figure 6C and 6D; Figure S10C). However, LV densities in hepatic tissue and ear dermal sheets were comparable (Figure S12). We further examined the accumulation of CD11c⁺ dendritic cells and CD3⁺ T cells in atherosclerotic lesions; however, no significant differences were observed between the 2 groups of mice (Figure S13). Collectively, these data suggest that LEC-specific *Cd47* deletion promotes proresolving macrophage phenotype and augments adventitial LV density in hypercholesterolemic mice.

DISCUSSION

The lymphatic vasculature is indispensable for the removal of extracellular fluid, extravasated leukocytes, and inflammatory cytokines/lipids from the inflamed tissue like arterial wall to the surrounding lymph nodes and back to the systemic circulation, thereby contributing to the attenuation of inflammatory processes, resolution of inflammation, and reduced atherosclerosis.^{2,3,32} In addition, hypercholesterolemia—a causative factor for atherosclerosis development—has been shown to degenerate LV and impair lymphatic drainage, which consequently leads to the accumulation of inflammatory cells in peripheral tissue.^{33,34} In the present study, we explored the role of the TSP1-CD47 signaling axis in regulating lymphangiogenesis and atherosclerosis development. TSP1 expression was observed remarkably elevated in atherosclerotic arteries, particularly in the plaque area and adventitial layer. We found that TSP1 inhibits lymphangiogenesis via CD47-mediated blockade of lymphangiogenic signaling. Moreover, utilizing the novel LEC-specific *Cd47*-deficient

mice, we report that deletion of *Cd47* from LEC in mice suppresses atherosclerosis, which is accompanied by increased adventitial LV density. Collectively, these findings suggest that LEC CD47 activation limits lymphangiogenesis and promotes atherosclerosis.

Various types of vascular cells and immune cells including VSMCs (vascular smooth muscle cells), endothelial cells, adventitial fibroblasts, macrophages, and dendritic cells, which are present in the arterial and periarterial regions, express TSP1.^{35–37} Arterial TSP1 levels upregulate in various vascular pathologies such as abdominal aortic aneurysm, injury-induced neointima formation, and atherosclerosis,^{10–12,38} and conflicting roles of TSP1 in atherosclerosis development have been reported.^{14,15} Further, TSP1 is a potent endogenous inhibitor of angiogenesis,³⁹ and yet knowledge of its ability to inhibit arterial lymphangiogenesis and the signaling mechanisms involved remains scant. To dissect the role of TSP1 in atherosclerotic lesion formation and manipulating arterial lymphangiogenesis, we first investigated the expression and localization of TSP1 protein in murine atherosclerotic vessels. Consistent with increased TSP1 expression in other vascular diseases, upregulated TSP1 protein expression was observed in mouse atherosclerotic arteries compared with nonatherosclerotic vascular tissue.^{11,40,41} We noted elevated TSP1 expression in the intraplaque CD68-positive areas and the adventitial layer of the arterial wall. Increased TSP1 expression has a pathophysiological significance as TSP1 treatment directly inhibited VEGF-C-stimulated LEC proliferation, migration, and both in vitro and in vivo lymphangiogenesis. A previous study reported that TSP1 via binding to CD36 present on macrophages suppresses the expression of lymphangiogenic factors, VEGF-C and VEGF-D, and consequently reduces inflammation-induced corneal lymphangiogenesis.⁴² However, the physiological importance of TSP1-CD36 signaling in lymphangiogenesis has not been explored. Cifarelli et al⁴³ demonstrated reduced VEGF-C-induced lymphangiogenesis by *CD36*-silenced LEC compared with control cells. Conversely, the deletion of CD36 abrogated oxidized LDL (low-density lipoprotein)-induced inhibition of lymphangiogenesis.³⁴ These findings predict the differential effects of LEC CD36 deficiency depending on the presence and concentration of various prolymphangiogenic and antilymphangiogenic factors in the microenvironment.

CD47 serves as a cognate receptor of TSP1, and its expression is upregulated in atherosclerotic arteries.¹⁸ Our results demonstrated CD47 as a major TSP1 receptor in LEC compared with CD36, and knockdown

Figure 4 Continued. executed. **F**, Representative Western blot images are shown. **G** through **I**, Bar diagrams represent pAKT/total AKT (**G**), peNOS/total eNOS (**H**), and pERK1/2/total ERK1/2 (**I**; n=3). **J**, Control or *CD47*-silenced LECs were pretreated with TSP1 (22 nM, 4 h), stimulated with VEGF-C (100 ng/mL, 1 h) and analyzed for NO (nitric oxide) production using DAF-FM diacetate. **K**, Control or *CD47*-silenced LECs were pretreated with vehicle or TSP1 (1 h), incubated with H2DCFDA solution, and fluorescence analyzed using flow cytometry. Statistical analyses were performed using 1-way ANOVA (**B–D**), 2-way ANOVA with Tukey test for multiple comparisons (**G–I** and **K**), and 2-tailed unpaired Student *t* test (**J**). Data represent mean±SEM.




Figure 5. Lymphatic endothelial cell (LEC)-specific *Cd47* deficiency reduces atherosclerotic lesion formation.

A through **M**, Male *Cd47^{WT}* and *Cd47^{ΔLEC}* mice were injected with AAV (adeno-associated virus) 8-*PCSK9* IP, fed a Western diet for 16 weeks and atherosclerosis analyzed. **A**, Representative in situ images of aortic arch (yellow arrowheads point to atherosclerotic lesions). Scale bar, 2 mm. **B**, Representative en face oil red O (ORO) staining of aorta. Scale bar, 5 mm. **C**, Quantification of plaque area in aorta ($n=13-14$). **D** through **G**, Bar diagrams show plasma total cholesterol (**D**), plasma triglycerides (**E**), fasting blood glucose levels (**F**), and body composition (*Continued*)

of *CD47* in LEC and treatment with CD47-blocking antibody prevented the effects of TSP1 on lymphangiogenesis. Similarly, Hawighorst et al⁴⁴ showed little-to-no expression of CD36 receptor in LYVE-1⁺ LEC present in murine skin tumor tissue and human skin. However, we believe that LECs also have significant expression of CD36 as shown by us³⁴ and other groups^{43,45} using both in vitro and in vivo studies. CD36 is a scavenger receptor, which helps in lipid internalization by various cell types including LECs. It has been shown by Wong et al⁴⁵ that LECs mainly utilize fatty acid oxidation for their energy needs. LEC fatty acid oxidation promotes LEC proliferation as it provides acetyl-CoA, which helps to sustain the Krebs cycle and dNTP synthesis for cell division. Additional experiments performed using control- and *CD47*-siRNA-treated cells suggested that TSP1 suppresses VEGF-C-stimulated lymphangiogenic signaling involving AKT and eNOS phosphorylation in control cells but not in *CD47*-silenced LECs. Exposure to TSP1 led to a significant increase in the expression of cell cycle inhibitors including p21 and p53. These data indicate that TSP1 is a potent lymphangiogenesis inhibitor and induces LEC senescence. Similarly, Gao et al⁴⁶ reported that TSP1 accelerates brain endothelial cell senescence in a CD47-dependent manner. It is possible that blockade of TSP1-CD47 signaling in LEC via CD47 deletion/blocking antibodies may contribute to LEC self-renewal via upregulating the expression of c-myc.⁴⁷ However, we cannot exclude the engagement of other TSP1's receptors like CD36 or integrins in TSP1-mediated inhibition of LEC proliferation as a significant decrease in cell growth was observed in *CD47*-silenced cells following TSP1 treatment. TSP1 treatment downregulated VEGFR3 expression in LECs without affecting *FLT4* (*VEGFR3*) mRNA and epsin 1 protein levels. TSP1 exposure may elevate the expression of other epsin proteins, which cause VEGFR3 degradation in the Golgi compartment,²⁶ and promotes proteasomal- and autophagy-mediated VEGFR3 degradation.⁴⁸ Oxidative stress via reducing NO bioavailability suppresses lymphangiogenesis.²⁶ Moreover, *CD47* deficiency in LECs improved NO production following TSP1 exposure and reduced TSP1-induced ROS production. Consistently, previous studies have shown that TSP1 reduces NO production in vascular endothelial cells and promotes ROS production in VSMCs, endothelial cells, and macrophages.^{49–51} Besides, it has been shown that TSP1 via stimulating ROS production increases p53 levels and inhibits cell growth.⁵² However, it is not known whether treatment

with antioxidants or NO donors can prevent TSP1-induced inhibition of lymphangiogenesis.

CD47 is a ubiquitously expressed protein and has pleiotropic effects in various cell types via interacting with multiple binding partners including TSP1, SIRP α , and integrins.⁵³ Global CD47-deficient mice and atheroprone mice treated with CD47-blocking antibodies are protected from atherosclerosis; however, myeloid cell-specific CD47 loss augments lesion formation suggesting the significance of cell-specific CD47 blockade in atherosclerosis.^{18,20} We investigated the role of lymphatic endothelium *Cd47* in atherosclerosis via generating novel LEC-specific *Cd47* knockout mice and showed that LEC CD47 contributes to atherosclerotic lesion formation without affecting the metabolic profile. Upregulated CD47 expression in atherosclerotic arteries inhibits phagocytosis, induces necrotic core formation, increases inflammation, and exacerbates atherosclerosis.^{18,54} Nonetheless, comparable necrotic core areas in *Cd47*^{WT} and *Cd47*^{LEEC} mice were observed, which demonstrates no role of LEC CD47 in regulating efferocytosis. Additionally, no differences in plasma proinflammatory cytokine levels were detected. Interestingly, we found reduced Il-10 levels in plasma of *Cd47*^{LEEC} mice. Further experiments revealed reduced arterial inflammation in *Cd47*^{LEEC} mice compared with control mice. CD47-SIRP α interaction plays important roles in regulating cell adhesion and migration. Deletion of *Cd47* in LECs may affect transmigration of immune cells into LV.

LVs play an important role in the maintenance of tissue homeostasis and are induced by inflammation to aid in the drainage of inflammatory cells, cytokines, and fluids, thereby promoting the resolution of inflammation. During atherogenesis, lymphangiogenesis is induced due to an increased concentration of lymphangiogenic factors like VEGF-C and VEGF-D,⁵⁵ but elevated levels of anti-lymphangiogenic factors in the advanced plaques prevent LV formation or regress them.⁵⁶ Our immunostaining analysis performed on aortic root cross-sections showed increased adventitial LV density with LEC-specific *Cd47* deletion without affecting the accumulation of intraplaque dendritic cells and T cells. LVs need to be functional to play their beneficial role in inflammatory diseases like atherosclerosis. None of the currently available methodologies allow the direct measurement of LV function in the adventitial layer of murine aortic arch inner curvature (area of increased atherosclerosis due to disturbed flow) and the aortic sinus. We speculate that there may be an improvement in lymphatic function with LEC *Cd47* loss. It would be interesting to determine the number of immune

Figure 5 Continued. (fat, lean, and fluid mass; **G**; n=8–13). **H**, Representative images of aortic root cross-sections stained with H&E (neointima area), ORO (lipid accumulation), Masson trichrome (collagen content), CD68 (cluster of differentiation; macrophage burden), and necrotic area (encircled in red). Scale bar, 200 μ m. **I** through **M**, Bar diagrams show neointima area (**I**), lipid deposition (**J**), collagen content (**K**), macrophage accumulation (**L**), and necrotic area (**M**) in aortic root sections (n=5–9). Statistical analyses were performed using a 2-tailed unpaired Mann-Whitney *U* test (**C**, **E**, and **M**), 2-tailed unpaired Student *t* test (**D**, **F**, and **I–L**), and 2-way ANOVA followed by Sidak post hoc test (**G**). Data represent mean \pm SEM.




Figure 6. Lymphatic endothelial cell (LEC)-specific deletion of *Cd47* (cluster of differentiation) in mice increases arterial lymphatic vessel (LV) density.

A through **D**, Aortic root cross-sections from male AAV (adeno-associated virus) 8-*PCSK9*-injected $Cd47^{WT}$ and $Cd47^{ALEC}$ mice (16-wk Western diet) were immunostained for CD68, iNOS (inducible NO [nitric oxide] synthase), Arg1 (arginase 1), and LYVE-1 (lymphatic vessel endothelial hyaluronan receptor-1). Nuclei were counterstained with DAPI (4',6-diamidino-2-phenylindole; blue). Representative confocal images of iNOS (green; **A**) and CD68 (red); Arg1 (green; **B**) and CD68 (red; scale bar, 50 μ m); and LYVE-1 staining (red; **C**; scale bar, 20 μ m) are shown ($n=5-6$). Statistical analyses were performed using a 2-tailed unpaired Student *t* test (**A** and **B**) and a Mann-Whitney *U* test (**C**). Data represent mean \pm SEM. A indicates adventitia; M, media; and P, plaque.

cells in the proximity of adventitial LV as an indirect measure of lymphatic function. Most of the previous studies evaluated lymphatic function either in dermal tissue or the peritoneal cavity and associated it with RCT and atherosclerosis development. Martel et al² using aorta transplantation experiments showed reduced RCT in *ApoE*^{-/-} mice treated with antilymphangiogenic VEGFR3–blocking antibody compared with control antibody–treated mice, suggesting the critical role of lymphatics in macrophage RCT from the arterial wall. Yeo et al³ recently devised a novel method to investigate the role of abdominal periaortic lymphatic drainage in descending aorta atherosclerosis and reported increased atherosclerosis with impaired drainage. Future functional studies using new innovative tools are required to investigate lymphatic function in the different regions of aortic arch and medium-sized arteries and determine its association with atherosclerosis development. Further, to fully understand the role of TSP1-CD47 signaling in regulating lymphatic function in mice, it will be important to quantify the levels of TSP1 in lymph fluid. As Lyve-1 is also expressed in a subset of M2-like aortic macrophages and Lyve-1⁺ macrophages affect arterial wall remodeling, it remains to be investigated whether deletion of *Cd47* utilizing *Lyve-1* Cre mice alters arterial stiffness and collagen deposition.⁵⁷ It is possible that in vivo deletion of *Cd47* from M2-like macrophages in the murine models utilized in the present study increased anti-inflammatory capacity of macrophages via blocking TSP1-CD47 signaling. Though we did not see any differences in the expression of CD47 in lesional macrophages between *Cd47*^{WT} and *Cd47*^{ΔLEC} mice, the role of macrophage CD47 may need to be ruled out using other LEC-specific Cre mouse line. Another limitation of the present study includes not determining the activation of LEC eNOS and AKT in our mouse models.

In summary, the present study suggests that TSP1 via activation of LEC CD47 restricts lymphangiogenesis, and deletion of *Cd47* in lymphatic endothelium increases arterial lymphangiogenesis and attenuates atherosclerosis. These findings demonstrate LEC CD47 inhibition as a potential therapeutic target to promote arterial lymphangiogenesis and reduce atherosclerosis.

ARTICLE INFORMATION

Received December 15, 2022; accepted May 16, 2023.

Affiliations

Department of Pharmaceutical Sciences, College of Pharmacy, The University of Tennessee Health Science Center, Memphis (B.S., R.V.A., N.P., I.K., M.S., F.A.E., F.P., M.G., U.P.S., S.K.). Vascular Biology Center, Medical College of Georgia, Augusta University (W.A., M.C.-S.).

Acknowledgments

The authors are very thankful to Dr Gabor Csanyi, Vascular Biology Center, Augusta University, for his feedback on the study design. In addition, we thank Esther Magdalena Marquez Wilkins, Neuroscience Imaging Center, UTHSC, for her help with confocal microscopy. B. Singla designed the study, performed most of the experiments, analyzed data, and wrote and edited the manuscript. R.V.

Aithbathula, N. Pervaiz, I. Kathuria, and W. Ahn assisted with data analysis. M. Swanson and F. Park helped with Matrigel plug paraffin embedding, sectioning, and further processing. F.A. Ekuban and M. Gyamfi performed plasma triglyceride quantitation. M. Cherian-Shaw measured animal weight every week. U.P. Singh and S. Kumar provided feedback on the manuscript.

Sources of Funding

This work was supported by the National Institutes of Health (NIH) grants (K99HL146954 and R00HL146954), American Heart Association postdoctoral fellowship (17POST33661254), and UTHSC College of Pharmacy Research Seed Grant awarded to B. Singla and NIH grant (R01AA028806) given to M. Gyamfi.

Disclosures

None.

Supplemental Material

Supplemental Materials and Methods

Figures S1–S13

Tables S1–S2

Major Resources Table

REFERENCES

- Lawton JS, Tamis-Holland JE, Bangalore S, Bates ER, Beckie TM, Bischoff JM, Bittl JA, Cohen MG, DiMaio JM, Don CW, et al. 2021 ACC/AHA/SCAI guideline for coronary artery revascularization: executive summary: a report of the American College of Cardiology/American Heart Association joint committee on clinical practice guidelines. *Circulation*. 2022;145:e4–e17. doi: 10.1161/CIR.0000000000001039
- Martel C, Li W, Fulp B, Platt AM, Gautier EL, Westerterp M, Bittman R, Tall AR, Chen SH, Thomas MJ, et al. Lymphatic vasculature mediates macrophage reverse cholesterol transport in mice. *J Clin Invest*. 2013;123:1571–1579. doi: 10.1172/JCI63685
- Yeo KP, Lim HY, Thiam CH, Azhar SH, Tan C, Tang Y, See WQ, Koh XH, Zhao MH, Phua ML, et al. Efficient aortic lymphatic drainage is necessary for atherosclerosis regression induced by ezetimibe. *Sci Adv*. 2020;6:eabc2697. doi: 10.1126/sciadv.abc2697
- Rademakers T, van der Vorst EP, Daissormont IT, Otten JJ, Theodorou K, Theelen TL, Gijbels M, Anisimov A, Nurmi H, Lindeman JH, et al. Adventitial lymphatic capillary expansion impacts on plaque T cell accumulation in atherosclerosis. *Sci Rep*. 2017;7:45263. doi: 10.1038/srep45263
- Vuorio T, Nurmi H, Moulton K, Kurkipuro J, Robciuc MR, Ohman M, Heino SE, Samaranyake H, Heikura T, Alitalo K, et al. Lymphatic vessel insufficiency in hypercholesterolemic mice alters lipoprotein levels and promotes atherogenesis. *Arterioscler Thromb Vasc Biol*. 2014;34:1162–1170. doi: 10.1161/ATVBAHA.114.302528
- Milasan A, Smaani A, Martel C. Early rescue of lymphatic function limits atherosclerosis progression in *Ldlr*^{-/-} mice. *Atherosclerosis*. 2019;283:106–119. doi: 10.1016/j.atherosclerosis.2019.01.031
- Bornstein P. Thrombospondins function as regulators of angiogenesis. *J Cell Commun Signal*. 2009;3:189–200. doi: 10.1007/s12079-009-0060-8
- Bornstein P. Thrombospondins as matricellular modulators of cell function. *J Clin Invest*. 2001;107:929–934. doi: 10.1172/JCI12749
- Isenberg JS, Romeo MJ, Abu-Asab M, Tsokos M, Oldenborg A, Pappan L, Wink DA, Frazier WA, Roberts DD. Increasing survival of ischemic tissue by targeting CD47. *Circ Res*. 2007;100:712–720. doi: 10.1161/01.RES.0000259579.357874e
- Raugi GJ, Mullen JS, Bark DH, Okada T, Mayberg MR. Thrombospondin deposition in rat carotid artery injury. *Am J Pathol*. 1990;137:179–185.
- Sajid M, Hu Z, Guo H, Li H, Stouffer GA. Vascular expression of integrin-associated protein and thrombospondin increase after mechanical injury. *J Invest Med*. 2001;49:398–406. doi: 10.2310/6650.2001.33784
- Riessen R, Kearney M, Lawler J, Isner JM. Immunolocalization of thrombospondin-1 in human atherosclerotic and restenotic arteries. *Am Heart J*. 1998;135:357–364. doi: 10.1016/s0002-8703(98)70105-x
- Stenina OI, Topol EJ, Plow EF. Thrombospondins, their polymorphisms, and cardiovascular disease. *Arterioscler Thromb Vasc Biol*. 2007;27:1886–1894. doi: 10.1161/ATVBAHA.107.141713
- Moura R, Tjwa M, Vandervoort P, Van Kerckhoven S, Holvoet P, Hoylaerts MF. Thrombospondin-1 deficiency accelerates atherosclerotic plaque maturation in *ApoE*^{-/-} mice. *Circ Res*. 2008;103:1181–1189. doi: 10.1161/CIRCRESAHA.108.185645

15. Ganguly R, Khanal S, Mathias A, Gupta S, Lallo J, Sahu S, Ohyanan V, Patel A, Storm K, Datta S, et al. TSP-1 (thrombospondin-1) deficiency protects ApoE(-/-) mice against leptin-induced atherosclerosis. *Arterioscler Thromb Vasc Biol.* 2021;41:e112–e127. doi: 10.1161/ATVBAHA.120.314962
16. Isenberg JS, Ridnour LA, Perruccio EM, Espey MG, Wink DA, Roberts DD. Thrombospondin-1 inhibits endothelial cell responses to nitric oxide in a cGMP-dependent manner. *Proc Natl Acad Sci USA.* 2005;102:13141–13146. doi: 10.1073/pnas.0502977102
17. Lahdenranta J, Hagendoorn J, Padera TP, Hoshida T, Nelson G, Kashiwagi S, Jain RK, Fukumura D. Endothelial nitric oxide synthase mediates lymphangiogenesis and lymphatic metastasis. *Cancer Res.* 2009;69:2801–2808. doi: 10.1158/0008-5472.CAN-08-4051
18. Kojima Y, Volkmer JP, McKenna K, Civelek M, Lusis AJ, Miller CL, Direnzo D, Nanda V, Ye J, Connolly AJ, et al. CD47-blocking antibodies restore phagocytosis and prevent atherosclerosis. *Nature.* 2016;536:86–90. doi: 10.1038/nature18935
19. Jarr KU, Nakamoto R, Doan BH, Kojima Y, Weissman IL, Advani RH, Iagaru A, Leeper NJ. Effect of CD47 blockade on vascular inflammation. *N Engl J Med.* 2021;384:382–383. doi: 10.1056/NEJMc2029834
20. Singla B, Lin HP, Ahn W, Xu J, Ma Q, Sghayyer M, Dong K, Cherian-Shaw M, Zhou J, Huo Y, et al. Loss of myeloid cell-specific SIRPalpha, but not CD47, attenuates inflammation and suppresses atherosclerosis. *Cardiovasc Res.* 2022;118:3097–3111. doi: 10.1093/cvr/cvab369
21. Gubbels Bupp MR. Sex, the aging immune system, and chronic disease. *Cell Immunol.* 2015;294:102–110. doi: 10.1016/j.cellimm.2015.02.002
22. Bourassa PA, Milos PM, Gaynor BJ, Breslow JL, Aiello RJ. Estrogen reduces atherosclerotic lesion development in apolipoprotein E-deficient mice. *Proc Natl Acad Sci USA.* 1996;93:10022–10027. doi: 10.1073/pnas.93.19.10022
23. Kumar S, Kang DW, Rezvan A, Jo H. Accelerated atherosclerosis development in C57Bl6 mice by overexpressing AAV-mediated PCSK9 and partial carotid ligation. *Lab Invest.* 2017;97:935–945. doi: 10.1038/labinvest.2017.47
24. Singla B, Lin HP, Chen A, Ahn W, Ghoshal P, Cherian-Shaw M, White J, Stansfield BK, Csanyi G. Role of R-spondin 2 in arterial lymphangiogenesis and atherosclerosis. *Cardiovasc Res.* 2021;117:1489–1509. doi: 10.1093/cvr/cvaa244
25. McCrohan MB, Huang SW, Sleasman JW, Klein PA, Kao KJ. Plasma thrombospondin as an indicator of intravascular platelet activation in patients with vasculitis. *Thromb Haemost.* 1987;58:850–852.
26. Wu H, Rahman HNA, Dong Y, Liu X, Lee Y, Wen A, To KH, Xiao L, Birsner AE, Bazinet L, et al. Epsin deficiency promotes lymphangiogenesis through regulation of VEGFR3 degradation in diabetes. *J Clin Invest.* 2018;128:4025–4043. doi: 10.1172/JCI96063
27. Song L, Ding S, Ge Z, Zhu X, Qiu C, Wang Y, Lai E, Yang W, Sun Y, Chow SA, et al. Nucleoside/nucleotide reverse transcriptase inhibitors attenuate angiogenesis and lymphangiogenesis by impairing receptor tyrosine kinases signalling in endothelial cells. *Br J Pharmacol.* 2018;175:1241–1259. doi: 10.1111/bph.14036
28. Roberts DD, Miller TW, Rogers NM, Yao M, Isenberg JS. The matricellular protein thrombospondin-1 globally regulates cardiovascular function and responses to stress via CD47. *Matrix Biol.* 2012;31:162–169. doi: 10.1016/j.matbio.2012.01.005
29. Adams JC, Lawler J. The thrombospondins. *Cold Spring Harb Perspect Biol.* 2011;3:a009712. doi: 10.1101/cshperspecta.a009712
30. Makinen T, Veikkola T, Mustjoki S, Karpanen T, Catimel B, Nice EC, Wise L, Mercer A, Kowalski H, Kerjaschki D, et al. Isolated lymphatic endothelial cells transduce growth, survival and migratory signals via the VEGF-C/D receptor VEGFR-3. *EMBO J.* 2001;20:4762–4773. doi: 10.1093/emboj/20.17.4762
31. Dotto GP, p21(WAF1/Cip1): more than a break to the cell cycle? *Biochim Biophys Acta.* 2000;1471:M43–M56. doi: 10.1016/s0304-419x(00)00019-6
32. Oliver G, Kipnis J, Randolph GJ, Harvey NL. The lymphatic vasculature in the 21(st) century: novel functional roles in homeostasis and disease. *Cell.* 2020;182:270–296. doi: 10.1016/j.cell.2020.06.039
33. Lim HY, Rutkowski JM, Helft J, Reddy ST, Swartz MA, Randolph GJ, Angeli V. Hypercholesterolemic mice exhibit lymphatic vessel dysfunction and degeneration. *Am J Pathol.* 2009;175:1328–1337. doi: 10.2353/ajpath.2009.080963
34. Singla B, Lin HP, Ahn W, White J, Csanyi G. Oxidatively modified LDL suppresses lymphangiogenesis via CD36 signaling. *Antioxidants (Basel).* 2021;10:331. doi: 10.3390/antiox10020331
35. Sid B, Sartelet H, Bellon G, El Btaouri H, Rath G, Delorme N, Haye B, Martiny L. Thrombospondin 1: a multifunctional protein implicated in the regulation of tumor growth. *Crit Rev Oncol Hematol.* 2004;49:245–258. doi: 10.1016/j.critrevonc.2003.09.009
36. Armstrong LC, Bornstein P. Thrombospondins 1 and 2 function as inhibitors of angiogenesis. *Matrix Biol.* 2003;22:63–71. doi: 10.1016/s0945-053x(03)00005-2
37. Doyen V, Rubio M, Braun D, Nakajima T, Abe J, Saito H, Delespesse G, Sarfati M. Thrombospondin 1 is an autocrine negative regulator of human dendritic cell activation. *J Exp Med.* 2003;198:1277–1283. doi: 10.1084/jem.20030705
38. Liu Z, Morgan S, Ren J, Wang Q, Annis DS, Mosher DF, Zhang J, Sorenson CM, Sheibani N, Liu B. Thrombospondin-1 (TSP1) contributes to the development of vascular inflammation by regulating monocytic cell motility in mouse models of abdominal aortic aneurysm. *Circ Res.* 2015;117:129–141. doi: 10.1161/CIRCRESAHA.117.305262
39. Good DJ, Poverini PJ, Rastinejad F, Le Beau MM, Lemons RS, Frazier WA, Bouck NP. A tumor suppressor-dependent inhibitor of angiogenesis is immunologically and functionally indistinguishable from a fragment of thrombospondin. *Proc Natl Acad Sci USA.* 1990;87:6624–6628. doi: 10.1073/pnas.87.17.6624
40. Yang H, Zhou T, Sorenson CM, Sheibani N, Liu B. Myeloid-derived TSP1 (thrombospondin-1) contributes to abdominal aortic aneurysm through suppressing tissue inhibitor of metalloproteinases-1. *Arterioscler Thromb Vasc Biol.* 2020;40:e350–e366. doi: 10.1161/ATVBAHA.120.314913
41. Rogers NM, Sharifi-Sanjani M, Yao M, Ghimire K, Bienes-Martinez R, Mutchler SM, Knupp HE, Baust J, Novelli EM, Ross M, et al. TSP1-CD47 signaling is upregulated in clinical pulmonary hypertension and contributes to pulmonary arterial vasculopathy and dysfunction. *Cardiovasc Res.* 2017;113:15–29. doi: 10.1093/cvr/cvw218
42. Cursiefen C, Maruyama K, Bock F, Saban D, Sadrai Z, Lawler J, Dana R, Masli S. Thrombospondin 1 inhibits inflammatory lymphangiogenesis by CD36 ligation on monocytes. *J Exp Med.* 2011;208:1083–1092. doi: 10.1084/jem.20092277
43. Cifarelli V, Appak-Baskoy S, Peche VS, Kluzak A, Shew T, Narendran R, Pietka KM, Cella M, Walls CW, Czepielewski R, et al. Visceral obesity and insulin resistance associate with CD36 deletion in lymphatic endothelial cells. *Nat Commun.* 2021;12:3350. doi: 10.1038/s41467-021-23808-3
44. Hawighorst T, Oura H, Streit M, Janes L, Nguyen L, Brown LF, Oliver G, Jackson DG, Detmar M. Thrombospondin-1 selectively inhibits early-stage carcinogenesis and angiogenesis but not tumor lymphangiogenesis and lymphatic metastasis in transgenic mice. *Oncogene.* 2002;21:7945–7956. doi: 10.1038/sj.onc.1205956
45. Wong BW, Wang X, Zecchin A, Thienpont B, Cornelissen I, Kalucka J, Garcia-Caballero M, Missiaen R, Huang H, Bruning U, et al. The role of fatty acid beta-oxidation in lymphangiogenesis. *Nature.* 2017;542:49–54. doi: 10.1038/nature21028
46. Gao Q, Chen K, Gao L, Zheng Y, Yang YG. Thrombospondin-1 signaling through CD47 inhibits cell cycle progression and induces senescence in endothelial cells. *Cell Death Dis.* 2016;7:e2368. doi: 10.1038/cddis.2016.155
47. Kaur S, Soto-Pantoja DR, Stein EV, Liu C, Elkahoun AG, Pendrak ML, Nicolae A, Singh SP, Nie Z, Levens D, et al. Thrombospondin-1 signaling through CD47 inhibits self-renewal by regulating c-Myc and other stem cell transcription factors. *Sci Rep.* 2013;3:1673. doi: 10.1038/srep01673
48. Lecker SH, Goldberg AL, Mitch WE. Protein degradation by the ubiquitin-proteasome pathway in normal and disease states. *J Am Soc Nephrol.* 2006;17:1807–1819. doi: 10.1681/ASN.2006010083
49. Isenberg JS, Frazier WA, Roberts DD. Thrombospondin-1: a physiological regulator of nitric oxide signaling. *Cell Mol Life Sci.* 2008;65:728–742. doi: 10.1007/s00018-007-7488-x
50. Csanyi G, Yao M, Rodriguez Al, Al Ghoulah I, Sharifi-Sanjani M, Frazziano G, Huang X, Kelley EE, Isenberg JS, Pagano PJ. Thrombospondin-1 regulates blood flow via CD47 receptor-mediated activation of NADPH oxidase 1. *Arterioscler Thromb Vasc Biol.* 2012;32:2966–2973. doi: 10.1161/ATVBAHA.112.300031
51. Csanyi G, Feck DM, Ghoshal P, Singla B, Lin H, Nagarajan S, Meijles DN, Al Ghoulah I, Cantu-Medellin N, Kelley EE, et al. CD47 and Nox1 mediate dynamic fluid-phase macropinocytosis of native LDL. *Antioxid Redox Signal.* 2017;26:886–901. doi: 10.1089/ars.2016.6834
52. Meijles DN, Sahoo S, Al Ghoulah I, Amaral JH, Bienes-Martinez R, Knupp HE, Attaran S, Sembrat JC, Nouraei SM, Rojas MM, et al. The matricellular protein TSP1 promotes human and mouse endothelial cell senescence through CD47 and Nox1. *Sci Signal.* 2017;10:eaaj1784. doi: 10.1126/scisignal.aaj1784
53. Sick E, Jeanne A, Schneider C, Dedieu S, Takeda K, Martiny L. CD47 update: a multifaceted actor in the tumour microenvironment of potential therapeutic interest. *Br J Pharmacol.* 2012;167:1415–1430. doi: 10.1111/j.1476-5381.2012.02099.x
54. Hansson GK, Libby P, Tabas I. Inflammation and plaque vulnerability. *J Intern Med.* 2015;278:483–493. doi: 10.1111/joim.12406

-
55. Grzegorek I, Drozd K, Chmielewska M, Gomulkiewicz A, Jablonska K, Piotrowska A, Karczewski M, Janczak D, Podhorska-Okolow M, Dziegiel P, et al. Arterial wall lymphangiogenesis is increased in the human iliac atherosclerotic arteries: involvement of CCR7 receptor. *Lymphat Res Biol*. 2014;12:222–231. doi: 10.1089/lrb.2013.0048
56. Taher M, Nakao S, Zandi S, Melhorn MI, Hayes KC, Hafezi-Moghadam A. Phenotypic transformation of intimal and adventitial lymphatics in atherosclerosis: a regulatory role for soluble VEGF receptor 2. *FASEB J*. 2016;30:2490–2499. doi: 10.1096/fj.201500112
57. Lim HY, Lim SY, Tan CK, Thiam CH, Goh CC, Carbajo D, Chew SHS, See P, Chakarov S, Wang XN, et al. Hyaluronan receptor LYVE-1-expressing macrophages maintain arterial tone through hyaluronan-mediated regulation of smooth muscle cell collagen. *Immunity*. 2018;49:326–341.e7. doi: 10.1016/j.immuni.2018.06.008