

# Dressed or undressed? How to measure children's body weight in overweight surveillance?

Laura Censi<sup>1,\*</sup>, Angela Spinelli<sup>2</sup>, Romana Roccaldo<sup>1</sup>, Noemi Bevilacqua<sup>1</sup>, Anna Lamberti<sup>2</sup>, Veronica Angelini<sup>1</sup>, Paola Nardone<sup>2</sup> and Giovanni Baglio<sup>2</sup>

<sup>1</sup>Agricultural Research Council, Food and Nutrition Research Centre (CRA-NUT), Via Ardeatina 546, 00178 Rome, Italy; <sup>2</sup>National Centre for Epidemiology, Surveillance and Health Promotion, National Institute of Health (Istituto Superiore di Sanità), Rome, Italy

Submitted 7 January 2013; Final revision received 16 July 2013; Accepted 8 October 2013; First published online 15 November 2013

## Abstract

*Objective:* To simplify body weight measurement and, particularly, to encourage children and their parents to participate in the Italian nutritional surveillance system OKkio alla SALUTE, children were measured with clothes and then the weight was corrected for the estimated weight of the clothes. In the present study we compared the children's weight measured in underwear, as recommended by the WHO ( $W_{\text{WHO}}$ ), with that obtained using the OKkio alla SALUTE protocol ( $W_{\text{OK}}$ ) and investigated how the latter affects the calculation of BMI and the assessment of overweight and obesity prevalence.

*Design:* Weight (twice in close sequence, with and without clothing) and height were measured. A checklist was used to describe the type of clothing worn. The estimated weight of clothing was subtracted from the  $W_{\text{OK}}$ . BMI was calculated considering both values of weight and height; ponderal status was defined using both the International Obesity Task Force and WHO BMI cut-offs.

*Setting:* Thirty-seven third grade classes of thirteen primary schools in Rome and in two towns in the Lazio Region were recruited.

*Subjects:* The anthropometric measurements were taken on 524 children aged 8–9 years.

*Results:* The error in the calculation of BMI from  $W_{\text{OK}}$  was very low, 0.005 kg/m<sup>2</sup> (95% CI -0.185, 0.195 kg/m<sup>2</sup>); the agreement between the percentages of overweight (not including obesity) and obese children calculated with the two methods was very close to 1 ( $\kappa = 0.98$ ).

*Conclusions:* The error in BMI and in nutritional classification can be considered minor in a surveillance system for monitoring overweight/obesity, but eases the procedure for measuring children.

**Keywords**  
Overweight  
Obesity  
Nutritional surveillance  
Body weight measure  
Children

The prevalence of obesity is increasing so rapidly worldwide that this phenomenon is referred to as a 'global epidemic' by the WHO and is a major public health problem in industrialized countries<sup>(1)</sup>. According to the WHO Regional Office for Europe, this trend is particularly alarming in children and adolescents because the prevalence is increasing constantly and is now ten times higher than that found in the 1970s. Childhood obesity is a condition that, once it has developed, tends to persist into adulthood and predisposes to a series of pathological alterations in both the short and long term<sup>(2,3)</sup>. Over 60% of children overweight before puberty will be overweight in early adulthood; this reduces the average age of onset of non-communicable diseases and significantly influences the burden on health services, increasing the number of medical contacts and treatment during adulthood<sup>(2,4)</sup>. BMI is the most commonly used

measure of ponderal status in a population and is calculated as body weight (in kilograms) divided by height (in metres) squared<sup>(1,5)</sup>. It may be an appropriate index to define overweight in children and adolescents<sup>(6)</sup> and to identify groups that are at increased risk for overweight and obesity-related health problems.

A national surveillance system to monitor the prevalence of overweight/obesity is fundamental to plan successful preventive health interventions. Surveillance of childhood overweight and obesity should be more widespread in order to monitor its trend and to permit inter-country comparisons. The important elements of a surveillance system are simplicity, acceptability, representativeness and cost<sup>(7)</sup>. In Italy the national survey of the prevalence of childhood overweight and obesity (OKkio alla SALUTE) was developed and started in 2008 and involves a representative sample of about 40 000 schoolchildren

\*Corresponding author: Email laura.censi@entecra.it

aged 8–9 years every two years<sup>(8,9)</sup>. OKkio alla SALUTE is part of the Childhood Obesity Surveillance Initiative (COSI) established by the WHO Regional Office for Europe<sup>(10)</sup>.

The WHO anthropometric evaluation procedure<sup>(5)</sup> specifies that weight and height must be measured on a subject wearing only light underwear. On the other hand, the surveillance system OKkio alla SALUTE decided to measure body weight with clothing and then to rectify the data according to the estimated weight of the clothes. The assumption is that the OKkio alla SALUTE procedure is more practical and quick and it may increase the participation rate of children and their parents.

The aim of the present study was to compare the weight obtained according to WHO recommendations<sup>(5)</sup>, that is measuring the weight of the children wearing only underwear, with that of clothed children adjusted to take account of the weight of their clothes, as for the OKkio alla SALUTE protocol. The second aim was to assess how the latter method to measure body weight affects the calculation of BMI and the assessment of overweight and obesity prevalence in schoolchildren.

## Methods

Thirty-seven third grade classes of thirteen primary schools in the municipality of Rome and in two towns in the Lazio Region were recruited. All students of the selected classes were enrolled on a voluntary basis. This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the ethical committee of the National Institute of Health. Prior to their acceptance, the children's parents or other caregivers were fully informed about the objectives and methods of the study and signed a consent form.

Anthropometric measurements were taken on 524 children between 8 and 9 years of age, with a higher percentage of boys (57.3%), due to a greater number of males in the sampled classes. Measurements were made by two researchers, trained and standardized according to WHO recommendations<sup>(5)</sup>. Body weight was measured to the nearest 50 g using an electronic scale SECA 872<sup>TM</sup> (Hamburg, Germany), with the child in fasting condition (or after a light breakfast) and after emptying the bladder. Height was measured to the nearest 0.1 cm by a stadiometer SECA 214<sup>TM</sup>, with the child without shoes, standing up, weight being equally distributed on both feet, with his/her head in the Frankfort plane and the back of the head, buttocks and heels (if possible) in contact with the vertical board. Anthropometric measurements were taken at school in the morning, one child at a time in a quiet room, in the presence of the teacher, ensuring confidentiality for each child. Data of weight, height, date of birth and gender were collected and transcribed on the measurement card, as stated in the OKkio alla SALUTE protocol.

**Table 1** The OKkio alla SALUTE clothing checklist and estimated weight of the clothes (kg)

| Clothing item                          | Estimated weight (kg) |
|--|-----------------------|
| Long sleeve shirt                      | 0.15850               |
| T-shirt                                | 0.10850               |
| Long denim trousers                    | 0.37500               |
| Short denim trousers                   | 0.32500               |
| Long trousers/winter gym suit trousers | 0.29075               |
| Long trousers/summer gym suit trousers | 0.20500               |
| Shorts                                 | 0.17400               |
| Cloth skirt                            | 0.17000               |
| Denim skirt                            | 0.23750               |
| Dress                                  | 0.17400               |
| Pantyhose                              | 0.06600               |

The weight of each child was measured twice in close sequence by the same surveyor and using the same scale:

1. The first time, the children were wearing their clothes according to the OKkio alla SALUTE protocol. They were asked to remove their shoes, jackets, heavy sweaters and belts. Children were also asked to empty their pockets and remove any watch, bracelets, etc. The remaining clothes that were worn during weight measurement were marked on the OKkio alla SALUTE measurement card and, in the analysis, the weight was adjusted by the estimated weight of each item of clothing used in OKkio alla SALUTE (Table 1).
2. The second time, the children were asked to get undressed, wearing only their underwear, as recommended by the WHO<sup>(5)</sup>.

BMI was calculated: first using the weight measured with the clothes, in accordance with the protocol of OKkio alla SALUTE ( $BMI_{OK}$ ), and second as indicated by WHO ( $BMI_{WHO}$ ). To classify the ponderal status of the children, BMI values (both  $BMI_{OK}$  and  $BMI_{WHO}$ ) were compared with those of the international reference tables, specific for age and sex, suggested by the International Obesity Task Force (IOTF)<sup>(11)</sup>. In these tables, the threshold values of reference BMI are listed at exact half-year ages; in our study we applied the cut-offs of BMI linearly interpolated to the child's age in months. As the European Childhood Obesity Group has recently recommended to use both IOTF<sup>(11)</sup> and WHO definitions<sup>(12,13)</sup> for the assessment of the prevalence of childhood overweight and obesity<sup>(14)</sup> to permit comparisons between studies, the ponderal status of the children was also classified according to the WHO definition<sup>(12,13)</sup>.

## Statistical analysis

Children's weight wearing underwear ( $W_{WHO}$ ) was subtracted from that obtained with their clothes on, corrected for the estimated weight of the clothes ( $W_{OK}$ ). How this difference (error) biases BMI and evaluation of the children's ponderal status classification was subsequently appraised.

The Bland–Altman plot<sup>(15)</sup> was applied to compare: (i) the children's weight measured with clothing rectified according to the estimated weight of the clothes ( $W_{OK}$ ) with their weight measured without clothing ( $W_{WHO}$ ); and (ii) the BMI calculated from each of the two children's weight measurements ( $BMI_{WHO}$  and  $BMI_{OK}$ ).

Cohen's kappa index<sup>(16)</sup> was used to assess the agreement in overweight and obesity classification according to the IOTF<sup>(11)</sup> reference and the WHO<sup>(12,13)</sup> definition from each of the two BMI values obtained ( $BMI_{WHO}$  and  $BMI_{OK}$ ).

The statistical analyses were carried out using the statistical software package IBM SPSS Statistics version 18.0.

**Results**

Table 2 shows the characteristics of the sample examined in the present study: the 524 studied children had mean age of 8.4 (SD 0.4) years, weight  $W_{WHO}$  of 31.3 (SD 6.8) kg, weight  $W_{OK}$  of 31.3 (SD 6.8) kg, height of 131.5 (SD 5.9) cm and  $BMI_{WHO}$  of 17.9 (SD 3.0) kg/m<sup>2</sup>.

Table 3 shows the difference between the weight measurement with clothes, corrected for the weight of the clothes ( $W_{OK}$ ), and the weight measured only with underwear ( $W_{WHO}$ ); this difference can also be expressed as the difference between the estimated weight of the clothes and that measured. The mean difference between the two weights was -0.013 (SD 0.169) kg, with a 95% CI of -0.027, 0.002 kg, which is not statistically significant ( $P = 0.08$ ).

The Bland–Altman plot showed that the error between the real weight and the estimated weight was -0.013 kg and between -0.344 kg and 0.318 kg. The plot showed a tendency to overestimate clothes with a lower weight and underestimate those with a higher weight (Spearman's  $R = -0.60$ ;  $P = 0.00$ ; Fig. 1).

The mean error in the calculation of BMI through  $W_{OK}$  procedure was 0.005 (SD 0.097) kg/m<sup>2</sup> and it was included between -0.185 kg/m<sup>2</sup> and 0.195 kg/m<sup>2</sup>, with a 95% CI of -0.0029, 0.014 kg/m<sup>2</sup>. There was no systematic error (Fig. 2).

By the IOTF definition<sup>(11)</sup>, the percentages of overweight and obese children calculated with the two methods were: 25.6% overweight (not including obesity)

and 10.7% obese with  $W_{OK}$  and 25.4% overweight and 10.3% obese with  $W_{WHO}$  (Table 4). The analysis of the agreement in the overweight and obesity status classification by  $BMI_{WHO}$  and that made from  $BMI_{OK}$  produced a  $\kappa$  index very close to 1 (0.98): an extremely high agreement between the two ways of categorizing through BMI. None of the measured overweight and obese children dropped into the thinness/normal weight class. Three children with normal weight were incorrectly classified as overweight; only two overweight children were erroneously classified as obese (Table 4).

By the WHO definition<sup>(12,13)</sup>, the percentages of overweight and obese children calculated with the two methods of measuring body weight were: 23.5% overweight (not including obesity) and 21.4% obese with  $W_{OK}$  and 23.3% overweight and 21.8% obese with  $W_{WHO}$  (Table 5). Similarly to the results obtained using the IOTF definition, the agreement in the overweight and obesity status classification by  $BMI_{WHO}$  and that from  $BMI_{OK}$  was very high, with the  $\kappa$  index equal to 0.98. Two overweight children dropped into the thinness/normal weight class. One thinness/normal weight child was classified as overweight and two obese children were categorized as overweight (Table 5).

**Discussion**

The validation analysis, carried out on 524 children aged 8.4 (SD 0.4) years, suggests that measuring children's weight with their clothes on and then rectifying the obtained weight with the estimated weight of the clothes, according to the protocol of the surveillance system

**Table 3** Difference between the corrected clothed weight/BMI and the weight/BMI measured/calculated without clothes among 524 children aged 8–9 years


| Difference                                  | Min    | Max   | Mean   | SD    |
|---|--------|-------|--------|-------|
| $W_{WHO} - W_{OK}$ (kg)                     | -0.666 | 0.466 | -0.013 | 0.169 |
| $BMI_{WHO} - BMI_{OK}$ (kg/m <sup>2</sup> ) | -0.283 | 0.321 | 0.005  | 0.097 |

$W_{WHO}$ , weight without clothes according to the WHO protocol;  $W_{OK}$ , weight with clothes corrected by the estimated weight of the clothes according to the OKkio alla SALUTE protocol;  $BMI_{WHO}$ , BMI calculated from  $W_{WHO}$ ;  $BMI_{OK}$ , BMI calculated from  $W_{OK}$ .


**Table 2** Characteristics of the study sample (524 children aged 8–9 years)

|       | n   | Age (years) |     | $W_{WHO}$ (kg) |     | $W_{OK}$ (kg) |     | Height (cm) |     | $BMI_{WHO}$ (kg/m <sup>2</sup> ) |     |
|-------|-----|-------------|-----|----------------|-----|---------------|-----|-------------|-----|----------------------------------|-----|
|       |     | Mean        | SD  | Mean           | SD  | Mean          | SD  | Mean        | SD  | Mean                             | SD  |
| Girls | 224 | 8.4         | 0.4 | 31.1           | 7.0 | 31.1          | 7.0 | 131.1       | 6.3 | 18.0                             | 3.1 |
| Boys  | 300 | 8.4         | 0.4 | 31.4           | 6.6 | 31.4          | 6.6 | 131.9       | 5.6 | 17.9                             | 2.9 |
| Total | 524 | 8.4         | 0.4 | 31.3           | 6.8 | 31.3          | 6.8 | 131.5       | 5.9 | 17.9                             | 3.0 |

$W_{WHO}$ , weight without clothes according to the WHO protocol;  $W_{OK}$ , weight with clothes corrected by the estimated weight of the clothes according to the OKkio alla SALUTE protocol;  $BMI_{WHO}$ , BMI calculated from  $W_{WHO}$ .



**Fig. 1** Bland–Altman plot showing agreement between the real and estimated weight of clothes: the difference between the real and estimated weight of clothes (kg) is plotted against the mean of the real and estimated weight of clothes (kg) for 524 children aged 8–9 years



**Fig. 2** Bland–Altman plot showing agreement between the BMI values calculated from the two weight measurements: the difference in BMI ( $\text{kg}/\text{m}^2$ ) calculated from children's weight obtained using the OKkio alla SALUTE protocol ( $W_{\text{OK}}$ ) and BMI calculated from children's weight measured by the WHO protocol ( $W_{\text{WHO}}$ ) is plotted against the mean BMI ( $\text{kg}/\text{m}^2$ ) calculated using  $W_{\text{OK}}$  and  $W_{\text{WHO}}$  for 524 children aged 8–9 years

OKkio alla SALUTE, leads to a slight error in the estimation of body weight that implies a small miscalculation of BMI and a negligible difference in the prevalence of overweight and obesity assessment. Considering the advantage of the significant simplification of measurement procedures in ponderal status surveillance, that are generally carried out on a large scale, this error can be regarded as negligible, especially in view of its very limited impact in evaluation of the ponderal status in

children using international BMI reference values specific for age and sex (both IOTF<sup>(11)</sup> and WHO<sup>(12,13)</sup>). In fact, to be sustainable a population surveillance system should be acceptable, simple and not expensive<sup>(7)</sup>. To measure children without asking them to wear only their underwear makes the procedure quicker for the data collectors, less invasive for the children and parents and can lead to a higher participation rate. This is an essential point to consider in population studies, because the participation

**Table 4** Agreement between the assessment of ponderal status according to the International Obesity Task Force<sup>(11)</sup> by BMI<sub>WHO</sub> and BMI<sub>OK</sub> among 524 children aged 8–9 years

|                        | BMI <sub>WHO</sub>     |      |            |      |          |      |          |       |
|------------------------|------------------------|------|------------|------|----------|------|----------|-------|
|                        | Thinness/normal weight |      | Overweight |      | Obese    |      | Total    |       |
|                        | <i>n</i>               | %    | <i>n</i>   | %    | <i>n</i> | %    | <i>n</i> | %     |
| BMI <sub>OK</sub>      |                        |      |            |      |          |      |          |       |
| Thinness/normal weight | 334                    | 63.7 | 0          | –    | 0        | –    | 334      | 63.7  |
| Overweight             | 3                      | 0.6  | 131        | 25.0 | 0        | –    | 134      | 25.6  |
| Obese                  | 0                      | –    | 2          | 0.4  | 54       | 10.3 | 56       | 10.7  |
| Total                  | 337                    | 64.3 | 133        | 25.4 | 54       | 10.3 | 524      | 100.0 |

BMI<sub>WHO</sub>, BMI calculated from W<sub>WHO</sub>; BMI<sub>OK</sub>, BMI calculated from W<sub>OK</sub>; W<sub>WHO</sub>, weight without clothes according to the WHO protocol; W<sub>OK</sub>, weight with clothes corrected by the estimated weight of the clothes according to the OKkio alla SALUTE protocol.

**Table 5** Agreement between the assessment of ponderal status according to the WHO<sup>(12,13)</sup> by BMI<sub>WHO</sub> and BMI<sub>OK</sub> among 524 children aged 8–9 years

|                        | BMI <sub>WHO</sub>     |      |            |      |          |      |          |       |
|------------------------|------------------------|------|------------|------|----------|------|----------|-------|
|                        | Thinness/normal weight |      | Overweight |      | Obese    |      | Total    |       |
|                        | <i>n</i>               | %    | <i>n</i>   | %    | <i>n</i> | %    | <i>n</i> | %     |
| BMI <sub>OK</sub>      |                        |      |            |      |          |      |          |       |
| Thinness/normal weight | 287                    | 54.8 | 2          | 0.4  | 0        | –    | 289      | 55.2  |
| Overweight             | 1                      | 0.2  | 120        | 22.9 | 2        | 0.4  | 123      | 23.5  |
| Obese                  | 0                      | –    | 0          | –    | 112      | 21.4 | 112      | 21.4  |
| Total                  | 288                    | 55.0 | 122        | 23.3 | 114      | 21.8 | 524      | 100.0 |

BMI<sub>WHO</sub>, BMI calculated from W<sub>WHO</sub>; BMI<sub>OK</sub>, BMI calculated from W<sub>OK</sub>; W<sub>WHO</sub>, weight without clothes according to the WHO protocol; W<sub>OK</sub>, weight with clothes corrected by the estimated weight of the clothes according to the OKkio alla SALUTE protocol.

rate may affect the calculated prevalence figures. During the past 30 years the participation rates for epidemiological studies have been declining, with even steeper declines in recent years<sup>(17)</sup>. The participation rate had a slight but significant positive association with the estimated prevalence of obese children aged 10–11 years in the National Child Measurement Programme (NCMP) for England in 2007/08. This was due to obese children being less likely to participate in the NCMP than other children<sup>(18)</sup>. Thus a lower participation rate may lead to an underestimate of the prevalence of obesity. The choice of measuring the children with their clothes on may have been one of the factors which led to a low refusal to participate (3% in 2008 and in 2010) in the surveillance system OKkio alla SALUTE<sup>(8,9)</sup>.

Beyond the strengths of the simplified procedure of measuring body weight used in this surveillance system, it should be stressed that measuring dressed children does not enable waist circumference to be measured accurately. This may be a limit to be taken into account on the basis of the evidence that, in terms of morbidity, waist circumference and the ratio of waist circumference to height (WC:Ht) are good indices for selecting children with the highest cardiometabolic risk<sup>(19,20)</sup>. This limitation is important because WC:Ht does not need population-specific, gender-specific or age-specific references as BMI does. Considering the availability of age-, sex- and ethnicity-independent cut-offs for WC:Ht, it may be

advisable in future large population samples to obtain also waist circumference and not just weight and height.

In conclusion, in view of the big advantage of not asking children to remove their clothes, the small error in the evaluation of BMI and of ponderal status classification can be considered minor in surveillance systems of overweight and obesity.

## Acknowledgements

*Sources of funding:* Funding for this study was provided by the Italian Ministry of Health/Centre for Disease Prevention and Control, Chapter 4393/2005–CCM. *Conflict of interest:* The authors have no conflict of interest. *Ethical information:* The study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving human subjects were approved by the ethical committee of the National Institute of Health on 20 April 2009 (Ref. PRE/182/09 p. CE-ISS09/259). *Authors' contributions:* A.S., L.C. and G.B. designed the study. R.R. and N.B. collected the data. L.C., V.A., A.S. and G.B. analysed the data. A.L. and P.N. contributed to the interpretation of the results. All authors were involved in writing the paper and approved the final manuscript. *Acknowledgements:* The authors gratefully acknowledge Professor John Frederick Osborn for carefully reading the manuscript, and for his suggestions. They also wish to

thank the members of the OKkio alla SALUTE Technical Committee, including Giulia Cairella, Marcello Caputo, Margherita Caroli, Chiara Cattaneo, Franco Cavallo, Amleto D'Amicis, Amalia De Luca, Barbara De Mei, Daniela Galeone, Giordano Giostra, Gianfranco Mazzarella, Giuseppe Perri, Maria Teresa Silani, Anna Rita Silvestri and Lorenzo Spizzichino. Finally, they gratefully thank the teachers, children and families who participated in this study.

## References

- World Health Organization (2000) *Obesity: Preventing and Managing the Global Epidemic. Report on a WHO Consultation. WHO Technical Report Series* no. 894. Geneva: WHO.
- Branca F, Nikogosian H & Lobstein T (editors) (2007) *The Challenge of Obesity in the WHO European Region and the Strategies for Response: Summary*. Copenhagen: WHO Regional Office for Europe.
- Dietz W (1998) Health consequences of obesity in youth: childhood predictors of adult disease. *Pediatrics* **101**, 518–525.
- Claire YW, McPherson K, Marsh T *et al.* (2011) Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet* **378**, 815–825.
- World Health Organization (1995) *Physical Status: The Use and Interpretation of Anthropometry. Report of a WHO Expert Committee. WHO Technical Report Series* no. 854. Geneva: WHO.
- Bellizzi MC & Dietz WH (1999) Workshop on childhood obesity: summary of the discussion. *Am J Clin Nutr* **70**, issue 1, 173S–175S.
- Declich S & Carter AO (1994) Public health surveillance: historical origins, methods and evaluation. *Bull World Health Organ* **72**, 285–304.
- Binkin N, Fontana G, Lamberti A *et al.* (2010) A national survey of the prevalence of childhood overweight and obesity in Italy. *Obes Rev* **11**, 2–10.
- Spinelli A, Lamberti A, Nardone P *et al.* (editors) (2012) *Sistema di sorveglianza OKkio alla SALUTE: risultati 2010. Rapporti ISTISAN 12/14*. Roma: Istituto Superiore di Sanità.
- Wijnhoven TM, van Raaij JM, Spinelli A *et al.* (2013) WHO European Childhood Obesity Surveillance Initiative 2008: weight, height and body mass index in 6–9-year-old children. *Pediatr Obes* **8**, 79–97.
- Cole TJ, Bellizzi MC, Flegal KM *et al.* (2000) Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* **320**, 1240–1243.
- de Onis M, Onyango AW, Borghi E *et al.* (2007) Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* **85**, 660–667.
- Blössner M, Siyam A, Borghi E *et al.* (2009) *WHO AnthroPlus for Personal Computers Manual: Software for Assessing Growth of the World's Children and Adolescents*. Geneva: WHO; available at [http://www.who.int/growthref/tools/who\\_anthroplus\\_manual.pdf](http://www.who.int/growthref/tools/who_anthroplus_manual.pdf) (accessed December 2012).
- Rolland-Cachera MF for the European Childhood Obesity Group (2011) Childhood obesity: current definitions and recommendations for their use. *Int J Pediatr Obes* **6**, 325–331.
- Bland JM & Altman DG (1986) Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* **1**, 307–310.
- Cohen J (1960) A coefficient of agreement for nominal scales. *Educ Psychol Meas* **20**, 37–46.
- Galea S & Melissa T (2007) Participation rates in epidemiologic studies. *Ann Epidemiol* **17**, 643–653.
- The NHS Information Centre, Lifestyle Statistics (2009) National Child Measurement Programme: England, 2008/09 school year. December 2009. [http://www.ic.nhs.uk/webfiles/publications/ncmp/ncmp0809/NCMP\\_England\\_2008\\_09\\_school\\_year\\_report\\_2.pdf](http://www.ic.nhs.uk/webfiles/publications/ncmp/ncmp0809/NCMP_England_2008_09_school_year_report_2.pdf) (accessed December 2012).
- Maffei C, Banzato C, Talamini G *et al.* (2008) Waist-to-height ratio, a useful index to identify high metabolic risk in overweight children. *J Pediatr* **152**, 207–213.
- Katzmarzyk PT, Shen W, Baxter-Jones A *et al.* (2012) Adiposity in children and adolescents: correlates and clinical consequences of fat stored in specific body depots. *Pediatr Obes* **7**, e42–e61.