



## Correction: Separation of long-stranded RNAs by RP-HPLC using an octadecyl-based column with super-wide pores

Tomomi Kuwayama<sup>1</sup> · Makoto Ozaki<sup>1</sup> · Motoshi Shimotsuma<sup>1</sup> · Tsunehisa Hirose<sup>1</sup> 

Published online: 2 March 2023

© The Author(s), under exclusive licence to The Japan Society for Analytical Chemistry 2023

**Correction: Analytical Sciences (2022) 39:417–425**  
<https://doi.org/10.1007/s44211-022-00253-w>

The original article has been corrected.

In this article ref. 23 and 26 were incorrect and should have been as below.

23. Z. Huang, S. Jayaseelan, J. Hebert, H. Seo, L. Niu, *Anal. Biochem.* 435, 35–43 (2013)

26. Y. Yamauchi, M. Taoka, Y. Nobe, K. Izumikawa, N. Takahashi, H. Nakayama, T. Isobe, *J. Chromatogr. A.* 1312, 87–92 (2013)

---

The original article can be found online at <https://doi.org/10.1007/s44211-022-00253-w>.

---

✉ Tsunehisa Hirose  
[hirose-t@nacalai.co.jp](mailto:hirose-t@nacalai.co.jp)

<sup>1</sup> Nacalai Tesque, Inc., Ishibashi Kaide-Cho, Muko, Kyoto 617-0004, Japan