



Cochrane
Library

Cochrane Database of Systematic Reviews

Medical treatment of eosinophilic esophagitis (Review)

Franciosi JP, Gordon M, Sinopoulou V, Dellon ES, Gupta SK, Reed CC, Gutiérrez-Junquera C, Venkatesh RD, Erwin EA, Egiz A, Elleithy A, Mougey EB

Franciosi JP, Gordon M, Sinopoulou V, Dellon ES, Gupta SK, Reed CC, Gutiérrez-Junquera C, Venkatesh RD, Erwin EA, Egiz A, Elleithy A, Mougey EB.

Medical treatment of eosinophilic esophagitis.

Cochrane Database of Systematic Reviews 2023, Issue 7. Art. No.: CD004065.

DOI: [10.1002/14651858.CD004065.pub4](https://doi.org/10.1002/14651858.CD004065.pub4).

www.cochranelibrary.com

Medical treatment of eosinophilic esophagitis (Review)

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

WILEY

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	3
SUMMARY OF FINDINGS	5
BACKGROUND	32
OBJECTIVES	33
METHODS	33
RESULTS	36
Figure 1.	37
Figure 2.	43
Figure 3.	46
DISCUSSION	57
AUTHORS' CONCLUSIONS	60
ACKNOWLEDGEMENTS	60
REFERENCES	62
CHARACTERISTICS OF STUDIES	80
DATA AND ANALYSES	193
Analysis 1.1. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 1: Clinical improvement at study endpoint (dichotomous)	198
Analysis 1.2. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 2: Clinical improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	199
Analysis 1.3. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 3: Clinical improvement at study endpoint (dichotomous), sensitivity analysis, validated instruments	199
Analysis 1.4. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 4: Clinical improvement at study endpoint (dichotomous), subgrouped by age	200
Analysis 1.5. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 5: Clinical improvement at study endpoint (dichotomous), subgrouped by type of steroid	201
Analysis 1.6. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 6: Clinical improvement at study endpoint (dichotomous), subgrouped by delivery method	202
Analysis 1.7. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 7: Clinical improvement at study endpoint (continuous)	202
Analysis 1.8. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 8: Clinical improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	203
Analysis 1.9. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 9: Clinical improvement at study endpoint (continuous), sensitivity analysis, validated instruments	203
Analysis 1.10. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 10: Clinical improvement at study endpoint (continuous), subgrouped by age	204
Analysis 1.11. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 11: Clinical improvement at study endpoint (continuous), subgrouped by type of steroid	204
Analysis 1.12. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 12: Clinical improvement at study endpoint (continuous), subgrouped by delivery method	205
Analysis 1.13. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 13: Histological improvement at study endpoint (dichotomous)	205
Analysis 1.14. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 14: Histological improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	206
Analysis 1.15. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 15: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold < 15 eos/hpf	206
Analysis 1.16. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 16: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 6 eos/hpf	207
Analysis 1.17. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 17: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 1 eos/hpf	207
Analysis 1.18. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 18: Histological improvement at study endpoint (dichotomous), subgrouped by age	208
Analysis 1.19. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 19: Histological improvement at study endpoint (dichotomous), subgrouped by type of steroid	209

Analysis 1.20. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 20: Histological improvement at study endpoint (dichotomous), subgrouped by delivery method	210
Analysis 1.21. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 21: Histological improvement at study endpoint (continuous)	211
Analysis 1.22. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 22: Histological improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	211
Analysis 1.23. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 23: Histological improvement at study endpoint (continuous), subgrouped by age	212
Analysis 1.24. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 24: Histological improvement at study endpoint (continuous), subgrouped by type of steroid	212
Analysis 1.25. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 25: Histological improvement at study endpoint (continuous), subgrouped by delivery method	213
Analysis 1.26. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 26: Endoscopic improvement at study endpoint (dichotomous)	213
Analysis 1.27. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 27: Endoscopic improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	214
Analysis 1.28. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 28: Endoscopic improvement at study endpoint (dichotomous), sensitivity analysis, validated instruments	214
Analysis 1.29. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 29: Endoscopic improvement at study endpoint (dichotomous), subgrouped by age	215
Analysis 1.30. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 30: Endoscopic improvement at study endpoint (dichotomous), subgrouped by delivery method	216
Analysis 1.31. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 31: Endoscopic improvement at study endpoint (continuous)	216
Analysis 1.32. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 32: Endoscopic improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	217
Analysis 1.33. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 33: Endoscopic improvement at study endpoint (continuous), sensitivity analysis, validated instruments	217
Analysis 1.34. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 34: Endoscopic improvement at study endpoint (continuous), subgrouped by age	218
Analysis 1.35. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 35: Endoscopic improvement at study endpoint (continuous), subgrouped by type of steroid	218
Analysis 1.36. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 36: Withdrawals due to adverse events	219
Analysis 1.37. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 37: Withdrawals due to adverse events, sensitivity analysis, fixed-effect	220
Analysis 1.38. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 38: Withdrawals due to adverse events, subgrouped by age	221
Analysis 1.39. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 39: Withdrawals due to adverse events, subgrouped by type of steroid	222
Analysis 1.40. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 40: Withdrawals due to adverse events, subgrouped by delivery method	223
Analysis 1.41. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 41: Serious adverse events	224
Analysis 1.42. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 42: Total adverse events	225
Analysis 1.43. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 43: Quality of life at study endpoint (continuous)	225
Analysis 2.1. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 1: Clinical improvement at study endpoint (dichotomous)	226
Analysis 2.2. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 2: Clinical improvement at study endpoint (continuous)	227
Analysis 2.3. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 3: Histological improvement at study endpoint (dichotomous)	227
Analysis 2.4. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 4: Histological improvement at study endpoint (continuous)	228
Analysis 2.5. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 5: Endoscopic improvement at study endpoint (continuous)	228

Analysis 2.6. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 6: Withdrawals due to adverse events	229
Analysis 2.7. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 7: Serious adverse events	229
Analysis 2.8. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 8: Total adverse events	230
Analysis 2.9. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 9: Quality of life at study endpoint (continuous)	230
Analysis 3.1. Comparison 3: Biologics vs placebo for induction of remission, Outcome 1: Clinical improvement at study endpoint (dichotomous)	235
Analysis 3.2. Comparison 3: Biologics vs placebo for induction of remission, Outcome 2: Clinical improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	235
Analysis 3.3. Comparison 3: Biologics vs placebo for induction of remission, Outcome 3: Clinical improvement at study endpoint (dichotomous), sensitivity analysis, validated instruments	236
Analysis 3.4. Comparison 3: Biologics vs placebo for induction of remission, Outcome 4: Clinical improvement at study endpoint (dichotomous), subgrouped by age	236
Analysis 3.5. Comparison 3: Biologics vs placebo for induction of remission, Outcome 5: Clinical improvement at study endpoint (dichotomous), subgrouped by mechanism	237
Analysis 3.6. Comparison 3: Biologics vs placebo for induction of remission, Outcome 6: Clinical improvement at study endpoint (continuous)	237
Analysis 3.7. Comparison 3: Biologics vs placebo for induction of remission, Outcome 7: Clinical improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	238
Analysis 3.8. Comparison 3: Biologics vs placebo for induction of remission, Outcome 8: Clinical improvement at study endpoint (continuous), sensitivity analysis, peer-reviewed manuscripts	238
Analysis 3.9. Comparison 3: Biologics vs placebo for induction of remission, Outcome 9: Clinical improvement at study endpoint (continuous), sensitivity analysis, less than high risk of bias	239
Analysis 3.10. Comparison 3: Biologics vs placebo for induction of remission, Outcome 10: Clinical improvement at study endpoint (continuous), sensitivity analysis, validated instruments	239
Analysis 3.11. Comparison 3: Biologics vs placebo for induction of remission, Outcome 11: Clinical improvement at study endpoint (continuous), subgrouped by mechanism	240
Analysis 3.12. Comparison 3: Biologics vs placebo for induction of remission, Outcome 12: Histological improvement at study endpoint (dichotomous)	241
Analysis 3.13. Comparison 3: Biologics vs placebo for induction of remission, Outcome 13: Histological improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	241
Analysis 3.14. Comparison 3: Biologics vs placebo for induction of remission, Outcome 14: Histological improvement at study endpoint (dichotomous), sensitivity analysis, peer-reviewed manuscripts	242
Analysis 3.15. Comparison 3: Biologics vs placebo for induction of remission, Outcome 15: Histological improvement at study endpoint (dichotomous), sensitivity analysis, less than high risk of bias	242
Analysis 3.16. Comparison 3: Biologics vs placebo for induction of remission, Outcome 16: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold < 15 eos/hpf	243
Analysis 3.17. Comparison 3: Biologics vs placebo for induction of remission, Outcome 17: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 6 eos/hpf	243
Analysis 3.18. Comparison 3: Biologics vs placebo for induction of remission, Outcome 18: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 1 eos/hpf	244
Analysis 3.19. Comparison 3: Biologics vs placebo for induction of remission, Outcome 19: Histological improvement at study endpoint (dichotomous), subgrouped by age	245
Analysis 3.20. Comparison 3: Biologics vs placebo for induction of remission, Outcome 20: Histological improvement at study endpoint (dichotomous), subgrouped by mechanism	246
Analysis 3.21. Comparison 3: Biologics vs placebo for induction of remission, Outcome 21: Histological improvement at study endpoint (continuous)	247
Analysis 3.22. Comparison 3: Biologics vs placebo for induction of remission, Outcome 22: Histological improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	247
Analysis 3.23. Comparison 3: Biologics vs placebo for induction of remission, Outcome 23: Histological improvement at study endpoint (continuous), sensitivity analysis, less than high risk of bias	248
Analysis 3.24. Comparison 3: Biologics vs placebo for induction of remission, Outcome 24: Histological improvement at study endpoint (continuous), subgrouped by age	248
Analysis 3.25. Comparison 3: Biologics vs placebo for induction of remission, Outcome 25: Histological improvement at study endpoint (continuous), subgrouped by mechanism	249

Analysis 3.26. Comparison 3: Biologics vs placebo for induction of remission, Outcome 26: Endoscopic improvement at study endpoint (dichotomous)	249
Analysis 3.27. Comparison 3: Biologics vs placebo for induction of remission, Outcome 27: Endoscopic improvement at study endpoint (continuous)	250
Analysis 3.28. Comparison 3: Biologics vs placebo for induction of remission, Outcome 28: Endoscopic improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	250
Analysis 3.29. Comparison 3: Biologics vs placebo for induction of remission, Outcome 29: Endoscopic improvement at study endpoint (continuous), sensitivity analysis, less than high risk of bias	250
Analysis 3.30. Comparison 3: Biologics vs placebo for induction of remission, Outcome 30: Withdrawals due to adverse events ..	251
Analysis 3.31. Comparison 3: Biologics vs placebo for induction of remission, Outcome 31: Withdrawals due to adverse events, sensitivity analysis, fixed-effect	251
Analysis 3.32. Comparison 3: Biologics vs placebo for induction of remission, Outcome 32: Withdrawals due to adverse events, sensitivity analysis, less than high risk of bias	252
Analysis 3.33. Comparison 3: Biologics vs placebo for induction of remission, Outcome 33: Withdrawals due to adverse events, sensitivity analysis, peer reviewed manuscripts	252
Analysis 3.34. Comparison 3: Biologics vs placebo for induction of remission, Outcome 34: Withdrawals due to adverse events, subgrouped by age	253
Analysis 3.35. Comparison 3: Biologics vs placebo for induction of remission, Outcome 35: Withdrawals due to adverse events, subgrouped by mechanism	254
Analysis 3.36. Comparison 3: Biologics vs placebo for induction of remission, Outcome 36: Serious adverse events	255
Analysis 3.37. Comparison 3: Biologics vs placebo for induction of remission, Outcome 37: Total adverse events	255
Analysis 3.38. Comparison 3: Biologics vs placebo for induction of remission, Outcome 38: Quality of life at study endpoint (continuous)	256
Analysis 4.1. Comparison 4: Cromolyn sodium vs placebo, Outcome 1: Clinical improvement at study endpoint (continuous) ..	256
Analysis 4.2. Comparison 4: Cromolyn sodium vs placebo, Outcome 2: Histological improvement at study endpoint (continuous)	257
Analysis 4.3. Comparison 4: Cromolyn sodium vs placebo, Outcome 3: Withdrawals due to adverse events	257
Analysis 4.4. Comparison 4: Cromolyn sodium vs placebo, Outcome 4: Serious adverse events	257
Analysis 5.1. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 1: Clinical improvement at study endpoint (continuous)	258
Analysis 5.2. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 2: Histological improvement at study endpoint (continuous)	258
Analysis 5.3. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 3: Endoscopic improvement at study endpoint (continuous)	259
Analysis 5.4. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 4: Withdrawals due to adverse events	259
Analysis 5.5. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 5: Serious adverse events	259
Analysis 6.1. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 1: Clinical improvement at study endpoint (dichotomous)	260
Analysis 6.2. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 2: Histological improvement at study endpoint (dichotomous)	261
Analysis 6.3. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 3: Histological improvement at study endpoint (continuous)	261
Analysis 6.4. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 4: Endoscopic improvement at study endpoint (dichotomous)	261
Analysis 6.5. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 5: Withdrawals due to adverse events	262
Analysis 6.6. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 6: Serious adverse events	262
Analysis 6.7. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 7: Total adverse events	263
Analysis 7.1. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 1: Clinical improvement at study endpoint (continuous)	264
Analysis 7.2. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 2: Histological improvement at study endpoint (dichotomous)	264
Analysis 7.3. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 3: Histological improvement at study endpoint (continuous)	264
Analysis 7.4. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 4: Endoscopic improvement at study endpoint (continuous)	265

Analysis 7.5. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 5: Withdrawals due to adverse events	265
Analysis 7.6. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 6: Serious adverse events	265
Analysis 7.7. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 7: Total adverse events	266
Analysis 8.1. Comparison 8: Esomeprazole vs fluticasone, Outcome 1: Clinical improvement at study endpoint (continuous)	267
Analysis 8.2. Comparison 8: Esomeprazole vs fluticasone, Outcome 2: Histological improvement at study endpoint (dichotomous)	267
Analysis 8.3. Comparison 8: Esomeprazole vs fluticasone, Outcome 3: Histological improvement at study endpoint (continuous)	268
Analysis 8.4. Comparison 8: Esomeprazole vs fluticasone, Outcome 4: Withdrawals due to adverse events	268
Analysis 8.5. Comparison 8: Esomeprazole vs fluticasone, Outcome 5: Serious adverse events	269
Analysis 8.6. Comparison 8: Esomeprazole vs fluticasone, Outcome 6: Total adverse events	269
Analysis 9.1. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 1: Clinical improvement at study endpoint (continuous)	270
Analysis 9.2. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 2: Histological improvement at study endpoint (dichotomous)	270
Analysis 9.3. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 3: Endoscopic improvement at study endpoint (continuous)	271
Analysis 9.4. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 4: Withdrawals due to adverse events	271
Analysis 9.5. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 5: Serious adverse events	272
Analysis 9.6. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 6: Total adverse events	272
Analysis 9.7. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 7: Quality of life at study endpoint (continuous)	273
Analysis 10.1. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 1: Clinical improvement at study endpoint (continuous)	274
Analysis 10.2. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 2: Histological improvement at study endpoint (dichotomous)	274
Analysis 10.3. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 3: Histological improvement at study endpoint (continuous)	274
Analysis 10.4. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 4: Endoscopic improvement at study endpoint (continuous)	275
Analysis 10.5. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 5: Withdrawals due to adverse events	275
Analysis 10.6. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 6: Serious adverse events	276
Analysis 10.7. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 7: Total adverse events	276
Analysis 10.8. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 8: Quality of life at study endpoint (continuous)	277
Analysis 11.1. Comparison 11: Four-food elimination diet with omeprazole vs omeprazole, Outcome 1: Histological improvement at study endpoint (dichotomous)	277
Analysis 11.2. Comparison 11: Four-food elimination diet with omeprazole vs omeprazole, Outcome 2: Histological improvement at study endpoint (continuous)	278
Analysis 11.3. Comparison 11: Four-food elimination diet with omeprazole vs omeprazole, Outcome 3: Withdrawals due to adverse events	278
Analysis 12.1. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 1: Clinical improvement at study endpoint (continuous)	279
Analysis 12.2. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 2: Histological improvement at study endpoint (dichotomous)	279
Analysis 12.3. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 3: Histological improvement at study endpoint (continuous)	280
Analysis 12.4. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 4: Endoscopic improvement at study endpoint (continuous)	280
Analysis 12.5. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 5: Withdrawals due to adverse events	280
Analysis 12.6. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 6: Serious adverse events	281

Analysis 12.7. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 7: Total adverse events	281
Analysis 13.1. Comparison 13: Nebulized budesonide vs viscous budesonide, Outcome 1: Clinical improvement at study endpoint (continuous)	282
Analysis 13.2. Comparison 13: Nebulized budesonide vs viscous budesonide, Outcome 2: Histological improvement at study endpoint (continuous)	282
Analysis 13.3. Comparison 13: Nebulized budesonide vs viscous budesonide, Outcome 3: Serious adverse events	283
Analysis 13.4. Comparison 13: Nebulized budesonide vs viscous budesonide, Outcome 4: Total adverse events	283
Analysis 14.1. Comparison 14: Viaskin milk patch vs placebo, Outcome 1: Clinical improvement at study endpoint (continuous)	284
Analysis 14.2. Comparison 14: Viaskin milk patch vs placebo, Outcome 2: Histological improvement at study endpoint (continuous)	284
Analysis 14.3. Comparison 14: Viaskin milk patch vs placebo, Outcome 3: Endoscopic improvement at study endpoint (continuous)	285
Analysis 14.4. Comparison 14: Viaskin milk patch vs placebo, Outcome 4: Withdrawals due to adverse events	285
Analysis 14.5. Comparison 14: Viaskin milk patch vs placebo, Outcome 5: Serious adverse events	285
Analysis 14.6. Comparison 14: Viaskin milk patch vs placebo, Outcome 6: Total adverse events	286
Analysis 14.7. Comparison 14: Viaskin milk patch vs placebo, Outcome 7: Quality of life at study endpoint (continuous)	286
Analysis 15.1. Comparison 15: Leukotriene receptor antagonist vs placebo for maintenance of remission, Outcome 1: Clinical improvement at study endpoint (dichotomous)	287
Analysis 15.2. Comparison 15: Leukotriene receptor antagonist vs placebo for maintenance of remission, Outcome 2: Withdrawals due to adverse events	287
Analysis 15.3. Comparison 15: Leukotriene receptor antagonist vs placebo for maintenance of remission, Outcome 3: Serious adverse events	287
Analysis 15.4. Comparison 15: Leukotriene receptor antagonist vs placebo for maintenance of remission, Outcome 4: Total adverse events	288
Analysis 16.1. Comparison 16: Mepolizumab 10 mg/kg vs mepolizumab 0.55 mg/kg, Outcome 1: Histological improvement at study endpoint (dichotomous)	288
Analysis 16.2. Comparison 16: Mepolizumab 10 mg/kg vs mepolizumab 0.55 mg/kg, Outcome 2: Withdrawals due to adverse events	289
Analysis 17.1. Comparison 17: Mepolizumab 2.5 mg/kg vs mepolizumab 0.55 mg/kg, Outcome 1: Histological improvement at study endpoint (dichotomous)	289
Analysis 17.2. Comparison 17: Mepolizumab 2.5 mg/kg vs mepolizumab 0.55 mg/kg, Outcome 2: Withdrawals due to adverse events	290
Analysis 18.1. Comparison 18: Mepolizumab 10 mg/kg vs mepolizumab 2.5 mg/kg, Outcome 1: Histological improvement at study endpoint (dichotomous)	290
Analysis 18.2. Comparison 18: Mepolizumab 10 mg/kg vs mepolizumab 2.5 mg/kg, Outcome 2: Withdrawals due to adverse events	291
ADDITIONAL TABLES	291
APPENDICES	361
WHAT'S NEW	363
HISTORY	365
CONTRIBUTIONS OF AUTHORS	365
DECLARATIONS OF INTEREST	365
SOURCES OF SUPPORT	366
DIFFERENCES BETWEEN PROTOCOL AND REVIEW	366
INDEX TERMS	367

[Intervention Review]

Medical treatment of eosinophilic esophagitis

James P Franciosi^{1,2a}, Morris Gordon^{3a}, Vassiliki Sinopoulou³, Evan S Dellon⁴, Sandeep K Gupta⁵, Craig C Reed⁴, Carolina Gutiérrez-Junquera⁶, Rajitha D Venkatesh⁷, Elizabeth A Erwin⁸, Abdullah Egiz³, Assem Elleithy³, Edward B Mougey⁹

¹Division of Gastroenterology, Hepatology, and Nutrition, Nemours Children's Hospital, Orlando, FL, USA. ²College of Medicine, University of Central Florida, Orlando, USA. ³School of Medicine, University of Central Lancashire, Preston, UK. ⁴Department of Medicine, Division of Gastroenterology and Hepatology, Center for Esophageal Diseases and Swallowing, University of North Carolina School of Medicine, Chapel Hill, NC, USA. ⁵Division of Pediatric Gastroenterology, Hepatology and Nutrition, University of Illinois College of Medicine at Peoria and Children's Hospital of Illinois, Peoria, IN, USA. ⁶Pediatric Gastroenterology, University Hospital Puerta de Hierro Majadahonda. Autonomous University of Madrid, Madrid, Spain. ⁷Pediatrics, Gastroenterology & Hepatology & Nutrition, Nationwide Children's Hospital, Columbus, OH, USA. ⁸Pediatric Allergy, Nationwide Children's Hospital, Columbus, OH, USA. ⁹Clinical Pharmacogenomics and Translational Research, Nemours Children's Health System, Jacksonville, FL, USA

^aThese authors should be considered joint first author

Contact: James P Franciosi, james.franciosi@nemours.org.

Editorial group: Cochrane Gut Group.

Publication status and date: New search for studies and content updated (conclusions changed), published in Issue 7, 2023.

Citation: Franciosi JP, Gordon M, Sinopoulou V, Dellon ES, Gupta SK, Reed CC, Gutiérrez-Junquera C, Venkatesh RD, Erwin EA, Egiz A, Elleithy A, Mougey EB. Medical treatment of eosinophilic esophagitis. *Cochrane Database of Systematic Reviews* 2023, Issue 7. Art. No.: CD004065. DOI: [10.1002/14651858.CD004065.pub4](https://doi.org/10.1002/14651858.CD004065.pub4).

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration. This is an open access article under the terms of the [Creative Commons Attribution Licence](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background

Eosinophilic esophagitis (EoE) is a chronic antigen-mediated eosinophilic inflammatory disease isolated to the esophagus. As a clinicopathologic disorder, a diagnosis of EoE requires a constellation of clinical symptoms of esophageal dysfunction and histologic findings (at least 15 eosinophils/high-powered microscope field (eos/hpf)). Current guidelines no longer require the failure of response to proton pump inhibitor medications to establish a diagnosis of EoE, but continue to suggest the exclusion of other etiologies of esophageal eosinophilia.

The treatment goals for EoE are improvement in clinical symptoms, resolution of esophageal eosinophilia and other histologic abnormalities, endoscopic improvement, improved quality of life, improved esophageal function, minimized adverse effects of treatment, and prevention of disease progression and subsequent complications.

Currently, there is no cure for EoE, making long-term treatment necessary. Standard treatment modalities include dietary modifications, esophageal dilation, and pharmacologic therapy. Effective pharmacologic therapies include corticosteroids, rapidly emerging biological therapies, and proton pump inhibitor medications.

Objectives

To evaluate the efficacy and safety of medical interventions for people with eosinophilic esophagitis.

Search methods

We searched CENTRAL, MEDLINE, Embase, ClinicalTrials.gov, and WHO ICTRP to 3 March 2023.

Medical treatment of eosinophilic esophagitis (Review)

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Selection criteria

Randomized controlled trials (RCTs) comparing any medical intervention or food elimination diet for the treatment of eosinophilic esophagitis, either alone or in combination, to any other intervention (including placebo).

Data collection and analysis

Pairs of review authors independently selected studies and conducted data extraction and risk of bias assessment. We expressed outcomes as a risk ratio (RR) and as the mean or standardized mean difference (MD/SMD) with 95% confidence interval (CI). We assessed the certainty of the evidence using GRADE.

Our primary outcomes were: clinical, histological, and endoscopic improvement, and withdrawals due to adverse events. Secondary outcomes were: serious and total adverse events, and quality of life.

Main results

We included 41 RCTs with 3253 participants. Eleven studies included pediatric patients while the rest recruited both children and adults. Four studies were in patients with inactive disease while the rest were in patients with active disease. We identified 19 intervention comparisons. In this abstract we present the results of the primary outcomes for the two main comparisons: corticosteroids versus placebo and biologics versus placebo, based on the prespecified outcomes defined of the primary studies.

Fourteen studies compared corticosteroids to placebo for induction of remission and the risk of bias for these studies was mostly low.

Corticosteroids may lead to slightly better clinical improvement (20% higher), measured dichotomously (risk ratio (RR) 1.74, 95% CI 1.08 to 2.80; 6 studies, 583 participants; number needed to treat for an additional beneficial outcome (NNTB) = 4; low certainty), and may lead to slightly better clinical improvement, measured continuously (standard mean difference (SMD) 0.51, 95% CI 0.17 to 0.85; 5 studies, 475 participants; low certainty).

Corticosteroids lead to a large histological improvement (63% higher), measured dichotomously (RR 11.94, 95% CI 6.56 to 21.75; 12 studies, 978 participants; NNTB = 3; high certainty), and may lead to histological improvement, measured continuously (SMD 1.42, 95% CI 1.02 to 1.82; 5 studies, 449 participants; low certainty).

Corticosteroids may lead to little to no endoscopic improvement, measured dichotomously (RR 2.60, 95% CI 0.82 to 8.19; 5 studies, 596 participants; low certainty), and may lead to endoscopic improvement, measured continuously (SMD 1.33, 95% CI 0.59 to 2.08; 5 studies, 596 participants; low certainty).

Corticosteroids may lead to slightly fewer withdrawals due to adverse events (RR 0.64, 95% CI 0.43 to 0.96; 14 studies, 1032 participants; low certainty).

Nine studies compared biologics to placebo for induction of remission.

Biologics may result in little to no difference in clinical improvement, measured dichotomously (RR 1.14, 95% CI 0.85 to 1.52; 5 studies, 410 participants; low certainty), and may result in better clinical improvement, measured continuously (SMD 0.50, 95% CI 0.22 to 0.78; 7 studies, 387 participants; moderate certainty).

Biologics result in better histological improvement (55% higher), measured dichotomously (RR 6.73, 95% CI 2.58 to 17.52; 8 studies, 925 participants; NNTB = 2; moderate certainty). We could not draw conclusions for this outcome when measured continuously (SMD 1.01, 95% CI 0.36 to 1.66; 6 studies, 370 participants; very low certainty).

Biologics may result in little to no difference in endoscopic improvement, measured dichotomously (effect not estimable, low certainty). We cannot draw conclusions for this outcome when measured continuously (SMD 2.79, 95% CI 0.36 to 5.22; 1 study, 11 participants; very low certainty).

There may be no difference in withdrawals due to adverse events (RR 1.55, 95% CI 0.88 to 2.74; 8 studies, 792 participants; low certainty).

Authors' conclusions

Corticosteroids (as compared to placebo) may lead to clinical symptom improvement when reported both as dichotomous and continuous outcomes, from the primary study definitions. Corticosteroids lead to a large increase in histological improvement (dichotomous outcome) and may increase histological improvement (continuous outcome) when compared to placebo. Corticosteroids may or may not increase endoscopic improvement (depending on whether the outcome is measured dichotomously or continuously). Withdrawals due to adverse events (dichotomous outcome) may occur less frequently when corticosteroids are compared to placebo.

Biologics (as compared to placebo) may not lead to clinical symptom improvement when reported as a dichotomous outcome and may lead to an increase in clinical symptom improvement (as a continuous outcome), from the primary study definitions. Biologics lead to a large increase in histological improvement when reported as a dichotomous outcome, but this is uncertain when reported as a continuous outcome, as compared to placebo. Biologics may not increase endoscopic improvement (dichotomous outcome), but this is uncertain

Medical treatment of eosinophilic esophagitis (Review)

when measured as a continuous outcome. Withdrawals due to adverse events as a dichotomous outcome may occur as frequently when biologics are compared to placebo.

PLAIN LANGUAGE SUMMARY

Medical treatments for eosinophilic esophagitis

Key messages

We found that while corticosteroids may improve patients' symptoms, they certainly reduce the amount of allergic cells (eosinophils) and they may improve what the disease looks like under visual inspection (endoscopy), when compared to placebo, for children and adults with eosinophilic esophagitis. They may be just as safe as a placebo (dummy treatment).

We found that biologics (a type of treatment that uses substances made from living organisms to treat disease) may improve patients' symptoms, that they certainly reduce the amount of allergic cells, and that they may be no different in terms of what the disease looks like under visual inspection, when compared to placebo, for children and adults with eosinophilic esophagitis. They may be just as safe as a placebo.

What is eosinophilic esophagitis?

Eosinophilic esophagitis is a long-term allergic condition in which the esophagus becomes inflamed (sore), which can lead to difficulty swallowing, vomiting, heartburn, and chest and stomach pain. Particles in foods or the air cause the immune system to have an allergic reaction and produce immune cells, which are called eosinophils. These build up in the esophagus, the tube that connects the mouth with the stomach. Eosinophilic esophagitis was first identified in the 1990s and since then it has been recognized as a major digestive illness. It is not known what causes it, but it might be related to genetics combined with environmental triggers. People with eosinophilic esophagitis tend to have other allergies as well. Currently, there is no cure for eosinophilic esophagitis, making long-term treatment necessary. Standard treatments include diets, stretching of the esophagus (dilation), and drugs such as corticosteroids, biological medications, and proton pump inhibitor medications.

What did we want to find out?

We wanted to find out if the available medical treatments for eosinophilic esophagitis work for improving patients' symptoms, reducing the amount of allergic cells when measured under a microscope, and improving what the disease looks like under visual examination. We also wanted to find out how safe they are and if they improve quality of life.

What did we do?

We searched for randomized controlled trials (studies where people are assigned to one of two or more treatment groups using a random method) comparing any medical treatment for eosinophilic esophagitis with any other medical treatment, in both adults and children.

What did we find?

We found 41 studies with 3253 participants. Eleven studies were in children only while the rest were in a mix of children and adults. We identified 19 comparisons. In this summary, we present the results of the two main comparisons: corticosteroids compared to placebo and biologics compared to placebo.

We found that corticosteroids may be better than placebo at improving patients' symptoms. We are highly certain that corticosteroids are better than placebo at reducing the amount of eosinophils (allergic cells) when measured under a microscope. Corticosteroids may be better than placebo at improving what the disease looks like under visual examination (endoscopy). We also found that people taking corticosteroids may be less likely to leave a study due to unwanted or harmful effects (side effects), and that they probably experience a similar number of both serious side effects and side effects in total, compared to placebo. There may be no difference between corticosteroids and placebo in the improvement of quality of life.

We found that biologics may be better than placebo at improving patients' symptoms. It is likely that biologics are better than placebo at reducing the amount of allergic cells when measured under a microscope. Biologics may be no different to placebo at improving what the disease looks like under visual examination. We also found that people on biologics may be equally likely to leave a study due to side effects, or have serious side effects, and may experience similar numbers of total side effects, compared to placebo. There may be no difference between biologics and placebo in the improvement of quality of life.

What are the limitations of the evidence?

The evidence in children only was quite limited and we do not know if the conclusions above can definitely apply to children specifically. Another limitation of the evidence is that the outcomes were measured in many different ways, which may have weakened our conclusions. Other treatments used by the participants were also something that varied a lot between people, and may have affected our conclusions.

Medical treatment of eosinophilic esophagitis (Review)

Finally, we were limited in the conclusions we could make about the effects of sex, age, extent of disease, dosage, and type of corticosteroid or biologic.

How up-to-date is this review?

This review is up-to-date to 3 March 2023.

SUMMARY OF FINDINGS

Summary of findings 1. Corticosteroids compared to placebo for induction of remission

Corticosteroids compared to placebo for induction of remission

Patient or population: active EoE patients

Setting: medical centers

Intervention: corticosteroids

Comparison: placebo

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with placebo**	Risk difference with corticosteroids	
Clinical improvement (dichotomous) 2 to 12 weeks	583 (6 studies)	⊕⊕⊕⊕ Low^a	RR 1.74 (1.08 to 2.80)	Study population 350 per 1000	259 more per 1000 (28 more to 378 more)	—
Clinical improvement (continuous) 2 to 12 weeks	475 (5 studies)	⊕⊕⊕⊕ Low^a	—	—	SMD 0.51 higher (0.17 higher to 0.85 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
Histological improvement (dichotomous) 2 to 12 weeks	978 (12 studies)	⊕⊕⊕⊕ High	RR 11.94 (6.56 to 21.75)	Study population 31 per 1000	339 more per 1000 (172 more to 643 more)	NNTB = 3
Histological improvement (continuous) 2 to 12 weeks	449 (5 studies)	⊕⊕⊕⊕ Low^b	—	—	SMD 1.42 higher (1.02 higher to 1.82 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
Endoscopic improvement (dichotomous) 6 to 12 weeks	102 (3 studies)	⊕⊕⊕⊕ Low^c	RR 2.60 (0.82 to 8.19)	Study population 136 per 1000	218 more per 1000 (24 less to 978 more)	—
Endoscopic improvement (continuous)	596 (5 studies)	⊕⊕⊕⊕ Low^d	—	—	SMD 1.33 higher (0.59 higher to 2.08 higher)	As a rule of thumb, 0.2 SMD represents a small

6 to 12 weeks						difference, 0.5 a moderate, and 0.8 a large effect.
Withdrawals due to adverse events (2 to 12 weeks)	1032 (14 studies)	⊕⊕⊕⊕ Low^e	RR 0.64 (0.43 to 0.96)	Study population	—	
				124 per 1000	45 fewer per 1000 (71 fewer to 5 fewer)	

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

** **The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded once due to inconsistency ($I^2 = 72\%$ and $I^2 = 55\%$ respectively) and once due to imprecision.

^bDowngraded once due to inconsistency ($I^2 = 50\%$) and once due to risk of bias across multiple domains.

^cDowngraded twice due to serious imprecision.

^dDowngraded twice due to serious inconsistency ($I^2 = 92\%$).

^eDowngraded once due to imprecision and once due to risk of bias across multiple domains.

Summary of findings 2. Corticosteroids compared to placebo for maintenance of remission

Corticosteroids compared to placebo for maintenance of remission

Patient or population: inactive EoE patients

Setting: medical centers

Intervention: corticosteroids

Comparison: placebo

Outcomes	N ^o of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)	Comments
----------	--	-----------------------------------	--------------------------	--	----------

				Risk with placebo	Risk difference with corti-costeroids	
Clinical improvement (dichotomous) 12 to 48 weeks	252 (2 studies)	⊕⊕⊕⊕ Very low^a	RR 2.17 (0.75 to 6.27)	Study population 297 per 1000	— 347 more per 1000 (74 fewer to 1000 more)	
Clinical improvement (continuous) 12 to 50 weeks	269 (3 studies)	⊕⊕⊕⊕ Very low^a	—	—	SMD 0.51 higher (0.49 lower to 1.52 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
Histological improvement (dichotomous) 12 to 50 weeks	280 (3 studies)	⊕⊕⊕⊕ Moderate^b	RR 4.58 (1.66 to 12.62)	Study population 133 per 1000	— 476 more per 1000 (88 more to 1000 more)	NNTB = 3
Histological improvement (continuous) 12 to 50 weeks	269 (3 studies)	⊕⊕⊕⊕ Moderate^c	—	—	SMD 1.26 higher (0.74 higher to 1.78 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
Endoscopic improvement at study endpoint (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement (continuous) 12 to 48 weeks	240 (2 studies)	⊕⊕⊕⊕ Very low^a	—	—	SMD 1.34 higher (0.27 lower to 2.95 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
Withdrawals due to adverse events 12 to 50 weeks	280 (3 studies)	⊕⊕⊕⊕ Low^d	RR 0.37 (0.16 to 0.87)	Study population 552 per 1000	— 348 fewer per 1000 (464 fewer to 72 fewer)	

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious inconsistency ($I^2 = 97%$, $I^2 = 91%$, and $I^2 = 95%$ respectively) and imprecision.

^bDowngraded once due to imprecision

^cDowngraded once due to inconsistency ($I^2 = 60%$).

^dDowngraded once due to inconsistency ($I^2 = 69%$) and once due to imprecision.

Summary of findings 3. Biologics compared to placebo for induction of remission

Biologics compared to placebo for induction of remission

Patient or population: active EoE patients

Setting: medical centers

Intervention: biologics

Comparison: placebo

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with placebo	Risk difference with biologics	
Clinical improvement (dichotomous) 12 to 44 weeks	410 (5 studies)	⊕⊕⊕⊕ Low^a	RR 1.14 (0.85 to 1.52)	Study population 504 per 1000	71 more per 1000 (76 fewer to 262 more)	—
Clinical improvement (continuous) 12 to 24 weeks	387 (7 studies)	⊕⊕⊕⊕ Moderate^b	—	—	SMD 0.50 higher (0.22 higher to 0.78 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
Histological improvement (dichotomous) 12 to 44 weeks	925 (8 studies)	⊕⊕⊕⊕ Moderate^b	RR 6.73 (2.58 to 17.52)	Study population		NNTB = 2

				115 per 1000	659 more (182 more to 1000 more)	
Histological improvement (continuous)	370 (6 studies)	⊕⊕⊕⊕ Very low^c	—	—	SMD 1.01 higher (0.36 higher to 1.66 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
Endoscopic improvement (dichotomous)	11 (1 study)	⊕⊕⊕⊕ Low^d	Not estimable	Study population		Both groups had zero patients with endoscopic improvement.
13 weeks				Not estimable	Not estimable	
Endoscopic improvement (continuous)	197 (3 studies)	⊕⊕⊕⊕ Very low^c	—	—	SMD 2.79 higher (0.36 higher to 5.22 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
12 to 24 weeks						
Withdrawals due to adverse events	792 (8 studies)	⊕⊕⊕⊕ Low^d	RR 1.55 (0.88 to 2.74)	Study population		—
12 to 44 weeks				58 per 1000	32 more per 1000 (7 fewer to 101 more)	

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

** **The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to imprecision.

^bDowngraded once due to imprecision.

^cDowngraded twice due to serious inconsistency ($I^2 = 83\%$ and $I^2 = 97\%$ respectively) and once due to imprecision.

^dDowngraded twice due to serious imprecision.

Summary of findings 4. Cromolyn sodium compared to placebo

Cromolyn sodium compared to placebo

Patient or population: active EoE pediatric patients

Setting: medical center

Intervention: cromolyn sodium

Comparison: placebo

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with placebo	Risk difference with cromolyn sodium	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement (continuous) 8 weeks	14 (1 study)	⊕⊕⊕⊕ Low^a	—	—	MD 4.70 higher (12.09 lower to 21.49 higher)	Measured on the Pediatric Eosinophilic Esophagitis Symptom Score (PEESS)
Histological improvement (dichotomous)	—	—	—	—	—	No data
Histological improvement (continuous) 8 weeks	15 (1 study)	⊕⊕⊕⊕ Low^a	—	—	MD 14.20 higher (36.90 lower to 65.30 higher)	Measured as change in peak eos/hpf from baseline
Endoscopic improvement (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement (continuous)	—	—	—	—	—	No data
Withdrawals due to adverse events at 8 weeks	16 (1 study)	⊕⊕⊕⊕ Low^a	RR 0.27 (0.01 to 5.70)	Study population 143 per 1000	104 less per 1000 (141 less to 672 more)	—

*The risk in the intervention group (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **eos/hpf:** eosinophils/high-power field; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision.

Summary of findings 5. PGD2R antagonist OC000459 compared to placebo

PGD2R antagonist OC000459 compared to placebo

Patient or population: active EoE patients

Setting: medical center

Intervention: PGD2R antagonist OC000459

Comparison: placebo

Outcomes	N° of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with placebo	Risk difference with PGD2R antagonist OC000459	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement (continuous) 8 weeks	26 (1 study)	⊕○○○ Very low^a	—	—	MD 1.06 lower (6.80 lower to 4.68 higher)	Measured as combined post-treatment means of several questionnaires
Histological improvement (dichotomous)	—	—	—	—	—	No data



Histological improvement (continuous) 8 weeks	26 (1 study)	⊕○○○ Very low^a	—	—	MD 26.21 higher (23.78 lower to 76.20 higher)	Measured as post-treatment eosinophil load
Endoscopic improvement (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement (continuous) 8 weeks	26 (1 study)	⊕○○○ Very low^a	—	—	MD 0.49 lower (2.05 lower to 1.07 higher)	Measured on a 10-point visual analogue scale
Withdrawals due to adverse events	26 (1 study)	⊕○○○ Very low^a	Not estimable	Study population	—	—
				0 per 1000	0 per 1000	

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

** **The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and once due to risk of bias for unclear allocation concealment and blinding of outcome assessment.

Summary of findings 6. Swallowed fluticasone compared to oral prednisone

Swallowed fluticasone compared to oral prednisone

Patient or population: active EoE pediatric patients

Setting: medical center

Intervention: swallowed fluticasone

Comparison: oral prednisone

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with comparator	Risk difference with corticosteroid	
Clinical improvement at 4 weeks (dichotomous)	80 (1 study)	⊕⊕⊕⊕ Very low^a	RR 1.09 (0.90 to 1.33)	Study population 800 per 1000	72 more per 1000 (80 fewer to 200 more)	—
Clinical improvement at 4 weeks (continuous)	—	—	—	—	—	No data
Histological improvement at 4 weeks (dichotomous)	80 (1 study)	⊕⊕⊕⊕ Very low^a	RR 1.1 (0.87 to 1.38)	Study population 750 per 1000	75 more per 1000 (98 fewer to 285 more)	—
Histological improvement at 4 weeks (continuous)	68 (1 study)	⊕⊕⊕⊕ Very low^a	—	—	MD 4.45 lower (9.08 lower to 0.18 higher)	Measured as mean peak eosinophils
Endoscopic improvement at 4 weeks (dichotomous)	80 (1 study)	⊕⊕⊕⊕ Very low^a	RR 1.13 (0.91 to 1.41)	Study population 750 per 1000	97 more per 1000 (68 fewer to 308 more)	—
Endoscopic improvement (continuous)	—	—	—	—	—	No data
Withdrawals due to adverse events at 4 weeks	80 (1 study)	⊕⊕⊕⊕ Very low^a	RR 0.50 (0.16 to 1.53)	Study population 200 per 1000	100 fewer per 1000 (168 fewer to 106 more)	—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

** **The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and once due to risk of bias for blinding of participants and personnel.

Summary of findings 7. Oral viscous budesonide compared to swallowed fluticasone

Oral viscous budesonide compared to swallowed fluticasone

Patient or population: active EoE patients

Setting: medical center

Intervention: oral viscous budesonide

Comparison: swallowed fluticasone

Outcomes	№ of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with swallowed fluticasone	Risk difference with oral viscous budesonide	
Clinical improvement at study endpoint (dichotomous)	—	—	—	—	—	No data
Clinical improvement at 8 weeks (continuous)	84 (1 study)	⊕⊕⊕⊕ Low^a	—	—	MD 0.6 lower (3.78 lower to 2.58 higher)	Measured on the Dysphagia Score Questionnaire
Histological improvement at study endpoint (dichotomous)	129 (1 study)	⊕⊕⊕⊕ Low^a	RR 1.13 (0.84 to 1.51)	Study population		—
				547 per 1000	71 more per 1000 (88 fewer to 279 more)	—
Histological improvement at 8 weeks (continuous)	111 (1 study)	⊕⊕⊕⊕ Low^a	—	—	MD 6.2 higher (5.63 lower to 18.03 higher)	Measured as eosinophils per high-power field

Endoscopic improvement at study end-point (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement at 8 weeks (continuous)	111 (1 study)	⊕⊕⊕⊕ Low^a	—	—	MD 0.7 higher (0.03 lower to 1.43 higher)	Measured on the endoscopic reference score
Withdrawals due to adverse events at 8 weeks	129 (1 study)	⊕⊕⊕⊕ Low^a	RR 0.98 (0.42 to 2.32)	Study population		—
				141 per 1000	3 fewer per 1000 (82 fewer to 186 more)	—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision.

Summary of findings 8. Esomeprazole compared to fluticasone

Esomeprazole compared to fluticasone

Patient or population: active EoE patients

Setting: medical centers

Intervention: esomeprazole

Comparison: fluticasone

Outcomes	Nº of participants	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)	Comments
----------	--------------------	-----------------------------------	--------------------------	--	----------

	(studies)			Risk with fluticasone	Risk difference with esomeprazole	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement at 8 weeks (continuous)	67 (2 studies)	⊕⊕⊕⊕ Very low^a	—	—	SMD 0.28 higher (0.2 lower to 0.76 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
Histological improvement at 8 weeks (dichotomous)	72 (2 studies)	⊕⊕⊕⊕ Very low^b	RR 1.62 (0.77 to 3.41)	Study population 222 per 1000	151 more per 1000 (42 fewer to 551 more)	—
Histological improvement at 8 weeks (continuous)	70 (2 studies)	⊕⊕⊕⊕ Very low^b	—	—	SMD 0.28 higher (0.20 lower to 0.76 higher)	As a rule of thumb, 0.2 SMD represents a small difference, 0.5 a moderate, and 0.8 a large effect.
Endoscopic improvement (dichotomous)	—	—	—	—	—	The studies reported data on specific endoscopic findings, which can be found in Table 1 .
Endoscopic improvement (continuous)	—	—	—	—	—	No data
Withdrawals due to adverse events at 8 weeks	72 (2 studies)	⊕⊕⊕⊕ Very low^b	RR 0.95 (0.07 to 13.38)	Study population 83 per 1000	4 fewer per 1000 (77 fewer to 1000 more)	—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded once for inconsistency ($I^2 = 81\%$), once for imprecision, and once for risk of bias

^bDowngraded twice for serious imprecision and once for risk of bias for blinding of participants, personnel, and outcome assessment, and selective reporting.

Summary of findings 9. One-food elimination diet compared to four-food elimination diet

One-food elimination diet compared to four-food elimination diet

Patient or population: active EoE pediatric patients

Setting: medical centers

Intervention: one-food elimination diet

Comparison: four-food elimination diet

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects [†] (95% CI)		Comments
				Risk with four-food elimination diet	Risk difference with one-food elimination diet	
Clinical improvement (dichotomous)	—	—	—	—	—	—
Clinical improvement at 12 weeks (continuous)	50 (1 study)	⊕○○○ Very low^a	—	—	MD 7.5 lower (16.28 lower to 1.28 higher)	Measured on the EoE Symptom Activity Index
Histological improvement at 12 weeks (dichotomous)	63 (1 study)	⊕○○○ Very low^a	RR 2.26 (1.15 to 4.43)	Study population 280 per 1000	353 more per 1000 (42 more to 960 more)	—
Histological improvement at study endpoint (continuous)	—	—	—	—	—	No data
Endoscopic improvement at study endpoint (dichotomous)	—	—	—	—	—	No data

Endoscopic improvement at 12 weeks (continuous)	34 (1 study)	⊕○○○ Very low^a	—	—	MD 0.6 lower (2.15 lower to 0.95 higher)	Measured on the endoscopic reference score
Withdrawals due to adverse events at 12 weeks	63 (1 study)	⊕○○○ Very low^a	RR 0.33 (0.11 to 0.98)	Study population	320 per 1000	214 fewer per 1000 (285 fewer to 6 fewer)

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and once due to risk of bias for unclear blinding of participants and personnel, and attrition.

Summary of findings 10. One-food elimination diet compared to six-food elimination diet

One-food elimination diet compared to six-food elimination diet

Patient or population: active EoE pediatric patients

Setting: medical centers

Intervention: one-food elimination diet

Comparison: six-food elimination diet

Outcomes	N° of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with six-food elimination diet	Risk difference with one-food elimination diet	

Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement at 6 weeks (continuous)	129 (1 study)	⊕⊕⊕⊕ Very low^a	—	—	MD 5.2 lower (11.06 lower to 0.66 higher)	Measured on the EoE Symptom Activity Index
Histological improvement at 6 weeks (dichotomous)	129 (1 study)	⊕⊕⊕⊕ Very low^a	RR 0.85 (0.54 to 1.33)	Study population	—	—
				403 per 1000	60 fewer per 1000 (185 fewer to 133 more)	—
Histological improvement at 6 weeks (continuous)	129 (1 study)	⊕⊕⊕⊕ Very low^a	—	—	MD 6.8 higher (10.4 lower to 24 higher)	Measured as changes in the EoE Histologic Scoring System
Endoscopic improvement at study endpoint (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement at 6 weeks (continuous)	129 (1 study)	⊕⊕⊕⊕ Very low^a	—	—	MD 0.42 lower (1.67 lower to 0.83 higher)	Measured on the endoscopic reference score
Withdrawals due to adverse events	129 (1 study)	⊕⊕⊕⊕ Very low^a	RR 0.62 (0.11 to 3.57)	Study population	—	—
				403 per 1000	60 fewer per 1000 (185 fewer to 133 more)	—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and risk of bias for blinding of participants and personnel, and other bias.

Summary of findings 11. Four-food elimination diet with omeprazole compared to omeprazole

Four-food elimination diet with omeprazole compared to omeprazole

Patient or population: active EoE patients

Setting: medical centers

Intervention: four-food elimination diet with omeprazole

Comparison: omeprazole

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with omeprazole	Risk difference with four-food elimination diet with omeprazole	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement (continuous)	—	—	—	—	—	No data
Histological improvement at 8 to 12 weeks (dichotomous)	64 (1 study)	⊕⊕⊕⊕ Very low^a	RR 1.57 (0.99 to 2.48)	Study population 438 per 1000	250 more per 1000 (4 fewer to 648 more)	—
Histological improvement at 8 to 12 weeks (continuous)	58 (1 study)	⊕⊕⊕⊕ Very low^a	—	—	MD 9.50 higher (11.18 lower to 30.18 higher)	—
Endoscopic improvement at study endpoint (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement at study endpoint (continuous)	—	—	—	—	—	No data
Withdrawals due to adverse events at 8 to 12 weeks	64 (1 study)	⊕⊕⊕⊕ Very low^a	RR 5.00 (0.62 to 40.44)	Study population 31 per 1000	124 more per 1000 (12 fewer to 1000 more)	—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and once due to risk of bias for blinding of participants, personnel, and outcome assessment, unclear attrition, and selective reporting.

Summary of findings 12. Four-food elimination and amino acid formula compared to four-food elimination diet

Four-food elimination and amino acid formula compared to four-food elimination diet

Patient or population: active EoE patients

Setting: medical center

Intervention: four-food elimination and amino acid formula

Comparison: four-food elimination diet

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with four-food elimination diet	Risk difference with four-food elimination and amino acid formula	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement (continuous)	41 (1 study)	⊕○○○ Very low^a	—	—	MD 0.50 lower (2.41 lower to 1.41 higher)	—
Histological improvement at 6 weeks (dichotomous)	41 (1 study)	⊕○○○ Very low^a	RR 1.90 (0.79 to 4.60)	Study population		—
				250 per 1000	225 more per 1000 (53 fewer to 900 more)	—

Histological improvement at 6 weeks (continuous)	41 (1 study)	⊕⊕⊕⊕ Very low^a	—	—	MD 13.8 higher (9.5 lower to 37.1 higher)	Measured as peak eosinophil count
Endoscopic improvement (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement at study endpoint (continuous)	41 (1 study)	⊕⊕⊕⊕ Very low^a	—	—	MD 1.00 lower (2.83 lower to 0.83 higher)	Measured on the endoscopic reference score
Withdrawals due to adverse events at 6 weeks	41 participants (1 study)	⊕⊕⊕⊕ Very low^a	RR 0.95 (0.06 to 14.22)	Study population	—	—
				50 per 1000	3 fewer per 1000 (47 fewer to 661 more)	—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

** **The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and once due to risk of bias for unclear randomization, blinding of participants, personnel, and outcome assessment.

Summary of findings 13. Nebulized budesonide compared to viscous budesonide

Nebulized budesonide compared to viscous budesonide

Patient or population: active EoE patients

Setting: medical center

Intervention: nebulized budesonide

Comparison: viscous budesonide

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with vis-cous budes-onide	Risk difference with nebu-lized budesonide	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement at 8 weeks (continuous)	22 (1 study)	⊕○○○ Very low^a	—	—	MD 6.00 lower (18.3 lower to 6.3 higher)	—
Histological improvement at 8 weeks (dichotomous)	—	—	—	—	—	No data
Histological improvement at 8 weeks (continuous)	22 (1 study)	⊕○○○ Very low^a	—	—	MD 78.00 higher (20.81 higher to 135.19 higher)	—
Endoscopic improvement (dichotomous)	25 (1 study)	⊕○○○ Very low^a	—	—	—	The study reported specific endoscopic characteristics which can be found in Table 1
Endoscopic improvement (continuous)	—	—	—	—	—	No data
Withdrawals due to adverse events at 8 weeks	25 (1 study)	⊕○○○ Very low^a	Not estimable	Study population 0 per 1000 0 fewer per 1000 (0 fewer to 0 fewer)		—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

** **The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to imprecision and once due to risk of bias for blinding of participants and personnel.

Summary of findings 14. Viaskin milk patch compared to placebo

Viaskin milk patch compared to placebo

Patient or population: active EoE pediatric patients

Setting: medical center

Intervention: Viaskin milk patch

Comparison: placebo

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with placebo	Risk difference with Viaskin milk patch	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement at 44 weeks (continuous)	9 (1 study)	⊕⊕⊕⊕ Low^a	—	—	MD 1.29 higher (0.83 lower to 3.41 higher)	Measured on the eosinophilic esophagitis symptom score
Histological improvement (dichotomous)	—	—	—	—	—	No data
Histological improvement at 44 weeks (continuous)	9 (1 study)	⊕⊕⊕⊕ Low^a	—	—	MD 69.43 higher (21.75 lower to 160.61 higher)	Measured as change in maximum esophageal eosinophil count from baseline to end of study
Endoscopic improvement (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement at 44 weeks (continuous)	20 (1 study)	⊕⊕⊕⊕ Low^a	—	—	MD 0.33 lower (2 lower to 1.34 higher)	Measured on the endoscopic reference score

Withdrawals due to adverse events at 44 weeks	20 (1 study)	⊕⊕○○ Low^a	RR 1.12 (0.05 to 23.99)	Study population	—
				0 per 1000	66 fewer per 1000 (372 fewer to 660 more)

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

** **The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision.

Summary of findings 15. Leukotriene receptor antagonist compared to placebo for maintenance of remission

Leukotriene receptor antagonist compared to placebo for maintenance of remission

Patient or population: inactive EoE patients

Setting: medical center

Intervention: leukotriene receptor antagonist

Comparison: placebo

Outcomes	№ of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with placebo	Risk difference with leukotriene receptor antagonist	
Clinical improvement at 26 weeks (dichotomous)	41 (1 study)	⊕○○○ Very low^a	RR 1.68 (0.66 to 4.28)	Study population 238 per 1000	162 more per 1000	—

					(81 fewer to 781 more)	
Clinical improvement (continuous)	—	—	—	—	—	No data
Histological improvement (dichotomous)	—	—	—	—	—	No data
Histological improvement (continuous)	—	—	—	—	—	No data
Endoscopic improvement (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement (continuous)	—	—	—	—	—	No data
Withdrawals due to adverse events at 26 weeks	41 (1 study)	⊕⊕⊕⊕ Very low^a	RR 2.10 (0.21 to 21.39)	Study population		—
				48 per 1000	53 more per 1000	—
					(38 fewer to 979 more)	

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

** **The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and once due to risk of bias for unclear selective reporting.

Summary of findings 16. Mepolizumab 10 mg/kg compared to mepolizumab 0.55 mg/kg

Mepolizumab 10 mg/kg compared to mepolizumab 0.55 mg/kg

Patient or population: active EoE pediatric patients

Setting: medical centers

Intervention: mepolizumab 10 mg/kg

Comparison: mepolizumab 0.55 mg/kg

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with viscous mepolizumab 0.55 mg/kg	Risk difference with mepolizumab 10 mg/kg	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement at (continuous)	—	—	—	—	—	No data
Histological improvement at 12 weeks (dichotomous)	39 (1 study)	⊕⊕⊕⊕ Very low^a	RR 1.19 (0.37 to 3.77)	Study population 211 per 1000	40 more per 1000 (133 fewer to 584 more)	—
Histological improvement (continuous)	—	—	—	—	—	No data
Endoscopic improvement (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement (continuous)	—	—	—	—	—	No data
Withdrawals due to adverse events at 8 weeks	39 (1 study)	⊕⊕⊕⊕ Very low^a	RR 0.63 (0.12 to 3.38)	Study population 158 per 1000	48 more per 1000 (139 fewer to 376 more)	—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

** **The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and once due to risk of bias.

Summary of findings 17. Mepolizumab 2.5 mg/kg compared to mepolizumab 0.55 mg/kg

Mepolizumab 2.5 mg/kg compared to mepolizumab 0.55 mg/kg

Patient or population: active EoE pediatric patients

Setting: medical centers

Intervention: mepolizumab 2.5 mg/kg

Comparison: mepolizumab 0.55 mg/kg

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with viscous mepolizumab 0.55 mg/kg	Risk difference with mepolizumab 2.5 mg/kg	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement at (continuous)	—	—	—	—	—	No data
Histological improvement at 12 weeks (dichotomous)	39 (1 study)	⊕⊕⊕⊕ Very low^a	RR 2.14 (0.79 to 5.79)	Study population 211 per 1000	241 more per 1000 (44 fewer to 1000 more)	—
Histological improvement (continuous)	—	—	—	—	—	No data
Endoscopic improvement (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement (continuous)	—	—	—	—	—	No data
Withdrawals due to adverse events at 8 weeks	39 (1 study)	⊕⊕⊕⊕ Very low^a	RR 0.32 (0.04 to 2.79)	Study population 158 per 1000	107 fewer per 1000 (152 fewer to 283 more)	—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and once due to risk of bias for unclear randomization, unclear blinding of outcome assessment, and selective reporting.

Summary of findings 18. Mepolizumab 10 mg/kg compared to mepolizumab 2.5 mg/kg

Mepolizumab 10 mg/kg compared to mepolizumab 2.5 mg/kg

Patient or population: active EoE pediatric patients

Setting: medical centers

Intervention: mepolizumab 10 mg/kg

Comparison: mepolizumab 2.5 mg/kg

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Anticipated absolute effects* (95% CI)		Comments
				Risk with viscous mepolizumab 2.5 mg/kg	Risk difference with mepolizumab 10 mg/kg	
Clinical improvement (dichotomous)	—	—	—	—	—	No data
Clinical improvement at (continuous)	—	—	—	—	—	No data
Histological improvement at 12 weeks (dichotomous)	40 (1 study)	⊕○○○ Very low^a	RR 0.56 (0.23 to 1.37)	Study population 450 per 1000	198 fewer per 1000 (347 fewer to 167 more)	—
Histological improvement (continuous)	—	—	—	—	—	No data

Endoscopic improvement (dichotomous)	—	—	—	—	—	No data
Endoscopic improvement (continuous)	—	—	—	—	—	No data
Withdrawals due to adverse events at 8 weeks	40 (1 study)	⊕⊕⊕⊕ Very low^a	RR 2.00 (0.20 to 20.33)	Study population	—	—
				50 per 1000	50 more per 1000 (40 fewer to 967 more)	—

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to serious imprecision and once due to risk of bias for unclear randomization, unclear blinding of outcome assessment, and selective reporting.

Summary of findings 19. Six-food elimination diet compared to swallowed fluticasone compared to swallowed budesonide compared to oral viscous budesonide

Six-food elimination diet compared to swallowed fluticasone compared to swallowed budesonide compared to oral viscous budesonide

Patient or population: active EoE pediatric patients

Setting: medical center

Comparison: six-food elimination diet versus swallowed fluticasone versus swallowed budesonide versus oral viscous budesonide

Outcomes	Nº of participants (studies)	Certainty of the evidence (GRADE)	Relative effect (95% CI)	Reported results	Comments
Clinical improvement (dichotomous)	—	—	—	—	No data

Clinical improvement (continuous)	—	—	—	—	No data
Histological improvement at 8 weeks (dichotomous)	64 (1 study)	⊕⊕⊕⊕ Very low^a		69% of participants in the six-food elimination diet achieved histological improvement, 67% in the swallowed fluticasone group, 75% in the swallowed budesonide group, and 85% in the oral viscous budesonide group.	—
Histological improvement (continuous)	—	—	—	—	No data
Endoscopic improvement (dichotomous)	—	—	—	—	No data
Endoscopic improvement (continuous)	—	—	—	—	No data
Withdrawals due to adverse events at 26 weeks	—	—	—	—	No data

***The risk in the intervention group** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

**** The risk for the control intervention has been calculated by dividing the number of cases to the number of randomized participants, using the numbers of our analyses. The risk for the comparison intervention has been calculated by multiplying the control risk with the RR and CI limits. The risk difference has been calculated by subtracting the control risk from the comparison intervention risk.**

CI: confidence interval; **EoE:** eosinophilic esophagitis; **MD:** mean difference; **NNTB:** number needed to treat for an additional beneficial outcome; **RR:** risk ratio; **SMD:** standardized mean difference

GRADE Working Group grades of evidence

High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.

Moderate certainty: we are moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.

Low certainty: our confidence in the effect estimate is limited: the true effect may be substantially different from the estimate of the effect.

Very low certainty: we have very little confidence in the effect estimate: the true effect is likely to be substantially different from the estimate of effect.

^aDowngraded twice due to risk of bias and once due to imprecision for unclear randomization, unclear blinding of outcome assessment, and selective reporting.

BACKGROUND

Description of the condition

Eosinophilic esophagitis (EoE) is a chronic type 2 antigen-mediated inflammatory disorder of the esophagus, causing upper gastrointestinal symptoms and characterized by increased esophageal infiltration with intraepithelial eosinophils (Liacouras 2011; Rothenberg 2015). It was originally described during the 1970s in adults with symptoms of esophagitis, who often had allergies, and who had high esophageal eosinophil counts. The diagnostic criteria for eosinophilic esophagitis are clinical symptoms of esophageal disease, a histological abnormality of 15 or more intraepithelial eosinophils per high-power field (hpf) on endoscopy, the exclusion of gastroesophageal reflux disease (GERD), and consistent endoscopic findings (Furuta 2007). Eosinophilic esophagitis affects young infants to adults. In young children, the symptoms may be associated with feeding difficulties, vomiting, weight loss, and abdominal pain, however dysphagia and food impaction occur more often among teenagers and adults (Croese 2003; Liacouras 1998; Liacouras 2011; Orenstein 2000). In eosinophilic esophagitis, the mucosa may look normal macroscopically, however thickening, ringing, furrowing, and erosion have been reported (Dellon 2018; Furuta 2015; Hassall 1996; Khan 2003; Orenstein 2000).

Currently, there is no cure for eosinophilic esophagitis, making long-term treatment critical. Treatment goals for eosinophilic esophagitis include improvement in clinical symptoms, resolution of esophageal eosinophilia and other histologic abnormalities, endoscopic improvement, improved quality of life, improved esophageal function, minimized adverse effects of treatment, and prevention of disease progression and subsequent complications (Muir 2021). Standard treatment modalities for eosinophilic esophagitis include dietary, pharmacologic, and endoscopic interventions. Dietary therapy involves empiric food elimination (most commonly milk protein) through dietary elimination and/or formula. Although effective, this approach can result in several endoscopies initially and be challenging to sustain long-term (Kelly 1995; Markowitz 2003). Pharmacologic therapy includes proton pump inhibitors, topical glucocorticoids, and rapidly emerging biologics, including the first (US) Food & Drug Administration (FDA) approved medication for eosinophilic esophagitis (Greuter 2017; Muir 2021; Faubion 1998; Liacouras 1998; Teitelbaum 2002). Endoscopic intervention such as esophageal dilation is effective in relieving symptoms such as dysphagia, however it does not alter the underlying inflammation. This therapy is often used in conjunction with diet and/or medication. Eosinophilic esophagitis is a chronic disease that requires ongoing therapy and long-term monitoring. If left untreated, eosinophilic esophagitis can result in complications such as fibrostenosis and strictures of the esophagus (Muir 2021; Schoepfer 2013).

Description of the intervention

Medical management of eosinophilic esophagitis includes pharmacological therapy and dietary elimination.

Steroids are anti-inflammatory drugs that have been used for the induction and maintenance of remission of eosinophilic esophagitis (Rank 2020). Budesonide and fluticasone have been the most frequently used, and to a lesser extent prednisone, beclomethasone, mometasone, and ciclesonide. Although initially

administered systemically, swallowed topical administration has been the most frequent way of delivery, either adapted from asthma formulations (swallowed metered dose or nebulized solutions) or with compound viscous formulations. More recently esophageal-specific topical steroid formulations, either oral suspension or oro-dispersible tablets, have been developed.

Diverse biological therapies have been studied for the treatment of eosinophilic esophagitis including anti-IL5 (mepolizumab, reslizumab), anti-IgE (omalizumab), anti-IL4r (dupilumab), anti-IL13 (RPC4046, alias cendakimab; QAX576, alias dectrekumab), and anti-sialic acid binding Ig-like lectin 8 (Siglec-8) (lirentelimab) (Nhu 2022).

There is very limited evidence of the efficacy of other drugs including mast cell inhibitors (sodium cromoglycate), leukotriene receptor antagonists (montelukast), and chemoattractant receptor-homologous molecule on Th2 cells (CRTH2) antagonist (OC000459) for the treatment of eosinophilic esophagitis (Straumann 2013).

The role of proton pump inhibitors (PPIs) has evolved from a tool to diagnose eosinophilic esophagitis, by excluding other entities associated with esophageal eosinophilia, to a true treatment for the condition (Dellon 2018). Different PPI drugs, such as omeprazole, esomeprazole, pantoprazole, and lansoprazole have been used to induce and maintain remission (Franciosi 2022).

Elimination diets have been used since the description of the disease, providing evidence that eosinophilic esophagitis is predominantly triggered by food antigens (Kelly 1995). Dietary strategies for treatment of eosinophilic esophagitis comprise elemental diet, allergy testing-directed elimination diet, and empiric elimination diets, avoiding the most frequent food triggers (milk, wheat, egg, soy/legumes, nuts, and fish/seafood).

How the intervention might work

Inflammation and the resulting symptoms of esophageal dysfunction in eosinophilic esophagitis are thought to result from penetration of the esophageal mucosa by food or aero-antigens resulting in cellular response and symptoms of esophageal dysfunction. A breach in the integrity of the esophageal epithelium, potentially facilitated by gastric acid exposure and/or carriage of genetic variants that compromise epithelial barrier function, allows ingress of food or aeroallergens leading to initiation of an immune response. Interleukins produced by activated Th2 cells can act directly to recruit eosinophils to the esophagus (IL-5), or can stimulate the epithelium to express inflammatory genes (IL-4/IL-13), including eotaxin-3, by activation of cell surface receptors that signal through a pathway involving JAKs and STAT6. Esophageal eosinophilia in eosinophilic esophagitis is driven largely by STAT6-dependent local expression of eotaxin-3.

Medical therapies for eosinophilic esophagitis have targeted esophageal inflammation broadly (corticosteroids), or targeted biologic mediators, including anti-IL5 (mepolizumab, reslizumab), anti-IgE (omalizumab), anti-IL4r (dupilumab), anti-IL13 (RPC4046, alias cendakimab; QAX576, alias dectrekumab), and anti-sialic acid binding Ig-like lectin 8 (lirentelimab) (Nhu 2022). The proposed mechanism of PPI therapy is by gastric acid suppression leading to a restoration of esophageal barrier function and unrelated PPI-mediated anti-inflammatory effects. Elimination diets have been used since the initial description of the disease, providing evidence

Medical treatment of eosinophilic esophagitis (Review)

that eosinophilic esophagitis is predominantly triggered by food antigens. Dietary strategies for treating eosinophilic esophagitis include the elemental diet, the allergy testing-directed elimination diet, and empiric elimination diets, which avoid the most frequent food triggers (milk, wheat, egg, soy/legumes, nuts, and fish/seafood).

Why it is important to do this review

There is no universally accepted treatment for eosinophilic esophagitis. Topical corticosteroids, hypoallergenic diets, proton pump inhibitors, biologics, and dilation have all been used to treat eosinophilic esophagitis (Muir 2021). In 2010, a previous version of this review was published that included only three randomized controlled trials (RCTs). The number of published RCTs in pediatric and adult eosinophilic esophagitis that meet our inclusion criteria has grown substantially, with a rapid pace of clinical trial publications and changing outcome metrics. The purpose of this review is to review the evidence from RCTs evaluating non-surgical interventions for eosinophilic esophagitis.

OBJECTIVES

To evaluate the efficacy and safety of medical interventions for people with eosinophilic esophagitis.

METHODS

Criteria for considering studies for this review

Types of studies

We considered all types of randomized controlled trials (RCTs) for inclusion. We excluded quasi-randomized trials (using no or non-appropriate randomization).

Types of participants

People of any age with a diagnosis of eosinophilic esophagitis, either with active disease (increased number of eosinophils (at least 15 eos/hpf) on esophageal biopsy and with symptoms of esophageal dysfunction) or inactive disease.

We applied no restrictions on sex, disease duration, or previous medication exposure.

We considered studies with only a subset of eligible participants for inclusion. If the subset had been planned for a subgroup analysis, we explored its impact through the methods described in [Subgroup analysis and investigation of heterogeneity](#). If a subgroup analysis had not been planned, the authoring team liaised to discuss the effect this may have on the planned outcomes and whether further subgroup analysis was necessary.

Types of interventions

Studies comparing any medical intervention (e.g. topical corticosteroid, biologic therapy, systemic corticosteroid, leukotriene receptor antagonist, mast cell stabilizer, epicutaneous immunotherapy, proton pump inhibitor) or food elimination diet (e.g. empiric elimination diet, elemental diet), either alone or in combination, to any other intervention.

Types of outcome measures

We included both dichotomous and continuous outcomes. When multiple thresholds were prespecified by the study, we chose the most inclusive threshold for the analysis. Study outcomes were not relevant for determining study eligibility.

Primary outcomes

- Clinical symptom improvement - defined by the study, either as a clinically successful improvement based on achieving a prespecified threshold on a symptom scoring scale (dichotomous), or absolute or relative symptom scores (continuous) - as measured by the authors at study end and all available intermediate study time points.
- Histological improvement - defined by the study, using a recognized histological grading system (continuous), achieving a prespecified eosinophil count threshold measured per high-powered microscope field (dichotomous), or absolute or relative eosinophil counts per high-powered microscope field (continuous) - as measured by the authors at study end and all available intermediate study time points.
- Endoscopic improvement - defined by the study, achieving a prespecified threshold on an endoscopic scoring scale (dichotomous), or absolute or relative endoscopic assessment scores (continuous) - as measured by the authors at study end and all available intermediate study time points.
- Withdrawals due to adverse events - (dichotomous) - as measured by the authors at study end.

Secondary outcomes

- Participants with serious adverse events as defined by the study (dichotomous) - as measured by the authors at study end.
- Total number of participants with adverse events as defined by the study (dichotomous) - as measured by the authors at study end.
- Quality of life (QOL) improvement as defined by the study, either as an improvement in quality of life based on achieving a prespecified threshold on a QOL scoring scale (dichotomous), or absolute or relative QOL scores (continuous) - as measured by the authors at study end and all available intermediate study time points.

Search methods for identification of studies

Electronic searches

On 24 October 2021 and 3 March 2023, the Cochrane Gut Information Specialist searched the following sources:

- Cochrane Central Register of Controlled Trials (CENTRAL) via the Cochrane Library (until search date; [Appendix 1](#));
- MEDLINE via Ovid SP (1946 to March 02, 2023; [Appendix 2](#));
- Embase via Ovid SP (1974 to 2023 Week 08; [Appendix 3](#));
- ClinicalTrials.gov (until search date; [Appendix 4](#));
- World Health Organization International Clinical Trials Registry Platform (WHO ICTRP) (until search date; [Appendix 5](#)).

There were no limitations to publication date, language, status, or document type in this search.

Searching other resources

We handsearched reference lists from the trials identified by electronic searching to identify further relevant trials. Other sources that we searched included reference lists of textbooks, reviews (*Cochrane Database of Systematic Reviews* and others), previous trials, and conference proceedings.

Data collection and analysis

Selection of studies

Pairs of authors independently assessed publications identified by the search strategy to determine eligibility based on the above inclusion criteria using *Covidence*, initially as titles/abstracts, followed by full-text assessments. Any disagreement was resolved by discussion and consensus among the authors. If consensus could not be reached, a third author was consulted. We documented the results of this process in a flow diagram (*PRISMA 2020*).

Data extraction and management

We collected data from the included studies using a piloted data collection form. Pairs of authors independently extracted data. Disagreements were resolved by discussion and consensus. A third author was consulted when consensus was not reached.

The extracted data included the following:

- General information (title, journal, year, publication type).
- Study information (design, setting, dates, single- or multi-center; RCT duration and endpoints; study outcomes; funding source; conflicts of interest).
- Participant information (disease activity; diagnostic criteria; inclusion and exclusion criteria; age; sex; concomitant medications).
- Intervention and control (type, dose, method of delivery of medication).
- Eligibility (total number of patients randomized and reaching end of study).
- Review outcomes (continuous scoring system or dichotomous success definition; outcome data at study endpoints).

For studies requiring translation we used online translation software, and if this was not adequate we sought translations by speakers of the relevant languages.

Assessment of risk of bias in included studies

Using the Cochrane risk of bias tool (*Higgins 2011*), pairs of authors independently assessed the risk of bias of each included study. We assessed the following factors:

- sequence generation (i.e. randomization method);
- allocation sequence concealment;
- blinding;
- incomplete outcome data (i.e. methods used by investigators to deal with attrition);
- selective outcome reporting (i.e. investigators reported all outcomes); and
- other potential sources of bias (i.e. anything else that could have increased bias).

We judged studies to be of high, low, or unclear risk of bias. Disagreement was resolved by consensus via discussion. A third author resolved cases where consensus was not reached.

Measures of treatment effect

We analyzed all data using Review Manager Web (*RevMan Web 2022*). For dichotomous outcomes, we expressed the treatment effect as a risk ratio (RR) with corresponding 95% confidence interval (CI). For continuous outcomes, we expressed the treatment effect as a mean or standardized mean difference (MD or SMD) with 95% CI.

Unit of analysis issues

The participant was the unit of analysis. For studies comparing more than two intervention groups, we made multiple pairwise comparisons between all possible pairs of intervention groups. In multiple-arm studies comparing different medication dosages to a comparator, we combined the different dosage groups into one. When we analyzed multiple treatment groups separately (e.g. different interventions within medication class groups), we divided the placebo group across the treatment groups.

To deal with repeated observations of participants, we divided shared intervention groups evenly among the comparisons. For dichotomous outcomes, we divided both the number of events and the total number of participants. To deal with events that may re-occur (e.g. adverse events), we reported on the proportion of participants who experienced at least one event. For continuous outcomes, we only divided the total number of participants, and left the means and standard deviations unchanged. We included cross-over studies, but we only pooled their data if they were reported separately before and after cross-over, and we only used pre-cross-over data.

In the case of cluster-RCTs we planned to use study data only if the authors used appropriate statistical methods in taking the clustering effect into account. We would also exclude cluster-RCTs from a sensitivity analysis to assess their impact on the results.

Dealing with missing data

Where data were missing, we contacted the corresponding authors of included studies to supply any unreported or unclear data. For all outcomes, we carried out analyses on an intention-to-treat (ITT) basis; that is, we included all participants randomized to each group in the analyses, and we analyzed all participants in the group to which they were allocated regardless of whether they received the allocated intervention.

For dichotomous efficacy outcomes we used the numbers randomized as denominators. As numerators, we used the numbers as reported by the authors. We assumed participants with missing or unclear data to be treatment failures. For safety outcomes, we considered participants with missing or unclear withdrawal data as withdrawals due to adverse events. The denominators used for this outcome were as reported by the authors. For serious and total adverse events we used the numbers of events per participants, as reported by the authors. Outcome data reported for mixes of randomized and non-randomized participants or post hoc data were discarded and not used for analysis.

For our dichotomous improvement outcomes, we scored an event when the prespecified threshold defined by the study was achieved. In studies that included threshold definitions for both partial and complete improvement, the total number of dichotomous events recorded reflects the sum of both the partial and complete events.

For missing continuous data, we estimated standard deviations from other available data, such as standard errors, or we imputed them using the methods suggested in [Higgins 2021b](#). We conducted analyses for continuous outcomes based on participants completing the trial, in line with available case analysis; this assumes that data were missing at random. If there was a discrepancy between the number randomized and the number analyzed in each treatment group, we calculated and reported the percentage lost to follow-up in each group.

We attempted to convert data presented in graphic from only to numerical data by digitizing them. When it was not possible to obtain missing data or gain clarity from the study authors, we recorded this in our risk of bias assessments, and rated it for bias based on the extent to which the missing data could bias our outcomes. Data that could not be used in our meta-analyses due to inadequate reporting (e.g. data not presented per intervention group, no available variance measures, data presented in graphic format which we could not convert) have been presented narratively in the additional tables.

Some studies may have reported data for more than one definition/threshold of a given outcome. We have reported which outcome definitions/thresholds we used in our meta-analyses in the description of included studies of the results section.

We employed the same methods in our subgroup and sensitivity analyses.

Assessment of heterogeneity

We scrutinized studies to ensure that they were clinically homogenous in terms of participants, interventions, comparators, and outcomes. To test for statistical heterogeneity, we used a Chi^2 test. A P value of less than 0.1 gave an indication of the presence of heterogeneity. We quantified and represented inconsistency using the I^2 statistic. We interpreted the thresholds as follows ([Higgins 2021a](#)):

- 0% to 40%: might not be important;
- 30% to 60%: may represent moderate heterogeneity;
- 50% to 90%: may represent substantial heterogeneity;
- 75% to 100%: considerable heterogeneity.

In the case of considerable statistical heterogeneity, we investigated whether this could be explained on clinical grounds or by risk of bias, in which case we conducted sensitivity analyses. If we could not find reasons for considerable statistical heterogeneity, we presented the results narratively, in detail.

Assessment of reporting biases

Our use of an inclusive search strategy minimized most reporting biases. We investigated publication bias using a funnel plot for outcomes with 10 or more studies and determined the magnitude of publication bias by visual inspection of the asymmetry of the

funnel plot or other methods mentioned in the *Cochrane Handbook for Systematic Reviews of Interventions* ([Egger 1997](#); [Higgins 2021a](#)).

Data synthesis

We combined data from individual studies for meta-analysis when we deemed the interventions, patient groups, and outcomes to be sufficiently similar (determined by consensus). We calculated the pooled RR and corresponding 95% CI for dichotomous outcomes. We calculated the pooled MD and corresponding 95% CI for continuous outcomes that were measured using the same units. We calculated the pooled standardized mean difference (SMD) and 95% CI when different scales were used to measure the same underlying construct. We carried out meta-analysis using a random-effects model.

Subgroup analysis and investigation of heterogeneity

We carried out subgroup analyses on the primary outcomes to further study the effects of a number of variables on the outcomes, when there were enough studies ([Deeks 2021](#)), using the formal test for subgroup differences in [RevMan Web 2022](#). Our planned subgroup analyses were decided by our review team as the characteristics most likely to have an impact on outcomes, and were:

1. age of participants (children < 18 years old, adults > 18 years old or mixed age populations);
2. specific interventions within categories (within the category of biologics these were grouped by mechanism including anti-IL13/anti-IL4R, anti-IL5, anti-IgE, anti-sialic acid binding Ig-like lectin 8; for others by specific intervention);
3. routes of delivery (specific to corticosteroids): esophageal-specific or not esophageal-specific through an inhaled route referred to as adapted asthma.

Any planned subgroup analyses that were ultimately not performed were due to no data being available or the original analysis comprising three or fewer studies.

The statistical methods described previously also applied to the subgroup analyses.

Sensitivity analysis

We conducted sensitivity analyses for the primary outcomes based on the following:

1. fixed-effect instead of random-effects model;
2. removing outcome data from studies that employed non-validated measures;
3. removing outcome data from non-peer-reviewed studies;
4. removing outcome data from studies judged to be at high risk for any risk of bias domain;
5. dichotomous histological reporting thresholds of < 15 eos/hpf, which was the first threshold used at the emergence of the field ([Konikoff 2006](#)) and currently employed;
6. dichotomous histological reporting thresholds of < 6 eos/hpf, which is currently advised by the FDA ([Reed 2018](#));
7. dichotomous histological reporting thresholds of < 1 eos/hpf, which signifies full or complete remission ([Greuter 2017](#)).

Any planned sensitivity analyses that were ultimately not performed were due to no data being available or the original analysis comprising three or fewer studies.

The statistical methods described previously also applied to the sensitivity analyses.

Summary of findings and assessment of the certainty of the evidence

We have presented summary of findings tables and GRADE decisions for all comparisons for all of our dichotomous and continuous primary outcomes. We assessed the overall certainty of evidence supporting the primary and secondary outcomes using the GRADE approach (Schünemann 2021). Evidence retrieved from RCTs is usually regarded as high-certainty. However, the certainty rating may be downgraded as a result of:

- risk of bias;
- indirect evidence;
- inconsistency (unexplained heterogeneity);
- imprecision;
- publication bias.

GRADE Working Group grades of evidence:

- High certainty: we are very confident that the true effect lies close to that of the estimate of the effect.
- Moderate certainty: we are moderately confident in the effect estimate; the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different.
- Low certainty: our confidence in the effect estimate is limited; the true effect may be substantially different from the estimate of the effect.

- Very low certainty: we have very little confidence in the effect estimate; the true effect is likely to be substantially different from the estimate of effect.

For the abstract of this review, we decided to focus on the two main comparisons of corticosteroids against placebo and biologics against placebo for induction of remission, reflecting what is most commonly used in current practice.

RESULTS

Description of studies

Results of the search


The literature search identified 3103 records through database searching and alternative sources. After removal of duplicates, 3089 unique records remained. Examination of the titles and abstracts left 294 records for full-text screening. After assessing all 294 records, we identified 208 records of 41 studies that met the inclusion criteria and these were included in the review ([Characteristics of included studies](#) table).

We excluded 39 studies (45 records; [Characteristics of excluded studies](#) table).

We identified 30 ongoing studies (39 records) ([Characteristics of ongoing studies](#) table). We categorized two studies (two records) as awaiting classification ([Characteristics of studies awaiting classification](#) table).

The results of the search are presented in the PRISMA flow diagram ([Figure 1](#)).

Figure 1. Study flow diagram.



Included studies

Setting

All studies took place in hospitals and medical centers, in North America, Europe, and Australia.

Twenty-three studies were multi-center studies (Assa'ad 2011; Butz 2014; Clayton 2014; Dellon 2017; Dellon 2022; Dellon 2021b; Dellon 2022a; Dellon 2022b; Gupta 2015; Heine 2019; Hirano 2019; Hirano 2020; Hirano 2020f; Hirano 2021; Kliever 2019; Kliever 2021; Lucendo 2019; Miehlike 2016; Rothenberg 2015; Rothenberg 2022; Spergel 2012; Straumann 2020; Tylor 2021).

Eighteen studies were single-center studies (Alexander 2012; Alexander 2017; Bhardwaj 2017; Dellon 2012; Dellon 2019; De Rooij 2022; Dohil 2010; Konikoff 2006; Lieberman 2018; Moawad 2013; Oliva 2018; Peterson 2010; Schaefer 2008; Spergel 2020; Straumann 2010a; Straumann 2010b; Straumann 2011; Straumann 2013).

Participants

The 41 RCTs included 3253 participants.

Eleven studies were in pediatric patients (Assa'ad 2011; Dohil 2010; Gupta 2015; Heine 2019; Kliever 2019; Konikoff 2006; Lieberman 2018; Oliva 2018; Schaefer 2008; Spergel 2012; Spergel 2020), while the rest were in both children and adults.

Four studies were in patients with inactive disease (Alexander 2017; Dellon 2021b; Straumann 2011; Straumann 2020), while the rest were in patients with active disease.

The use of add-on therapies per included study can be found in Table 2.

Interventions

For induction of remission

1. Fourteen studies compared corticosteroids to placebo for induction of remission (Alexander 2012; Bhardwaj 2017; Butz 2014; Dellon 2017; Dellon 2022a; Dohil 2010; Gupta 2015; Hirano 2020f; Hirano 2021; Konikoff 2006; Lucendo 2019; Miehlike 2016; Straumann 2010b; Tylor 2021).
2. Nine studies compared biologics to placebo for induction of remission (Clayton 2014; Dellon 2022; Dellon 2022b; Hirano 2019; Hirano 2020; Rothenberg 2015; Rothenberg 2022; Spergel 2012; Straumann 2010a).
3. One study compared cromolyn sodium to placebo for induction of remission (Lieberman 2018).
4. One study compared PGD2R antagonist OC000459 to placebo for induction of remission (Straumann 2013).
5. One study compared swallowed fluticasone to oral prednisone for induction of remission (Schaefer 2008).
6. One study compared oral viscous budesonide to swallowed fluticasone for induction of remission (Dellon 2019).
7. Two studies compared esomeprazole to fluticasone for induction of remission (Moawad 2013; Peterson 2010).
8. One study compared a one-food elimination diet to a four-food elimination diet for induction of remission (Kliever 2019).
9. One study compared a one-food elimination diet to a six-food elimination diet for induction of remission (Kliever 2021).

10. One study compared a four-food elimination diet with omeprazole to omeprazole for induction of remission (Heine 2019).
11. One study compared a four-food elimination diet with amino acid formula to a four-food elimination diet for induction of remission (De Rooij 2022).
12. One study compared nebulized swallowed budesonide to viscous swallowed budesonide (Dellon 2012).
13. One study compared Viaskin milk patch to placebo (Spergel 2020).
14. One study compared a low dose of the biologic mepolizumab (0.55 mg/kg) to a medium dose (2.5 mg/kg) and to a high dose (10 mg/kg) (Assa'ad 2011).
15. One study compared a six-food elimination diet to swallowed fluticasone to swallowed budesonide and to oral viscous budesonide (Oliva 2018).

The duration of induction RCTs ranged from two weeks (Miehlike 2016; Straumann 2010b) to 44 weeks (Spergel 2020).

For maintenance of remission

1. Three studies compared corticosteroids to placebo for maintenance of remission (Dellon 2021b; Straumann 2011; Straumann 2020).
2. One study compared leukotriene receptor antagonist to placebo for maintenance of remission (Alexander 2017).

The duration of maintenance RCTs ranged from 36 weeks (Dellon 2021b) to 50 weeks (Straumann 2011).

The earliest RCT was published in 2006 (Konikoff 2006), and there has been an exponential growth in RCTs in recent years.

Outcomes

Definitions of dichotomous clinical improvement thresholds

1. Three studies derived a dichotomous clinical outcome from the Mayo Dysphagia Questionnaire (two-week recall; Peloquin 2006). In Alexander 2012 and Alexander 2017, complete symptom response was defined as an answer of "no" to the question, "In the past 2 weeks, have you had trouble swallowing, not associated with other cold symptoms (such as strep throat or mononucleosis)?" on the Mayo dysphagia questionnaire two-week version. A partial symptom response was defined as an answer of "yes" to the earlier-described question and a decrease in the severity of at least two levels (or to a level of "Doesn't bother me at all"), or a decrease in the frequency of at least one level. In Rothenberg 2015, the sum change of Mayo Dysphagia Questionnaire items 1, 2, 4, 9, 10, 13, 14, 16, 20, and 21 from baseline was computed at end of therapy. A positive sum change was scored as an improvement, a negative sum change was scored as a worsening, and a zero-sum change was scored as unchanged.
2. Two studies derived a dichotomous clinical outcome from the Dysphagia Symptom Questionnaire (DSQ; Hudgens 2017). In Dellon 2021b, efficacy was defined as maintenance of the $\geq 30\%$ reduction in DSQ score from baseline that was achieved during the induction phase. Improvement as defined by Dellon 2021b, was scored as a dichotomous event. In Hirano 2021, efficacy was defined as a $\geq 30\%$ reduction in DSQ score from baseline at end of therapy.

Medical treatment of eosinophilic esophagitis (Review)

3. Two studies derived a dichotomous clinical outcome from the Eosinophilic Esophagitis Activity Index (EESAI; [Schoepfer 2014](#)). In [Hirano 2020](#), efficacy was defined as a $\geq 40\%$ improvement in EESAI score from baseline to end of therapy. In [Straumann 2020](#), efficacy was defined as score of ≤ 20 on the EESAI at end of therapy.
 4. One study derived a dichotomous clinical outcome from the Dysphagia Symptom Score (DSS; [Straumann 2010](#)). In [Straumann 2010b](#), frequency and intensity of dysphagia events were scored on a scale of 1 to 4 and 1 to 5, respectively. The Dysphagia Symptom Score was the sum of the two. A clinical response was defined as a decrease in the Dysphagia Symptom Score of at least three points compared with baseline.
 5. One study derived a dichotomous clinical outcome from the EoE Clinical Symptom Score (EoE CSS; [Dohil 2010](#)). In [Gupta 2015](#), efficacy was defined as a $\geq 50\%$ reduction in the EoE CSS from baseline to end of therapy.
 6. One study derived a dichotomous clinical outcome from an esophagus-related symptom score ([Straumann 2003](#)). In [Straumann 2010a](#), clinical efficacy was defined as an improvement of ≥ 1 grade of the esophagus-related symptom score from baseline to end of therapy.
 7. One study derived a dichotomous clinical outcome from a pair of numerical rating scales (0 to 10) that individually assessed dysphagia and odynophagia ([Lucendo 2019](#)). In [Lucendo 2019](#), clinical efficacy was defined as a score of ≤ 2 on each rating scale on each day of the week before end of therapy.
 8. One study derived a dichotomous clinical outcome from the frequency of dysphagia and/or heartburn ([Bhardwaj 2017](#)). In [Bhardwaj 2017](#), clinical efficacy was defined as a decrease in the frequency of dysphagia and/or heartburn from baseline to the end of therapy.
 9. One study derived a dichotomous clinical outcome from the presence or absence of the presenting symptoms by patient/guardian report and by physician assessment ([Schaefer 2008](#)). In [Schaefer 2008](#), clinical efficacy was defined as the absence of presenting symptoms at end of therapy.
 10. One study derived a dichotomous clinical outcome from a Physician's Eosinophilic Esophagitis Global Assessment Score (PGA; [Schoepfer 2014](#)). In [Spergel 2012](#), clinical efficacy was defined as an improvement of ≥ 1 level on the Physician's Eosinophilic Esophagitis Global Assessment Score from baseline to end of therapy.
 11. Twenty-seven studies did not define a dichotomous clinical outcome or did not publish data for a dichotomous clinical outcome in a form that could be used by this review ([Assa'ad 2011](#); [Butz 2014](#); [Clayton 2014](#); [Dellon 2012](#); [Dellon 2017](#); [Dellon 2019](#); [Dellon 2022](#); [Dellon 2022a](#); [Dellon 2022b](#); [De Rooij 2022](#); [Dohil 2010](#); [Heine 2019](#); [Hirano 2019](#); [Hirano 2020f](#); [Kliewer 2019](#); [Kliewer 2021](#); [Konikoff 2006](#); [Lieberman 2018](#); [Miehlke 2016](#); [Moawad 2013](#); [Oliva 2018](#); [Peterson 2010](#); [Rothenberg 2022](#); [Spergel 2020](#); [Straumann 2011](#); [Straumann 2013](#); [Tytor 2021](#)).
- Scales used for clinical improvement continuous measurement**
1. Six studies reported a continuous clinical outcome based on the Dysphagia Symptom Questionnaire (DSQ; [Hudgens 2017](#)). [Dellon 2019](#) and [Hirano 2021](#) reported mean Dysphagia Symptom Questionnaire scores at end of therapy. [Dellon 2017](#), [Dellon 2022](#), [Dellon 2021b](#), and [Rothenberg 2022](#) reported mean change in Dysphagia Symptom Questionnaire score from baseline to end of therapy.
 2. Five studies reported a continuous clinical outcome based on the Straumann Dysphagia Instrument (SDI; [Straumann 2010](#)). [Straumann 2010b](#) and [Straumann 2011](#) reported the mean Straumann Dysphagia Instrument score at end of therapy. [Miehlke 2016](#), [Hirano 2020](#), and [De Rooij 2022](#) reported the mean change in Straumann Dysphagia Instrument score from baseline to end of therapy.
 3. Two studies reported a continuous clinical outcome based on the Pediatric Eosinophilic Esophagitis Symptom Score version 2.0 (PEESS V2.0; [Franciosi 2011](#)). [Kliewer 2019](#) and [Spergel 2020](#) reported mean Pediatric Eosinophilic Esophagitis Symptom Scores version 2.0 at end of therapy.
 4. Two studies reported a continuous clinical outcome based on the Mayo Dysphagia Questionnaire (two-week recall; [Peloquin 2006](#)). [Moawad 2013](#) and [Rothenberg 2015](#) reported mean Mayo Dysphagia Questionnaire (two-week recall) scores at end of therapy.
 5. One study reported a continuous clinical outcome based on the Pediatric Eosinophilic Esophagitis Symptom Score version 1.0 (PEESS V1.0; [Pentiuk 2009](#)). [Lieberman 2018](#) reported mean Pediatric Eosinophilic Esophagitis Symptom Scores version 1.0 at end of therapy.
 6. Two studies reported a continuous clinical outcome based on the Eosinophilic Esophagitis Activity Index (EESAI; [Schoepfer 2014](#)). [Kliewer 2021](#) and [Straumann 2020](#) reported mean Eosinophilic Esophagitis Activity Index scores at end of therapy.
 7. One study reported a continuous clinical outcome based on the proportion of days that participants reported difficulty in swallowing averaged over the seven days prior to the clinic visit ([Straumann 2010a](#)). [Straumann 2010a](#) reported the mean proportion of dysphagia-free days in the week prior to the last clinic visit at end of therapy.
 8. One study reported a continuous clinical outcome based on the Dysphagia Scale ([DiSario 2002](#)). [Peterson 2010](#) reported mean Dysphagia Scale score at end of therapy.
 9. One study reported a continuous clinical outcome based on the Watson Dysphagia Score (WDS; [Dakkak 1992](#)). [Tytor 2021](#) reported mean change in Watson Dysphagia Score from baseline to end of therapy.
 10. One study reported a mean continuous clinical outcome, at end of therapy, based on the following criteria: 0 = no dysphagia; 1 = solid food dysphagia monthly; 2 = solid food dysphagia < weekly; 3 = solid food dysphagia > weekly and < daily; 4 = solid food dysphagia daily; 5 = solid food dysphagia with every meal; and 6 = dysphagia for solid and liquid food ([Clayton 2014](#)).
 11. One study reported a continuous clinical outcome based on the Mayo Dysphagia Questionnaire (30-day recall; [McElhiney 2009](#)). [Dellon 2012](#) reported mean Mayo Dysphagia Questionnaire (30-day recall) scores at end of therapy.
 12. One study reported a continuous clinical outcome based on the sum of scores from a visual dysphagia questionnaire (VDQ; [Straumann 2013](#)) and chest pain as recorded by the Straumann Dysphagia Instrument (SDI; [Straumann 2010](#)). [Straumann 2013](#) reported mean composite visual dysphagia questionnaire/ Straumann Dysphagia Instrument chest pain scores at end of therapy.

Medical treatment of eosinophilic esophagitis (Review)

13. One study reported a continuous clinical outcome based on a symptom scoring tool (Aceves 2009). Dohil 2010 reported the mean symptom score at end of therapy.
14. One study reported a continuous clinical outcome based on Daily Dysphagia Symptom Diary (DSD) scores (DSD; Hirano 2019). Hirano 2019 reported mean change in Daily Dysphagia Symptom Diary scores from baseline to end of therapy.
15. Fifteen studies did not report a continuous clinical outcome or did not publish data for a clinical continuous outcome in a form that could be used by this review (Alexander 2012; Alexander 2017; Assa'ad 2011; Bhardwaj 2017; Butz 2014; Dellon 2022a; Dellon 2022b; Gupta 2015; Heine 2019; Hirano 2020f; Konikoff 2006; Lucendo 2019; Oliva 2018; Schaefer 2008; Spergel 2012).

Definitions of dichotomous histological improvement thresholds

1. Three studies reported a dichotomous histological threshold of < 20 mean peak eos/hpf at end of therapy (Dohil 2010; Straumann 2010b; Straumann 2011).
2. Fifteen studies reported a dichotomous histological threshold of < 15 mean peak eos/hpf at end of therapy (Butz 2014; Dellon 2012; Dellon 2019; Dellon 2022; Dellon 2021b; De Rooij 2022; Hirano 2019; Hirano 2020; Hirano 2020f; Hirano 2021; Kliewer 2019; Kliewer 2021; Peterson 2010; Spergel 2012; Straumann 2020).
3. One study reported a dichotomous histological threshold of ≤ 7 mean peak eos/hpf at end of therapy (Moawad 2013).
4. Eight studies reported a dichotomous histological threshold of ≤ 6 mean peak eos/hpf at end of therapy (Assa'ad 2011; Dellon 2017; Dellon 2022a; Dellon 2022b; Gupta 2015; Konikoff 2006; Rothenberg 2022; Straumann 2010a).
5. Two studies reported a dichotomous histological threshold of < 5 mean peak eos/hpf at end of therapy (Lucendo 2019; Miehlike 2016).
6. One study defined a dichotomous complete histologic response as a decrease in the mean eosinophil level of 90% from baseline to end of therapy. A partial response was defined as a decrease of more than 50% from baseline to end of therapy (Alexander 2012).
7. One study defined dichotomous histologic response as a decrease in mean peak eos/hpf of 75% from baseline to end of therapy (Rothenberg 2015).
8. One study defined dichotomous complete histologic response as "normal biopsy specimens" at end of therapy (Schaefer 2008).
9. Nine studies did not report a dichotomous histologic response or did not publish data for a dichotomous histologic outcome in a form that could be used by this review (Alexander 2017; Bhardwaj 2017; Clayton 2014; Heine 2019; Lieberman 2018; Oliva 2018; Spergel 2020; Straumann 2013; Tytor 2021).

Scales used for histological improvement continuous measurement

1. Twenty-three studies reported continuous histologic outcomes based on mean peak counts of eosinophils per high-powered microscope field. Six studies reported a continuous histologic outcome as the change in mean peak eos/hpf from baseline to end of therapy (Dellon 2017; De Rooij 2022; Hirano 2019; Hirano 2020; Kliewer 2021; Straumann 2020). Seventeen studies reported a continuous histologic outcome as mean peak eos/hpf at end of therapy (Bhardwaj 2017; Clayton 2014; Dellon 2012; Dellon 2019; Dohil 2010; Hirano 2021; Lieberman 2018;

- Moawad 2013; Peterson 2010; Rothenberg 2015; Schaefer 2008; Spergel 2012; Spergel 2020; Straumann 2010a; Straumann 2010b; Straumann 2011; Straumann 2013).
2. Eighteen studies did not report a continuous histologic response or did not publish data for a continuous histologic outcome in a form that could be used by this review (Alexander 2012; Alexander 2017; Assa'ad 2011; Butz 2014; Dellon 2022; Dellon 2021b; Dellon 2022a; Dellon 2022b; Gupta 2015; Heine 2019; Hirano 2020f; Kliewer 2019; Konikoff 2006; Lucendo 2019; Miehlike 2016; Oliva 2018; Rothenberg 2022; Tytor 2021).

Definitions of dichotomous endoscopic improvement thresholds

1. Two studies defined a dichotomous endoscopic outcome from histologic grading (Schaefer 2008). In Schaefer 2008 and Straumann 2010a, improvement of ≥ 1 histologic grade was scored as a dichotomous endoscopic event.
2. One study derived a dichotomous endoscopic outcome from the Endoscopic Reference Score (EREFS; Hirano 2013). In Hirano 2020f, efficacy was defined as the sign of the change in EREFS score from baseline to end of therapy.
3. One study derived a dichotomous endoscopic outcome from esophageal furrows (Konikoff 2006). In Konikoff 2006, the lack of esophageal furrows at end of therapy was scored as a dichotomous endoscopic event.
4. One study defined a dichotomous endoscopic outcome from endoscopic findings (Alexander 2012). In Alexander 2012, resolution of all endoscopic findings was scored as a dichotomous endoscopic event.
5. Thirty-five studies did not report a dichotomous endoscopic response or did not publish data for a dichotomous endoscopic outcome in a form that could be used by this review (Alexander 2017; Assa'ad 2011; Bhardwaj 2017; Butz 2014; Clayton 2014; Dellon 2012; Dellon 2017; Dellon 2019; Dellon 2022; Dellon 2021b; Dellon 2022a; Dellon 2022b; Dohil 2010; Gupta 2015; Heine 2019; Hirano 2019; Hirano 2020; Hirano 2021; Kliewer 2019; Kliewer 2021; Lieberman 2018; Lucendo 2019; Miehlike 2016; Moawad 2013; Oliva 2018; Peterson 2010; Rothenberg 2015; Rothenberg 2022; Spergel 2012; Spergel 2020; Straumann 2010b; Straumann 2011; Straumann 2013; Straumann 2020; Tytor 2021).

Scales used for endoscopic improvement continuous measurement

1. Thirteen studies reported continuous endoscopic outcomes based on the Endoscopic Reference Score (EREFS; Hirano 2013). Eight studies reported the mean change in Endoscopic Reference Score from baseline to end of therapy (Dellon 2017; Dellon 2022; Dellon 2021b; Dellon 2022a; Hirano 2020; Kliewer 2019; Kliewer 2021; Spergel 2020). Five studies reported mean Endoscopic Reference Score at end of therapy (Dellon 2019; Hirano 2019; Hirano 2021; Lucendo 2019; Straumann 2020).
2. One study reported continuous endoscopic outcomes based on an Endoscopy Scoring Tool (EST, Aceves 2009). In Dohil 2010, the mean score from the Endoscopy Scoring Tool was reported at end of therapy.
3. One study reported continuous endoscopic outcomes based on endoscopic findings (Straumann 2013). In Straumann 2013, the mean endoscopic findings score was reported at end of therapy.
4. Twenty-five studies did not report a continuous endoscopic response or did not publish data for a continuous endoscopic outcome in a form that could be used by this review (Alexander

Medical treatment of eosinophilic esophagitis (Review)

2017; Assa'ad 2011; Bhardwaj 2017; Butz 2014; Clayton 2014; Dellon 2012; Dellon 2022b; De Rooij 2022; Gupta 2015; Heine 2019; Hirano 2020f; Konikoff 2006; Lieberman 2018; Miehlke 2016; Moawad 2013; Oliva 2018; Peterson 2010; Rothenberg 2015; Rothenberg 2022; Schaefer 2008; Spergel 2012; Straumann 2010a; Straumann 2010b; Straumann 2011; Tytor 2021).

Withdrawals due to adverse events

1. Information about withdrawals due to adverse events was reported for 39 studies (Alexander 2012; Alexander 2017; Assa'ad 2011; Bhardwaj 2017; Butz 2014; Clayton 2014; Dellon 2012; Dellon 2017; Dellon 2019; Dellon 2022; Dellon 2021b; Dellon 2022a; Dellon 2022b; De Rooij 2022; Dohil 2010; Gupta 2015; Heine 2019; Hirano 2019; Hirano 2020; Hirano 2020f; Hirano 2021; Kliewer 2019; Kliewer 2021; Konikoff 2006; Lieberman 2018; Lucendo 2019; Miehlke 2016; Moawad 2013; Peterson 2010; Rothenberg 2015; Schaefer 2008; Spergel 2012; Spergel 2020; Straumann 2010a; Straumann 2010b; Straumann 2011; Straumann 2013; Straumann 2020; Tytor 2021).
2. Two studies did not report information about withdrawals due to adverse events or did not publish data for withdrawals due to adverse events in a form that could be used by this review (Oliva 2018; Rothenberg 2022).

Serious adverse events

1. Thirty-seven studies reported information about serious adverse events (as defined by the study) (Alexander 2012; Alexander 2017; Assa'ad 2011; Bhardwaj 2017; Butz 2014; Clayton 2014; Dellon 2012; Dellon 2017; Dellon 2019; Dellon 2022; Dellon 2021b; Dellon 2022a; Dellon 2022b; De Rooij 2022; Dohil 2010; Gupta 2015; Hirano 2019; Hirano 2020; Hirano 2020f; Hirano 2021; Kliewer 2019; Kliewer 2021; Konikoff 2006; Lieberman 2018; Lucendo 2019; Miehlke 2016; Moawad 2013; Peterson 2010; Rothenberg 2015; Schaefer 2008; Spergel 2012; Spergel 2020; Straumann 2010a; Straumann 2010b; Straumann 2011; Straumann 2013; Straumann 2020; Tytor 2021).
2. Four studies did not report information about serious adverse events (as defined by the study) or did not publish data for serious adverse events in a form that could be used by this review (Dellon 2022a; Heine 2019; Oliva 2018; Rothenberg 2022).

Total adverse events

1. Thirty-three studies reported information about adverse events (as defined by the study) (Alexander 2012; Alexander 2017; Assa'ad 2011; Butz 2014; Clayton 2014; Dellon 2012; Dellon 2017; Dellon 2019; Dellon 2021b; Dellon 2022a; Dellon 2022b; De Rooij 2022; Dohil 2010; Gupta 2015; Hirano 2019; Hirano 2020; Hirano 2020f; Hirano 2021; Kliewer 2019; Kliewer 2021; Konikoff 2006; Lucendo 2019; Miehlke 2016; Moawad 2013; Peterson 2010; Rothenberg 2015; Spergel 2012; Spergel 2020; Straumann 2010a; Straumann 2010b; Straumann 2011; Straumann 2020; Tytor 2021).
2. Eight studies did not report information about adverse events (as defined by the study) or did not publish data for adverse events in a form that could be used by this review (Bhardwaj 2017; Dellon 2022; Heine 2019; Lieberman 2018; Oliva 2018; Rothenberg 2022; Schaefer 2008; Straumann 2013).

Quality of life

1. Four studies reported continuous quality of life outcomes based on the Adult Eosinophilic Esophagitis Quality of Life questionnaire (EoE QoL-A; Taft 2011). Lucendo 2019 and Straumann 2020 reported mean Adult Eosinophilic Esophagitis Quality of Life questionnaire scores at end of therapy. Hirano 2020 and Kliewer 2021 reported mean change in Adult Eosinophilic Esophagitis Quality of Life questionnaire scores from baseline to end of therapy.
2. One study reported continuous quality of life outcomes based on the Pediatric Quality of Life 4.0, Eosinophilic Esophagitis Module (PedsQL 4.0 EoE; Franciosi 2013). Kliewer 2019 reported mean change in Pediatric Quality of Life 4.0, Eosinophilic Esophagitis from baseline to end of therapy.
3. Thirty-six studies did not report a continuous quality of life score or did not publish data for quality of life in a form that could be used by this review (Alexander 2012; Alexander 2017; Assa'ad 2011; Bhardwaj 2017; Butz 2014; Clayton 2014; Dellon 2012; Dellon 2017; Dellon 2019; Dellon 2022; Dellon 2021b; Dellon 2022a; Dellon 2022b; De Rooij 2022; Dohil 2010; Gupta 2015; Heine 2019; Hirano 2019; Hirano 2020f; Hirano 2021; Konikoff 2006; Lieberman 2018; Miehlke 2016; Moawad 2013; Oliva 2018; Peterson 2010; Rothenberg 2015; Rothenberg 2022; Schaefer 2008; Spergel 2012; Spergel 2020; Straumann 2010a; Straumann 2010b; Straumann 2011; Straumann 2013; Tytor 2021).

Contact with authors

We contacted authors of 35 studies with requests for data and clarification where risk of bias was unclear (Alexander 2012; Alexander 2017; Assa'ad 2011; Bhardwaj 2017; Butz 2014; Clayton 2014; Dellon 2012; Dellon 2017; Dellon 2019; Dellon 2021b; Dellon 2022; Dellon 2022a; Dellon 2022b; De Rooij 2022; Dohil 2010; Gupta 2015; Heine 2019; Hirano 2019; Hirano 2020; Hirano 2020f; Hirano 2021; Kliewer 2019; Kliewer 2021; Lieberman 2018; Miehlke 2016; Moawad 2013; Oliva 2018; Peterson 2010; Rothenberg 2015; Rothenberg 2022; Spergel 2012; Spergel 2020; Straumann 2010b; Straumann 2011; Straumann 2013; Straumann 2020; Tytor 2021). We received responses from all except 11 (Assa'ad 2011; Bhardwaj 2017; Clayton 2014; De Rooij 2022; Heine 2019; Hirano 2020f; Hirano 2021; Oliva 2018; Straumann 2010b; Straumann 2011; Straumann 2013).

Funding sources and conflicts of interest

Twenty-six studies received funding from pharmaceutical companies (Alexander 2017; Assa'ad 2011; Clayton 2014; Dellon 2012; Dellon 2017; Dellon 2022; Dellon 2021b; Dellon 2022a; Dellon 2022b; Dohil 2010; Gupta 2015; Hirano 2019; Hirano 2020; Hirano 2020f; Hirano 2021; Lucendo 2019; Miehlke 2016; Rothenberg 2015; Rothenberg 2022; Spergel 2012; Spergel 2020; Straumann 2010a; Straumann 2010b; Straumann 2011; Straumann 2013; Straumann 2020).

Thirteen studies received funding from universities, foundations, medical associations or research institutions, and no funding from pharmaceutical companies (Alexander 2012; Bhardwaj 2017; Butz 2014; Dellon 2019; Heine 2019; Kliewer 2019; Kliewer 2021; Konikoff 2006; Lieberman 2018; Moawad 2013; Peterson 2010; Schaefer 2008; Tytor 2021).

All studies that reported conflicts of interest had authors with conflicts of interest, except three (Bhardwaj 2017; Clayton 2014, Moawad 2013).

Medical treatment of eosinophilic esophagitis (Review)

Two studies did not report on their funding or conflicts of interest (De Rooij 2022; Oliva 2018). Six studies did not report on conflicts of interest (Heine 2019; Kliewer 2019; Kliewer 2021; Konikoff 2006; Peterson 2010; Schaefer 2008).

More details about the funding and conflicts of interest of the included studies can be found in the [Characteristics of included studies](#) tables.

Table 2 is a summary of key characteristics of the included studies.

Excluded studies

We excluded 39 studies (45 records) for the reasons presented in the [Characteristics of excluded studies](#) table, summarized below.

- Twenty-eight studies were excluded due to the wrong study design (Ceves 2005; Della 2017; Eluri 2017; Eluri 2017a; EUCTR2014-002465-30-IT 2014; Francis 2012; Helou 2008; Hudgens 2017; JPRN-UMIN000021041 2016; JPRN-UMIN000026704 2017; Kagalwalla 2006; Kruszewski 2016;

Kuzumoto 2021; Molina-Infante 2017; NCT01498497 2011; NTR4892 2014; Safroneeva 2015; Safroneeva 2018; Safroneeva 2018a; Savarino 2015; Song 2020; Spengel 2002; Spengel 2005; Syverson 2020; Vazquez-Elizondo 2013; Wang 2017; Warners 2016; Wechsler 2017)

- Seven studies were excluded due to the wrong population (Braathen 2006; Comer 2017; Dellon 2020c; Hefner 2016; Tripp 2017; Wright 2020; Wright 2021).
- One study was excluded for the wrong intervention (Kavitt 2016).
- Three studies were abandoned RCTs without results (NCT01458418 2011; NCT01702701 2012; NCT01821898 2013).

Risk of bias in included studies

The results of our risk of bias assessments are presented in [Figure 2](#) and the risk of bias tables in the [Characteristics of included studies](#) table. We conducted our initial assessment using the information presented in the published papers. In studies where the risk of bias assessment was unclear, we sought clarification from at least one author or contact person (or both) per study. Where we received responses, we adapted our initial assessment accordingly.

Figure 2. Risk of bias summary: review authors' judgments about each risk of bias item for each included study.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias): All outcomes	Blinding of outcome assessment (detection bias): All outcomes	Incomplete outcome data (attrition bias): All outcomes	Selective reporting (reporting bias)	Other bias
Alexander 2012	+	+	+	+	+	+	+
Alexander 2017	+	+	+	+	+	?	+
Assa'ad 2011	?	+	+	?	+	-	+
Bhardwaj 2017	+	+	+	+	?	?	+
Butz 2014	+	+	+	+	+	?	+
Clayton 2014	+	+	?	+	+	?	+
Dellon 2012	+	+	-	+	+	+	+
Dellon 2017	+	+	+	+	+	+	+
Dellon 2019	+	+	+	+	+	+	+
Dellon 2021b	+	+	+	+	+	+	+
Dellon 2022	+	+	+	+	+	+	+
Dellon 2022a	+	+	+	+	+	+	+
Dellon 2022b	+	+	+	+	+	+	+
De Rooij 2022	?	+	?	?	+	+	+
Dohil 2010	+	+	?	+	+	+	+
Gupta 2015	+	+	+	+	+	+	+
Heine 2019	+	+	-	-	?	-	+

Figure 2. (Continued)

Heine 2019	+	+	-	-	?	-	+
Hirano 2019	+	+	+	+	+	+	+
Hirano 2020	+	+	+	+	+	+	+
Hirano 2020f	+	?	+	+	+	?	+
Hirano 2021	+	?	+	+	+	+	+
Kliewer 2019	+	+	-	+	?	+	+
Kliewer 2021	+	+	-	+	+	+	?
Konikoff 2006	+	+	+	+	+	+	+
Lieberman 2018	+	+	+	+	+	+	+
Lucendo 2019	+	+	+	+	+	+	+
Miehlke 2016	+	+	+	+	+	?	+
Moawad 2013	+	+	-	+	+	+	+
Oliva 2018	?	?	-	-	?	?	?
Peterson 2010	+	+	-	-	+	-	+
Rothenberg 2015	+	+	+	+	+	+	+
Rothenberg 2022	+	+	+	+	+	?	+
Schaefer 2008	+	+	-	+	+	+	+
Spergel 2012	+	+	+	+	+	+	+
Spergel 2020	+	+	+	+	+	+	+
Straumann 2010a	+	+	+	+	+	+	+
Straumann 2010b	+	?	+	?	+	+	+
Straumann 2011	+	?	+	?	+	+	+
Straumann 2013	+	?	+	?	+	+	+
Straumann 2020	+	+	+	+	+	+	+
Tytor 2021	+	+	+	+	+	+	+

Allocation

Three studies did not sufficiently describe randomization and therefore were at unclear risk of bias (Assa'ad 2011; De Rooij 2022; Oliva 2018). We rated all other studies at low risk of bias for randomization.

We rated six studies at unclear risk of bias for allocation concealment, as they did not provide enough information about their selection and allocation concealment processes (Hirano 2020f; Hirano 2021; Oliva 2018; Straumann 2010b; Straumann 2011; Straumann 2013). We rated all other studies at low risk of bias for allocation concealment.

Blinding

We rated eight studies at high risk of performance bias for not blinding participants and/or personnel (Dellon 2012; Heine 2019; Kliewer 2019; Kliewer 2021; Moawad 2013; Oliva 2018; Peterson

2010; Schaefer 2008), and three studies at unclear risk, as not enough information was available (Clayton 2014; De Rooij 2022; Dohil 2010). We rated all other studies at low risk for performance bias.

We rated detection bias as high risk in three studies, for not blinding outcome assessors (Heine 2019; Oliva 2018; Peterson 2010), and five studies at unclear risk (Assa'ad 2011; De Rooij 2022; Straumann 2010b; Straumann 2011; Straumann 2013). We rated all other studies at low risk for detection bias.

Incomplete outcome data

We rated attrition bias unclear in four studies where it was not possible to judge whether incomplete outcome data had affected outcomes (Bhardwaj 2017; Heine 2019; Kliewer 2019; Oliva 2018). We rated all other studies at low risk for attrition bias.

Medical treatment of eosinophilic esophagitis (Review)

Selective reporting

We rated the risk of bias for selective reporting as high in three studies (Assa'ad 2011; Heine 2019; Peterson 2010), and as unclear in eight studies (Alexander 2017; Bhardwaj 2017; Butz 2014; Clayton 2014; Hirano 2020f; Miehlke 2016; Oliva 2018; Rothenberg 2022). We rated all other studies at low risk of reporting bias.

Other potential sources of bias

We rated two studies as unclear for other potential sources of bias (Kliwer 2021; Oliva 2018). We rated all other studies at low risk of other bias.

Effects of interventions

See: **Summary of findings 1** Corticosteroids compared to placebo for induction of remission; **Summary of findings 2** Corticosteroids compared to placebo for maintenance of remission; **Summary of findings 3** Biologics compared to placebo for induction of remission; **Summary of findings 4** Cromolyn sodium compared to placebo; **Summary of findings 5** PGD2R antagonist OC000459 compared to placebo; **Summary of findings 6** Swallowed fluticasone compared to oral prednisone; **Summary of findings 7** Oral viscous budesonide compared to swallowed fluticasone; **Summary of findings 8** Esomeprazole compared to fluticasone; **Summary of findings 9** One-food elimination diet compared to four-food elimination diet; **Summary of findings 10** One-food elimination diet compared to six-food elimination diet; **Summary of findings 11** Four-food elimination diet with omeprazole compared to omeprazole; **Summary of findings 12** Four-food elimination and amino acid formula compared to four-food elimination diet; **Summary of findings 13** Nebulized budesonide compared to viscous budesonide; **Summary of findings 14** Viaskin milk patch compared to placebo; **Summary of findings 15** Leukotriene receptor antagonist compared to placebo for maintenance of remission; **Summary of findings 16** Mepolizumab 10 mg/kg compared to mepolizumab 0.55 mg/kg; **Summary of findings 17** Mepolizumab 2.5 mg/kg compared to mepolizumab 0.55 mg/kg; **Summary of findings 18** Mepolizumab 10 mg/kg compared to mepolizumab 2.5 mg/kg; **Summary of findings 19** Six-food elimination diet compared to swallowed fluticasone compared to swallowed budesonide compared to oral viscous budesonide

All outcome data used in our analyses can be found in Table 3; Table 4; Table 1; Table 5; Table 6.

Corticosteroids versus placebo for induction of remission

Fourteen studies compared corticosteroids to placebo for induction of remission (Alexander 2012; Bhardwaj 2017; Butz 2014; Dellon 2017; Dellon 2022a; Dohil 2010; Gupta 2015; Hirano 2020f; Hirano 2021; Konikoff 2006; Lucendo 2019; Miehlke 2016; Straumann 2010b; Tytor 2021).

Primary outcomes

Clinical improvement

Six studies compared corticosteroids to placebo for clinical improvement as a dichotomous outcome (Alexander 2012; Bhardwaj 2017; Gupta 2015; Hirano 2021; Lucendo 2019; Straumann 2010b).

Corticosteroids (n = 210/380) may lead to slightly better clinical improvement compared to placebo (n = 71/203), measured as a dichotomous outcome (risk ratio (RR) 1.74, 95% confidence interval (CI) 1.08 to 2.80). The results are of low certainty due to inconsistency and imprecision (Analysis 1.1; Summary of findings 1).

Sensitivity analyses using a fixed-effect model (RR 1.54, 95% CI 1.25 to 1.89; Analysis 1.2), and for validated instruments (RR 1.39, 95% CI 1.08 to 1.79; Analysis 1.3), showed similar results.

The subgroup analyses provided only limited explanation for the variation in treatment effect across the studies. While there was a statistically significant difference in the interaction test for subgroup analysis by age (Analysis 1.4), this was based on only one study in children younger than 18 years old (Gupta 2015), and five studies in mixed adult and child participants (Alexander 2012; Bhardwaj 2017; Hirano 2021; Lucendo 2019; Straumann 2010b). The remaining subgroup analyses were based on type of corticosteroid (one beclomethasone study (Bhardwaj 2017), four budesonide studies (Gupta 2015; Hirano 2021; Lucendo 2019; Straumann 2010b), and one fluticasone study (Alexander 2012)), and based on corticosteroid delivery method (three studies using an adapted asthma delivery method (Alexander 2012; Bhardwaj 2017; Straumann 2010b), and three studies using an esophageal-specific method (Gupta 2015; Hirano 2021; Lucendo 2019)). There was insufficient evidence in these subgroup analyses to determine whether there were any differences in the subgroup effects (see Analysis 1.5 and Analysis 1.6).

Five studies compared corticosteroids to placebo for clinical improvement as a continuous outcome (Dellon 2017; Dohil 2010; Hirano 2021; Straumann 2010b; Tytor 2021).

Corticosteroids (n = 302) may lead to slightly better clinical improvement compared to placebo (n = 173), measured as a continuous outcome (standardized mean difference (SMD) 0.51, 95% CI 0.17 to 0.85). The results are of low certainty due to inconsistency and imprecision (Analysis 1.7; Summary of findings 1).

Sensitivity analyses using a fixed-effect model (SMD 0.37, 95% CI 0.18 to 0.56; Analysis 1.8), and for validated instruments (SMD 0.35, 95% CI 0.07 to 0.64; Analysis 1.9), showed similar results.

Again, the subgroup analyses provided limited explanation for the variation in treatment effect. The test for subgroup differences by age did not show a statistical difference (Analysis 1.10), however it was only based on one study in children younger than 18 years old (Dohil 2010), and four studies on mixed adult and child participants (Dellon 2017; Hirano 2021; Straumann 2010b; Tytor 2021). Similarly, we observed no difference for the type of corticosteroid (Analysis 1.11), based on four studies of budesonide (Dellon 2017; Dohil 2010; Hirano 2021; Straumann 2010b), and one of mometasone (Tytor 2021). However, we observed a difference between subgroups for corticosteroid delivery method (Analysis 1.12), based on one study using an adapted asthma delivery method (Straumann 2010b), and four using an esophageal-specific method (Dellon 2017; Dohil 2010; Hirano 2021; Tytor 2021).

Histological improvement

Twelve studies compared corticosteroids to placebo for histological improvement as a dichotomous outcome (Alexander 2012; Butz

Medical treatment of eosinophilic esophagitis (Review)


2014; Dellon 2017; Dellon 2022a; Dohil 2010; Gupta 2015; Hirano 2020f; Hirano 2021; Konikoff 2006; Lucendo 2019; Miehlike 2016; Straumann 2010b).

Corticosteroids (n = 428/652) lead to a large histological improvement compared to placebo (n = 10/326), measured as a

dichotomous outcome (RR 11.94, 95% CI 6.56 to 21.75, NNTB = 3). These results are of high certainty (Analysis 1.13; Summary of findings 1).

Funnel plot inspection did not reveal publication bias (Figure 3).

Figure 3.



A sensitivity analysis using a fixed-effect model showed similar results but a higher magnitude of effect (RR 18.87, 95% CI 10.57 to 33.71; Analysis 1.14).

Sensitivity analyses for histological thresholds of < 15 eos/hpf (RR 18.47, 95% CI 4.45 to 76.72; Analysis 1.15), ≤ 6 eos/hpf (RR 14.03, 95% CI 6.73 to 29.26; Analysis 1.16), and ≤ 1 eos/hpf (RR 10.97, 95% CI 3.12 to 38.55; Analysis 1.17) showed similar results.

A subgroup analysis for participant age, based on three studies in children younger than 18 years old (Dohil 2010; Gupta 2015; Konikoff 2006), and eight studies in mixed adult and child participants did not reveal differences (Alexander 2012; Butz 2014; Dellon 2017; Dellon 2022a; Hirano 2021; Lucendo 2019; Miehlike 2016; Straumann 2010b) (Analysis 1.18).

A subgroup analysis for the type of corticosteroid, with seven studies of budesonide (Dellon 2017; Dohil 2010; Gupta 2015; Hirano 2021; Lucendo 2019; Miehlike 2016; Straumann 2010b), and four studies of fluticasone (Alexander 2012; Butz 2014; Dellon 2022a; Konikoff 2006), did not reveal differences (Analysis 1.19).

A subgroup analysis for corticosteroid delivery method, based on seven studies using an esophageal-specific method (Dellon 2017; Dellon 2022a; Dohil 2010; Gupta 2015; Hirano 2021; Lucendo 2019; Miehlike 2016), and four using an adapted asthma delivery method (Alexander 2012; Butz 2014; Konikoff 2006; Straumann 2010b), did not reveal differences (Analysis 1.20).

Five studies compared corticosteroids to placebo for histological improvement as a continuous outcome (Bhardwaj 2017; Dellon 2017; Dohil 2010; Hirano 2021; Straumann 2010b).

Corticosteroids (n = 287) may lead to histological improvement compared to placebo (n = 162), measured as a continuous outcome (SMD 1.42, 95% CI 1.02 to 1.82). The results are of low certainty, due to inconsistency and risk of bias (Analysis 1.21; Summary of findings 1).

A sensitivity analysis using a fixed-effect model showed similar results (SMD 1.33, 95% CI 1.12 to 1.55; Analysis 1.22).

Little insight could be gained from the subgroup analyses. We observed no differences between subgroups for the following: participant age (Analysis 1.23), based on only one study in children

younger than 18 years old (Dohil 2010), and four studies in mixed adult and child participants (Bhardwaj 2017; Dellon 2017; Hirano 2021; Straumann 2010b); the type of steroid (Analysis 1.24), based on only one study of beclomethasone (Bhardwaj 2017), and four studies of budesonide (Dellon 2017; Dohil 2010; Hirano 2021; Straumann 2010b); and corticosteroid delivery method (Analysis 1.25), based on two studies using an adapted asthma delivery method (Bhardwaj 2017; Straumann 2010b), and three studies using an esophageal-specific method (Dellon 2017; Dohil 2010; Hirano 2021).

Endoscopic improvement

Three studies compared corticosteroids to placebo for endoscopic improvement as a dichotomous outcome (Alexander 2012; Hirano 2020f; Konikoff 2006).

Corticosteroids ($n = 25/58$) may lead to little to no endoscopic improvement compared to placebo ($n = 6/44$), measured as a dichotomous outcome (RR 2.60, 95% CI 0.82 to 8.19). The results are of low certainty due to serious imprecision (Analysis 1.26; Summary of findings 1).

A sensitivity analysis using a fixed-effect model showed similar results (RR 2.73, 95% CI 1.27 to 5.86; Analysis 1.27).

We cannot draw any conclusions from a sensitivity analysis based on validated instruments (RR 5.87, 95% CI 1.11 to 31.02; Analysis 1.28). The results are of very low certainty due to serious imprecision and risk of bias.

Subgroup analyses for participant age (Analysis 1.29), and corticosteroid delivery method (Analysis 1.30), provided limited explanation for the variation in treatment effect. They were based on only one study in children younger than 18 years old (Konikoff 2006), and two studies in mixed adult and child participants (Alexander 2012; Hirano 2020f), and two studies using an adapted asthma delivery method (Alexander 2012; Konikoff 2006), and one study using an esophageal-specific method (Hirano 2020f).

Five studies compared corticosteroids to placebo for endoscopic improvement as a continuous outcome (Dellon 2017; Dellon 2022a; Dohil 2010; Hirano 2021; Lucendo 2019).

Corticosteroids ($n = 409$) may lead to endoscopic improvement compared to placebo ($n = 187$), measured as a continuous outcome (SMD 1.33, 95% CI 0.59 to 2.08). The results are of low certainty due to serious inconsistency (Analysis 1.31; Summary of findings 1).

A sensitivity analysis using a fixed-effect model showed similar results (SMD 0.93, 95% CI 0.74 to 1.11; Analysis 1.32).

We cannot draw any conclusions from a sensitivity analysis based on validated instruments (SMD 1.31, 95% CI 0.46 to 2.17; Analysis 1.33).

A subgroup analysis for participant age, based on one study in children younger than 18 years old (Dohil 2010), and four studies in mixed adult and child participants (Dellon 2017; Dellon 2022a; Hirano 2021; Lucendo 2019), revealed no differences between subgroups and there was no statistical difference from placebo in both groups. This was based on limited data for children (Analysis 1.34).

A subgroup analysis for type of steroid, based on four studies of budesonide (Dellon 2017; Dohil 2010; Hirano 2021; Lucendo 2019), and one study of fluticasone (Dellon 2022a), showed similar statistically significant differences from placebo in both groups (Analysis 1.35).

Withdrawals due to adverse events

Fourteen studies provided data on withdrawals due to adverse events when corticosteroids were compared to placebo (Alexander 2012; Bhardwaj 2017; Butz 2014; Dellon 2017; Dellon 2022a; Dohil 2010; Gupta 2015; Hirano 2020f; Hirano 2021; Konikoff 2006; Lucendo 2019; Miehke 2016; Straumann 2010b; Tytor 2021).

Corticosteroids ($n = 59/678$) may lead to slightly fewer withdrawals due to adverse events compared to placebo ($n = 44/354$) (RR 0.64, 95% CI 0.43 to 0.96; Analysis 1.36). The results are of low certainty due to imprecision and risk of bias (Summary of findings 1).

A sensitivity analysis using a fixed-effect model showed similar effects (RR 0.65, 95% CI 0.44 to 0.94; Analysis 1.37).

Subgroup analysis for the following did not reveal any differences between subgroups (Analysis 1.38; Analysis 1.39; Analysis 1.40): participant age, based on three studies in children younger than 18 years old (Dohil 2010; Gupta 2015; Konikoff 2006), and 11 studies in mixed adult and child participants (Alexander 2012; Bhardwaj 2017; Butz 2014; Dellon 2017; Dellon 2022a; Hirano 2020f; Hirano 2021; Lucendo 2019; Miehke 2016; Straumann 2010b; Tytor 2021); type of steroid, based on one study of beclomethasone (Bhardwaj 2017), seven of budesonide (Dellon 2017; Dohil 2010; Gupta 2015; Hirano 2021; Lucendo 2019; Miehke 2016; Straumann 2010b), five of fluticasone (Alexander 2012; Butz 2014; Dellon 2022a; Hirano 2020f; Konikoff 2006), and one of mometasone (Tytor 2021); and corticosteroid delivery method, based on five studies using an adapted asthma method (Alexander 2012; Bhardwaj 2017; Butz 2014; Konikoff 2006; Straumann 2010b), and nine studies using an esophageal-specific method (Dellon 2017; Dellon 2022a; Dohil 2010; Gupta 2015; Hirano 2020f; Hirano 2021; Lucendo 2019; Miehke 2016; Tytor 2021).

Secondary outcomes

Serious adverse events

Fourteen studies provided data on serious adverse events when corticosteroids were compared to placebo (Alexander 2012; Bhardwaj 2017; Butz 2014; Dellon 2017; Dellon 2022a; Dohil 2010; Gupta 2015; Hirano 2020f; Hirano 2021; Konikoff 2006; Lucendo 2019; Miehke 2016; Straumann 2010b; Tytor 2021).

Corticosteroids ($n = 11/678$) may result in little to no difference in serious adverse events compared to placebo ($n = 14/354$) (RR 0.35, 95% CI 0.17 to 0.73; Analysis 1.41). The results are of low certainty due to serious imprecision.

Total adverse events

Thirteen studies provided data on total adverse events when corticosteroids were compared to placebo (Alexander 2012; Butz 2014; Dellon 2017; Dellon 2022a; Dohil 2010; Gupta 2015; Hirano 2020f; Hirano 2021; Konikoff 2006; Lucendo 2019; Miehke 2016; Straumann 2010b; Tytor 2021).

Corticosteroids ($n = 290/669$) probably result in little to no difference in total adverse events compared to placebo ($n =$

111/345) (RR 1.14, 95% CI 0.94 to 1.40; [Analysis 1.42](#)). The results are of moderate certainty due to imprecision.

Quality of life

One study provided continuous data on quality of life when corticosteroids were compared to placebo ([Lucendo 2019](#)).

Corticosteroids (n = 59) may result in little to no difference in quality of life compared to placebo (n = 29) (mean difference (MD) 0.20, 95% CI -0.14 to 0.54; [Analysis 1.43](#)). The results are of low certainty due to serious imprecision.

Corticosteroids versus placebo for maintenance of remission

Three studies compared corticosteroids to placebo for maintenance of remission ([Dellon 2021b](#); [Straumann 2011](#); [Straumann 2020](#)).

Primary outcomes

Clinical improvement

Two studies compared corticosteroids to placebo for clinical improvement as a dichotomous outcome ([Dellon 2021b](#); [Straumann 2020](#)).

We cannot draw any conclusions on the effects of corticosteroids (n = 118/161) on clinical improvement for maintenance of remission compared to placebo (n = 20/91), measured as a dichotomous outcome (RR 2.17, 95% CI 0.75 to 6.27). The results are of very low certainty due to serious inconsistency and imprecision ([Analysis 2.1](#); [Summary of findings 2](#)).

Three studies compared corticosteroids to placebo for clinical improvement as a continuous outcome ([Dellon 2021b](#); [Straumann 2011](#); [Straumann 2020](#)).

We cannot draw any conclusions on the effects of corticosteroids (n = 169) on clinical improvement for maintenance of remission compared to placebo (n = 100), measured as a continuous outcome (SMD 0.51, 95% CI -0.49 to 1.52). The results are of very low certainty due to serious inconsistency and imprecision ([Analysis 2.2](#); [Summary of findings 2](#)).

Histological improvement

Three studies compared corticosteroids to placebo for histological improvement as a dichotomous outcome ([Dellon 2021b](#); [Straumann 2011](#); [Straumann 2020](#)).

Corticosteroids (n = 146/175) probably result in histological improvement for maintenance of remission compared to placebo (n = 14/105), measured as a dichotomous outcome (RR 4.58, 95% CI 1.66 to 12.62, NNTB = 3). The results are of moderate certainty, downgraded once due to imprecision ([Analysis 2.3](#); [Summary of findings 2](#)).

Three studies compared corticosteroids to placebo for histological improvement as a continuous outcome ([Dellon 2021b](#); [Straumann 2011](#); [Straumann 2020](#)).

Corticosteroids (n = 169) probably result in large histological improvement for maintenance of remission compared to placebo (n = 100), measured as a continuous outcome (SMD 1.26, 95% CI 0.74 to 1.78). The results are of moderate certainty due to inconsistency ([Analysis 2.4](#); [Summary of findings 2](#)).

Endoscopic improvement

Two studies compared corticosteroids to placebo for clinical improvement as a continuous outcome ([Dellon 2021b](#); [Straumann 2020](#)).

We cannot draw any conclusions on the effects of corticosteroids (n = 154) on endoscopic improvement for maintenance of remission compared to placebo (n = 86), measured as a continuous outcome (SMD 1.34, 95% CI -0.27 to 2.95). The results are of very low certainty due to serious inconsistency and imprecision ([Analysis 2.5](#); [Summary of findings 2](#)).

Withdrawals due to adverse events

Three studies provided data on withdrawals due to adverse events when corticosteroids were compared to placebo for maintenance of remission ([Dellon 2021b](#); [Straumann 2011](#); [Straumann 2020](#)).

Corticosteroids (n = 26/175) may lead to fewer withdrawals due to adverse events for maintenance of remission compared to placebo (n = 58/105) (RR 0.37, 95% CI 0.16 to 0.87). The results are of low certainty due to inconsistency and imprecision ([Analysis 2.6](#); [Summary of findings 2](#)).

Secondary outcomes

Serious adverse events

Three studies provided data on serious adverse events when corticosteroids were compared to placebo for maintenance of remission ([Dellon 2021b](#); [Straumann 2011](#); [Straumann 2020](#)).

We cannot draw any conclusions on the effects of corticosteroids (n = 4/175) on serious adverse events for maintenance of remission compared to placebo (n = 1/105) (RR 1.27, 95% CI 0.09 to 18.03; [Analysis 2.7](#)). The results are of low certainty due to serious imprecision.

Total adverse events

Three studies provided data on total adverse events when corticosteroids were compared to placebo for maintenance of remission ([Dellon 2021b](#); [Straumann 2011](#); [Straumann 2020](#)).

We cannot draw any conclusions on the effects of corticosteroids (n = 133/175) on total adverse events for maintenance of remission compared to placebo (n = 75/105) (RR 1.10, 95% CI 0.75 to 1.62; [Analysis 2.8](#)). The results are of very low certainty due to serious inconsistency and imprecision.

Quality of life

One study provided continuous data on quality of life when corticosteroids were compared to placebo for maintenance of remission ([Straumann 2020](#)).

Corticosteroids (n = 136) may lead to improved quality of life compared to placebo (n = 68), measured as a continuous outcome (MD 0.60, 95% CI 0.40 to 0.80; [Analysis 2.9](#)). The results are of low certainty due to serious imprecision.

Biologics versus placebo for induction of remission

Nine studies compared biologics to placebo for induction of remission ([Clayton 2014](#); [Dellon 2022](#); [Dellon 2022b](#); [Hirano 2019](#); [Hirano 2020](#); [Rothenberg 2015](#); [Rothenberg 2022](#); [Spergel 2012](#); [Straumann 2010a](#)).

Medical treatment of eosinophilic esophagitis (Review)

Primary outcomes

Clinical improvement

Five studies compared biologics to placebo for clinical improvement as a dichotomous outcome (Hirano 2019; Hirano 2020; Rothenberg 2015; Spergel 2012; Straumann 2010a).

Biologics (n = 169/281) may result in little to no difference in clinical improvement compared to placebo (n = 65/129), measured as a dichotomous outcome (RR 1.14, 95% CI 0.85 to 1.52). The results are of low certainty due to inconsistency and imprecision (Analysis 3.1; Summary of findings 3).

Sensitivity analyses using a fixed-effect model (RR 1.10, 95% CI 0.92 to 1.32; Analysis 3.2), and based on validated instruments (RR 1.37, 95% CI 1.02 to 1.85; Analysis 3.3), showed similar results.

Subgroup analysis for participant age, based on only one study in children younger than 18 years old (Spergel 2012), and four studies in mixed adult and child participants (Hirano 2019; Hirano 2020; Rothenberg 2015; Straumann 2010a), showed differences between the two age groups, with the mixed age group showing a statistically significant effect for the intervention, while no effect was observed for the children. Subgroup analysis for biologic mechanism, based on three studies of anti-IL-13 or anti-IL-4ra (Hirano 2019; Hirano 2020; Rothenberg 2015), and two studies of anti-IL-5 (Spergel 2012; Straumann 2010a), showed differences between the two groups, with the IL-13 group showing a statistically significant effect for the intervention, while no effect was observed for the IL-5 group (Analysis 3.4; Analysis 3.5).

Seven studies compared biologics to placebo for clinical improvement as a continuous outcome (Clayton 2014; Dellon 2022; Hirano 2019; Hirano 2020; Rothenberg 2015; Rothenberg 2022; Straumann 2010a).

Biologics (n = 195) may result in better clinical improvement compared to placebo (n = 192), measured as a continuous outcome (SMD 0.50, 95% CI 0.22 to 0.78). The results are of moderate certainty due to imprecision (Analysis 3.6; Summary of findings 3).

Sensitivity analyses using a fixed-effect model (SMD 0.48, 95% CI 0.28 to 0.69; Analysis 3.7), risk of bias (SMD 0.61, 95% CI 0.40 to 0.82; Analysis 3.9), and for validated instruments (SMD 0.62, 95% CI 0.37 to 0.88; Analysis 3.10) showed similar results.

A sensitivity analysis for peer-reviewed manuscripts showed slightly different results (SMD 0.36, 95% CI -0.09 to 0.81; Analysis 3.8).

A subgroup analysis for biologic mechanism, based on one study of anti-IgE (Clayton 2014), five studies of anti-IL-13/anti-IL-4ra (Dellon 2022; Hirano 2019; Hirano 2020; Rothenberg 2015; Rothenberg 2022), and one study of anti-IL-5 (Straumann 2010a), showed differences between the three groups, with statistically significant difference from placebo for anti-IL13/anti-IL-4r, however the data are very limited for the other categories (Analysis 3.11).

Histological improvement

Eight studies compared biologics to placebo for histological improvement as a dichotomous outcome (Dellon 2022; Dellon 2022b; Hirano 2019; Hirano 2020; Rothenberg 2015; Rothenberg 2022; Spergel 2012; Straumann 2010a).

Biologics (n = 394/586) may result in better histological improvement compared to placebo (n = 39/339), measured as a dichotomous outcome (RR 6.73, 95% CI 2.58 to 17.52, NNTB = 2). The results are of moderate certainty, downgraded once due to imprecision (Analysis 3.12; Summary of findings 3).

Sensitivity analyses using a fixed-effect model (RR 5.12, 95% CI 3.86 to 6.78; Analysis 3.13), peer-reviewed manuscripts (RR 6.13, 95% CI 1.67 to 22.51; Analysis 3.14), risk of bias (RR 6.73, 95% CI 2.58 to 17.52; Analysis 3.15), and for different histological thresholds showed similar results (RR 5.61, 95% CI 1.00 to 31.32; Analysis 3.16; RR 8.85, 95% CI 5.73 to 13.67; Analysis 3.17; RR 18.01, 95% CI 7.24 to 44.83; Analysis 3.18).

A subgroup analysis for participant age, based on two studies in children younger than 18 years old (Dellon 2022b; Spergel 2012), six studies in mixed adult and children participants (Dellon 2022; Hirano 2019; Hirano 2020; Rothenberg 2015; Rothenberg 2022; Straumann 2010a), and one study that provided separate data on children and adults (Dellon 2022b), revealed no evidence of a difference between the age subgroups (Analysis 3.19).

A subgroup analysis for biologic mechanism (Analysis 3.20), based on one study on anti-sialic acid binding Ig-like lectin 8 (Dellon 2022b), five studies of anti-IL-13/anti-IL-4ra (Dellon 2022; Hirano 2019; Hirano 2020; Rothenberg 2015; Rothenberg 2022), and two studies of anti-IL-5 (Spergel 2012; Straumann 2010a), showed significant differences between all categories, but it was based on limited data.

Six studies compared biologics to placebo for histological improvement as a continuous outcome (Clayton 2014; Hirano 2019; Hirano 2020; Rothenberg 2015; Spergel 2012; Straumann 2010a).

We cannot draw any conclusions on the effects of biologics (n = 240) on histological improvement compared to placebo (n = 130) measured as a continuous outcome (SMD 1.01, 95% CI 0.36 to 1.66). The results are of very low certainty due to serious inconsistency and imprecision (Analysis 3.21; Summary of findings 3).

A sensitivity analysis using a fixed-effect model showed that biologics (n = 240) may lead to better histological improvement when compared to placebo (n = 130), measured as a continuous outcome (SMD 1.25, 95% CI 1.01 to 1.49; Analysis 3.22). A sensitivity analysis for risk of bias showed that biologics (n = 224) probably result in better histological improvement compared to placebo (n = 116), measured as a continuous outcome (SMD 1.39, 95% CI 1.01 to 1.77; Analysis 3.23).

A subgroup analysis for participant age, based on one study in children younger than 18 years old (Spergel 2012), and five studies in mixed adult and child participants (Clayton 2014; Hirano 2019; Hirano 2020; Rothenberg 2015; Straumann 2010a), revealed no differences between age groups, however this was based on limited data (Analysis 3.24).

A subgroup analysis for biologic mechanism, based on one study of anti-IgE (Clayton 2014), three studies of anti-IL-13/anti-IL-4ra (Hirano 2019; Hirano 2020; Rothenberg 2015), and two studies of anti-IL-5 (Spergel 2012; Straumann 2010a), showed differences between groups, with statistically significant effects for anti-IL-13 and anti-IL-5, however this was also based on limited data (Analysis 3.25).

Medical treatment of eosinophilic esophagitis (Review)

Endoscopic improvement

One study compared biologics to placebo for endoscopic improvement, measured as a dichotomous outcome (Straumann 2010a).

Biologics (n = 0/5) may result in little to no difference in endoscopic improvement compared to placebo (n = 0/6), measured as a dichotomous outcome (effect not estimable). The results are of low certainty due to serious imprecision (Analysis 3.26; Summary of findings 3).

Three studies compared biologics to placebo for endoscopic improvement, measured as a continuous outcome (Dellon 2022; Hirano 2019; Hirano 2020).

We cannot draw any conclusions about the effects of biologics (n = 115) on endoscopic improvement compared to placebo (n = 82), measured as a continuous outcome (SMD 2.79, 95% CI 0.36 to 5.22; Analysis 3.27). The results are of very low certainty due to serious inconsistency and imprecision.

A sensitivity analysis using a fixed-effect model showed similar results (SMD 1.20, 95% CI 0.86 to 1.55; Analysis 3.28). A sensitivity analysis for risk of bias showed that biologics (n = 80) may result in slightly better endoscopic improvement compared to placebo (n = 56), measured as a continuous outcome (SMD 0.82, 95% CI 0.42 to 1.21; Analysis 3.29). The results are of low certainty due to serious imprecision.

Withdrawals due to adverse events

Eight studies provided data on withdrawals due to adverse events when biologics were compared to placebo (Clayton 2014; Dellon 2022; Dellon 2022b; Hirano 2019; Hirano 2020; Rothenberg 2015; Spergel 2012; Straumann 2010a).

There may be no difference between biologics (n = 54/518) and placebo (n = 16/274) in withdrawals due to adverse events (RR 1.55, 95% CI 0.88 to 2.74). The results are of low certainty due to serious imprecision (Analysis 3.30; Summary of findings 3).

Sensitivity analyses using a fixed-effect model (RR 1.53, 95% CI 0.89 to 2.64; Analysis 3.31), risk of bias (RR 1.55, 95% CI 0.88 to 2.74; Analysis 3.32), and peer-reviewed publications (RR 1.55, 95% CI 0.88 to 2.74; Analysis 3.33) showed similar results.

Subgroup analyses for the following did not reveal differences between the subgroups (Analysis 3.34; Analysis 3.35): participant age, based on one study in children younger than 18 years old (Spergel 2012), and seven studies in mixed adult and child participants (Clayton 2014; Dellon 2022; Dellon 2022b; Hirano 2019; Hirano 2020; Rothenberg 2015; Straumann 2010a); and biologic mechanism, based on one study of anti-IgE (Clayton 2014), one study of anti-sialic acid-binding Ig-like lectin 8 (Dellon 2022b), four studies of anti-IL-13 (Dellon 2022; Hirano 2019; Hirano 2020; Rothenberg 2015), and two studies of anti-IL-5 (Spergel 2012; Straumann 2010a).

Secondary outcomes

Serious adverse events

Six studies provided data on serious adverse events when biologics were compared to placebo (Dellon 2022b; Hirano 2019; Hirano 2020; Rothenberg 2015; Spergel 2012; Straumann 2010a).

There may be no difference between biologics (n = 10/464) and placebo (n = 6/221) in serious adverse events (RR 0.70, 95% CI 0.25 to 1.97; Analysis 3.36). The results are of low certainty due to serious imprecision.

Total adverse events

Six studies provided data on total adverse events when biologics were compared to placebo (Dellon 2022b; Hirano 2019; Hirano 2020; Rothenberg 2015; Spergel 2012; Straumann 2010a).

There is probably no difference between biologics (n = 310/464) and placebo (n = 136/221) in total adverse events (RR 1.07, 95% CI 0.94 to 1.23; Analysis 3.37). The results are of moderate certainty due to imprecision.

Quality of life

One study provided continuous data on quality of life when biologics were compared to placebo (Hirano 2020).

There may be little to no difference between biologics (n = 23) and placebo (n = 24) in quality of life (MD 0.33, 95% CI -0.06 to 0.72; Analysis 3.38). The results are of low certainty due to serious imprecision.

Cromolyn sodium versus placebo

One study compared cromolyn sodium to placebo for induction of remission (Lieberman 2018).

Primary outcomes

Clinical improvement

Cromolyn sodium (n = 8) may result in little to no difference in clinical improvement compared to placebo (n = 6), measured as a continuous outcome (MD 4.70, 95% CI -12.09 to 21.49). The results are of low certainty due to serious imprecision (Analysis 4.1; Summary of findings 4).

Histological improvement

Cromolyn sodium (n = 9) may result in little to no difference in histological improvement compared to placebo (n = 6), measured as a continuous outcome (MD 14.20, 95% CI -36.90 to 65.30). The results are of low certainty due to serious imprecision (Analysis 4.2; Summary of findings 4).

Endoscopic improvement

There were no data for meta-analysis for this outcome.

Withdrawals due to adverse events

Cromolyn sodium (n = 0/9) may result in little to no difference in withdrawals due to adverse events compared to placebo (n = 1/7) (RR 0.27, 95% CI 0.01 to 5.70). The results are of low certainty due to serious imprecision (Analysis 4.3; Summary of findings 4).

Secondary outcomes

Serious adverse events

Cromolyn sodium (n = 0/9) may result in little to no difference in serious adverse events compared to placebo (n = 0/7) (effect not estimable; Analysis 4.4). The results are of low certainty due to serious imprecision

Medical treatment of eosinophilic esophagitis (Review)

Total adverse events

There were no data for this outcome.

Quality of life

There were no data for this outcome.

PGD2R antagonist OC000459 versus placebo

One study compared PGD2R antagonist OC000459 to placebo for induction of remission ([Straumann 2013](#)).

Primary outcomes

Clinical improvement

We cannot draw any conclusions on the effect of PGD2R antagonist (n = 14) on clinical improvement compared to placebo (n = 12), measured as a continuous outcome (MD -1.06, 95% CI -6.80 to 4.68). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 5.1](#); [Summary of findings 5](#)).

Histological improvement

We cannot draw any conclusions on the effect of PGD2R antagonist (n = 14) on histological improvement compared to placebo (n = 12), measured as a continuous outcome (MD 26.21, 95% CI -23.78 to 76.20). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 5.2](#); [Summary of findings 5](#)).

Endoscopic improvement

We cannot draw any conclusions on the effect of PGD2R antagonist (n = 14) on histological improvement compared to placebo (n = 12), measured as a continuous outcome (MD -0.49, 95% CI -2.05 to 1.07). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 5.3](#); [Summary of findings 5](#)).

Withdrawals due to adverse events

We cannot draw any conclusions on the effect of PGD2R antagonist (n = 0/14) on withdrawals due to adverse events compared to placebo (n = 0/12) (effect not estimable). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 5.4](#); [Summary of findings 5](#)).

Secondary outcomes

Serious adverse events

We cannot draw any conclusions on the effect of PGD2R antagonist (n = 1/14) on serious adverse events compared to placebo (n = 0/12) (RR 2.60, 95% CI 0.12 to 58.48; [Analysis 5.5](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Total adverse events

There were no data for this outcome.

Quality of life

There were no data for this outcome.

Swallowed fluticasone versus oral prednisone

One study compared swallowed fluticasone to oral prednisone for induction of remission ([Schaefer 2008](#)).

Primary outcomes

Clinical improvement

We cannot draw any conclusions on the effects of swallowed fluticasone (n = 35/40) on clinical improvement compared to oral prednisone (n = 32/40), measured as dichotomous outcome (RR 1.09, 95% CI 0.90 to 1.33). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 6.1](#); [Summary of findings 6](#)).

Histological improvement

We cannot draw any conclusions on the effects of swallowed fluticasone (n = 33/40) on histological improvement compared to oral prednisone (n = 30/40), measured as a dichotomous outcome (RR 1.10, 95% CI 0.87 to 1.38). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 6.2](#); [Summary of findings 6](#)).

We cannot draw any conclusions on the effects of swallowed fluticasone (n = 36) on histological improvement compared to oral prednisone (n = 32/40), measured as a continuous outcome (MD -4.45, 95% CI -9.08 to 0.18). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 6.3](#); [Summary of findings 6](#)).

Endoscopic improvement

We cannot draw any conclusions on the effects of swallowed fluticasone (n = 34/40) on endoscopic improvement compared to oral prednisone (n = 30/40), measured as dichotomous outcome (RR 1.13, 95% CI 0.91 to 1.41). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 6.4](#); [Summary of findings 6](#)).

Withdrawals due to adverse events

We cannot draw any conclusions on the effects of swallowed fluticasone (n = 4/40) on withdrawals due to adverse events compared to oral prednisone (n = 8/40) (RR 0.50, 95% CI 0.16 to 1.53). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 6.5](#); [Summary of findings 6](#)).

Secondary outcomes

Serious adverse events

We cannot draw any conclusions on the effects of swallowed fluticasone (n = 0/40) on serious adverse events compared to oral prednisone (n = 3/40) (RR 0.14, 95% CI 0.01 to 2.68; [Analysis 6.6](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Total adverse events

We cannot draw any conclusions on the effects of swallowed fluticasone (n = 6/40) on total adverse events compared to oral prednisone (n = 16/40) (RR 0.38, 95% CI 0.16 to 0.86; [Analysis 6.7](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Quality of life

There were no data for this outcome.

Medical treatment of eosinophilic esophagitis (Review)

Oral viscous budesonide versus swallowed fluticasone

One study compared oral viscous budesonide to swallowed fluticasone for induction of remission ([Dellon 2019](#)).

Primary outcomes

Clinical improvement

Oral viscous budesonide (n = 46) may result in little to no difference in clinical improvement compared to swallowed fluticasone (n = 38), measured as a continuous outcome (MD -0.60, 95% CI -3.78 to 2.58). The results are of low certainty due to serious imprecision ([Analysis 7.1](#); [Summary of findings 7](#)).

Histological improvement

Oral viscous budesonide (n = 40/65) may result in little to no difference in histological improvement compared to swallowed fluticasone (n = 35/64), measured as a dichotomous outcome (RR 1.13, 95% CI 0.84 to 1.51). The results are of low certainty due to serious imprecision ([Analysis 7.2](#); [Summary of findings 7](#)).

Oral viscous budesonide (n = 56) may result in little to no difference in histological improvement compared to swallowed fluticasone (n = 55), measured as a continuous outcome (MD 6.20, 95% CI -5.63 to 18.03). The results are of low certainty due to serious imprecision ([Analysis 7.3](#); [Summary of findings 7](#)).

Endoscopic improvement

Oral viscous budesonide (n = 56) may result in little to no difference in endoscopic improvement compared to swallowed fluticasone (n = 55), measured as a continuous outcome (MD 0.70, 95% CI -0.03 to 1.43). The results are of low certainty due to serious imprecision ([Analysis 7.4](#); [Summary of findings 7](#)).

Withdrawals due to adverse events

Oral viscous budesonide (n = 9/65) may result in little to no difference in withdrawals due to adverse events compared to swallowed fluticasone (n = 9/64) (RR 0.98, 95% CI 0.42 to 2.32). The results are of low certainty due to serious imprecision ([Analysis 7.5](#); [Summary of findings 7](#)).

Secondary outcomes

Serious adverse events

Oral viscous budesonide (n = 0/65) may result in little to no difference in serious adverse events compared to swallowed fluticasone (n = 1/64) (RR 0.33, 95% CI 0.01 to 7.91; [Analysis 7.6](#)). The results are of low certainty due to serious imprecision.

Total adverse events

Oral viscous budesonide (n = 10/65) may result in little to no difference in total adverse events compared to swallowed fluticasone (n = 15/64) (RR 0.66, 95% CI 0.32 to 1.35; [Analysis 7.7](#)). The results are of low certainty due to serious imprecision.

Quality of life

There were no data for this outcome.

Esomeprazole versus fluticasone

Two studies compared esomeprazole to fluticasone for induction of remission ([Moawad 2013](#); [Peterson 2010](#)).

Primary outcomes

Clinical improvement

Both studies provided continuous data on clinical improvement ([Moawad 2013](#); [Peterson 2010](#)).

We cannot draw any conclusions on the effects of esomeprazole (n = 32) on clinical improvement compared to fluticasone (n = 31), measured as a continuous outcome (SMD 0.32, 95% CI -0.88 to 1.52). The results are of very low certainty due to inconsistency, imprecision, and risk of bias ([Analysis 8.1](#); [Summary of findings 8](#)).

Histological improvement

Both studies provided dichotomous and continuous data on histological improvement ([Moawad 2013](#); [Peterson 2010](#)).

We cannot draw any conclusions on the effects of esomeprazole (n = 13/36) on histological improvement compared to fluticasone (n = 8/36), measured as a dichotomous outcome (RR 1.62, 95% CI 0.77 to 3.41). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 8.2](#); [Summary of findings 8](#)).

We cannot draw any conclusions on the effects of esomeprazole (n = 33) on histological improvement compared to fluticasone (n = 34), measured as a continuous outcome (SMD 0.28, 95% CI -0.20 to 0.76). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 8.3](#); [Summary of findings 8](#)).

Endoscopic improvement

No studies provided meta-analysis data for this outcome.

They reported data on specific endoscopic findings, which can be found in [Table 1](#).

Withdrawals due to adverse events

Both studies provided data on withdrawals due to adverse events ([Moawad 2013](#); [Peterson 2010](#)).

We cannot draw any conclusions on the effects of esomeprazole (n = 3/36) on withdrawals due to adverse events compared to fluticasone (n = 3/36) (RR 0.95, 95% CI 0.07 to 13.38). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 8.4](#); [Summary of findings 8](#)).

Secondary outcomes

Serious adverse events

We cannot draw any conclusions on the effects of esomeprazole (n = 0/36) on serious adverse events compared to fluticasone (n = 0/36) (effect not estimable; [Analysis 8.5](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Total adverse events

We cannot draw any conclusions on the effects of esomeprazole (n = 0/33) on total adverse events compared to fluticasone (n = 3/34) (RR 0.14, 95% CI 0.01 to 2.61; [Analysis 8.6](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Quality of life

There were no data for this outcome.

Medical treatment of eosinophilic esophagitis (Review)

One-food elimination diet versus four-food elimination diet

One study compared a one-food elimination diet to a four-food elimination diet for induction of remission (Kliewer 2019)

Primary outcomes

Clinical improvement

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 33) on clinical improvement compared to a four-food elimination diet (n = 17), measured as a continuous outcome (MD -7.50, 95% CI -16.28 to 1.28). The results are of very low certainty due to serious imprecision and risk of bias (Analysis 9.1; Summary of findings 9).

Histological improvement

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 24/38) on histological improvement compared to a four-food elimination diet (n = 7/25), measured as a dichotomous outcome (RR 2.26, 95% CI 1.15 to 4.43). The results are of very low certainty due to serious imprecision and risk of bias (Analysis 9.2; Summary of findings 9).

Endoscopic improvement

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 22) on endoscopic improvement compared to a four-food elimination diet (n = 12), measured as a continuous outcome (MD -0.60, 95% CI -2.15 to 0.95). The results are of very low certainty due to serious imprecision and risk of bias (Analysis 9.3; Summary of findings 9).

Withdrawals due to adverse events

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 4/38) on withdrawals due to adverse events compared to a four-food elimination diet (n = 8/25) (RR 0.33, 95% CI 0.11 to 0.98). The results are of very low certainty due to serious imprecision and risk of bias (Analysis 9.4; Summary of findings 9).

Secondary outcomes

Serious adverse events

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 1/38) on serious adverse events compared to a four-food elimination diet (n = 1/25) (RR 0.66, 95% CI 0.04 to 10.04; Analysis 9.5). The results are of very low certainty due to serious imprecision and risk of bias.

Total adverse events

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 5/38) on total adverse events compared to a four-food elimination diet (n = 8/25) (RR 0.41, 95% CI 0.15 to 1.11; Analysis 9.6). The results are of very low certainty due to serious imprecision and risk of bias.

Quality of life

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 38) on quality of life compared to a four-food elimination diet (n = 25), measured as a continuous outcome (MD 0.10, 95% CI -6.49 to 6.69; Analysis 9.7). The results are of very low certainty due to serious imprecision and risk of bias.

One-food elimination diet versus six-food elimination diet

One study compared a one-food elimination diet to a six-food elimination diet for induction of remission (Kliewer 2021).

Primary outcomes

Clinical improvement

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 67) on clinical improvement compared to a six-food elimination diet (n = 62), measured as a continuous outcome (MD -5.20, 95% CI -11.06 to 0.66). The results are of very low certainty due to serious imprecision and risk of bias (Analysis 10.1; Summary of findings 10).

Histological improvement

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 23/67) on histological improvement compared to a six-food elimination diet (n = 25/62), measured as a dichotomous outcome (RR 0.85, 95% CI 0.54 to 1.33). The results are of very low certainty due to serious imprecision and risk of bias (Analysis 10.2; Summary of findings 10).

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 67) on histological improvement compared to a six-food elimination diet (n = 62), measured as a continuous outcome (MD 6.80, 95% CI -10.40 to 24.00). The results are of very low certainty due to serious imprecision and risk of bias (Analysis 10.3; Summary of findings 10).

Endoscopic improvement

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 67) on endoscopic improvement compared to a six-food elimination diet (n = 62), measured as a continuous outcome (MD -0.42, 95% CI -1.67 to 0.83). The results are of very low certainty due to serious imprecision and risk of bias (Analysis 10.4; Summary of findings 10).

Withdrawals due to adverse events

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 2/67) on withdrawals due to adverse events compared to a six-food elimination diet (n = 3/62) (RR 0.62, 95% CI 0.11 to 3.57). The results are of very low certainty due to serious imprecision and risk of bias (Analysis 10.5; Summary of findings 10).

Secondary outcomes

Serious adverse events

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 0/67) on serious adverse events compared to a six-food elimination diet (n = 0/62) (effect not estimable; Analysis 10.6). The results are of very low certainty due to serious imprecision and risk of bias.

Total adverse events

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 1/67) on total adverse events compared to a six-food elimination diet (n = 2/62) (RR 0.46, 95% CI 0.04 to 4.98; Analysis 10.7). The results are of very low certainty due to serious imprecision and risk of bias.

Medical treatment of eosinophilic esophagitis (Review)

Quality of life

We cannot draw any conclusions on the effects of a one-food elimination diet (n = 67) on quality of life compared to a six-food elimination diet (n = 62), measured as a continuous outcome (MD 0.57, 95% CI -3.25 to 4.39; [Analysis 10.8](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Four-food elimination diet with omeprazole versus omeprazole

One study compared a four-food elimination diet with omeprazole to omeprazole for induction of remission ([Heine 2019](#)).

Primary outcomes

Clinical improvement

There were no data for this outcome.

Histological improvement

We cannot draw any conclusions on the effects of a four-food elimination diet with omeprazole (n = 22/32) on histological improvement compared to omeprazole (n = 14/32), measured as a dichotomous outcome (RR 1.57, 95% CI 0.99 to 2.48). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 11.1](#); [Summary of findings 11](#)).

We cannot draw any conclusions on the effects of a four-food elimination diet with omeprazole (n = 27) on histological improvement compared to omeprazole (n = 31), measured as a continuous outcome (MD 9.50, 95% CI -11.18 to 30.18). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 11.2](#); [Summary of findings 11](#)).

Endoscopic improvement

There were no data for this outcome.

Withdrawals due to adverse events

We cannot draw any conclusions on the effects of a four-food elimination diet with omeprazole (n = 5/32) on withdrawals due to adverse events compared to omeprazole (n = 1/32) (RR 5.00, 95% CI 0.62 to 40.44). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 11.3](#); [Summary of findings 11](#)).

Secondary outcomes

There were no data for any of the secondary outcomes.

Four-food elimination diet and amino acid formula versus four-food elimination diet

One study compared a four-food elimination diet with amino acid formula to a four-food elimination diet for induction of remission ([De Rooij 2022](#)).

Primary outcomes

Clinical improvement

We cannot draw any conclusions on the effects of a four-food elimination diet with amino acid formula (n = 21) on clinical improvement compared to a four-food elimination diet (n = 20), measured as a continuous outcome (MD -0.50, 95% CI -2.41 to 1.41). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 12.1](#); [Summary of findings 12](#)).

Histological improvement

We cannot draw any conclusions on the effects of a four-food elimination diet with amino acid formula (n = 10/21) on histological improvement compared to a four-food elimination diet (n = 5/20), measured as a dichotomous outcome (RR 1.90, 95% CI 0.79 to 4.60). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 12.2](#); [Summary of findings 12](#)).

We cannot draw any conclusions on the effects of a four-food elimination diet with amino acid formula (n = 21) on histological improvement compared to a four-food elimination diet (n = 20), measured as a continuous outcome (MD 13.80, 95% CI -9.50 to 37.10). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 12.3](#); [Summary of findings 12](#)).

Endoscopic improvement

We cannot draw any conclusions on the effect of a four-food elimination diet with amino acid formula (n = 21) on endoscopic improvement compared to a four-food elimination diet (n = 20), measured as a continuous outcome (MD -1.00, 95% CI -2.83 to 0.83). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 12.4](#); [Summary of findings 12](#)).

Withdrawals due to adverse events

We cannot draw any conclusions on the effects of a four-food elimination diet with amino-acid formula (n = 1/21) compared to a four-food elimination diet (n = 1/20) on withdrawals due to adverse events (RR 0.95, 95% CI 0.06 to 14.22). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 12.5](#); [Summary of findings 12](#)).

Secondary outcomes

Serious adverse events

We cannot draw any conclusions on the effects of a four-food elimination diet with amino-acid formula (n = 0/21) compared to a four-food elimination diet (n = 0/20) on serious adverse events (effect not estimable; [Analysis 12.6](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Total adverse events

We cannot draw any conclusions on the effects of a four-food elimination diet with amino-acid formula (n = 1/21) compared to a four-food elimination diet (n = 0/20) on total adverse events (RR 2.86, 95% CI 0.12 to 66.44). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 12.7](#)).

Quality of life

There were no data for this outcome.

Nebulized budesonide versus viscous budesonide

One study compared nebulized swallowed budesonide to viscous swallowed budesonide for induction of remission ([Dellon 2012](#)).

Primary outcomes

Clinical improvement

We cannot draw any conclusions on the effects of nebulized swallowed budesonide (n = 11) on clinical improvement compared to viscous swallowed budesonide (n = 11), measured as a continuous outcome (MD -6.00, 95% CI -18.30 to 6.30). The results

Medical treatment of eosinophilic esophagitis (Review)

are of very low certainty due to serious imprecision and risk of bias ([Analysis 13.1](#); [Summary of findings 13](#)).

Histological improvement

We cannot draw any conclusions on the effects of nebulized swallowed budesonide ($n = 11$) on histological improvement compared to viscous swallowed budesonide ($n = 11$), measured as a continuous outcome (MD 78.00, 95% CI 20.81 to 135.19). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 13.2](#); [Summary of findings 13](#)).

Endoscopic improvement

There were no data for this outcome.

Withdrawals due to adverse events

We cannot draw any conclusions on the effects of nebulized swallowed budesonide ($n = 0/13$) compared to viscous swallowed budesonide ($n = 0/12$) on withdrawals due to adverse events (effect not estimable). The results are of very low certainty due to serious imprecision and risk of bias ([Summary of findings 12](#)).

Secondary outcomes

Serious adverse events

We cannot draw any conclusions on the effects of nebulized swallowed budesonide ($n = 0/13$) compared to viscous swallowed budesonide ($n = 0/12$) on serious adverse events (effect not estimable; [Analysis 13.3](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Total adverse events

We cannot draw any conclusions on the effects of nebulized swallowed budesonide ($n = 2/13$) compared to viscous swallowed budesonide ($n = 2/12$) on total adverse events (RR 0.92, 95% CI 0.15 to 5.56; [Analysis 13.4](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Quality of life

There were no data for this outcome.

Viaskin milk patch versus placebo

One study compared Viaskin milk patch to placebo for induction of remission ([Spergel 2020](#)).

Primary outcomes

Clinical improvement

Viaskin milk patch ($n = 7$) may result in little to no difference in clinical improvement compared to placebo ($n = 2$), measured as a continuous outcome (MD 1.29, 95% CI -0.83 to 3.41). The results are of low certainty due to serious imprecision ([Analysis 14.1](#); [Summary of findings 14](#)).

Histological improvement

Viaskin milk patch ($n = 7$) may result in little to no difference in histological improvement compared to placebo ($n = 2$), measured as a continuous outcome (MD 69.43, 95% CI -21.75 to 160.61). The results are of low certainty due to serious imprecision ([Analysis 14.2](#); [Summary of findings 14](#)).

Endoscopic improvement

Viaskin milk patch ($n = 15$) may result in little to no difference in endoscopic improvement compared to placebo ($n = 5$), measured as a continuous outcome (MD -0.33, 95% CI -2.00 to 1.34). The results are of low certainty due to serious imprecision ([Analysis 14.3](#); [Summary of findings 14](#)).

Withdrawals due to adverse events

Viaskin milk patch ($n = 1/15$) may result in little to no difference in withdrawals due to adverse events compared to placebo ($n = 0/5$) (RR 1.12, 95% CI 0.05 to 23.99). The results are of low certainty due to serious imprecision ([Analysis 14.4](#); [Summary of findings 14](#)).

Secondary outcomes

Serious adverse events

Viaskin milk patch ($n = 0/15$) may result in little to no difference in serious adverse events compared to placebo ($n = 1/5$) (RR 0.13, 95% CI 0.01 to 2.67; [Analysis 14.5](#)). The results are of low certainty due to serious imprecision.

Total adverse events

Viaskin milk patch ($n = 15/15$) may result in little to no difference in total adverse events compared to placebo ($n = 5/5$) (RR 1.00, 95% CI 0.77 to 1.29; [Analysis 14.6](#)). The results are of low certainty due to serious imprecision.

Quality of life

Viaskin milk patch ($n = 7$) may result in little to no difference in quality of life compared to placebo ($n = 2$), measured as a continuous outcome (MD 13.60, 95% CI -16.12 to 43.32; [Analysis 14.7](#)). The results are of low certainty due to serious imprecision.

Leukotriene receptor antagonist versus placebo for maintenance of remission

One study compared leukotriene receptor antagonist to placebo for maintenance of remission ([Alexander 2017](#)).

Primary outcomes

Clinical improvement

We cannot draw any conclusions on the effects of leukotriene receptor antagonist ($n = 8/20$) compared to placebo ($n = 5/21$) on clinical improvement for maintenance of remission, measured as a dichotomous outcome (RR 1.68, 95% CI 0.66 to 4.28). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 15.1](#); [Summary of findings 15](#)).

Histological improvement

There were no data for meta-analysis for this outcome.

Endoscopic improvement

There were no data for meta-analysis for this outcome.

Withdrawals due to adverse events

We cannot draw any conclusions on the effects of leukotriene receptor antagonist ($n = 2/20$) on withdrawals due to adverse events compared to placebo ($n = 1/21$) for maintenance of remission (RR 2.10, 95% CI 0.21 to 21.39). The results are of very low certainty due

to serious imprecision and risk of bias ([Analysis 15.2](#); [Summary of findings 15](#)).

Secondary outcomes

Serious adverse events

We cannot draw any conclusions on the effects of leukotriene receptor antagonist (n = 0/20) on serious adverse events compared to placebo (n = 0/21) for maintenance of remission (effect not estimable; [Analysis 15.3](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Total adverse events

We cannot draw any conclusions on the effects of leukotriene receptor antagonist (n = 0/20) on total adverse events compared to placebo (n = 0/21) for maintenance of remission (effect not estimable; [Analysis 15.4](#)). The results are of very low certainty due to serious imprecision and risk of bias.

Quality of life

There were no data for this outcome.

Mepolizumab high-dose (10 mg/kg) versus low-dose (0.55 mg/kg)

One study compared a low dose of the biologic mepolizumab (0.55 mg/kg) to a medium dose (2.5 mg/kg) and to a high dose (10 mg/kg) ([Assa'ad 2011](#)). The study did not include any control groups other than mepolizumab.

Primary outcomes

Clinical improvement

There were no data for this outcome.

Histological improvement

We cannot draw any conclusions about the effects of mepolizumab 10 mg/kg (n = 5/20) compared to mepolizumab 0.55 mg/kg (n = 4/19) on histological improvement, measured as a dichotomous outcome (RR 1.19, 95% CI 0.37 to 3.77). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 16.1](#); [Summary of findings 16](#)).

Endoscopic improvement

There were no data for this outcome.

Withdrawals due to adverse events

We cannot draw any conclusions about the effects of mepolizumab 10 mg/kg (n = 2/20) compared to mepolizumab 0.55 mg/kg (n = 3/19) on withdrawals due to adverse events (RR 0.63, 95% CI 0.12 to 3.38). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 16.2](#); [Summary of findings 16](#)).

Secondary outcomes

Serious adverse events

The study reported three serious adverse events but did not specify in which group they occurred. No conclusions can be drawn. The results are of very low certainty due to serious risk of bias and imprecision.

Total adverse events

The study reported a total of 51/59 participants with more than one adverse event, but did not give specifics per group. The results are of very low certainty due to serious risk of bias and imprecision.

Quality of life

There were no data for this outcome.

Mepolizumab medium-dose (2.5 mg/kg) versus low-dose (0.55 mg/kg)

One study compared a low dose of the biologic mepolizumab (0.55 mg/kg) to a medium dose (2.5 mg/kg) and to a high dose (10 mg/kg) ([Assa'ad 2011](#)). The study did not include any control groups other than mepolizumab.

Primary outcomes

Clinical improvement

There were no data for this outcome.

Histological improvement

We cannot draw any conclusions about the effects of mepolizumab 2.5 mg/kg (n = 9/20) compared to mepolizumab 0.55 mg/kg (n = 4/19) on histological improvement, measured as a dichotomous outcome (RR 2.14, 95% CI 0.79 to 5.79). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 17.1](#); [Summary of findings 17](#)).

Endoscopic improvement

There were no data for this outcome.

Withdrawals due to adverse events

We cannot draw any conclusions about the effects of mepolizumab 2.5 mg/kg (n = 1/20) compared to mepolizumab 0.55 mg/kg (n = 3/19) on withdrawals due to adverse events (RR 0.32, 95% CI 0.04 to 2.79). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 17.2](#); [Summary of findings 17](#)).

Secondary outcomes

Serious adverse events

The study reported three serious adverse events but did not specify in which group they occurred. No conclusions can be drawn. The results are of very low certainty due to serious risk of bias and imprecision.

Total adverse events

The study reported a total of 51/59 participants with more than one adverse event, but did not give specifics per group. The results are of very low certainty due to serious risk of bias and imprecision.

Quality of life

There were no data for this outcome.

Mepolizumab high-dose (10 mg/kg) versus medium-dose (2.5 mg/kg)

One study compared a low dose of the biologic mepolizumab (0.55 mg/kg) to a medium dose (2.5 mg/kg) and to a high dose (10 mg/kg) ([Assa'ad 2011](#)). The study did not include any control groups other than mepolizumab.

Primary outcomes

Clinical improvement

There were no data for this outcome.

Histological improvement

We cannot draw any conclusions about the effects of mepolizumab 10 mg/kg (n = 5/20) compared to mepolizumab 2.5 mg/kg (n = 9/20) on histological improvement, measured as a dichotomous outcome (RR 0.56, 95% CI 0.23 to 1.37). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 18.1](#); [Summary of findings 18](#)).

Endoscopic improvement

There were no data for this outcome.

Withdrawals due to adverse events

We cannot draw any conclusions about the effects of mepolizumab 10 mg/kg (n = 2/20) compared to mepolizumab 2.5 mg/kg (n = 1/20) on withdrawals due to adverse events (RR 2.00, 95% CI 0.20 to 20.33). The results are of very low certainty due to serious imprecision and risk of bias ([Analysis 18.2](#); [Summary of findings 18](#)).

Secondary outcomes

Serious adverse events

The study reported three serious adverse events but did not specify in which group they occurred. No conclusions can be drawn. The results are of very low certainty due to serious risk of bias and imprecision.

Total adverse events

The study reported a total of 51/59 participants with more than one adverse event, but did not give specifics per group. The results are of very low certainty due to serious risk of bias and imprecision.

Quality of life

There were no data for this outcome.

Six-food elimination diet versus swallowed fluticasone versus swallowed budesonide versus oral viscous budesonide

One study compared a six-food elimination diet to swallowed fluticasone to swallowed budesonide and to oral viscous budesonide ([Oliva 2018](#)). This was an abstract report without any data that could be used for meta-analysis of any outcome.

Primary outcomes

Clinical improvement

There were no data for this outcome.

Histological improvement

The study only reported percentages, with the numbers of patients in each group not reported. At eight weeks, 69% of participants in the six-food elimination diet group achieved histological improvement, 67% in the swallowed fluticasone group, 75% in the swallowed budesonide group, and 85% in the oral viscous budesonide group. No conclusions can be drawn. These results are of very low certainty due to serious risk of bias and imprecision.

Endoscopic improvement

There were no data for this outcome.

Withdrawals due to adverse events

There were no data for this outcome.

Secondary outcomes

There were no data for any of the secondary outcomes.

DISCUSSION

Summary of main results

In the studies comparing corticosteroids and placebo for induction, corticosteroids may lead to clinical symptom improvement when reported as both dichotomous and continuous outcomes. Corticosteroids lead to a large increase in histological improvement (dichotomous outcome) and may increase histological improvement (continuous outcome) when compared to placebo. Corticosteroids may increase endoscopic improvement (dichotomous and continuous outcome). Withdrawals due to adverse events as a dichotomous outcome may be lower for corticosteroids when compared to placebo.

In the studies comparing corticosteroids and placebo for maintenance, corticosteroids probably lead to a large increase in histological improvement (dichotomous outcome) and probably increase histological improvement (continuous outcome) when compared to placebo. No conclusions can be drawn for any other outcomes due to very low-certainty evidence.

In the studies comparing biologics and placebo, biologics may lead to little to no clinical symptom improvement when reported as a dichotomous outcome, and may lead to an increase in clinical symptom improvement when reported as a continuous outcome. Biologics may result in increased histological improvement when reported as a dichotomous outcome, but this is uncertain when reported as a continuous outcome when compared to placebo. Biologics may increase endoscopic improvement when measured as a dichotomous outcome, but this is uncertain when measured as a continuous outcome. Withdrawals due to adverse events as a dichotomous outcome may occur as frequently when biologics are compared to placebo.

In the study comparing cromolyn sodium to placebo, cromolyn sodium may lead to clinical symptom and histological improvement, measured as a continuous outcome. There may be no difference between cromolyn sodium and placebo in withdrawals due to adverse events and serious adverse events. Other outcomes were not reported.

In the study comparing PGD2R antagonist to placebo, we could not reach any conclusions for any primary outcome or serious adverse events due to the very low certainty of the results. Other secondary outcomes were not reported.

In the study comparing swallowed fluticasone to oral prednisone, we could not reach any conclusions for any primary outcome, or serious or total adverse events, due to the very low certainty of the results. Quality of life was not reported.

In the study comparing oral viscous budesonide to swallowed fluticasone there may be little to no difference for all primary

Medical treatment of eosinophilic esophagitis (Review)

outcomes, or serious and total adverse events. Quality of life was not reported.

In the two studies comparing esomeprazole to fluticasone we cannot draw any conclusions for clinical and histological improvement, withdrawals due to adverse events, or serious and total adverse events, due to the very low certainty of the results. Endoscopic improvement and quality of life were not reported.

In the study comparing a one-food elimination diet to a four-food elimination diet we cannot draw any conclusions for any outcome, due to the very low certainty of the results.

In the study comparing a one-food elimination diet to a six-food elimination diet we cannot draw any conclusions for any outcome, due to the very low certainty of the results.

In the study comparing a four-food elimination diet with omeprazole to omeprazole we cannot draw any conclusions for histological improvement or withdrawals due to adverse events, due to the very low certainty of the results. No other outcomes were reported.

In the study comparing a four-food elimination diet with amino acid formula to a four-food elimination diet, we could not reach any conclusions for endoscopic or histological improvement, or serious or total adverse events, due to the very low certainty of the results. Clinical improvement and quality of life were not reported.

In the study comparing nebulized swallowed budesonide to viscous swallowed budesonide we cannot draw any conclusions for clinical and histological improvement, withdrawals due to adverse events, or serious and total adverse events, due to the very low certainty of the results. Endoscopic improvement and quality of life were not reported.

In the study comparing Viaskin milk patch to placebo, Viaskin milk patch may result in little to no difference for all outcomes.

In the study comparing leukotriene receptor antagonist to placebo for maintenance of remission, we cannot draw any conclusions for clinical improvement, withdrawals due to adverse events, or serious and total adverse events, due to the very low certainty of the results. Other outcomes were not reported.

In the study comparing a low dose of the biologic mepolizumab (0.55 mg/kg) to a medium dose (2.5 mg/kg) and to a high dose (10 mg/kg), we cannot draw any conclusions for all combinations of comparisons, for histological improvement, withdrawals due to adverse events, or serious and total adverse events, due to the very low certainty of the results. Other outcomes were not reported.

In the study comparing a six-food elimination diet to swallowed fluticasone to swallowed budesonide and to oral viscous budesonide, no meta-analyses were possible for any combination of comparisons. We cannot draw any conclusions for histological improvement due to the very low certainty of the results. Other outcomes were not reported.

Overall completeness and applicability of evidence

In a number of our primary outcomes in key groups of treatments, high- and moderate-certainty evidence has been synthesized. However, there are several issues with the overall evidence from

the included studies, which limits our ability to draw convergent conclusions to inform clinical practice and decision-making.

There are limited studies specifically in children, often with high clinical and methodological heterogeneity. Most studies used mixed groups of adolescents and adults, but the data were combined. This not only limits the applicability to younger children, but it limits the completeness of the evidence in the adolescent age group.

The first primary outcome was clinical symptom improvement as defined by the individual study, but the assessments varied widely, with a lack of consistent validated, patient-reported outcomes. This limited the scope for meta-analysis and in turn reduced certainty, particularly in the pediatric setting. The limited data set in children makes it difficult to determine whether a lack of efficacy for these outcomes reflects issues with different symptomatology, the validity and reliability of such measures or instruments in children, or underlying poor efficacy. A similar point can be made about the adult studies. This is a key clinical question across all patient ages and so limits the applicability of these outcomes. Ideally, a single validated clinical outcome would be used across studies in order to facilitate data pooling and comparison of results.

There were also a range of therapies used within each of the two main groups seen, corticosteroids and biologics. The subgroup analysis does suggest that certain specific subclasses of medications within each of these groups have higher efficacy on a range of outcomes, but as the numbers of studies reduces substantially in these analyses, the certainty of the evidence is also lower. This is another key clinical area and until further targeted studies increase certainty, it represents an area of the evidence that is incomplete.

Histologic outcomes were also heterogenous, with different scoring systems, different thresholds, and the use of both dichotomous and continuous outcomes. This limits the scope for analysis and as such the certainty of evidence. Debate remains in the field as to which dichotomous histologic threshold is the preferred outcome and there may be differences in use of such a threshold. There is good but not universal agreement on a cut-off threshold in clinical practice (i.e. < 15 eos/hpf in an appropriate clinical setting) or per FDA guidance (≤ 6 eos/hpf), or if so-called "deep remission" (≤ 1 eos/hpf) is desired. Additionally, while the eosinophil count has traditionally been the biomarker of interest for response assessment in clinical practice and trials, and is also the most visually evident and readily available one to use, there is growing recognition that it may not fully reflect the underlying disease process. As such, the EoE Histologic Scoring System (EoE-HSS; Collins 2017) has been used in some recent trials as a secondary or exploratory endpoint, but could not be included as an outcome in the present study. Future iterations of this review will hopefully be able to assess this outcome in more detail.

Reporting of endoscopic measures was less common and had similar impacts, limiting the completeness of evidence in this area. Quality of life was sparsely reported and represents another incomplete area of evidence.

The prior use of proton pump inhibitor (PPI) therapy before recruitment and the concomitant use of such therapy varied between studies. This clinical and methodological source of heterogeneity reduces the applicability of the findings.

This is a rapidly moving field with study designs changing constantly, moving targets for outcome measures, concomitant PPI use, and thresholds of treatment success. For many current studies prior PPI failure is no longer necessary at study inclusion. Issues like these further increase heterogeneity and reduce applicability.

The reporting of adverse events was inconsistent across the included studies, with a lack of uniformity on what constitutes an adverse event and a serious adverse event. While the outcome of withdrawal is a much more objective measure in this context, it can be argued that this is of less interest to patients and therefore does limit the application of these data.

Finally, the complex clinical and methodological heterogeneity issues limit the scope for meaningful subgroup analysis of key factors, such as gender, age, dosage, and type of corticosteroid or biologic, and extent of disease, ultimately limiting the completeness of the evidence.

Quality of the evidence

The certainty of the evidence for a number of primary outcomes comparing corticosteroids and biologics to placebo was high and moderate, but most other outcomes and comparisons were of low certainty, primarily due to inconsistency and issues with risk of bias. A number of the inconsistency issues were explained in subgroup analysis as due to age (corticosteroids) and mechanism of agent (biologics), but these investigations in turn increased imprecision due to smaller participant numbers.

Imprecision was seen throughout due to the pervasive issues with heterogeneity in the specific outcomes measures used, as already discussed above. Whilst the homogeneous deployment of outcome measures would increase the scope for analysis and in turn reduce issues with imprecision, there were a number of studies with very low participant numbers. We have previously published work highlighting how common this type of difficulty in reporting sample size estimation is (Iheozor-Ejiofor 2021), with a need for adequate sample size calculations using published resources (Gordon 2021).

Risk of bias within the primary studies was low for all judged criteria in a little under half of the studies, with the remainder exhibiting issues in a number of the areas.

Potential biases in the review process

Gaps in information to judge risk of bias were pervasive, as discussed above. Given the contemporaneous nature of the evidence, with an exponential increase in studies since the last published version of this review, the review team considered it prudent and likely fruitful to reach out to primary authors to request clarification or additional information. Many did respond and as such judgments could be changed from those we made based on the published study reports, this could be considered a source of bias. Conversely, for those where no response was received, the judgments are based on the published forms of the studies.

The team aimed to include data that may become available in future updates, but this could represent a source of bias in the review, with 23 ongoing studies identified in the review process. Conversely, the use of such unpublished data can also be seen as a source of bias.

We are aware of the possibility that industry funding may affect the validity of the results. Funding from manufacturing companies or any conflicts of interests from both the primary studies and the review team have been reported.

For the main comparisons within this review, there were potential biases within the extraction and analysis process. When multiple outcomes were reported, we made decisions to report those that were validated and homogenous, but this may have introduced bias. There were a number of circumstances when calculations were needed to convert measures such as standard error to standard deviation or to convert units for histological outcomes. These have all been reported within the review, but could introduce bias.

Finally, the significant clinical heterogeneity remained a challenge when using our pre-planned subgroups. We made consensus decisions on the classification of particular subgroup characteristics, but again this is a potential source of bias.

There was repeated heterogeneity in clinical, histologic, and endoscopic outcomes. Additionally, subgroups of children compared to adults, co-therapy especially with proton pump inhibitors, differences in drug delivery, differences in dosage, differences in frequency, and differences in mechanism were all identified as potential biases.

With varying and occasionally multiple histological thresholds reported in the studies, the analysis did not contain a single dichotomous histological threshold. To minimize bias and subjectivity, we elected to include all histological thresholds with most common being < 15 , ≤ 6 , and ≤ 1 . In addition, we performed sensitivity analysis amongst the various histological thresholds, but the decision to include all such thresholds in a primary analysis could be considered source of bias.

Agreements and disagreements with other studies or reviews

In 2010, a previous version of this review was published with only three randomized controlled trials (RCTs). As of 2022, the number of published RCTs for pediatric and adult eosinophilic esophagitis that meet our inclusion criteria has grown to 41, with approximately half completed in the past five years. Given the rapid pace of clinical trial publications for eosinophilic esophagitis and changing outcome metrics, prior publications appear outdated, even those as recent as 2020. As an example, a recent American Gastroenterological Association (AGA) technical review of eosinophilic esophagitis management reported nine corticosteroid trials (compared to 14) and four biologic clinical trials (compared to nine) (Rank 2020). Another example is the European guidelines, which included even fewer RCTs as they were published three years earlier (Lucendo 2017). The AGA technical review clearly employed GRADE and when RCTs were synthesized, the broad findings were similar, although the certainty broadly increased in our review in key highly studied areas (biologics and non-systemic steroids) due to enhanced quality of reporting and precision (Rank 2020).

A recent UK joint adult and pediatric national consensus guideline, which considered a similar body of evidence, did state alignment with GRADE methodology (Dhar 2022). However, no technical review details are presented and in several key areas of

management, the GRADE of evidence included in this review does not match what is reported. For example, several forms of diet therapy with different comparators are noted to have moderate or low GRADE certainty of evidence in their guideline. However, all of them are very low-certainty GRADE outcomes in our review for all primary outcome measures. It is difficult to explain this difference as the detailed technical review information is missing.

The focus of prior reviews has been primarily on histologic outcomes and adverse events. Governmental pharmaceutical regulatory agencies have emphasized that outcomes for eosinophilic esophagitis must focus on patient-reported symptom outcomes and the newer assessments of esophageal function. This current Cochrane Review includes dichotomous and continuous outcomes for symptom, histologic, and endoscopic outcomes.

AUTHORS' CONCLUSIONS

Implications for practice

The evidence from this review demonstrates that for induction of remission, corticosteroids improve histologic outcomes (high certainty) and that biologic anti-IL-13 and anti-IL-4r therapies may improve clinical outcomes (low to moderate certainty). Eleven studies included children up to 18 years and 30 studies included adolescents and adults.

Corticosteroid therapy compared to placebo may lead to slightly better clinical improvement, as a dichotomous outcome (low certainty), leads to a large histological improvement (high certainty, number needed to treat for an additional beneficial outcome (NNTB) = 3), and may lead to fewer adverse event withdrawals (low certainty). Biologic anti-IL-5 therapy may result in little to no difference in clinical improvement (low certainty), and may lead to slightly better histological improvement (low certainty). Anti-IL-13 and anti-IL-4r therapy may lead to slightly better clinical improvement (low certainty) and may result in better histological improvement (moderate certainty, NNTB = 3). For anti-sialic acid binding Ig-like lectin 8 therapy, compared to placebo, clinical outcomes could not be reported due to incomplete published data; for histologic dichotomous outcomes anti-sialic acid binding Ig-like lectin 8 therapy may lead to slightly better improvement (low certainty). For anti-IgE compared to placebo, no conclusions can be made regarding clinical improvement (very low certainty). In studies comparing cromolyn sodium or Viaskin milk patch to placebo there may be no difference in clinical improvement, histologic improvement, or adverse event withdrawals (low certainty).

We could not draw conclusions about clinical improvement, histological improvement, or adverse event outcomes for the following: active comparator studies with PGD2R antagonist OC000459 versus placebo, esomeprazole versus fluticasone, swallowed fluticasone versus oral prednisone, nebulized swallowed budesonide versus swallowed viscous budesonide, oral viscous budesonide versus swallowed fluticasone, anti-IL-5 (10 mg/kg versus 0.55 mg/kg), anti-IL-5 (2.5 mg/kg versus 0.55 mg/kg), anti-IL-5 (10 mg/kg versus 2.5 mg/kg), a one-food elimination diet versus a four-food elimination diet, a one-food elimination diet versus a six-food elimination diet, or a four-food elimination diet with an amino acid formula versus a four-food elimination diet (all low- or very low-certainty evidence).

The evidence from this review demonstrates that for maintenance of remission, corticosteroids probably result in histological improvement (moderate certainty), but no other conclusions can be drawn (very low certainty). No conclusions can be made regarding leukotriene receptor antagonists in achieving maintenance of remission (very low certainty).

There were no clinical trials that compared either proton pump inhibitor (PPI) medication or dietary elimination therapies to a placebo for induction or maintenance of remission.

Implications for research

As heterogeneity in reporting of outcomes and the thresholds for specific outcomes was pervasive in the evidence base, the use of validated tools and standardized thresholds for success is key for future research. These should align with regulatory guidelines that employ such defined and publicly available homogenous outcomes in all key areas, including histologic outcome systems (Collins 2017), validated patient-reported outcomes (PROs) of symptoms and health-related quality of life (Hudgens 2017), validated endoscopic outcomes (Ma 2022a), and validated assessments of esophageal function (e.g. esophageal distensibility testing; Donnan 2020). This could build on initial work done by the COREOS group, which proposed an initial consensus core outcomes set for eosinophilic esophagitis (Ma 2022b). Clinical outcome results for children are more heterogenous than adults, and pediatric patient-reported outcomes are an important area for future research and development. Studies should follow clear reporting guidelines in line with the CONSORT statement to reduce the risk of bias and enhance the certainty of the evidence base as a whole, regardless of findings.

There is a clear direction for future research of head-to-head comparisons of corticosteroids and biologics. Randomized, placebo-controlled clinical trials for proton pump inhibitor medications and dietary elimination are also needed. Future direction can also include personalized medicine clinical trials with precision medicine-focused therapies compared to conventional therapies.

In accordance with the findings of other recently published reviews, clinical guidelines, and regulatory statements within eosinophilic esophagitis, large-scale, long-term measurement of outcomes to include safety and adverse events is needed.

ACKNOWLEDGEMENTS

Editorial contributions

Cochrane Gut supported the authors in the development of this systematic review. The following people conducted the editorial process for this article:

Sign-off Editor (final editorial decision): Michael Brown, Michigan State University College of Human Medicine, USA; **Managing Editor** (selected peer reviewers, collated peer reviewer comments, provided editorial guidance to authors, edited the article): Marwah Anas El-Wegoud, Cochrane Central Editorial Service; **Editorial Assistant** (conducted editorial policy checks and supported editorial team): Lisa Wydrzynski, Cochrane Central Editorial Service; **Copy Editor** (copy editing and production): Jenny Bellorini, Cochrane Central Production Service

Peer reviewers (provided comments and recommended an editorial decision): Alexander Link, Head of Molecular Gastroenterology and microbiota-associated diseases; Otto-von-Guericke University Magdeburg, Germany (**clinical review**); Edoardo Vincenzo Savarino; Department of Surgery, Oncology and Gastroenterology, University of Padua, Padua, Italy (**clinical**

review); Alfretta Vanderheyden, consumer reviewer, UGPD (**consumer review**); Nuala Livingstone, Cochrane Evidence Production and Methods Department (**methods review**); Jo Platt, Information Specialist, Cochrane GNOC Review Group (**search review**). One additional peer reviewer provided clinical peer review but chose not to be publicly acknowledged.

REFERENCES

References to studies included in this review

Alexander 2012 {published data only}

Alexander J, Jung KW, Arora A, Enders F, Francis D, Kita H, et al. A randomized placebo-controlled trial of swallowed fluticasone as treatment of eosinophilic esophagitis 2010 ACG auxiliary award. *American Journal of Gastroenterology* 2010;**105** Suppl 1:S21-2. [DOI: [10.1038/ajg.2010.320-1](https://doi.org/10.1038/ajg.2010.320-1)]

* Alexander J, Jung KW, Arora A, Enders F, Katzka D, Kephart D, et al. Swallowed fluticasone improves histologic but not symptomatic response of adults with eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology* 2012;**10**(7):742-9. [DOI: [10.1016/j.cgh.2012.03.018](https://doi.org/10.1016/j.cgh.2012.03.018)] [PMID: 22475741]

NCT00275561. Topical steroid treatment for eosinophilic esophagitis [A randomized placebo-controlled trial of swallowed fluticasone in treatment of eosinophilic esophagitis]. <https://clinicaltrials.gov/ct2/show/NCT00275561> (first received 12 January 2006).

Alexander 2017 {published data only}

Alexander J, Karthik R, Enders F, Geno D, Kryzer L, Mara K, et al. A randomized placebo-controlled trial of montelukast in maintenance treatment of eosinophilic esophagitis. *Gastroenterology* 2016;**150**(4 Suppl 1):S19.

* Alexander J, Ravi K, Enders F, Geno D, Kryzer L, Mara K, et al. Montelukast does not maintain symptom remission after topical steroid therapy for eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology* 2017;**15**(2):214-21. [DOI: [10.1016/j.cgh.2016.09.013](https://doi.org/10.1016/j.cgh.2016.09.013)] [PMID: 27650328]

NCT00511316. Trial of montelukast in eosinophilic esophagitis [A randomized placebo-controlled trial of montelukast in maintenance therapy of asymptomatic eosinophilic esophagitis]. <https://clinicaltrials.gov/ct2/show/NCT00511316> (first received 3 August 2007).

Assa'ad 2011 {published data only}

Aceves S, Gupta S, Justinich C, Assa'ad A, Heath A, Perschy T, et al. Clinical characteristics of pediatric eosinophilic esophagitis subjects with inadequate response to or intolerance of prior therapies receiving mepolizumab in a multicenter clinical trial. *Journal of Allergy and Clinical Immunology* 2010;**125**(2 Suppl 1):AB159. [DOI: [10.1016/j.jaci.2009.12.625](https://doi.org/10.1016/j.jaci.2009.12.625)]

Aceves S, Newbury R, Chen D, Broide D, Bastian JF, Dohil R. Resolution of esophageal remodeling following oral viscous budesonide (OVB) therapy for eosinophilic esophagitis (EoE). *Gastroenterology* 2010;**138**(5 Suppl 1):S123.

Assa'ad A, Collins M, Aceves S, Gupta S, Drouin E, Jurgensen C, et al. The pharmacodynamic effects of mepolizumab, a humanized monoclonal antibody against IL-5, in pediatric patients with eosinophilic esophagitis: A randomized, double-blind, controlled clinical trial. *Journal of Allergy and Clinical Immunology* 2010;**125**(2 Suppl 1):AB129. [DOI: [10.1016/j.jaci.2009.12.509](https://doi.org/10.1016/j.jaci.2009.12.509)]

* Assa'ad AH, Collins MH, Gupta SK, Thomson M, Heath AT, Smith DA, et al. An antibody against IL-5 reduces numbers of esophageal intraepithelial eosinophils in children with eosinophilic esophagitis. *Gastroenterology* 2011;**141**(5):1593-604. [DOI: [10.1053/j.gastro.2011.07.044](https://doi.org/10.1053/j.gastro.2011.07.044)] [PMID: 21835135]

Collins M, Heath A, Jurgensen C. Effect of a monoclonal antibody (mepolizumab) against Interleukin-5 (IL-5) in reducing eosinophilic inflammation in children with eosinophilic esophagitis (EoE). *Laboratory Investigation* 2010;**90**(2):291. [DOI: [10.1038/labinvest.2009.151](https://doi.org/10.1038/labinvest.2009.151)]

Gupta S, Justinich C, Collins M, Assa'ad A, Kobak G, Gold B, et al. PK and PD of mepolizumab in pediatric subjects with eosinophilic esophagitis: A randomized, double-blind, controlled clinical trial. *Journal of Pediatric Gastroenterology and Nutrition* 2009;**49** Suppl 1:E65-6. [DOI: [10.1097/MPG.0b013e3181c64b4a](https://doi.org/10.1097/MPG.0b013e3181c64b4a)]

Gupta S, Kita H, Collins M, Perschy T, Heath A, Jurgensen C. Utility of eosinophil-derived neurotoxin as a biomarker in eosinophilic esophagitis: results from a randomized, double-blind, controlled trial of mepolizumab in pediatric patients. *Journal of Allergy and Clinical Immunology* 2010;**125**(2 Suppl 1):AB234. [DOI: [10.1016/j.jaci.2009.12.913](https://doi.org/10.1016/j.jaci.2009.12.913)]

NCT00358449. Intravenous mepolizumab in children with eosinophilic esophagitis [A randomized, double-blind, parallel group clinical trial to assess safety, tolerability, pharmacokinetics, and pharmacodynamics of intravenous mepolizumab (SB240563)(0.55mg/kg, 2.5mg/kg or 10mg/kg) in pediatric subjects with eosinophilic esophagitis, aged 2 to 17 years (study MEE103219)]. <https://clinicaltrials.gov/ct2/show/NCT00358449> (first received 31 July 2006).

Newbury RO, Beppu LY, Dohil R, Otani IM, Anilkumar AA, Bhagat M, et al. Anti-IL-5 therapy reduces mast cell and IL-9 cell numbers in pediatric patients with eosinophilic esophagitis. *Journal of Allergy and Clinical Immunology* 2013;**131**(6):1576. [DOI: [10.1016/j.jaci.2013.02.042](https://doi.org/10.1016/j.jaci.2013.02.042)]

Otani I M, Bhagat M, Newbury R O, Dohil R, Broide D H, Aceves S S. The effect of anti-IL-5 therapy on esophageal mastocytosis in pediatric eosinophilic esophagitis. *Journal of Allergy and Clinical Immunology* 2012;**129**(2 Suppl 1):AB202. [DOI: [10.1016/j.jaci.2011.12.180](https://doi.org/10.1016/j.jaci.2011.12.180)]

Bhardwaj 2017 {published data only}

* Bhardwaj N, Ishmael F, Lehman E, Bethards D, Ruggiero F, Ghaffari G. Effect of topical beclomethasone on inflammatory markers in adults with eosinophilic esophagitis: a pilot study. *Allergy Rhinol (Providence)* 2017;**8**(2):85-94. [DOI: [10.2500/ar.2017.8.0202](https://doi.org/10.2500/ar.2017.8.0202)] [PMID: 28583232]

Ghaffari G. A randomized double blind placebo controlled crossover study of the effect of swallowed beclomethasone dipropionate on inflammatory markers in adult patients with eosinophilic esophagitis: a pilot study. *Annals of Allergy, Asthma and Immunology* 2012;**109**(5 Suppl):A19-20. [DOI: [10.1016/j.anai.2012.09.018](https://doi.org/10.1016/j.anai.2012.09.018)]

Medical treatment of eosinophilic esophagitis (Review)

Ghaffari G. A randomized double blind placebo controlled crossover study of the effect of swallowed beclomethasone dipropionate on inflammatory markers in adult patients with eosinophilic esophagitis: a pilot study. *Annals of Allergy, Asthma and Immunology* 2011;**107**(5 Suppl 1):A99. [DOI: [10.1016/j.anai.2011.09.019](https://doi.org/10.1016/j.anai.2011.09.019)]

NCT01016223. A randomized double blind placebo controlled study of the effect of swallowed beclomethasone dipropionate on inflammatory markers in adult patients with eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT01016223> (first received 19 November 2009).

Butz 2014 {published data only}

* Butz B, Wen T, Gleich G, Furuta G, Spergel J, King E, et al. Efficacy, dose reduction, and resistance to high-dose fluticasone in patients with eosinophilic esophagitis. *Gastroenterology* 2014;**147**(2):324-33.e5. [DOI: [10.1053/j.gastro.2014.04.019](https://doi.org/10.1053/j.gastro.2014.04.019)] [PMID: 24768678]

NCT00426283. A study of Flovent in patients with eosinophilic esophagitis [A double blinded, randomized trial of swallowed 1760mcg fluticasone propionate versus placebo in the treatment of eosinophilic esophagitis]. <https://clinicaltrials.gov/ct2/show/NCT00426283> (first received 24 January 2007).

Clayton 2014 {published data only}

* Clayton F, Chen X, Emerson L, Fang J, Vinson L, Cox K, et al. Eosinophilic esophagitis in adults is associated with IgG4 and not mediated by IgE. *Gastroenterology* 2014;**147**(3):602-9. [DOI: [10.1053/j.gastro.2014.05.036](https://doi.org/10.1053/j.gastro.2014.05.036)] [PMID: 24907494]

Fang J, Hilden K, Gleich G, Emerson L, Ogorman M, Lowichik A, et al. A pilot study of the treatment of eosinophilic esophagitis with omalizumab. *Gastroenterology* 2011;**140**(5 Suppl 1):S235. [DOI: [10.1016/S0016-5085%2811%2960951-7](https://doi.org/10.1016/S0016-5085%2811%2960951-7)]

NCT00123630. A pilot study of the treatment of eosinophilic esophagitis with omalizumab. <https://clinicaltrials.gov/ct2/show/NCT00123630> (first received 25 July 2005).

Dellon 2012 {published data only}

Dellon E, Sheikh A, Speck O, Woodward K, Whitlow A, Hores J, et al. Nebulized/swallowed vs viscous/swallowed budesonide for treatment of eosinophilic esophagitis: results from a randomized, open-label, clinical trial. *Gastroenterology* 2012;**142**(5 Suppl 1):S180.

* Dellon E, Sheikh A, Speck O, Woodward K, Whitlow A, Hores J, et al. Viscous topical is more effective than nebulized steroid therapy for patients with eosinophilic esophagitis. *Gastroenterology* 2012;**143**(2):321-4. [DOI: [10.1053/j.gastro.2012.04.049](https://doi.org/10.1053/j.gastro.2012.04.049)] [PMID: 22561055]

Dellon E, Sheikh A, Whitlow A, Ivanovic M, Chau A, Woodward K, et al. Mucosal contact time determines histologic effectiveness of oral budesonide preparations in eosinophilic esophagitis: a randomized trial using scintigraphy. *Gastroenterology* 2012;**142**(5 Suppl 1):S180.

NCT00961233. Topical budesonide treatment for eosinophilic esophagitis (EoE) [Topical budesonide treatment for eosinophilic esophagitis (eoe): esophageal medication

distribution, treatment response, and effect on the adrenal axis]. <https://clinicaltrials.gov/ct2/show/NCT00961233> (first received 18 August 2009).

Dellon 2017 {published data only}

Collins M, Dellon E, Hirano I, Katzka D, Mukkada V, Williams J, et al. Baseline characterization of adolescent and adult patients with eosinophilic esophagitis (EOE) using the eoe histology scoring system: analyses from a randomized, placebo-controlled trial. *Journal of Pediatric Gastroenterology and Nutrition* 2019;**69** Suppl 2:S26. [DOI: [10.1097/MPG.0000000000002518](https://doi.org/10.1097/MPG.0000000000002518)]

Collins M, Dellon E, Hirano I, Katzka D, Williams J, Lan L, et al. Baseline characterization of patients with eosinophilic esophagitis (EOE) using the eoe histology scoring system and correlations with endoscopic and symptom outcomes: Analyses from a randomized, placebo-controlled trial. *American Journal of Gastroenterology* 2019;**114** Suppl:S229-30. [DOI: [10.14309/01.ajg.0000591100.32919.d0](https://doi.org/10.14309/01.ajg.0000591100.32919.d0)]

Collins M, Dellon E, Katzka D, Hirano I, Williams J, Lan L. Budesonide oral suspension significantly improves eosinophilic esophagitis histology scoring system results: analyses from a 12-week, phase 2, randomized, placebo-controlled trial. *American Journal of Surgical Pathology* 2019;**43**(11):1501-9. [DOI: [10.1097/PAS.0000000000001361](https://doi.org/10.1097/PAS.0000000000001361)] [PMID: 31498177]

Dellon E, Katzka D, Collins M, Gupta S, Lan L, Williams J, et al. Safety and efficacy of budesonide oral suspension maintenance therapy in patients with eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology* 2019;**17**(4):666-73. [DOI: [10.1016/j.cgh.2018.05.051](https://doi.org/10.1016/j.cgh.2018.05.051)] [PMID: 29902649]

Dellon E, Katzka D, Collins M, Hamdani M, Gupta S, Hirano I. Safety and efficacy of oral budesonide suspension for maintenance therapy in eosinophilic esophagitis: Results from a prospective open-label study of adolescents and adults. *Gastroenterology* 2016;**150**(4 Suppl 1):S188.

* Dellon E, Katzka D, Collins M, Hamdani M, Gupta S, Kagalwalla A, et al. Budesonide oral suspension improves symptomatic, endoscopic, and histologic parameters compared with placebo in patients with eosinophilic esophagitis. *Gastroenterology* 2017;**152**(4):776-86. [DOI: [10.1053/j.gastro.2016.11.021](https://doi.org/10.1053/j.gastro.2016.11.021)] [PMID: 27889574]

Dellon E, Katzka D, Collins M, Hamdani M, Hill M, Hirano I. Pain with swallowing is a frequent and under-appreciated symptom in eosinophilic esophagitis: analysis from a multicenter randomized placebo-controlled trial. *Gastroenterology* 2016;**150**(4 Suppl 1):S662-3.

Dellon E, Katzka D, Collins M, Hirano I. Oral budesonide suspension significantly improves dysphagia and esophageal eosinophilia: results from a multicenter randomized double-blind placebo-controlled trial in adolescents and adults with eosinophilic esophagitis. *Gastroenterology* 2015;**148**(4 Suppl 1):S157.

Dellon E, Katzka D, Collins M, Hirano I. The dysphagia symptom questionnaire is responsive and valid for measuring patient-reported outcomes in eosinophilic esophagitis: Analysis

Medical treatment of eosinophilic esophagitis (Review)

from a multicenter randomized placebo controlled trial. *Gastroenterology* 2015;**148**(4 Suppl 1):S792.

Dellon ES, Collins MH, Katzka DA, Hudgens S, Lan L, Williams J, et al. Improvements in dysphagia and pain with swallowing in patients with eosinophilic esophagitis receiving budesonide oral suspension. *Clinical Gastroenterology and Hepatology* 2021;**19**(4):699-706. [DOI: [10.1016/j.cgh.2020.03.060](https://doi.org/10.1016/j.cgh.2020.03.060)] [PMID: 32272243]

Gupta S, Mukkada V, Dellon E, Gold B, Collins M, Katzka D, et al. Efficacy and safety of budesonide oral suspension in a pediatric population: pooled data from a phase 2 and phase 3 trial in patients with eosinophilic esophagitis. *Journal of Allergy and Clinical Immunology* 2021;**147**(2 Suppl):AB91. [DOI: [10.1016/j.jaci.2020.12.348](https://doi.org/10.1016/j.jaci.2020.12.348)]

Hirano I, Dellon E, Collins M, Williams J, Lan L, Katzka D. Clinical features at baseline cannot predict symptom response to placebo in patients with eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology* 2019;**17**(10):2126-8. [DOI: [10.1016/j.cgh.2018.11.045](https://doi.org/10.1016/j.cgh.2018.11.045)] [PMID: 30502508]

Hirano I, Katzka D, Collins M, Dellon E. Randomized, double-blind, placebo controlled trial demonstrates the efficacy of oral budesonide suspension in improving endoscopically identified esophageal abnormalities in eosinophilic esophagitis. *Gastroenterology* 2015;**148**(4 Suppl 1):S29.

Hirano I, Williams J, Collins M, Dellon E, Uddin S, Katzka D. Clinical features at baseline are not clearly associated with symptomatic placebo response in adolescents and adults with eosinophilic esophagitis during a placebo run-in period of a double-blind, randomized, controlled trial of budesonide oral suspension. *Gastroenterology* 2017;**152**(5 Suppl 1):S854.

NCT01642212. OBS in adolescent and adults with EoE: a phase II, randomized, double-blind, placebo controlled, study with an open label extension. <https://clinicaltrials.gov/ct2/show/NCT01642212> (first received 17 July 2012).

Dellon 2019 {published data only}

Cotton C, Woosley J, Moist S, McGee S, Shaheen N, Dellon E. Determination of an endoscopic response threshold for the eosinophilic esophagitis endoscopic reference score: analysis of a randomized comparative clinical trial of topical steroids. *Gastroenterology* 2020;**158**(6 Suppl 1):S819-20. [DOI: [10.1016/S0016-5085\(20\)32728-1](https://doi.org/10.1016/S0016-5085(20)32728-1)]

Cotton C, Woosley J, Moist S, McGee S, Shaheen N, Dellon E. Dysphagia in eosinophilic esophagitis improves with topical steroid treatment independent of improvement in endoscopic and histologic measures of mucosal inflammation: analysis of a randomized comparative effectiveness clinical trial. *Gastroenterology* 2020;**158**(6 Suppl 1):S-820. [DOI: [10.1016/S0016-5085%2820%2932729-3](https://doi.org/10.1016/S0016-5085%2820%2932729-3)]

Dellon E, McGee S, Moist S, Shaheen N, Woosley J. Utility of major basic protein, eotaxin-3, and mast cell tryptase staining for prediction of response to topical steroid treatment in eosinophilic esophagitis: analysis of a randomized, double-blind, double dummy clinical trial. *Diseases of the Esophagus* 2020;**33**(6):1-8. [DOI: [10.1093/dote/daaa003](https://doi.org/10.1093/dote/daaa003)] [PMID: 32193541]

* Dellon E, Woosley J, Arrington A, McGee S, Covington J, Moist S, et al. Efficacy of budesonide vs fluticasone for initial treatment of eosinophilic esophagitis in a randomized controlled trial. *Gastroenterology* 2019;**157**(1):65-73. [DOI: [10.1053/j.gastro.2019.03.014](https://doi.org/10.1053/j.gastro.2019.03.014)] [PMID: 30872104]

Dellon E, Woosley J, Arrington A, McGee S, Covington J, Moist S, et al. Rapid recurrence of symptoms, endoscopic findings, and esophageal eosinophilia without maintenance therapy after successful initial treatment: results from the observation phase of a randomized, double-blind, double dummy clinical trial. *Gastroenterology* 2019;**156**(6 Suppl 1):713-4. [DOI: [10.1016/S0016-5085\(19\)38720-7](https://doi.org/10.1016/S0016-5085(19)38720-7)]

Dellon E, Woosley J, McGee S, Moist S, Shaheen N. Utility of major basic protein, eotaxin-3, and mast cell tryptase staining for prediction of response to topical steroid treatment in eosinophilic esophagitis: analysis of a randomized, double-blind, double-dummy clinical trial. *Gastroenterology* 2020;**158**(6 Suppl 1):S837-8. [DOI: [10.1016/S0016-5085\(20\)32771-2](https://doi.org/10.1016/S0016-5085(20)32771-2)]

Dellon E, Woosley J, Arrington A, McGee S, Covington J, Moist S, et al. Rapid recurrence of eosinophilic esophagitis activity after successful treatment in the observation phase of a randomized, double-blind, double-dummy trial. *Clinical Gastroenterology and Hepatology* 2020;**18**(7):1483-92. [DOI: [10.1016/j.cgh.2019.08.050](https://doi.org/10.1016/j.cgh.2019.08.050)] [PMID: 31499249]

Dellon ES, Woosley JT, Arrington A, McGee SJ, Covington J, Moist SE, et al. Oral viscous budesonide versus swallowed fluticasone inhaler for initial treatment of adolescents and adults with eosinophilic esophagitis: a randomized, double-blind, double-dummy clinical trial. *Gastroenterology* 2019;**156**(6 Suppl 1):72-3. [DOI: [10.1016/S0016-5085\(19\)36966-5](https://doi.org/10.1016/S0016-5085(19)36966-5)]

NCT02019758. Budesonide versus fluticasone for treatment of eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT02019758> (first received 24 December 2013).

Safroneeva E, Zwahlen M, Cotton C, Dellon E, Schoepfer A, Woosley J. Dilation modifies association between symptoms and esophageal eosinophilia in adult patients with eosinophilic esophagitis. *American Journal of Gastroenterology* 2020;**115**(12):2098-102. [DOI: [10.14309/ajg.0000000000000957](https://doi.org/10.14309/ajg.0000000000000957)] [PMID: 33009060]

Dellon 2021b {published data only}

* Dellon ES, Collins MH, Katzka DA, Mukkada VA, Falk GW, Morey R, et al. Long-term treatment of eosinophilic esophagitis with budesonide oral suspension. *Clinical Gastroenterology and Hepatology* 2021;**20**(7):P1488-98.E11. [DOI: [10.1016/j.cgh.2021.06.020](https://doi.org/10.1016/j.cgh.2021.06.020)]

Dellon ES, Collins MH, Katzka DA, Mukkada VA, Falk GW, Morey R, et al. Loss of histologic, symptom and endoscopic efficacy following treatment withdrawal in patients with eosinophilic esophagitis: a phase 3, randomized withdrawal study of budesonide oral suspension. *Gastroenterology* 2021;**160**(6):S-247-8.

NCT02736409. An extension study to evaluate maintenance of efficacy and long-term treatment effect of oral budesonide suspension (OBS) in adults and adolescents with eosinophilic

Medical treatment of eosinophilic esophagitis (Review)

esophagitis (EoE). <https://clinicaltrials.gov/ct2/show/NCT02736409> (first received 13 April 2016).

Dellon 2022 {published data only}

Dellon E, Rothenberg M, Bredenoord A, Lucendo A, Hirano I, Sun X et al. Dupilumab improves clinical, symptomatic, histologic, and endoscopic aspects of EoE up to 24 weeks: pooled results from parts A and B of phase 3 LIBERTY-EoE-TREET. *American Journal of Gastroenterology* 2022;**117**(Supplement):s387.

Dellon E, Rothenberg M, Hirano I, Chehade M, Bredenoord A, Spergel J, et al. Dupilumab improves health-related quality of life (HRQoL) and reduces symptom burden in patients with eosinophilic esophagitis (EoE): results from part A of a randomized, placebo-controlled three-part phase 3 study. *Journal of Allergy and Clinical Immunology* 2021;**147**(2 Suppl):AB91. [DOI: [10.1016/j.jaci.2020.12.347](https://doi.org/10.1016/j.jaci.2020.12.347)]

Dellon ES, Rothenberg ME, Bredenoord A, Collins MH, Hirano I, Chehade M, et al. Clinical and histological improvements with weekly dupilumab treatment in adult and adolescent patients with eosinophilic esophagitis at week 24: weekly and every 2 weeks; results from part b of the 3-part LIBERTY EoT TREET study. *Gastroenterology* 2022;**162**(7 Suppl):S-1391-2.

Dellon ES, Rothenberg ME, Bredenoord A, Collins MH, Hirano I, Chehade M, et al. Dupilumab improves histologic and clinical aspects of disease in adult and adolescent patients with eosinophilic esophagitis at week 24: results from part b of the 3-part LIBERTY EoT TREET Study. *Gastroenterology* 2022;**162**(7 Suppl):S-237-8.

Dellon ES, Rothenberg ME, Collins MH, Hirano I, Chehade M, Bredenoord A. A phase 3, randomized, 3-part study to investigate the efficacy and safety of dupilumab in adult and adolescent patients with eosinophilic esophagitis: results from part A. *American Journal of Gastroenterology* 2020;**115**:LB3.

Dellon ES, Rothenberg ME, Collins MH, Hirano I, Chehade M, Bredenoord A. Dupilumab efficacy and safety up to 52 weeks in adult and adolescent patients with eosinophilic esophagitis: results from part A and C of a randomized, placebo-controlled three-part, phase 3 LIBERTY EoE TREET Study. *United European Gastroenterology Journal* 2021;**9**(10):LB10. [DOI: <https://doi.org/10.1002/ueg2.12172>]

Dellon ES, Rothenberg ME, Collins MH, Hirano I, Chehade M, Bredenoord A. LIBERTY EoE TREET: Results from parts A and C of the phase 3, randomized, 3-part LIBERTY EoE TREET study to investigate the efficacy and safety of dupilumab in adult and adolescent patients with eosinophilic esophagitis up to 52-weeks. *American Journal of Gastroenterology* 2021;**116**:52.

Dellon ES, Rothenberg ME, Collins MH, Hirano I, Chehade M, Bredenoord AJ, et al. A phase 3, randomized, 3-part study to investigate the efficacy and safety of dupilumab in adult and adolescent patients with eosinophilic esophagitis: results from part A. In: <https://www.eventscribe.com/2020/ACG/fsPopup.asp?efp=REICSUFHUK02NDI2&PresentationID=783291&rnd=0.6661928&mode=7> American College of Gastroenterology, October 2020.

* Dellon ES, Rothenberg ME, Collins MH, Hirano I, Chehade M, Bredenoord AJ, et al. Dupilumab in adults and adolescents with eosinophilic esophagitis. *New England Journal of Medicine* 2022;**387**(25):2317-30.

EUCTR2018-000844-25-IT. A study to determine the efficacy and safety of dupilumab in adult and adolescent patients with eosinophilic esophagitis (EoE). <https://trialsearch.who.int/Trial2.aspx?TrialID=NCT03633617> (first received 24 September 2018).

NCT03633617. Study to determine the efficacy and safety of dupilumab in adult and adolescent patients With eosinophilic esophagitis (EoE) [A phase 3, randomized, 3-part study to investigate the efficacy and safety of dupilumab in adult and adolescent patients with eosinophilic esophagitis (EoE)]. <https://clinicaltrials.gov/ct2/show/NCT03633617> (first received 18 August 2018).

Dellon 2022a {published data only}

Dellon E, Falk G, Hirano I, Lucendo A, Schlag C, Schoepfer A, et al. An episode-based, patient-reported outcome measure of dysphagia accurately captures response to treatment with apt-1011 in a phase 2b, randomized, double-blind, placebo-controlled, clinical trial in adults with eosinophilic esophagitis. *Gastroenterology* 2020;**158**(6 Suppl 1):S817. [DOI: [10.1016/S0016-5085\(20\)32722-0](https://doi.org/10.1016/S0016-5085(20)32722-0)]

Dellon E, Lucendo A, Schlag C, Schoepfer A, Falk G, Richardson P, et al. Safety and efficacy of long-term treatment of EoE with fluticasone propionate orally disintegrating tablet (APT-1011): results from 40 weeks treatment in a phase 2b randomized double-blind placebo-controlled trial. *American Journal of Gastroenterology* 2020;**115** Suppl:S169-70. [DOI: [10.14309/01.ajg.0000703428.23468.41](https://doi.org/10.14309/01.ajg.0000703428.23468.41)]

Dellon E, Lucendo Villarin A, Schlag C, Schoepfer A, Knoop K, Richardson P, et al. Safety and efficacy of 4 dosage regimens of APT-1011: results from a phase 2B randomized double-blind placebo-controlled trial to select the best dosage regimen for further development. *United European Gastroenterology Journal* 2019;**7**(10):1420. [DOI: [10.1177/2050640619888859](https://doi.org/10.1177/2050640619888859)]

* Dellon ES, Lucendo AJ, Schlag C, Schoepfer AM, Falk GW, Eagle G, et al. Fluticasone Propionate Orally Disintegrating Tablet (APT-1011) for Eosinophilic Esophagitis: randomized Controlled Trial. *Clinical gastroenterology and hepatology* 2022;**20**(11):2485-2494.e15.

Dellon ES, Lucendo AJ, Schlag C, Schoepfer AM, Falk GW, Eagle G, et al. Fluticasone propionate orally disintegrating tablet (APT-1011) for eosinophilic esophagitis: randomized controlled trial. *Clinical Gastroenterology and Hepatology* 2022 Feb 15 [epub ahead of print]:S1542. [DOI: [10.1016/j.cgh.2022.02.013](https://doi.org/10.1016/j.cgh.2022.02.013)] [PMID: 35181572]

EUCTR2016-004749-10-BE. Study in subjects with eosinophilic esophagitis. <https://www.clinicaltrialsregister.eu/ctr-search/search?query=EUCTR2016-004749-10-BE> (first received 19 June 2016).

NCT03191864. Efficacy, safety, and pharmacokinetics of APT-1011 in subjects with eosinophilic esophagitis (EoE).

Medical treatment of eosinophilic esophagitis (Review)

<https://clinicaltrials.gov/ct2/show/NCT03633617> (first received 19 June 2017).

Turner R, Rohay JM, Sparling N, Whitsett J, Eagle G, Nezamis J, et al. Patient-reported symptom improvement following treatment with APT-1011 in patients with eosinophilic esophagitis as measured by the prose: results from flute, a phase 2b clinical trial. *American Journal of Gastroenterology* 2021;**116**(Suppl):S203.

Dellon 2022b {published and unpublished data}

Allakos. Allakos corporate update. <https://investor.allakos.com/static-files/74ad659a-8f45-4448-b1fb-97b647be1952> 2022.

* Dellon E, Chedhade M, Genta M, Leiman D, Peterson K, Spergel J. Results from KRYPTOS, a phase 2/3 study of lirentelimab (AK002) in adults and adolescents with EoE. *American journal of Gastroenterology* 2022;**117**(Suppl):S316.

De Rooij 2022 {published data only}

De Rooij W, Bredenoord A, Vlieg-Boerstra B, Warners M, Van Ampting M, Eussen S, et al. Effect of elemental nutrition added to four-food elimination in adult eosinophilic esophagitis patients: a randomized controlled trial. *United European Gastroenterology Journal* 2020;**8**(8S):56-7. [DOI: [10.1177/2050640620927344](https://doi.org/10.1177/2050640620927344)]

* De Rooij W, Vlieg-Boerstra B, Warners MJ, Van Ampting MTJ, Van Esch BCAM, Eussen SRBM, et al. Effect of amino acid-based formula added to four-food elimination in adult eosinophilic esophagitis patients: a randomized clinical trial. *Neurogastroenterology & Motility* 2022;**34**(7):e14291. [DOI: [10.1111/nmo.14291](https://doi.org/10.1111/nmo.14291)] [PMID: 34792264]

NTR6778. Effect of addition of elemental nutrition to an elimination diet on esophageal inflammation in adult eosinophilic esophagitis (EoE) patients: a randomized controlled trial. <http://www.who.int/trialsearch/Trial2.aspx?TrialID=NTR6778> (first received 20 October 2017).

de Rooij WE, Vlieg-Boerstra B, Warners MJ, Van Ampting MTJ, van Esch BCAM, Eussen SRBM, et al. Effect of amino acid-based formula added to four-food elimination in adult eosinophilic esophagitis patients: a randomized clinical trial. *Neurogastroenterology and Motility* 2022;**34**(7):e14291.

Dohil 2010 {published data only}

Dohil R, Newbury R, Fox L, Bastian J, Aceves S. A randomized, double-blind, placebo-controlled study of oral viscous budesonide (OVB) in children with eosinophilic esophagitis (EoE). *Gastroenterology* 2010;**138**(5 Suppl 1):S62-3.

* Dohil R, Newbury R, Fox L, Bastian J, Aceves S. Oral viscous budesonide is effective in children with eosinophilic esophagitis in a randomized, placebo-controlled trial. *American Journal of Gastroenterology* 2010;**139**(2):418-29. [DOI: [10.1053/j.gastro.2010.05.001](https://doi.org/10.1053/j.gastro.2010.05.001)] [PMID: 20457157]

NCT00638456. Use of topical budesonide in the treatment of eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT00638456> (first received 19 March 2008).

Gupta 2015 {published data only}

Collins M, Irani A, Ngo PD, Vitanza JM. Reduction of esophageal epithelial pathology and endoscopic abnormalities following treatment with oral budesonide suspension (OBS) in pediatric subjects with eosinophilic esophagitis (EoE). *Gastroenterology* 2011;**140**(5 Suppl 1):S237. [DOI: [10.1016/S0016-5085%2811%2960956-6](https://doi.org/10.1016/S0016-5085%2811%2960956-6)]

Gupta S, Collins M, Lewis J, Farber R. Efficacy and safety of oral budesonide suspension (OBS) in pediatric subjects with eosinophilic esophagitis (EoE): results from the double-blind, placebo-controlled PEER study. *Gastroenterology* 2011;**140**(5 Suppl 1):S179. [DOI: [10.1016/S0016-5085%2811%2960725-7](https://doi.org/10.1016/S0016-5085%2811%2960725-7)]

* Gupta S, Vitanza J, Collins M. Efficacy and safety of oral budesonide suspension in pediatric patients with eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology* 2015;**13**(1):66-76. [DOI: [10.1016/j.cgh.2014.05.021](https://doi.org/10.1016/j.cgh.2014.05.021)] [PMID: 24907502]

NCT00762073. Dose-ranging study of oral viscous budesonide in pediatrics with eosinophilic esophagitis [Oral viscous budesonide suspension (MB-7) in subjects with eosinophilic esophagitis: a randomized, placebo-controlled, dose-ranging study in children and adolescents]. <https://clinicaltrials.gov/ct2/show/NCT00762073> (first received 30 September 2008).

Heine 2019 {published data only}

ACTRN12613001210763. Effects of diet and omeprazole in patients with eosinophilic oesophagitis: a multi-centre, randomised controlled trial and investigation of novel disease mechanisms. <http://www.who.int/trialsearch/Trial2.aspx?TrialID=ACTRN12613001210763> (first received 4 November 2013).

* Heine R, Peters R, Cameron D, O-Loughlin E, Axelrad C, Allen K, et al. Effect of a 4-food elimination diet and omeprazole in children with eosinophilic esophagitis - a randomized controlled trial. *Journal of Allergy and Clinical Immunology* 2019;**143**(2 Suppl):AB309. [DOI: [10.1016/j.jaci.2018.12.940](https://doi.org/10.1016/j.jaci.2018.12.940)]

Hirano 2019 {published data only}

Dellon E S, Collins M H, Assouline-Dayana Y, Evans L, Gupta S, Schoepfer A, et al. Long-term efficacy and safety of RPC4046, an anti-interleukin-13 monoclonal antibody, in patients with eosinophilic esophagitis: Results from the openlabel extension of the heroes study. *United European Gastroenterology Journal* 2018;**6**(8 Suppl):A124-5. [DOI: [10.1177/2050640618792817](https://doi.org/10.1177/2050640618792817)]

Dellon E S, Collins M H, Assouline-Dayana Y, Evans L, Gupta S, Schoepfer A, et al. Long-term efficacy and safety of RPC4046, an anti-interleukin-13 monoclonal antibody, in patients with eosinophilic esophagitis: results from the open-label extension of the heroes study. *Allergy: European Journal of Allergy and Clinical Immunology* 2019;**74**(Suppl 106):308-9. [DOI: [10.1111/all.13961](https://doi.org/10.1111/all.13961)]

Dellon E S, Collins M H, Rothenberg M, Assouline-Dayana Y, Evans L, Gupta S, et al. Efficacy of RPC4046, an anti-interleukin-13 monoclonal antibody, in patients with active eosinophilic esophagitis: Analysis of the steroid-refractory subgroup from the heroes study. *Allergy: European Journal of*

Medical treatment of eosinophilic esophagitis (Review)

Allergy and Clinical Immunology 2019;**74**(Suppl 106):98. [DOI: [10.1111/all.13957](https://doi.org/10.1111/all.13957)]

Dellon E, Collins M, Assouline-Dayana Y, Evans L, Gupta S, Schoepfer A, et al. Long-term efficacy and safety of RPC4046, an anti-interleukin-13 monoclonal antibody, in patients with eosinophilic esophagitis: results from the open-label extension of the HEROES study. *Journal of Allergy and Clinical Immunology* 2019;**143**(2 Suppl):AB208. [DOI: [10.1016/j.jaci.2018.12.635](https://doi.org/10.1016/j.jaci.2018.12.635)]

Dellon E, Collins M, Rothenberg M, Assouline-Dayana Y, Evans L, Gupta S, et al. Efficacy of RPC4046, an anti-interleukin-13 monoclonal antibody, in patients with active eosinophilic esophagitis: Analysis of the steroidrefractory subgroup from the heroes study. *United European Gastroenterology Journal* 2018;**6**(8 Suppl):A51-2. [DOI: [10.1177/2050640618792817](https://doi.org/10.1177/2050640618792817)]

Dellon ES, Collins M, Assouline-Dayana Y, Evans L, Gupta SK, Schoepfer A, et al. A randomized, double-blind, placebo-controlled trial of a novel recombinant, humanized, anti-interleukin-13 monoclonal antibody (RPC4046) in patients with active eosinophilic esophagitis: Results of the heroes study. *American Journal of Gastroenterology* 2016;**111** Suppl 1:S186. [DOI: [10.1038/ajg.2016.355](https://doi.org/10.1038/ajg.2016.355)]

Dellon ES, Collins MH, Assouline-Dayana Y, Evans L, Gupta SK, Schoepfer AM, et al. Efficacy and safety of rpc4046, an anti-interleukin-13 monoclonal antibody, in patients with active eosinophilic esophagitis: analysis of the steroid-refractory subgroup from the HEROES study. *Gastroenterology* 2018;**154**(6 Suppl 1):S-244. [DOI: [10.1016/S0016-5085%2818%2931195-8](https://doi.org/10.1016/S0016-5085%2818%2931195-8)]

Dellon ES, Collins MH, Assouline-Dayana Y, Evans L, Gupta SK, Schoepfer AM, et al. Sustained long-term efficacy and safety of rpc4046, an anti-interleukin-13 monoclonal antibody, in patients with eosinophilic esophagitis: results from the open-label extension of the HEROES study. *Gastroenterology* 2018;**154**(6 Suppl 1):S174-5. [DOI: [10.1016/S0016-5085\(18\)30996-X](https://doi.org/10.1016/S0016-5085(18)30996-X)]

Dellon ES, Collins MH, Rothenberg ME, Assouline-Dayana Y, Evans L, Gupta S, et al. Long-term efficacy and tolerability of rpc4046 in an open-label extension trial of patients with eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology* 2021;**19**(3):473-83.e17. [DOI: [10.1016/j.cgh.2020.03.036](https://doi.org/10.1016/j.cgh.2020.03.036)]

Gann P, Deaton R, McMahon N, Collins M, Dellon E, Hirano I, et al. An anti-IL-13 antibody reverses epithelial-mesenchymal transition biomarkers in eosinophilic esophagitis: phase 2 trial results. *Journal of Allergy and Clinical Immunology* 2020;**146**(2):367. [DOI: [10.1016/j.jaci.2020.03.045](https://doi.org/10.1016/j.jaci.2020.03.045)]

Gann P, Deaton R, McMahon N, Collins M, Dellon E, Hirano I, et al. Markers of esophageal epithelial mesenchymal transition are significantly reduced in active eosinophilic esophagitis following 16 weeks of treatment with RPC4046, an anti-interleukin-13 monoclonal antibody. *United European Gastroenterology Journal* 2018;**6**(8 Suppl):A507-8. [DOI: [10.1177/2050640618792819](https://doi.org/10.1177/2050640618792819)]

Gann P, Deaton R, McMahon N, Collins M, Dellon E, Hirano I, et al. Markers of esophageal epithelial-mesenchymal transition are significantly reduced in active EoE following 16 weeks of

treatment with RPC4046, an anti-interleukin-13 monoclonal antibody. *Allergy: European Journal of Allergy and Clinical Immunology* 2019;**74**(Suppl 106):99. [DOI: [10.1111/all.13957](https://doi.org/10.1111/all.13957)]

Gann P, Deaton R, McMahon N, Collins M, Dellon E, Hirano I, et al. Markers of esophageal epithelial-mesenchymal transition are significantly reduced in active eosinophilic esophagitis following 16 weeks of treatment with RPC4046, an anti-interleukin-13 monoclonal antibody. *Journal of Allergy and Clinical Immunology* 2019;**143**(2 Suppl):AB135. [DOI: [10.1016/j.jaci.2018.12.409](https://doi.org/10.1016/j.jaci.2018.12.409)]

Gann P, Deaton R, McMahon N, Collins MH, Dellon ES, Hirano I, et al. Markers of esophageal epithelial-mesenchymal transition (emt) are significantly reduced in active eosinophilic esophagitis following 16 weeks of treatment with RPC4046, an anti-interleukin-13 monoclonal antibody. *Gastroenterology* 2018;**154**(6 Suppl 1):S243. [DOI: [10.1016/S0016-5085%2818%2931192-2](https://doi.org/10.1016/S0016-5085%2818%2931192-2)]

Hirano I, Collins M, Assouline-Dayana Y, Evans L, Gupta S, Schoepfer A, et al. A randomised, double-blind, placebo-controlled trial of a novel recombinant, humanised, antiinterleukin-13 monoclonal antibody (RPC4046) in patients with active eosinophilic oesophagitis: Results of the heroes study. *United European Gastroenterology Journal* 2016;**4**(5 Suppl 1):A127. [DOI: [10.1177/2050640616663688](https://doi.org/10.1177/2050640616663688)]

Hirano I, Collins M, Assouline-Dayana Y, Evans L, Gupta S, Schoepfer A, et al. A randomized, double-blind, placebo-controlled trial of anti-interleukin-13 monoclonal antibody RPC4046 in patients with eosinophilic esophagitis. *Annals of Allergy, Asthma and Immunology* 2018;**121**(5 Suppl):S15. [DOI: [10.1016/j.ana.2018.09.047](https://doi.org/10.1016/j.ana.2018.09.047)]

* Hirano I, Collins M, Assouline-Dayana Y, Evans L, Gupta S, Schoepfer A, et al. RPC4046, a monoclonal antibody against il13, reduces histologic and endoscopic activity in patients with eosinophilic esophagitis. *Gastroenterology* 2019;**156**(3):592-603. [DOI: [10.1053/j.gastro.2018.10.051](https://doi.org/10.1053/j.gastro.2018.10.051)] [PMID: 30395812]

Hirano I, Dellon E, Collins M, Assouline-Dayana Y, Evans L, Gupta S, et al. Efficacy of RPC4046, an anti-interleukin-13 monoclonal antibody, in patients with active eosinophilic esophagitis: analysis of the steroid-refractory subgroup from the HEROES study. *Journal of Allergy and Clinical Immunology* 2019;**143**(2 Suppl):AB308. [DOI: [10.1016/j.jaci.2018.12.939](https://doi.org/10.1016/j.jaci.2018.12.939)]

NCT02098473. Dose ranging study of RPC4046 in eosinophilic esophagitis [A phase2, multi-center, multi-national, randomized, double-blind, placebo-controlled parallel-group clinical trial to evaluate the efficacy and safety of RPC4046 in adult subjects with eosinophilic esophagitis]. <https://clinicaltrials.gov/ct2/show/NCT02098473> (first received 28 March 2014).

Hirano 2020 {published data only}

Collins M, Hirano I, Dellon E, Peterson K, Chehade M, Schoepfer A, et al. Baseline characteristics and correlation between dysphagia and disease activity in patients with eosinophilic esophagitis in a randomized, placebo-controlled, phase 2 dupilumab trial. *Gastroenterology* 2018;**154**(6 Suppl):S259. [DOI: [10.1016/S0016-5085\(18\)31232-0](https://doi.org/10.1016/S0016-5085(18)31232-0)]

Medical treatment of eosinophilic esophagitis (Review)

Collins M, Rothenberg M, Hamilton J, Lim W, Hamon S, Whipperman M, et al. The gene expression signature modulated by dupilumab is highly correlated with eosinophilic esophagitis histological severity. *United European Gastroenterology Journal* 2020;**8**(8 Suppl):57-8. [DOI: [10.1177/2050640620927344](https://doi.org/10.1177/2050640620927344)]

Corren J, Canonica GW, Graham NMH, Harel S, Chen Z, Amin N, et al. Dupilumab suppresses serum total IgE levels across multiple atopic, allergic diseases. *Allergy: European Journal of Allergy and Clinical Immunology* 2020;**75**(Suppl 109):64. [DOI: [10.1111/all.14504](https://doi.org/10.1111/all.14504)]

Hirano I, Collins M, Dellon E, Rothenberg M, Schoepfer A, Chegade M, et al. Endoscopic features are significantly associated with histologic measures of disease activity in patients with active eosinophilic esophagitis: a post HOC analysis of a randomized, placebo-controlled, phase 2 dupilumab trial. *Gastroenterology* 2018;**154**(6 Suppl 1):S244. [DOI: [10.1016/S0016-5085%2818%2931193-4](https://doi.org/10.1016/S0016-5085%2818%2931193-4)]

Hirano I, Dellon E S, Hamilton J D, Qing Z, Graham N M, Akinlade B, et al. Dupilumab efficacy and safety in adult patients with active eosinophilic oesophagitis: a randomised double-blind placebo-controlled phase 2 trial. *United European Gastroenterology Journal* 2017;**5**(8):1146-7. [DOI: [10.1177/2397847317743186](https://doi.org/10.1177/2397847317743186)]

* Hirano I, Dellon E, Hamilton J, Collins M, Peterson K, Chegade M, et al. Efficacy of dupilumab in a phase 2 randomized trial of adults with active eosinophilic esophagitis. *Gastroenterology* 2020;**158**(1):111-22. [DOI: [10.1053/j.gastro.2019.09.042](https://doi.org/10.1053/j.gastro.2019.09.042)] [PMID: 31593702]

Hirano I, Hamilton J, Collins M, Lin Z, Dellon E, Chen Z, et al. Correlation between esophageal distensibility and objective measures of disease in patients with active eosinophilic esophagitis: a post hoc analysis of a randomized, placebo-controlled, phase 2 dupilumab trial. *Gastroenterology* 2018;**154**(6 Suppl 1):S244. [DOI: [10.1016/S0016-5085%2818%2931194-6](https://doi.org/10.1016/S0016-5085%2818%2931194-6)]

NCT02379052. Study of dupilumab in adult participants with active eosinophilic esophagitis (EoE) [A randomized, double-blind, parallel, placebo-controlled study of the efficacy, safety and tolerability of dupilumab in adult patients with active eosinophilic esophagitis]. <https://clinicaltrials.gov/ct2/show/NCT02379052> (first received 19 March 2015).

Hirano 2020f {published data only}

* Hirano I, Safroneeva E, Roumet MC, Comer GM, Eagle G, Schoepfer A, et al. Randomised clinical trial: the safety and tolerability of fluticasone propionate orally disintegrating tablets versus placebo for eosinophilic oesophagitis. *Alimentary Pharmacology and Therapeutics* 2020;**51**(8):750-9. [DOI: [10.1111/apt.15670](https://doi.org/10.1111/apt.15670)] [PMID: 32149424]

Hirano I, Schoepfer A, Comer G, Safroneeva E, Meltzer B, Falk G. A randomized, double-blind, placebo-controlled trial of a fluticasone propionate orally disintegrating tablet in adult and adolescent patients with eosinophilic esophagitis: a phase 1/2A safety and tolerability study. *Gastroenterology* 2017;**152**(5 Suppl 1):S195.

NCT01386112. Safety and tolerability study of oral EUR-1100 to treat eosinophilic esophagitis [A multicenter, randomized, double-blind, placebo-controlled, safety and tolerability phase 1/2a study of two dosing regimens of EUR-1100 for oral use, in subjects with eosinophilic esophagitis (EoE)]. <https://clinicaltrials.gov/ct2/show/NCT01386112> (first received 30 June 2011).

Hirano 2021 {published data only}

Collins M, Hirano I, Katzka D, Cianferoni A, Williams J, Desai N, et al. Budesonide oral suspension improves histologic features in patients with eosinophilic esophagitis: results from a phase 3, randomized, double-blind, placebo-controlled trial. *Journal of Allergy and Clinical Immunology* 2020;**145**(2 Suppl):AB166. [DOI: [10.1016/j.jaci.2019.12.341](https://doi.org/10.1016/j.jaci.2019.12.341)]

Collins M, Mukkada V, Hirano I, Katzka D, Falk G, Williams J, et al. Improved histopathologic features in patients with eosinophilic esophagitis: results and analyses from a phase 3, randomized, placebo-controlled trial of budesonide oral suspension. *American Journal of Gastroenterology* 2020;**115** Suppl:S200-1. [DOI: [10.14309/01.ajg.0000703664.24718.78](https://doi.org/10.14309/01.ajg.0000703664.24718.78)]

Dellon E, Hirano I, Mukkada V, Katzka D, Gold B, Desai N, et al. Study design and baseline patient characteristics of a phase 3, randomized, placebo-controlled trial of budesonide oral suspension in patients with eosinophilic esophagitis. *Journal of Pediatric Gastroenterology and Nutrition* 2019;**69**(Suppl 2):283. [DOI: [10.1097/MPG.0000000000002518](https://doi.org/10.1097/MPG.0000000000002518)]

Goodwin B, Williams J, Desai N, Lan L. PGI1 estimation of the minimal important differences in eosinophilic esophagitis quality of life (EoE-QoL): analysis from a phase 3, randomized, placebo-controlled trial of budesonide oral suspension. *Value in Health* 2020;**23**(Suppl 1):S143. [DOI: [10.1016/j.jval.2020.04.361](https://doi.org/10.1016/j.jval.2020.04.361)]

Goodwin B, Williams J, Desai N, Lan L. PGI36 Estimation of the minimal important differences in euroqol-5 dimensions (EQ-5D) for patients with eosinophilic esophagitis: analysis from a phase 3, randomized, placebo-controlled trial of budesonide oral suspension. *Value in Health* 2020;**23** Suppl 1:S149. [DOI: [10.1016/j.jval.2020.04.390](https://doi.org/10.1016/j.jval.2020.04.390)]

* Hirano I, Collins M, Katzka D, Mukkada V, Falk G, Morey R, et al. Budesonide oral suspension improves outcomes in patients with eosinophilic esophagitis: results from a phase 3 trial. *Clinical Gastroenterology and Hepatology* 2021;**20**(3):525-34. [DOI: [10.1016/j.cgh.2021.04.022](https://doi.org/10.1016/j.cgh.2021.04.022)] [PMID: 33887475]

Hirano I, Collins M, Katzka D, Mukkada V, Falk G, Morey R, et al. Correction: Budesonide oral suspension improves outcomes in patients with eosinophilic esophagitis: results from a phase 3 trial (Clinical Gastroenterology and Hepatology (2022) 20(3) (525-534.e10), (S1542356521004560), (10.1016/j.cgh.2021.04.022)). *Clinical Gastroenterology and Hepatology* 2022;**20**(3):525-34.e10.

Hirano I, Collins M, Katzka D, Mukkada V, Falk G, Williams J, et al. 227 Budesonide oral suspension (bos) improves endoscopic activity in adolescents and adults with eosinophilic esophagitis: results and correlation analysis from a phase 3, randomized, placebo-controlled trial. *Clinical Gastroenterology*

Medical treatment of eosinophilic esophagitis (Review)

and *Hepatology* 2020;**158**(6 Suppl 1):S43. [DOI: [10.1016/S0016-5085\(20\)2930788-5](https://doi.org/10.1016/S0016-5085(20)2930788-5)]

Hirano I, Collins M, Mukkada V, Katzka D, Falk G, Johnston D, et al. Budesonide oral suspension treatment improves dysphagia symptoms in patients with eosinophilic esophagitis. *Annals of Allergy, Asthma and Immunology* 2019;**123**(5 Suppl):S56-7. [DOI: [10.1016/j.anai.2019.08.338](https://doi.org/10.1016/j.anai.2019.08.338)]

Hirano I, Collins M, Mukkada V, Katzka D, Falk G, Williams J, et al. Determination of the eosinophilic esophagitis endoscopic reference score associated with histologic response to therapy: analysis from a phase 3 placebo-controlled trial of budesonide oral suspension. *American Journal of Gastroenterology* 2020;**115** Suppl:S217-8. [DOI: [10.14309/01.ajg.0000703788.60508.8e](https://doi.org/10.14309/01.ajg.0000703788.60508.8e)]

Hirano I, Collins M, Mukkada V, Katzka D, Falk Gary W, Williams J, et al. Efficacy of budesonide oral suspension for eosinophilic esophagitis in adolescents and adults: Results from a phase 3, randomized, placebo-controlled trial. *American Journal of Gastroenterology* 2019;**114** Suppl:S205-6. [DOI: [10.14309/01.ajg.0000590932.88962.0b](https://doi.org/10.14309/01.ajg.0000590932.88962.0b)]

Hirano I, Katzka D A, Collins M, Mukkada V, Gold B, Falk G, et al. Predictors of response to corticosteroid therapy in patients with eosinophilic esophagitis: results from a phase 3, randomized, placebo-controlled trial of budesonide oral suspension. *Gastroenterology* 2020;**158**(6 Suppl 1):817-8. [DOI: [10.1016/S0016-5085\(20\)32723-2](https://doi.org/10.1016/S0016-5085(20)32723-2)]

NCT02605837. A placebo- controlled study in adolescents and adults with eosinophilic esophagitis (EoE) measuring histologic response and determine if reduction in dysphagia is achieved [Oral budesonide suspension (OBS) in adolescent and adult subjects (11 to 55 years of age, inclusive) with eosinophilic esophagitis: a phase 3 randomized, double-blind, placebo-controlled study]. <https://clinicaltrials.gov/ct2/show/NCT02605837> (first received 16 November 2015).

Kliewer 2019 {published data only}

* Kliewer K, Aceves SS, Atkins D, Bonis PA, Chehade M, Collins MH, et al. Efficacy of 1-food and 4-food elimination diets for pediatric eosinophilic esophagitis in a randomized multisite study. *Gastroenterology* 2019;**156**(6 Suppl 1):S-172-3.

Kliewer K, Gonsalves N, Dellon ES, Katzka DA, Aceves SS, Arva N, et al. Efficacy of one-food versus six-food elimination diet for treatment of eosinophilic esophagitis in adults: results from the multicenter randomized controlled sofeed trial. *Gastroenterology* 2021;**160**((6 Suppl)):S-109-10.

NCT02610816. Eosinophilic esophagitis intervention trial-1 food vs. 4 food elimination diet followed by swallowed glucocorticoids. <https://clinicaltrials.gov/ct2/show/NCT02610816> (first received 20 November 2015).

Kliewer 2021 {published data only}

* Kliewer K, Gonsalves N, Dellon E, Katzka DA, Seema S, Aceves NA, et al. 535 Efficacy of one-food versus six-food elimination diet for treatment of eosinophilic esophagitis in adults: results from the multicenter randomized controlled sofeed trial. *Gastroenterology* 2021;**160**(6 Suppl 1):S109. [DOI: [10.1016/S0016-5085\(19\)37223-3](https://doi.org/10.1016/S0016-5085(19)37223-3)]

NCT02778867. SOFEED: Six food vs. one food eosinophilic esophagitis diet study [Six food vs one food eosinophilic esophagitis elimination diet (SOFEED) followed by swallowed glucocorticoid trial]. <https://clinicaltrials.gov/ct2/show/NCT02778867> (first received 20 May 2016).

Konikoff 2006 {published data only}

* Konikoff MR, Noel RJ, Blanchard C, et al. A randomized, double-blind, placebo-controlled trial of fluticasone propionate for pediatric eosinophilic esophagitis. *Gastroenterology* 2006;**131**(5):1381-91. [DOI: [10.1053/j.gastro.2006.08.033](https://doi.org/10.1053/j.gastro.2006.08.033)] [PMID: 17101314]

NCT00266578. A study comparing swallowed Flovent and placebo in patients with eosinophilic disorders [Effect of swallowed fluticasone propionate on eosinophilic esophagitis; a prospective, randomized, placebo-controlled trial]. <https://www.clinicaltrials.gov/ct2/show/NCT00266578> (first received 19 December 2005).

Lieberman 2018 {published data only}

* Lieberman J, Whitworth J, Cavender C, Zhang J. A randomized, double-blinded, placebo-controlled study of the use of viscous oral cromolyn sodium for the treatment of eosinophilic esophagitis. *Annals of Allergy, Asthma and Immunology* 2018;**120**(5):527-31. [DOI: [10.1016/j.anai.2018.03.006](https://doi.org/10.1016/j.anai.2018.03.006)] [PMID: 29544738]

Lieberman J, Zhang J, Cavender C. Viscous oral cromolyn for the treatment of eosinophilic esophagitis: a double-blind, placebo-controlled trial. *Annals of Allergy, Asthma and Immunology* 2017;**119**(5 Suppl 1):S9.

NCT02371941. Oral cromolyn sodium for the treatment of eosinophilic esophagitis [A randomized, double-blind, placebo-controlled study of the use of oral cromolyn sodium for the treatment of eosinophilic esophagitis]. <https://clinicaltrials.gov/ct2/show/NCT02371941> (first received 26 February 2015).

Lucendo 2019 {published data only}

Lucendo A, Miehke S, Vieth M, Schlag C, Von A, Molina-Infante J, et al. Budesonide orodispersible tablets are highly effective for treatment of active eosinophilic esophagitis: results from a randomized, double-blind, placebo-controlled, pivotal multicenter trial (EOS-1). *Gastroenterology* 2017;**152**(5 Suppl 1):S207.

* Lucendo A, Miehke S, Schlag C, Vieth M, von Arnim U, Molina-Infante J, et al. Efficacy of budesonide orodispersible tablets as induction therapy for eosinophilic esophagitis in a randomized placebo-controlled trial. *Gastroenterology* 2019;**157**(1):74-86. [DOI: [10.1053/j.gastro.2019.03.025](https://doi.org/10.1053/j.gastro.2019.03.025)] [PMID: 30922997]

Schlag C, Miehke S, Lucendo A, Biedermann L, Santander C, Hartmann D, et al. Efficacy of budesonide orodispersible tablets for induction of remission in patients with active eosinophilic esophagitis: results from the 6-weeks open-label treatment phase of EOS-2 trial. *Gastroenterology* 2019;**156**(6 Suppl 1):S715. [DOI: [10.1016/S0016-5085\(20\)28199-2](https://doi.org/10.1016/S0016-5085(20)28199-2)]

Miehke 2016 {published data only}

EUCTR2009-016692-29-BE. A study to investigate the efficacy and tolerance of budesonide tablets and suspension in

Medical treatment of eosinophilic esophagitis (Review)

patients suffering from esophagitis over a 14 day period. <https://www.clinicaltrialsregister.eu/ctr-search/search?query=EUCTR2009-016692-29-BE> (first received 24 November 2011).

Hruz P, Mielhke S, Von Arnim U, Madisch A, Vieth M, Bussmann C, et al. A randomized, double-blind, double-dummy, placebo-controlled, European multicentre trial of two new budesonide formulations for treatment of active eosinophilic esophagitis. *United European Gastroenterology Journal* 2014;**2**(1 Suppl 1):A128. [DOI: [10.1177/2050640614548974](https://doi.org/10.1177/2050640614548974)]

* Mielhke S, Hruz P, Vieth M, Bussmann C, Von Arnim U, Bajbouj M, et al. A randomised, double-blind trial comparing budesonide formulations and dosages for short-term treatment of eosinophilic oesophagitis. *Gut* 2016;**65**(3):390-9. [DOI: [10.1136/gutjnl-2014-308815](https://doi.org/10.1136/gutjnl-2014-308815)] [PMID: 25792708]

Mielhke S, Hruz P, Von Ulrike A, Madisch A, Vieth M, Bussmann C, et al. Two new budesonide formulations are highly efficient for treatment of active eosinophilic esophagitis: Results from a randomized, double-blind, double-dummy, placebo-controlled multicenter trial. *Gastroenterology* 2014;**146**(5 Suppl 1):S16. [DOI: [10.1016/S0016-5085%2814%2960055-X](https://doi.org/10.1016/S0016-5085%2814%2960055-X)]

NCT02280616. Budesonide effervescent tablets vs. viscous budesonide suspension vs. placebo in eosinophilic esophagitis [Double-blind, double-dummy, randomized, placebo-controlled, phase iia study on the efficacy and tolerability of a 14-day treatment with budesonide effervescent tablets vs. viscous budesonide suspension vs. placebo in patients with eosinophilic esophagitis]. <https://clinicaltrials.gov/ct2/show/NCT02280616> (first received 31 October 2014).

Moawad 2013 {published data only}

Moawad F, Dias J, Veerappan G, Baker T, Maydonovitch C, Wong R. Comparison of aerosolized swallowed fluticasone to esomeprazole for the treatment of eosinophilic esophagitis. *American Journal of Gastroenterology* 2011;**106**:S12.

* Moawad FJ, Veerappan GR, Dias JA, Baker TP, Maydonovitch CL, Wong RK. Randomized controlled trial comparing aerosolized swallowed fluticasone to esomeprazole for esophageal eosinophilia. *American Journal of Gastroenterology* 2013;**108**(3):366-72. [DOI: [10.1038/ajg.2012.443](https://doi.org/10.1038/ajg.2012.443)] [PMID: 23399553]

NCT00895817. Fluticasone versus esomeprazole to treat eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT00895817> (first received 8 May 2009).

Oliva 2018 {published data only}

Oliva S, Ruggiero C, Rossetti D, Tinti V, Salimbene L, Sarli M, et al. A randomized controlled trial comparing six-food elimination diet vs topical steroids in inducing and maintaining remission of paediatric eosinophilic esophagitis. *Journal of Pediatric Gastroenterology and Nutrition* 2018;**66** Suppl 2:89.

* Oliva S, Ruggiero C, Rossetti D, Tinti V, Salimbene L, Sarli M, et al. A randomized controlled trial comparing six-food elimination diet vs topical steroids in inducing and maintaining remission of

pediatric eosinophilic esophagitis. *Gastroenterology* 2018;**154**(6 Suppl 1):S163-4. [DOI: [10.1016/S0016-5085\(18\)30957-0](https://doi.org/10.1016/S0016-5085(18)30957-0)]

Peterson 2010 {published data only}

NCT00123656. Comparison of esomeprazole to aerosolized, swallowed fluticasone for eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT00123656> (first received 25 July 2005).

* Peterson K, Thomas K, Hilden K, Emerson L, Wills J, Fang J. Comparison of esomeprazole to aerosolized, swallowed fluticasone for eosinophilic esophagitis. *Digestive Diseases and Sciences* 2010;**55**(5):1313-9. [DOI: [10.1007/s10620-009-0859-4](https://doi.org/10.1007/s10620-009-0859-4)] [PMID: 19533356]

Rothenberg 2015 {published data only}

NCT01022970. Efficacy and safety of QAX576 in patients with eosinophilic esophagitis [A double blinded randomized placebo-controlled trial of intravenous qax576 in the treatment of eosinophilic esophagitis (EoE)]. <https://clinicaltrials.gov/ct2/show/NCT01022970> (first received 1 December 2009).

* Rothenberg M, Wen T, Greenberg A, Alpan O, Enav B, Hirano I, et al. Intravenous anti-IL-13 mAb QAX576 for the treatment of eosinophilic esophagitis. *Journal of Allergy and Clinical Immunology* 2015;**135**(2):500-7. [DOI: [10.1016/j.jaci.2014.07.049](https://doi.org/10.1016/j.jaci.2014.07.049)] [PMID: 25226850]

Rothenberg 2022 {published data only}

* Dellon ES, Rothenberg ME, Collins MH, Hirano I, Chehade M, Bredenoord AJ, et al. Dupilumab in adults and adolescents with eosinophilic esophagitis. *New England Journal of Medicine* 2022;**387**(25):2317-30.

Rothenberg M, Dellon E, Bredenoord A, Collins M, Hirano I, Chehade M, et al. Dupilumab improves clinical and histologic aspects of disease in adult and adolescent patients with eosinophilic esophagitis at week 24: results from Part B of the 3-Part LIBERTY EoE TREET Study. *Journal of Allergy and Clinical Immunology* 2022;**149**(2 (Suppl)):AB312.

Schaefer 2008 {published data only}

Gupta SK, Fitzgerald JF, Davis MM, Croffie JM, Pfefferkorn, MD, Molleston JP, et al. Treatment of allergic eosinophilic esophagitis (AEE) with oral prednisone (P) and swallowed fluticasone (F): a randomized, prospective study in children. *Gastroenterology* 2003;**124**(4 Suppl 1):A19.

* Schaefer ET, Fitzgerald JF, Molleston JP, Croffie JM, Pfefferkorn MD, Corkins MR, et al. Comparison of oral prednisone and topical fluticasone in the treatment of eosinophilic esophagitis: a randomized trial in children. *Clinical Gastroenterology and Hepatology* 2008;**6**(2):165-73. [DOI: [10.1016/j.cgh.2007.11.008](https://doi.org/10.1016/j.cgh.2007.11.008)] [PMID: 18237866]

Spergel 2012 {published data only}

Markowitz J, Eke R, Frost C. Long-term safety and efficacy of reslizumab in children and adolescents with eosinophilic esophagitis: A review of 501 doses in 12 children over 8 years. *Journal of Pediatric Gastroenterology and Nutrition* 2016;**63**(Suppl 2):S143. [DOI: [10.1097/01.mpg.0000503536.79797.66](https://doi.org/10.1097/01.mpg.0000503536.79797.66)]

Medical treatment of eosinophilic esophagitis (Review)

Markowitz J, Jobe L, Miller M, Frost C, Eke R. Long-term safety and efficacy of reslizumab in children and adolescents with eosinophilic esophagitis: a review of 477 doses in 12 children over 7 years. *Journal of Pediatric Gastroenterology and Nutrition* 2016;**137**(2 Suppl 1):AB234.

Markowitz J, Jobe L, Miller M, Laney Z, Frost C, Eke R. Safety and efficacy of reslizumab for children and adolescents with eosinophilic esophagitis treated for 9 years. *Journal of Pediatric Gastroenterology and Nutrition* 2018;**66**(6):893-7. [DOI: [10.1097/MPG.0000000000001840](https://doi.org/10.1097/MPG.0000000000001840)] [PMID: 29176476]

NCT00538434. Efficacy and safety study of reslizumab to treat eosinophilic esophagitis in subjects aged 5 to 18 years [An efficacy and safety study of reslizumab (CTX55700) in the treatment of eosinophilic esophagitis in subjects aged 5 to 18 years]. <https://clinicaltrials.gov/ct2/show/NCT00538434> (first received 2 October 2007).

Spergel J, Rothenberg M, Collins M, Furuta G, Markowitz J, Fuchs G, et al. Reslizumab in children and adolescents with eosinophilic esophagitis: results of a double-blind, randomized, placebo-controlled study. *Gastroenterology* 2011;**140**(5 Suppl 1):S136.

* Spergel J, Rothenberg M, Collins M, Furuta G, Markowitz J, Fuchs G, et al. Reslizumab in children and adolescents with eosinophilic esophagitis: results of a double-blind, randomized, placebo-controlled trial. *Journal of Allergy and Clinical Immunology* 2012;**129**(2):456-63. [DOI: [10.1016/j.jaci.2011.11.044](https://doi.org/10.1016/j.jaci.2011.11.044)] [PMID: 22206777]

Spergel 2020 {published data only}

Burke D, Lewis M, Spergel J, Brown-Whitehorn T, Cianferoni A, Muir A, et al. Sustained milk consumption after 2 years post-milk epicutaneous immunotherapy for eosinophilic esophagitis. *Allergy: European Journal of Allergy and Clinical Immunology* 2021;**76**(5):1573-6. [DOI: [10.1111/all.14659](https://doi.org/10.1111/all.14659)] [PMID: 33202086]

NCT02579876. Milk patch for eosinophilic esophagitis [A double, blind placebo-controlled, randomized trial to study efficacy and safety of the Viaskin milk for treating milk induced eosinophilic esophagitis in children]. <https://clinicaltrials.gov/ct2/show/NCT02579876> 2015.

* Spergel J, Elci O, Muir A, Liacouras C, Wilkins B, Burke D, et al. Efficacy of epicutaneous immunotherapy in children with milk-induced eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology* 2020;**18**(2):328-36. [DOI: [10.1016/j.cgh.2019.05.014](https://doi.org/10.1016/j.cgh.2019.05.014)] [PMID: 31100455]

Straumann 2010a {published data only}

Conus S, Straumann A, Bettler E, Simon H U. Mepolizumab does not alter levels of eosinophils, T cells, and mast cells in the duodenal mucosa in eosinophilic esophagitis. *Journal of Allergy and Clinical Immunology* 2010;**126**(1):175-7. [DOI: [10.1016/j.jaci.2010.04.029](https://doi.org/10.1016/j.jaci.2010.04.029)] [DOI: [10.1016/j.jaci.2011.01.014](https://doi.org/10.1016/j.jaci.2011.01.014)] [PMID: 20542323]

EUCTR2005-006074-10-GB. A randomized, double-blind, parallel group clinical trial to assess safety, tolerability, pharmacokinetics, and pharmacodynamics of intravenous mepolizumab (SB240563) (0.55mg/kg, 2.5mg/kg or 10mg/

kg) in pediatric subjects with eosinophilic esophagitis, ag. <https://www.clinicaltrialsregister.eu/ctr-search/search?query=EUCTR2005-006074-10-GB> (first received 7 February 2006).

NCT00274703. An evaluation of mepolizumab in therapy of eosinophilic oesophagitis in adult patients [A randomised, double-blind, placebo-controlled, single-centre study to provide a preliminary evaluation of pharmacokinetics, pharmacodynamics, safety and tolerability of intravenous anti-human interleukin-5 (mepolizumab, 750mg and 1500mg) in the treatment of eosinophilic oesophagitis in adults]. <https://clinicaltrials.gov/ct2/show/NCT00274703> (first received January 2006).

Straumann A, Conus H, Kita G, Kephart C, Bussmann C, Beglinger J et al. Mepolizumab, a humanized monoclonal antibody to IL-5, for severe eosinophilic esophagitis in adults: a randomized, placebo-controlled double-blind trial. *Journal of Allergy and Clinical Immunology* 2008;**121**(2 Suppl 1):S44.

* Straumann A, Conus S, Grzonka P, Simon H U, Kita H, Kephart G, et al. Anti-interleukin-5 antibody treatment (mepolizumab) in active eosinophilic oesophagitis: a randomised, placebo-controlled, double-blind trial. *Gut* 2010;**59**(1):21-30. [DOI: [10.1136/gut.2009.178558](https://doi.org/10.1136/gut.2009.178558)] [PMID: 19828470]

Straumann 2010b {published data only}

Degen L, Bussmann C, Conus S, Thalmann C, Simon H, Straumann A. Budesonide as induction treatment for active eosinophilic esophagitis in adolescents and adults: a randomised, double-blind, placebo-controlled study (BEE-I trial). In: *Allergy: European Journal of Allergy and Clinical Immunology*. Vol. 65 Suppl 92. 2010:343-4. [DOI: [10.1111/j.1398-9995.2010.02393.x](https://doi.org/10.1111/j.1398-9995.2010.02393.x)]

* Straumann A, Conus S, Degen L, Felder S, Kummer M, Engel H, et al. Budesonide is effective in adolescent and adult patients with active eosinophilic esophagitis. *Gastroenterology* 2010;**139**(5):1526-37. [DOI: [10.1053/j.gastro.2010.07.048](https://doi.org/10.1053/j.gastro.2010.07.048)] [PMID: 20682320]

Straumann A, Degen L, Felder S, Conus S, Bussmann C, Thalmann C, et al. Budesonide as induction treatment for active eosinophilic esophagitis in adolescents and adults: a randomized, double-blind, placebo-controlled Study. *Journal of Allergy and Clinical Immunology* 2009;**123**(2 Suppl 1):S267-8.

Straumann 2011 {published data only}

NCT00271349. Budesonide for eosinophilic esophagitis [Budesonide for eosinophilic esophagitis: a randomized, placebo-controlled, double-blind treatment study]. <https://clinicaltrials.gov/ct2/show/NCT00271349> (first received 30 December 2005).

Straumann A, Conus S, Bussmann C, Degen L, Beglinger C, Simon H. Maintenance-treatment of eosinophilic esophagitis with budesonide (BEE-M trial). *Allergy: European Journal of Allergy and Clinical Immunology* 2010;**65** Suppl 92:342-3. [DOI: [10.1111/j.1398-9995.2010.02393.x](https://doi.org/10.1111/j.1398-9995.2010.02393.x)]

Medical treatment of eosinophilic esophagitis (Review)

* Straumann A, Conus S, Degen L, Frei C, Bussmann C, Beglinger C, et al. Long-term budesonide maintenance treatment is partially effective for patients with eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology* 2011;**9**(5):400-9. [DOI: [10.1016/j.cgh.2011.01.017](https://doi.org/10.1016/j.cgh.2011.01.017)] [PMID: 21277394]

Straumann A, Degen L, Bussmann C, Beglinger C, Schoepfer A, Thalman C, et al. Long-term treatment of eosinophilic esophagitis with budesonide. *Gastroenterology* 2010;**138**(5 Suppl 1):S123.

Straumann 2013 {published data only}

NCT01056783. Proof of concept study of OC000459 in eosinophilic esophagitis [A randomized, placebo-controlled, double-blind, single-centre proof-of-concept study of OC000459 in adult patients with active eosinophilic esophagitis]. <https://clinicaltrials.gov/ct2/show/NCT01056783> (first received 26 January 2010).

Straumann A, Bussmann C, Collins L P, Pettipher R, Hunter M, Steiner J, et al. Treatment of eosinophilic esophagitis with the CRTH2-antagonist oc000459: a novel therapeutic principle. *United European Gastroenterology Journal* 2013;**1**(1 Suppl 1):A123. [DOI: [10.1177/2050640613502899](https://doi.org/10.1177/2050640613502899)]

Straumann A, Bussmann C, Perkins M, Pearce C, Pettipher R, Hunter M, et al. Treatment of eosinophilic esophagitis with the CRTH2-antagonist OC000459: a novel therapeutic principle. *Swiss Medical Weekly* 2012;**142**:8S.

Straumann A, Bussmann C, Perkins MC, Collins LP, Pettipher R, Hunter M, et al. Treatment of eosinophilic esophagitis with the crth2-antagonist oc000459: a novel therapeutic principle. *Gastroenterology* May 2021;**142**(5 Suppl 1):S147. [DOI: [10.1016/S0016-5085\(12\)60552-6](https://doi.org/10.1016/S0016-5085(12)60552-6)]

* Straumann A, Hoesli S, Stuck M, Simon HU, Bussmann C, Perkins M, et al. Anti-eosinophil activity and clinical efficacy of the CRTH2 antagonist OC000459 in eosinophilic esophagitis. *Allergy: European Journal of Allergy and Clinical Immunology* 2013;**68**(3):375-85. [DOI: [10.1111/all.12096](https://doi.org/10.1111/all.12096)] [PMID: 23379537]

Straumann 2020 {published data only}

Biedermann L, Lucendo A, Miehke S, Schlag C, Santander Vaquero C, Ciriza De Los Rios C, et al. A novel budesonide orodispersible tablet formulation is highly effective to maintain endoscopic inflammatory remission and even complete endoscopic remission in adult patients with eosinophilic esophagitis: Results from the 48-weeks, double-blind, placebo. *United European Gastroenterology Journal* 2019;**7**:55.

EUCTR2014-001484-12-DE. Double-blind (neither physician nor patient knows of the actual treatment which can with or without active substance), randomized (patient will be allocated to a certain treatment group by chance), placebo-controlled (tested against tablets without active. <https://eudract.ema.europa.eu/> (first received 9 April 2015).

EUCTR2014-001485-99-DE. Double-blind (neither physician nor patient knows of the actual treatment which can with or without active substance), randomized (patient will be allocated to a certain treatment group by chance), phase

III study on the efficacy and tolerability of a 48-. <https://eudract.ema.europa.eu/> (first received 8 December 2015).

Lucendo A, Miehke S, Vieth M, Schlag C, Biedermann L, Santander C, et al. Budesonide orodispersible tablets are highly effective to maintain clinico-histological remission in adult patients with eosinophilic esophagitis: results from the 48-weeks, double-blind, placebo-controlled, pivotal eos-2 trial. *Gastroenterology* 2019;**156**:S-1509.

Lucendo A, Miehke S, Vieth M, Schlag C, von Arnim U, Molina-Infante J, et al. Prolongation of eosinophilic esophagitis treatment with budesonide orodispersible tablets for incomplete responder is effective and safe: results from a 6-weeks open-label treatment phase of the pivotal trial EOS-1. *Gastroenterology* 2018;**154**:S-75.

Lucendo A, Schlag C, Miehke S, Biedermann L, Santander Vaquero C, Hartmann D, et al. A novel oral budesonide formulation is highly effective for induction of remission in patients with active eosinophilic esophagitis: results from the 6-weeks open-label treatment phase of EOS-2 trial. *United European Gastroenterology Journal* 2019;**7**:54.

Lucendo A, Schlag C, Straumann A, Vieth M, Mueller R, Greinwald R, et al. Budesonide orodispersible tablets can effectively induce complete remission of endoscopic and histologic mucosal abnormalities and can induce deep disease remission in active eosinophilic esophagitis: results from a post-hoc analysis of the randomized, do. *Gastroenterology* 2019;**156**:715-6.

Miehke S, Lucendo A J, Vieth M, Schlag C, Von Arnim U, Molina-Infante J, et al. A randomized, double-blind, placebo controlled, pivotal multicenter trial with budesonide orodispersible tablets for treatment of active eosinophilic esophagitis (EOS-1). *Allergy: European Journal of Allergy and Clinical Immunology* 2017;**72**:78-9.

Miehke S, Lucendo A J, Vieth M, Schlag C, Von Arnim U, Molina-Infante J, et al. Predictive factors for early vs delayed response to budesonide orodispersible tablets in eosinophilic esophagitis: Results from the pivotal trial EOS-1. *Allergy: European Journal of Allergy and Clinical Immunology* 2018;**73**:37.

Miehke S, Schlag C, Straumann A, Vieth M, Muller R, Greinwald R, et al. A novel budesonide orodispersible tablet with a special esophageal-targeting can induce complete clinical, endoscopic and histologic remission in active Eosinophilic Esophagitis: results from a post-hoc analysis of the randomized, double-blind, placebo-controlled EOS-1 trial. *United European Gastroenterology Journal* 2019;**7**:423.

NCT02434029. Budesonide orodispersible tablet vs. placebo in active eosinophilic esophagitis. [ClinicalTrials.gov](https://clinicaltrials.gov) (first received 5 May 2015).

Schlag C, Biedermann L, Lucendo AJ, Miehke S, Santander Vaquero C, Ciriza De Los Rios C, et al. Budesonide orodispersible tablets are superior to maintain and even further improve quality of life in adult patients with eosinophilic esophagitis: Results from the 48-weeks, double-blind, placebo-controlled pivotal EOS-2 trial. *United European Gastroenterology Journal* 2019;**7**:705.

Medical treatment of eosinophilic esophagitis (Review)

Srinivasan S, Sharma P. In eosinophilic esophagitis, budesonide orodispersible tablets maintained remission at 48 wk. *Annals of Internal Medicine* 2021;**174**:JC44.

Straumann A, Lucendo A J, Greinwald R, Mueller R, Attwood S. Efficacy and safety of budesonide orodispersible tablets in active eosinophilic oesophagitis: Results from a randomised, double-blind, placebo-controlled, pivotal, European multicentre trial (EOS-1). *United European Gastroenterology Journal* 2017;**147**:10S.

* Straumann A, Lucendo A, Miehle S, Vieth M, Schlag C, Biedermann L, et al. Budesonide orodispersible tablets maintain remission in a randomized, placebo-controlled trial of patients with eosinophilic esophagitis. *Gastroenterology* 2020;**159**:1672-85. [PMID: 32721437]

Straumann A, Lucendo AJ, Biedermann L, Hruz P, Mueller R, Greinwald R, et al. Budesonide orodispersible tablets maintain remission in a randomized, placebo-controlled trial of patients with eosinophilic esophagitis. *Swiss Medical Weekly* 2021;**151**(Suppl 253):2S.

Straumann A, Lucendo V, Miehle S, Vieth M, Schlag C, Von Arnim U, et al. Efficacy and safety of budesonide orodispersible tablets in active eosinophilic oesophagitis: Results from a randomised, double blind, placebo-controlled, pivotal, European multicentre trial (EOS-1). *United European Gastroenterology Journal* 2017;**5**:A146-7.

Tytor 2021 {published data only}

EUCTR2012-005842-39-SE. Treatment of patients with eosinophilic esophagitis with mometasone furoate aerosol: a randomised, placebo-controlled phase II study. <https://www.clinicaltrialsregister.eu/ctr-search/search?query=EUCTR2012-005842-39-SE> (first received 5 November 2013).

NCT02113267. Mometasone-furoate for treatment of eosinophilic esophagitis - a randomized placebo controlled study [Treatment of eosinophilic esophagitis with mometasone furoate aerosol: a randomised, placebo-controlled phase ii study for evaluation of treatment effect on group level including symptom questionnaires]. <https://clinicaltrials.gov/ct2/show/NCT02113267> (first received 14 April 2014).

* Tytor J, Larsson H, Bove M, Johansson L, Bergquist H. Topically applied mometasone furoate improves dysphagia in adult eosinophilic esophagitis-results from a double-blind, randomized, placebo-controlled trial. *Scandinavian Journal of Gastroenterology* 2021;**56**(6):629-34. [PMID: 10.1080/00365521.2021.1906314] [PMID: 33831327]

References to studies excluded from this review

Braathén 2006 {published data only}

Braathén L, Straumann A, Wenk A, Spichtin H, Simon H U, Simon D. Eosinophilic esophagitis in adults - no clinical relevance of wheat and rye sensitizations. *Allergy: European Journal of Allergy and Clinical Immunology* 2006;**61**:1480-3.

Ceves 2005 {published data only}

Aceves SS, Dohil R, Newbury RO, Bastian JF. Topical viscous budesonide suspension for treatment of eosinophilic esophagitis. *Journal of Allergy and Clinical Immunology* 2005;**116**(3):705-6.

Comer 2017 {published data only}

Comer GM, Meltzer BA. A phase 1 study to assess the pharmacokinetics, safety, and tolerability of single dose APT-1011 administered under fed or fasted conditions or at bedtime in a randomised three-way, crossover design. *United European Gastroenterology Journal* 2017;**5**(5 Suppl 1):A351. [DOI: [10.1177/2050640617725676](https://doi.org/10.1177/2050640617725676)]

Della 2017 {published data only}

Della Coletta M, Bartolo O, Savarino E, Tolone S, De Bortoli N, Bodini G, et al. Different proton pump inhibitors are equally effective in inducing endoscopic and histologic remission in patients with proton pump inhibitor-response esophageal eosinophilia. *Neurogastroenterology and Motility* 2017;**29**:23-4.

Della Coletta M, Bartolo O, Savarino E, Tolone S, De Bortoli N, Bodini G, et al. Esomeprazole, rabeprazole and pantoprazole are equally effective in inducing endoscopic and histologic remission in patients with proton pump inhibitor-response esophageal eosinophilia. *Digestive and Liver Disease* 2018;**50**:e180-1.

Della Coletta M, Savarino E, Tolone S, De Bortoli N, Bartolo O, Bodini G, et al. Esomeprazole, rabeprazole and pantoprazole are equally effective in inducing endoscopic and histologic remission in patients with proton pump inhibitor-response esophageal eosinophilia. *UEG Journal* 2017;**5**(5 Suppl 1):A352.

Della Coletta M, Tolone S, De Bortoli N, Bartolo O, Bodini G, Marabotto E, et al. Different proton pump inhibitors are equally effective in inducing endoscopic and histologic remission in patients with proton pump inhibitor-response esophageal eosinophilia. *Gastroenterology* 2017;**152**(5 Suppl 1):S860-1.

Dellon 2020c {published data only}

Dellon ES, Peterson KA, Murray JA, Falk GW, Gonsalves N, Chehade M, et al. Histologic and symptomatic improvement across multiple forms of eosinophilic gastrointestinal diseases in enigma, a randomized, double-blind, placebo-controlled trial of antolimumab (AK002). *Gastroenterology* 2020;**158**(6 Suppl 1):S186.

NCT04322708. A study of antolimumab (AK002) in patients with active eosinophilic esophagitis. ClinicalTrials.gov 2020.

Eluri 2017 {published data only}

Eluri S, Selitsky S, Parker J, Hollyfield J, Perjar I, Rusin S, et al. Clinical and molecular predictors of histologic response to topical steroid treatment in eosinophilic esophagitis: Results from a prospective study. *Gastroenterology* 2017;**152**:S858.

Eluri 2017a {published data only}

Eluri S, Shaheen NJ, Dellon ES, Perjar I, Hollyfield J, Rusin S, et al. A multivariable model predicts complete histologic response to topical corticosteroid treatment in eosinophilic esophagitis. *Gastrointestinal Endoscopy* 2017;**85**(5 Suppl 1):AB121.

Medical treatment of eosinophilic esophagitis (Review)

EUCTR2014-002465-30-IT 2014 {published data only}

EUCTR2014-002465-30-IT. Spontaneous study on the efficacy and safety of an oral viscous suspension of budesonide administered to children affected by eosinophilic esophagitis. <https://eudract.ema.europa.eu> (first received 24 June 2014).

Francis 2012 {published data only}

Francis D, Arora A, Jensen K, Nord S, Alexander J, Romero Y, et al. Results of ambulatory pH monitoring do not reliably predict response to therapy in patients with eosinophilic oesophagitis. *Alimentary Pharmacology and Therapeutics* 2012;**35**:300-7.

Hefner 2016 {published data only}

Hefner JN, Howard RS, Massey R, Valencia M, Stocker DJ, Philla KQ et al. A randomized controlled comparison of esophageal clearance times of oral budesonide preparations. *Digestive Diseases and Sciences* 2016;**61**:1582-90.

Helou 2008 {published data only}

Helou EF, Simonson J, Arora AS. 3-yr-follow-up of topical corticosteroid treatment for eosinophilic esophagitis in adults. *American Journal of Gastroenterology* 2008;**9**:2194-9.

Hudgens 2017 {published data only}

Hudgens S, Evans C, Phillips E, Hill M. Psychometric validation of the dysphagia symptom questionnaire in patients with eosinophilic esophagitis treated with budesonide oral suspension. *Journal of Patient-reported Outcomes* 2017;**1**:3.

JPRN-UMIN000021041 2016 {published data only}

JPRN-UMIN000021041. A study investigating the therapeutic effect of vonoprazan on the patients with proton pump inhibitor resistant eosinophilic esophagitis. <https://trialsearch.who.int/?TrialID=JPRN-UMIN000021041> (first received 15 February 2016).

JPRN-UMIN000026704 2017 {published data only}

JPRN-UMIN000026704. A study investigating the effect of CYP2C19 polymorphism on the therapeutic efficacy in patients with eosinophilic esophagitis. https://center6.umin.ac.jp/cgi-open-bin/ctr_e/ctr_view.cgi?recptno=R000030656 (first received 27 March 2017).

Kagalwalla 2006 {published data only}

Kagalwalla AF, Sentongo TA, Ritz S, Hess T, Nelson SP et al. Effect of six-food elimination diet on clinical and histologic outcomes in eosinophilic esophagitis. *Clinical Gastroenterology and Hepatology* 2006;**4**:1097-102.

Kavitt 2016 {published data only}

Kavitt RT, Ates F, Slaughter JC, Higginbotham T, Shepherd BD, Sumner EL, et al. Randomized controlled trial comparing esophageal dilation to no dilation among adults with esophageal eosinophilia and dysphagia. *Diseases of the Esophagus* 2016;**29**:983-991.

Kruszewski 2016 {published data only}

Kruszewski P, Russo J, Franciosi J, Varni J, Platts-Mills T, Erwin E. Prospective, comparative effectiveness trial of cow's milk elimination and swallowed fluticasone for pediatric eosinophilic esophagitis. *Diseases of the Esophagus* 2016;**29**:377-84.

Kuzumoto 2021 {published data only}

Kuzumoto T, Tanaka F, Sawada A, Nadatani Y, Otani K, Hosomi S, et al. Vonoprazan shows efficacy similar to that of proton pump inhibitors with respect to symptomatic, endoscopic, and histological responses in patients with eosinophilic esophagitis. *Esophagus* 2021;**18**:372-9.

Molina-Infante 2017 {published data only}

Molina-Infante J, Gonzalez-Cordero P, Casabona-Frances S, Arias A, Alcedo J, Garcia-Romero R, et al. Step-up empiric elimination diet for pediatric and adult eosinophilic esophagitis: The 2-4-6 study. *Gastroenterology* 2017;**152**:S207.

NCT01458418 2011 {published data only}

NCT01458418. A trial of montelukast for maintenance therapy of eosinophilic esophagitis in children. <https://ClinicalTrials.gov> (first received 18 October 2011).

NCT01498497 2011 {published data only}

NCT01498497. A six month, safety follow-up study in eosinophilic esophagitis subjects who completed study PR-021. <https://ClinicalTrials.gov> (first received 16 December 2011).

NCT01702701 2012 {published data only}

NCT01702701. Eosinophilic esophagitis treatment: montelukast vs fluticasone. <https://ClinicalTrials.gov> (first received 4 October 2012).

NCT01821898 2013 {published data only}

NCT01821898. Eosinophilic esophagitis clinical therapy comparison trial. <https://ClinicalTrials.gov> (first received 27 March 2013).

NTR4892 2014 {published data only}

NTR4892. Effect of amino acid formula on esophageal inflammation and intestinal permeability in adult eosinophilic esophagitis patients. <https://trialsearch.who.int/Trial2.aspx?TrialID=NTR4892> (first received 5 November 2014).

Safroneeva 2015 {published data only}

Safroneeva E, Coslovsky M, Kuehni CE, Zwahlen M, Haas N, Panczak R, et al. Eosinophilic oesophagitis: relationship of quality of life with clinical, endoscopic and histological activity. *Alimentary Pharmacology and Therapeutics* 2015;**42**:1000-10.

Safroneeva 2018 {published data only}

Safroneeva E, Balsiger L, Hafner D, Zwahlen M, Kuehni C, Roumet M, et al. Short- and long-term therapeutic goals of adult patients with eosinophilic esophagitis. *Gastroenterology* 2018;**154**:S256.

Safroneeva 2018a {published data only}

Safroneeva E, Roumet MC, Coslovsky M, Kuehni C, Zwahlen M, Hirano I, et al. Both fibrotic and inflammatory endoscopic alterations contribute to symptom severity in adults with eosinophilic esophagitis. *Gastroenterology* 2018;**154**:S77.

Savarino 2015 {published data only}

Savarino E, De Cassan C, Ottavia B, Caccaro R, Galeazzi F, Nicoletti L, et al. Clinical and high resolution manometry data support the hypothesis that proton pump inhibitor-

Medical treatment of eosinophilic esophagitis (Review)

responsive esophageal eosinophilia represent a GERD-related phenomenon. *Gastroenterology* 2015;**148**:S415.

Song 2020 {published data only}

Song I, Finkelman R, Lan L. A pharmacokinetic bridging study to compare systemic exposure to budesonide between budesonide oral suspension and entocort ec in healthy individuals. *Drugs in R and D* 2020;**20**:359-67.

Spergel 2002 {published data only}

Spergel JM, Beausoleil JL, Mascarenhas M, Liacouras CA. The use of skin prick tests and patch tests to identify causative foods in eosinophilic esophagitis. *Journal of Allergy and Clinical Immunology* 2002;**109**:363-8.

Spergel 2005 {published data only}

Spergel JM, Andrews T, Brown-Whitehorn TF, Beausoleil JL, Liacouras CA. Treatment of eosinophilic esophagitis with specific food elimination diet directed by a combination of skin prick and patch tests. *Annals of Allergy, Asthma & Immunology* 2005;**95**:336-43.

Syversen 2020 {published data only}

Syversen E, Hait E, McDonald D, Rubinstein E, Goldsmith J, Ngo P, et al. Oral viscous mometasone is an effective treatment for eosinophilic esophagitis. *Journal of Allergy and Clinical Immunology* 2020;**8**:1107-9.

Tripp 2017 {published data only}

Tripp C, Cuff C, Campbell AL, Hendrickson BA, Voss J, Melim T, et al. RPC4046, a novel anti-interleukin-13 antibody, blocks IL-13 binding to IL-13 alpha 1 and alpha 2 receptors: a randomized, double-blind, placebo-controlled, dose-escalation first-in-human study. *Advances in Therapy* 2017;**34**(6):1364-81. [DOI: [10.1007/s12325-017-0525-8](https://doi.org/10.1007/s12325-017-0525-8)] [PMID: 28455782]

Tripp C, Cuff C, Miller R, Wu C, Campbell A, Hendrickson B, et al. The characterization of RPC4046, a novel anti-IL13 monoclonal antibody that blocks binding to IL-13 receptor alpha1 and alpha 2 in development for eosinophilic esophagitis: Results of a first-inhuman study. *Annals of Allergy, Asthma and Immunology* 2014;**113**(5 Suppl 1):A104-5.

Vazquez-Elizondo 2013 {published data only}

Vazquez-Elizondo G, Ngamruengphong S, Devault K R, Talley N, Achem S, Khrisna M. The outcome of patients with oesophageal eosinophilic infiltration after an eight-week trial of a proton pump inhibitor. *Alimentary Pharmacology and Therapeutics* 2013;**38**:1312-9.

Wang 2017 {published data only}

Wang R, Hirano I, Doerfler B, Zalewski A, Gonsalves N, Taft T. Assessing adherence and barriers to long term dietary elimination therapy in patients with eosinophilic esophagitis (EoE). *Digestive Diseases and Sciences* 2017;**152**:S864.

Wang R, Hirano I, Doerfler B, Zalewski A, Gonsalves N, Taft T. Assessing adherence and barriers to long-term elimination diet therapy in adults with eosinophilic esophagitis. *Digestive Diseases and Sciences* 2018;**63**:1756-62.

Warners 2016 {published data only}

Warners M, Vlieg-Boerstra B, Verheij J, Van Ampting M, Harthoorn L, De Jonge W, et al. Amino acid-based diet restores the esophageal mucosal integrity in eosinophilic esophagitis whereas duodenal mucosal integrity remains unaffected. *Gastroenterology* 2016;**150**:S18-9.

Wechsler 2017 {published data only}

Wechsler J, Amsden K, Schwartz S, Manuel-Rubio M, Sulkowski M, Johnson K, et al. Efficacy of a 4-food elimination diet for children with eosinophilic esophagitis. *Clin Gastroenterol Hepatol* 2017;**15**:1698.

Wright 2020 {published data only}

Wright B, Rank M, Kita H, Katzka D, Shim K, Doyle A, et al. Peanut oral immunotherapy induces gastrointestinal eosinophilia in a longitudinal randomized controlled trial. *Journal of Allergy and Clinical Immunology* 2020;**145**:AB84.

Wright 2021 {published data only}

Wright BL, Fernandez-Becker NQ, Kambham N, Purington N, Cao S, Tupa D, et al. Gastrointestinal eosinophil responses in a longitudinal, randomized trial of peanut oral immunotherapy. *Clinical Gastroenterology and Hepatology* 2021;**19**:1151-9.

References to studies awaiting assessment

Amini 2022 {published data only}

* Amini N, Khademian M, Momen T, Saneian H, Nasri P, Famouri F, et al. The effectiveness of synbiotic on the improvement of clinical symptoms in children with eosinophilic esophagitis. *International Journal of Pediatrics* 2022;**2022**:4211626.

NCT01846962 2012 {published data only}

NCT01846962. Dietetic versus topical steroids for pediatric eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT01846962> 2012;(first received 6 May 2013).

References to ongoing studies

ACTRN12619000141145 2019 {published data only}

ACTRN12619000141145. Two food group elimination diet versus swallowed fluticasone for the management of adult eosinophilic esophagitis, a single-centred randomised prospective study. <https://trialssearch.who.int/Trial2.aspx?TrialID=ACTRN12619000141145> (first received 30 January 2019).

ACTRN12621001406897 {published data only}

ACTRN12621001406897. Efficacy of Fucoidan for eosinophilic oesophagitis: a phase 2 pilot study. <https://trialssearch.who.int/Trial2.aspx?TrialID=ACTRN12621001406897> (first received 19 October 2021).

EUCTR2017-003516-39-ES 2021 {published data only}

EUCTR2017-003516-39-DE. A phase III clinical study in adult and adolescent patients with eosinophilic inflammation of the gutlet to prove superiority compared to placebo of an episodic and/or a continuous 48-week treatment with budesonide orodispersible tablets for maintaining remission. <https://>

Medical treatment of eosinophilic esophagitis (Review)

www.clinicaltrialsregister.eu/ctr-search/trial/2017-003516-39/DE (first received 21 December 2020).

EUCTR2017-003516-39-ES. A phase III clinical study in adult and adolescent patients with eosinophilic inflammation of the gullet to prove superiority compared to placebo of an episodic and/or a continuous 48-week treatment with budesonide orodispersible tablets for maintaining remission. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2017-003516-39/ES> (first received 21 December 2020).

EUCTR2017-003737-29-ES 2019 {published data only}

EUCTR2017-003737-29-DE. Double-blind (neither physician nor patient knows of the actual treatment which can be with or without active substance), randomized (patient will be allocated to a certain treatment group by chance), placebo-controlled (one of the treatment groups receive. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2017-003737-29/DE> (first received 17 December 2018).

EUCTR2017-003737-29-ES. Double-blind (neither physician nor patient knows of the actual treatment which can be with or without active substance), randomized (patient will be allocated to a certain treatment group by chance), placebo-controlled (one of the treatment groups receive. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2017-003737-29/ES> 2019.

EUCTR2019-002871-32-ES 2019 {published data only}

EUCTR2019-002871-32-ES. A study of benralizumab in patients with eosinophilic esophagitis. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2019-002871-32/ES> (first received 5 February 2019).

EUCTR2019-002871-32-NL. A study of benralizumab in patients with eosinophilic esophagitis. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2019-002871-32/NL> (first reviewed 2 March 2020).

EUCTR2019-004391-19-NL 2020 {published data only}

EUCTR2019-004391-19-NL. Study to evaluate the efficacy and safety of AK002 in patients with active eosinophilic esophagitis. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2019-004391-19/NL> (first received 2 March 2020).

EUCTR2020-000082-16-DE 2020 {published data only}

EUCTR2020-000082-16-DE. A study to investigate the efficacy and tolerability of the drug ESO-101 in adult patients with inflammation of the esophagus. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-000082-16/DE> (first received 29 July 2020).

NCT04849390. A study to investigate the efficacy and tolerability of ESO-101 in patients With eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT04849390> (first received 19 April 2021).

EUCTR2020-001314-37-DE 2020 {published data only}

EUCTR2020-001314-37-DE. clinical study to show equal clinical efficacy of two dosing regimen of budesonid orodispersible tablets (twice daily vs. once daily) for treatment of inflammation of the esophagus. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-001314-37/DE> (first received 27 November 2020).

[search/trial/2020-001314-37/DE](https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-001314-37/DE) (first received 27 November 2020).

EUCTR2020-003226-23-BE 2020 {published data only}

EUCTR2020-003226-23-BE. A study evaluating the efficacy and safety of etrasimod in the treatment of patients with eosinophilic esophagitis. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-003226-23/BE> (first received 23 November 2020).

NCT04682639. A study to assess the safety and efficacy of oral etrasimod in adult participants with eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT04682639> (first received 19 December 2020).

EUCTR2020-004336-16-DE 2021 {published data only}

EUCTR2020-004336-16-DE. A study to evaluate the efficacy and safety of CC-93538 in adult and adolescent patients who have eosinophilic esophagitis. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-004336-16/DE> (first received 29 September 2021).

NCT04753697. A study to evaluate the efficacy and safety of CC-93538 in adult and adolescent participants With eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT04753697> (first received 29 September 2021).

Henry 2019 {published data only}

* Henry S, Al'Atrash H, James F, Molle-Rios Z, Lee D, Suskind D, et al. A novel food additive removal diet for eosinophilic esophagitis (free study): conceptual design and clinical trial methods. *Journal of Pediatric Gastroenterology and Nutrition* 2019;**69**(Suppl 2):195. [DOI: <http://dx.doi.org/10.1097/MPG.0000000000002518>]

IRCT20171230038142N14 2020 {published data only}

IRCT20171230038142N14. evaluation of efficacy of synbiotics in children with eosinophilic esophagitis. <https://en.irct.ir/search/result?query=IRCT20171230038142N14> (first received 11 March 2020).

IRCT20191211045703N1 2020 {published data only}

IRCT20191211045703N1. Comparison of the efficacy and side effects of nebulized oral pulmicort and inhaler budesonide in patients with eosinophilic esophagitis. <https://en.irct.ir/search/result?query=IRCT20191211045703N1> (first received 24 June 2020).

JPRN-jRCT2051200140 {published data only}

JPRN-jRCT2051200140. A phase 3, multicenter, multinational, randomized, double-blind, placebo-controlled induction and maintenance study to evaluate the efficacy and safety of CC-93538 in adult and adolescent subjects with eosinophilic esophagitis. <https://trialsearch.who.int/Trial2.aspx?TrialID=JPRN-jRCT2051200140> (first received 19 February 2021).

NCT02873468 {published data only}

NCT02873468. Efficacy and safety of three doses of florence oral suspension in adults with eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT02873468> (first received 16 August 2016).

NCT03656380 2019 {published data only}

NCT03656380. Mepo for EoE study. <https://clinicaltrials.gov/ct2/show/NCT03656380> (first received 18 August 2018).

NCT03657771 2018 {published data only}

NCT03657771. A food additive removal diet for pediatric eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT03657771> (first received 31 August 2018).

NCT03781596 {published data only}

NCT03781596. EoE RCT fluticasone and omeprazole vs fluticasone alone. <https://clinicaltrials.gov/ct2/show/NCT03781596> (first received 20 December 2018).

NCT04248712 2020 {published data only}

NCT04248712. Antihistamines in eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT04248712> (first received 28 January 2020).

NCT04281108 2020 {published data only}

EUCTR2019-001521-27-DE. Study with eosinophilic esophagitis subjects. <https://www.clinicaltrialsregister.eu/ctr-search/trial/2019-001521-27/DE> (first received 3 August 2020).

NCT04281108. Efficacy and safety APT-1011 in adult subjects with eosinophilic esophagitis (EoE) (FLUTE-2). <https://clinicaltrials.gov/ct2/show/NCT04281108> (first received 20 February 2020).

NCT05083312. Efficacy and safety APT-1011 in adolescent subjects with eosinophilic esophagitis (EoE) - a sub-study of the FLUTE-2 Trial. <https://clinicaltrials.gov/ct2/show/NCT05083312> (first received 6 October 2021).

NCT04394351 2020 {published data only}

NCT04394351. Study to investigate the efficacy and safety of dupilumab in pediatric patients with active eosinophilic esophagitis (EoE). <https://clinicaltrials.gov/ct2/show/NCT04394351> (first received 1 September 2020).

NCT04543409 {published data only}

NCT04543409. A study of benralizumab in patients with eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT04543409> 2020.

NCT04593251 2020 {published data only}

NCT04593251. Dose escalation study to evaluate an experimental new treatment (CALY-002) in healthy subjects and subjects with celiac disease and eosinophilic esophagitis. <https://ClinicalTrials.gov> (first received 13 October 2020).

NCT04835168 2022 {published data only}

NCT04835168. Safety and pharmacokinetics of orodispersible BT-11 in active eosinophilic esophagitis. <https://clinicaltrials.gov/ct2/show/NCT04835168> (first received 8 April 2021).

NCT05084963 2021 {published data only}

NCT05084963. A study to assess the efficacy, safety and tolerability of IRL201104 in adults with active eosinophilic

esophagitis. <https://clinicaltrials.gov/ct2/show/NCT05084963> (first received 20 October 2021).

NCT05214599 {published data only}

NCT05214599. Pharmacokinetics, Efficacy, Tolerability and Safety of Different Budesonide Oral Gel Doses in Adults' Subjects of Both Genders With Eosinophilic Esophagitis (EoE). <https://ClinicalTrials.gov/show/NCT05214599> (first received 28 January 2022).

NCT05543512 {published data only}

NCT05543512. The Immune Directed Individualized Elimination Therapy (iDIET) Study. <https://ClinicalTrials.gov/show/NCT05543512> (first received 16 September 2022).

NCT05583227 {published data only}

EUCTR2022-001294-31. Efficacy and safety of tezepelumab in patients with eosinophilic esophagitis. <https://trialssearch.who.int/Trial2.aspx?TrialID=EUCTR2022-001294-31-DK-2022> (first received 4 October 2022).

NCT05583227. Efficacy and safety of tezepelumab in patients with eosinophilic esophagitis. <https://ClinicalTrials.gov/show/NCT05583227> (first received 17 October 2022).

NCT05634746 {published data only}

NCT05634746. 24-week induction study of APT-1011 in adult subjects with eosinophilic esophagitis (EoE) (FLUTE 3). <https://ClinicalTrials.gov/show/NCT05634746> (first received 2 December 2022).

NCT05695456 {published data only}

NCT05695456. Targeted elimination diet in EoE patients following identification of trigger nutrients using confocal laser endomicroscopy. <https://ClinicalTrials.gov/show/NCT05695456> (first received 25 January 2023).

Additional references

Aanen 2006

Ananen MC, Numans ME, Weusten BLAM, Smout AJPM. Diagnostic value of the Reflux Disease Questionnaire in general practice. *Digestion* 2006;**74**:162-8.

Aceves 2009

Aceves S, Newbury R, Dohil M, Bastian J, Dohil R. A symptom scoring tool identifies pediatric eosinophilic esophagitis patients and correlates with tissue inflammation. *Annals of Allergy Asthma and Immunology* 2009;**103**(5):401-6. [DOI: [10.1016/S1081-1206\(10\)60359-6](https://doi.org/10.1016/S1081-1206(10)60359-6)] [PMID: 19927538]

Collins 2017

Collins M, Martin L, Alexander E, Boyd J, Sheridan R, He H, et al. Newly developed and validated eosinophilic esophagitis histology scoring system and evidence that it outperforms peak eosinophil count for disease diagnosis and monitoring. *Diseases of the Esophagus* 2017;**30**:1-8.

Covidence [Computer program]

Covidence. Melbourne, Australia: Veritas Health Innovation, 2022. Available at www.covidence.org.

Medical treatment of eosinophilic esophagitis (Review)

Croese 2003

Croese J, Fairley SK, Masson JW, et al. Clinical and endoscopic features of eosinophilic esophagitis in adults. *Gastrointestinal Endoscopy* 2003;**58**:516-22.

Dakkak 1992

Dakkak M, Bennett JR. A new dysphagia score with objective validation. *Gastroenterology* 1992;**14**(2):99-100.

Deeks 2021

Deeks JJ, Higgins JPT, Altman DG, on behalf of the Cochrane Statistical Methods Group. Chapter 10: Analysing data and undertaking meta-analyses. In: Higgins JP, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, Welch VA, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 6.2 (updated February 2021). Cochrane, 2021. Available from training.cochrane.org/handbook.

Dellon 2018

Dellon ES, Liacouras CA, Molina-Infante J, Furuta GT, Spergel JM, Zevit N, et al. Updated International Consensus Diagnostic Criteria for Eosinophilic Esophagitis: Proceedings of the AGREE Conference. *Gastroenterology* 2018;**155**(4):1022-33.

Dhar 2022

Dhar A, Haboubi H, Attwood S, Auth M, Dunn J, Sweis R, et al. British Society of Gastroenterology (BSG) and British Society of Paediatric Gastroenterology, Hepatology and Nutrition (BSPGHAN) joint consensus guidelines on the diagnosis and management of eosinophilic oesophagitis in children and adults. *Gut* 2022;**71**:1459-87.

DiSario 2002

DiSario JA, Pedersen PJ, Bichiş-Canoutas C, Alder SC, Fang JC. Incision of recurrent distal esophageal (Schatzki) ring after dilation. *Gastrointestinal Endoscopy* 2002;**56**(2):544-8.

Donnan 2020

Donnan E, Pandolfino J. EndoFLIP in the esophagus: assessing sphincter function, wall stiffness, and motility to guide treatment. *Gastroenterology Clinics of North America* 2020;**49**:427-35.

Egger 1997

Egger M, Smith GD, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;**315**(7109):629-34.

Faubion 1998

Faubion WA Jr, Perrault J, Burgart LJ, Zein NN, Clawson M, Freese DK. Treatment of eosinophilic esophagitis with inhaled corticosteroids. *Journal of Pediatric Gastroenterology and Nutrition* 1998;**27**(1):90-3.

Flood 2008

Flood EM, Beusterien KM, Amonkar MM, Jurgensen CH, Dewit OE, Kahl LP, et al. Patient and caregiver perspective on pediatric eosinophilic esophagitis and newly developed symptom questionnaires. *Current Medical Research and Opinion* 2008;**24**(12):3369-81. [DOI: [10.1185/03007990802536900](https://doi.org/10.1185/03007990802536900)] [PMID: 19032119]

Franciosi 2011

Franciosi JP, Hommel KA, DeBrosse CW, Greenberg AB, Greenler AJ, Abonia JP, et al. Development of a validated patient-reported symptom metric for pediatric eosinophilic esophagitis: qualitative methods. *BMC Gastroenterology* 2011;**11**:126. [DOI: [10.1186/1471-230X-11-126](https://doi.org/10.1186/1471-230X-11-126)] [PMID: 22099448]

Franciosi 2013

Franciosi JP, Hommel KA, Bendo CB, King EC, Collins MH, Eby MD, et al. PedsQL™ Eosinophilic Esophagitis Module: feasibility, reliability and validity. *Journal of Pediatric Gastroenterology and Nutrition* 2013;**57**(1):57-66.

Franciosi 2022

Franciosi JP, Mougey EB, Dellon ES, Gutiérrez-Junquera C, Fernández-Fernández S, Venkatesh RD, et al. Proton pump inhibitor therapy for eosinophilic esophagitis: history, mechanisms, efficacy, and future directions. *Journal of Asthma and Allergy* 2022;**15**:281-302.

Furuta 2007

Furuta GT, Liacouras CA, Collins MH, et al. First International Gastrointestinal Eosinophil Research Symposium (FIGERS). *Gastroenterology* 2007;**133**:1342-63.

Furuta 2015

Furuta GT, Katzka DA. Eosinophilic esophagitis. *New England Journal of Medicine* 2015;**373**(17):1640-8.

Gordon 2021

Gordon M, Lakunina S, Sinopoulou V, et al. Minimum sample size estimates for trials in inflammatory bowel disease: a systematic review of a support resource. *World Journal of Gastroenterology* 2021;**27**:7572.

Greuter 2017

Greuter T, Bussmann C, Safroneeva E, Schoepfer AM, Biedermann L, Vavricka SR, et al. Long-term treatment of eosinophilic esophagitis with swallowed topical corticosteroids: development and evaluation of a therapeutic concept. *Official Journal of the American College of Gastroenterology* 2017;**112**(10):1527-35.

Hassall 1996

Hassall E. Macroscopic versus microscopic diagnosis of reflux esophagitis: erosions or eosinophils? *Journal of Pediatric Gastroenterology and Nutrition* 1996;**22**(3):321-5.

Higgins 2011

Higgins JPT, Green S, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 5.1.0 (updated March 2011). The Cochrane Collaboration, 2011. Available from training.cochrane.org/handbook/archive/v5.1/.

Higgins 2021a

Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 6.2 (updated February 2021). Cochrane, 2021. Available from www.training.cochrane.org/handbook.

Medical treatment of eosinophilic esophagitis (Review)

Higgins 2021b

Higgins JPT, Li T, Deeks JJ, editor(s). Chapter 6: Choosing effect measures and computing estimates of effect. In: Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al, editor(s). *Cochrane Handbook for Systematic Reviews of Interventions* Version 6.2 (updated February 2021). Cochrane, 2021. Available from www.training.cochrane.org/handbook.

Hirano 2013

Hirano I, Moy N, Heckman MG, Thomas CS, Gonsalves N, Achem SR. Endoscopic assessment of the oesophageal features of eosinophilic esophagitis: validation of a novel classification and grading system. *Gut* 2013;**62**(4):489-95.

Hudgens 2017

Hudgens S, Evans C, Phillips E, Hill M. Psychometric validation of the Dysphagia Symptom Questionnaire in patients with eosinophilic esophagitis treated with budesonide oral suspension. *Journal of Patient-Reported Outcomes* 2017;**1**(3):1-11. [DOI: [10.1186/s41687-017-0006-5](https://doi.org/10.1186/s41687-017-0006-5)] [PMID: 29757322]

Iheozor-Ejiofor 2021

Iheozor-Ejiofor Z, Lakunina S, Gordon M, Akintelure D, Sinopoulou V, Akobeng A. Sample-size estimation is not reported in 24% of randomised controlled trials of inflammatory bowel disease: a systematic review. *United European Gastroenterology Journal* 2021;**9**:47-53.

Kelly 1995

Kelly KJ, Lazenby AJ, Rowe PC, Yardley JH, Perman JA, Sampson HA. Eosinophilic esophagitis attributed to gastroesophageal reflux: improvement with an amino acid-based formula. *Gastroenterology* 1995;**109**(5):1503-12.

Khan 2003

Khan S, Orenstein SR, Di Lorenzo C, Kocoshis SA, Putnam PE, Sigurdsson L, et al. Eosinophilic Esophagitis: Strictures, Impaction, Dysphagia. *Digestive Diseases and Sciences* 2003;**48**(1):22-9.

Landgraf 2014

Landgraf JM. Child Health Questionnaire (CHQ). In: Michalos AC, editors(s). *Encyclopedia of Quality of Life and Well-Being Research*. Dordrecht: Springer, 2014:698–702. [DOI: [10.1007/978-94-007-0753-5_324](https://doi.org/10.1007/978-94-007-0753-5_324)]

Liacouras 1998

Liacouras CA, Wenner WJ, Brown K, Ruchelli E. Primary eosinophilic esophagitis in children: successful treatment with oral corticosteroids. *Journal of Pediatric Gastroenterology and Nutrition* 1998;**26**(4):380-5.

Liacouras 2011

Liacouras CA, Furuta GT, Hirano I, Atkins D, Attwood SE, Bonis PA, et al. Eosinophilic esophagitis: updated consensus recommendations for children and adults. *Journal of Allergy and Clinical Immunology* 2011;**128**(1):3-20.

Lucendo 2017

Lucendo A, Molina-Infante J, Arias A, Von Arnim U, Bredenoord A, Bussmann C, et al. Guidelines on eosinophilic esophagitis: evidence-based statements and recommendations for diagnosis and management in children and adults. *UEG Journal* 2017;**5**:335-58.

Ma 2022a

Ma C, Bredenoord A, Dellon E, Alexander J, Bierdermann L, Hogan M, et al. Reliability and responsiveness of endoscopic disease activity assessment in eosinophilic esophagitis. *Gastrointestinal Endoscopy* 2022;**95**:1126-37.

Ma 2022b

Ma C, Schoepfer A, Dellon E, Bredenoord A, Chehade M, Collins M, et al. Development of a core outcome set for therapeutic studies in eosinophilic esophagitis (COREOS). *Journal of Allergy and Clinical Immunology* 2022;**149**:659-70.

Markowitz 2003

Markowitz JE, Spergel JM, Ruchelli E, Liacouras CA. Elemental diet is an effective treatment for eosinophilic esophagitis in children and adolescents. *American Journal of Gastroenterology* 2003;**98**(4):777-82.

McElhiney 2009

McElhiney J, Lohse MR, Arora AS, Peloquin JM, Geno DM, Kuntz MM, et al. The Mayo Dysphagia Questionnaire-30: documentation of reliability and validity of a tool for interventional trials in adults with esophageal disease. *Dysphagia* 2009;**25**:221–30.

Muir 2021

Muir A, Falk GW. Eosinophilic esophagitis: a review. *JAMA* 2021;**326**(13):1310-8.

Nhu 2022

Nhu QM, Aceves SS. Current state of biologics in treating eosinophilic esophagitis. *Annals of Allergy, Asthma & Immunology* 2022;**22**:S1081-206.

Orenstein 2000

Orenstein SR, Shalaby TM, Di Lorenzo C, Putnam PE, Sigurdsson L, Kocoshis SA. The spectrum of pediatric eosinophilic esophagitis beyond infancy: a clinical series of 30 children. *American Journal of Gastroenterology* 2000;**95**(6):1422-30.

Peloquin 2006

Peloquin JM, Fredericksen M, Geno D, Abdalla AA, McElhiney J, Khan Y, et al. Mayo Dysphagia Questionnaire –2 week: validation of a symptom questionnaire. *American Journal of Gastroenterology* 2006;**101**:s418.

Pentiuk 2009

Pentiuk S, Putnam PE, Collins MH, Rothenberg ME. Dissociation between symptoms and histological severity in pediatric eosinophilic esophagitis. *Journal of Pediatric Gastroenterology and Nutrition* 2009;**48**:152-60.

PRISMA 2020

Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021;**372**:71.

Rank 2020

Rank MA, Sharaf RN, Furuta GT, Aceves AA, Greenhawt M, Spergel JM, et al. Technical review on the management of eosinophilic esophagitis: a report from the AGA institute and the joint task force on allergy-immunology practice parameters. *Annals of Allergy, Asthma & Immunology* 2020;**124**(5):424-40.

Reed 2018

Reed CC, Wolf WA, Cotton CC, Rusin S, Perjar I, Hollyfield J, et al. Optimal histologic cutpoints for treatment response in patients with eosinophilic esophagitis: analysis of data from a prospective cohort study. *Clinical Gastroenterology and Hepatology* 2018;**16**(2):226-33.

RevMan Web 2022 [Computer program]

Review Manager Web (RevMan Web). Version 4.1.0. The Cochrane Collaboration, 2022. Available at revman.cochrane.org.

Rothenberg 2015

Rothenberg ME. Molecular, genetic, and cellular bases for treating eosinophilic esophagitis. *Gastroenterology* 2015;**148**:1143-57.

Schoepfer 2013

Schoepfer AM, Safroneeva E, Bussmann C, et al. Delay in diagnosis of eosinophilic esophagitis increases risk for stricture formation in a time-dependent manner. *Gastroenterology* 2013;**145**(6):1230-6.

Schoepfer 2014

Schoepfer AM, Straumann A, Panczak R, Coslovsky M, Kuehni CE, Maurer E, et al. Development and validation of a symptom-based activity index for adults with eosinophilic esophagitis. *Gastroenterology* 2014;**147**(6):1255-66. [DOI: [10.1053/j.gastro.2014.08.028](https://doi.org/10.1053/j.gastro.2014.08.028)] [PMID: 25160980]

Schünemann 2021

Schünemann HJ, Higgins JPT, Vist GE, Glasziou P, Akl EA, Skoetz N, et al. Chapter 14: Completing 'Summary of findings' tables and grading the certainty of the evidence. In: Higgins

JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al, editor(s). Cochrane Handbook for Systematic Reviews of Interventions Version 6.2 (updated February 2021). Cochrane, 2021. Available from www.training.cochrane.org/handbook.

Straumann 2003

Straumann A, Spichtin HP, Grize L, et al. Natural history of primary eosinophilic esophagitis: a follow-up of 30 adult patients for up to 11.5 years. *Gastroenterology* 2003;**125**(6):1660-9. [DOI: [10.1053/j.gastro.2003.09.024](https://doi.org/10.1053/j.gastro.2003.09.024)] [PMID: 14724818]

Straumann 2010

Straumann A, Conus S, Degen L, Felder S, Kummer M, Engel H, et al. Budesonide is effective in adolescent and adult patients with active eosinophilic esophagitis. *Gastroenterology* 2010;**139**:1526-37.

Straumann 2013

Straumann A, Hoesli S, Bussmann CH, Stuck M, Perkins M, Collins LP, et al. Anti-eosinophil activity and clinical efficacy of the CRTH2 antagonist OC000459 in eosinophilic esophagitis. *Allergy* 2013;**68**(3):375-85.

Taft 2011

Taft TH, Kern E, Kwiatek MA, Hirano I, Gonsalves N, Keefer L. The adult eosinophilic oesophagitis quality of life questionnaire: a new measure of health-related quality of life. *Alimentary Pharmacology & Therapeutics* 2011;**34**(7):790-8.

Teitelbaum 2002

Teitelbaum JE, Fox VL, Twarog FJ, Nurko S, Antonilli D, Gleich G, et al. Eosinophilic oesophagitis in children: immunopathological analysis and response to fluticasone propionate. *Gastroenterology* 2002;**122**(5):1216-25.

References to other published versions of this review
Elliott 2010

Elliott EJ, Thomas D, Markowitz JE. Non-surgical interventions for eosinophilic esophagitis. *Cochrane Database of Systematic Reviews* 2010, Issue 3. Art. No: CD004065. [DOI: [10.1002/14651858.CD004065.pub3](https://doi.org/10.1002/14651858.CD004065.pub3)]

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES
Characteristics of included studies [ordered by study ID]
Alexander 2012
Study characteristics

Methods	RCT design and number of study arms: RCT; 2 arms
	Single-center or multi-center: single-center; Esophageal Clinic at Mayo Clinic Rochester Minnesota
	Countries: USA

Medical treatment of eosinophilic esophagitis (Review)

Alexander 2012 (Continued)

Study dates: October 2005 to December 2009

Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: ≥ 20 eosinophils/hpf (peak)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 18 to 65 years of age • Mayo Dysphagia Questionnaire: 1a) "Have you had trouble swallowing unrelated to a sore throat or cold?" - yes; 1c) Severity \geq moderate; 2) Frequency - at least once per week • ≥ 20 peak eosinophils/hpf on biopsies from mid esophagus (hpf = high-powered field) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Patients with endoscopic evidence of stricture, Schatzki ring, mass, or Los Angeles grade C or D esophagitis at endoscopy • Topical steroid therapy for EoE at any time in the past • Esophageal dilation at the index endoscopy or at any time in the past <p>Age at beginning of study per study group: fluticasone mean (range): 37 years (19 to 59); placebo mean (range): 38 (20 to 57)</p> <p>Sex (m/f) per study group: fluticasone (m/f): 18/3; placebo (m/f): 16/5</p> <p>Number randomized per study group: fluticasone: 21; placebo: 21</p> <p>Number reaching end of study per study group: fluticasone: 19; placebo: 15</p>
Interventions	<p>Study group 1 (placebo): placebo inhaler swallowed twice a day for 6 weeks</p> <p>Study group 2 (fluticasone): aerosolized swallowed fluticasone 880 μg twice a day for 6 weeks</p>
Outcomes	<p>Primary outcomes of the study: number of participants with complete response to dysphagia (time frame: 2 weeks), measured by the Mayo Dysphagia Questionnaire, a validated 28-item instrument: 0 = no dysphagia, higher levels indicate greater dysphagia severity. A complete symptom response was defined as an answer of "no" to the question "In the past 2 weeks, have you had trouble swallowing, not associated with other cold symptoms (such as strep throat or mono)?"</p> <p>Secondary outcomes of the study:</p> <ol style="list-style-type: none"> 1. Number of participants with complete response to dysphagia (time frame: 2 weeks), measured by the Mayo Dysphagia Questionnaire, a validated 28 item instrument; 0 = no dysphagia, higher levels indicate greater dysphagia severity. A partial symptom response was defined as an answer of "yes" to the question "In the past 2 weeks, have you had trouble swallowing, not associated with other cold symptoms (such as strep throat or mono)?" and a decrease in severity of at least 2 levels, or a decrease in frequency of at least 1 level. 2. Number of participants with complete histologic response (time frame: 2 weeks) <p>A complete histologic response was defined as a $> 90\%$ decrease in mean eosinophil count/high-powered field. A partial response was defined as a decrease of more than 50% from the pre-treatment value.</p>
Notes	<p>Funding source: Mayo Clinic</p> <p>Conflicts of interest: Jeffrey Alexander is a consultant for Meritage Pharmacia and has stock ownership in Meritage Pharmacia</p>
Risk of bias	
Bias	Authors' judgement Support for judgement

Medical treatment of eosinophilic esophagitis (Review)

Alexander 2012 (Continued)

Random sequence generation (selection bias)	Low risk	Computer-generated, coded randomization method.
Allocation concealment (selection bias)	Low risk	"The pharmacist and esophageal nurse saw the actual inhalers which were similar but not identical; all other study personnel were blinded to the study treatment allocation. Other than the inhaler education session, the esophageal nurse had no other subject contact or involvement with the study."
Blinding of participants and personnel (performance bias) All outcomes	Low risk	The author confirmed that both treatments "were inhalers with no particular taste" and caregivers were also blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The author confirmed that "the outcome assessors were blinded".
Incomplete outcome data (attrition bias) All outcomes	Low risk	2 participants from the fluticasone arm (1 travel, 1 scheduling) and 6 participants from the placebo arm (1 withdrawal by participant, 1 travel, 2 scheduling, 2 family issues) did not finish the study. Balanced and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Low risk	Protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	PPI use was continued during the study by 26.3% (5 of 19) of the fluticasone group and 0% (0 of 15) of the placebo group No major baseline differences between groups. No other concerns.

Alexander 2017
Study characteristics

Methods	<p>RCT design and number of study arms: RCT, 2 arms</p> <p>Single-center or multi-center: single-center, Esophageal Clinic at Mayo Clinic in Rochester Minnesota</p> <p>Countries: USA</p> <p>Study dates: April 2008 to February 2015</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: inactive</p> <p>EoE definition/diagnostic criteria: ≥ 20 eosinophils/hpf (peak); Los Angeles Grade A or B after steroid treatment</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 18 to 65 years of age • Mayo Dysphagia Questionnaire: (1a) "Have you had trouble swallowing unrelated to a sore throat or cold?" - yes; (1c) Severity \geq moderate; (2) Frequency \geq 1/week • ≥ 20 peak eosinophils/HPF on biopsies from mid-esophagus at baseline • Los Angeles Grade A or B after steroid treatment <p>Exclusion criteria:</p>

Medical treatment of eosinophilic esophagitis (Review)

Alexander 2017 (Continued)

- Patients with endoscopic evidence of stricture, Schatzki ring, mass, or Los Angeles grade C or D esophagitis at endoscopy
- Patients with other diseases known to be associated with esophageal eosinophilia
- Patients who underwent esophageal dilation within a year
- Patients without a complete symptomatic response to topical steroid therapy

Age at beginning of study per study group: montelukast mean (SD): 44.4 (13.7); placebo mean (SD): 40.4 (12.1)

Sex (m/f) per study group: montelukast m/f = 13/7 (65.0%/35%); placebo m/f = 12/9 (57.1%/42.9%)

Number randomized per study group: montelukast n = 20; placebo n = 21

Number reaching end of study per study group: montelukast n = 18; placebo n = 20

Interventions	<p>Study group 1: placebo tablets, similar-appearing</p> <p>Study group 2: montelukast 20 mg tablets for 26 weeks or until they developed symptomatic dysphagia</p>
Outcomes	<p>Primary outcomes of the study: symptomatic remission is defined as the absence of dysphagia as defined above</p> <p>Secondary outcomes of the study: unclear - outcomes not properly defined in the paper but defined in the protocol: "evaluate safety of montelukast in eosinophilic esophagitis"</p>
Notes	<p>Funding source: this work was supported by a grant from Merck (Kenilworth, NJ)</p> <p>Conflicts of interest: "These authors disclose the following: Jeffrey A. Alexander owns stock in Meritage Pharmacia/Shire, and has received research funding from Merck and Shire, and David A. Katzka has received research funding from Shire and Covidien. The remaining authors disclose no conflicts."</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated, coded randomization method.
Allocation concealment (selection bias)	Low risk	Placebo tablets were similar-appearing to the montelukast tablets. Only a pharmacist, who had no participant involvement, was unblinded to the participants.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	The study was blinded to all caregivers but was known only by the pharmacist who had no participant involvement.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The author confirmed that the outcome assessors were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Unclear risk	The inclusion criteria of ≥ 20 eosinophils/HPF in the full text is higher than that in the protocol (> 14 eosinophils). This retrospectively changed higher cut-off may have potentially helped the authors achieve a desired response/remission.

Medical treatment of eosinophilic esophagitis (Review)

Alexander 2017 (Continued)

The author responded that the change was made because the initial "was a criteria used very early in EoE. We wanted to limit as much as possible overlap with GERD often associated with mild eosinophilia and the plan was changed". As this is an appropriate a priori change, we have made the judgment of low risk.

Unclear definition of pre-planned secondary outcomes.

Other bias	Low risk	No major baseline differences between groups. No other concerns.
------------	----------	--

Assa'ad 2011
Study characteristics

Methods	<p>RCT design and number of study arms: RCT with 3 arms</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: USA and UK</p> <p>Study dates: 11 September 2006 to 25 November 2008</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active EoE</p> <p>EoE definition/diagnostic criteria: "pediatric patients with isolated EoE, defined as ≥ 20 eos/hpf" - hpf size = 0.3 mm^2</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age 2 to 17 • History of inadequate histopathologic response or intolerance to prior EoE therapy <p>Exclusion criteria: concurrent eosinophilic gastrointestinal enteropathy based on the baseline endoscopy; gastroesophageal reflux disease or other causes of esophagitis; and presence or history of hypereosinophilic syndromes or collagen vascular disease, vasculitis, allergic drug reaction with peripheral eosinophilia, graft-versus-host disease, chronic idiopathic inflammatory bowel disorders, or celiac disease</p> <p>Age at beginning of study per study group: mean age (SD) in the 3 study groups (low-, medium-, high-dose) was: 10.4 (4.3), 10.5 (5.2), and 10.4 (4.7)</p> <p>Sex (m/f) per study group: sex (m/f) in the 3 study groups (low-, medium-, high-dose) was: 16/3; 14/6; 17/3</p> <p>Number randomized per study group: 19/59, 20/59, 20/59 in the low-, medium-, and high-dose groups, respectively</p> <p>Number reaching end of study per study group: 15, 19, 18 in the low-, medium-, and high dose groups, respectively</p>
Interventions	<p>Study group 1 (control - low-dose): 0.55 mg/kg mepolizumab (note: "The lowest dose was expected to be minimally efficacious and serve as a comparator group. A placebo group was not included.")</p> <p>Study group 2 (medium-dose): 2.5 mg/kg</p> <p>Study group 3 (high-dose): 10 mg/kg</p>
Outcomes	<p>Primary outcomes of the study:</p> <ul style="list-style-type: none"> • The proportion of patients with a peak eosinophil count < 5 eos/hpf in the week 12 biopsy

Medical treatment of eosinophilic esophagitis (Review)

Assa'ad 2011 (Continued)

- Safety, tolerability, and pharmacokinetics of mepolizumab

Secondary outcomes of the study:

- Changes in peak and mean eosinophil counts
- Improvements in histopathologic findings
- Improvements in endoscopic findings
- Blood eosinophils counts
- Frequency and severity of EoE symptoms
- Proportion of partial responders (eosinophil counts 5 to 19 eos/hpf)

Notes

Funding source: GSK

Conflicts of interest: All investigators, including all non-GlaxoSmithKline authors, received funding from the study sponsor (to their institution and not personally) for recruiting patients and conducting the study at their respective sites. Margaret H. Collins received funding (to the institution and not personally) to process and interpret biopsy specimens as the central review pathologist for this study. Amal H. Assa'ad has been an advisory board member and consultant for GlaxoSmithKline. Sandeep K. Gupta has been a consultant for GlaxoSmithKline. Margaret H. Collins has been an advisory board member for GlaxoSmithKline and received funding (to the institution and not personally) to process and/or interpret biopsy specimens as central review pathologist for Ception Therapeutics (now Cephalon) and Meritage Pharmaceuticals and is a consultant for Sunovion. Mike Thomson discloses no conflicts. Amy T. Heath, Deborah A. Smith, Teresa L. Perschy, Cynthia H. Jurgensen, and Hector G. Ortega are employees of GlaxoSmithKline. Seema S. Aceves has been an advisory board member for GlaxoSmithKline and has financial arrangements with Meritage Pharmaceuticals. Editorial support in the form of outline development, collating author comments, grammatical editing, and referencing was provided by Elaine F. Griffin, DPhil, at Evidence Scientific Solutions and was funded by GlaxoSmithKline.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	A randomization number was given but no details on how it was generated. Authors contacted for further details in November 2022. No response received.
Allocation concealment (selection bias)	Low risk	Patients were assigned to study treatment in accordance with the randomization schedule generated and maintained by the study sponsor. Each participant was assigned a subject number, a randomization number, and a container number. All study personnel and patients were blinded to study group allocation.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Participants and personnel were blinded and randomization schedule maintained by the study sponsor. No details on how blinding was achieved, but since the study is stated to be "double-blind" and the infusions used can be easily masked, we chose to rate this domain as low risk.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	The authors state all personnel and patients were blinded but no further details were given. Authors contacted for further details in November 2022. No response received.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Low and balanced attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	High risk	Not all primary outcomes were reported.

Medical treatment of eosinophilic esophagitis (Review)

Assa'ad 2011 (Continued)

The pre-specified outcomes in the trial registration look different than those reported in the paper. Specifically, there are a number of outcomes missing related to exploratory outcomes not related to our review.

For the clinical outcomes, the authors did not report means and SDs.

For serious and total adverse events, authors did not specify the group in which they occurred.

Authors contacted for further details in November 2022. No response received.

Other bias

Low risk

No major baseline differences between groups. No other concerns.

Bhardwaj 2017
Study characteristics

Methods

RCT design and number of study arms: randomized, double-blind, placebo-controlled, cross-over study. 2 arms followed by washout, then 2 arms.

Single Center: Penn State College of Medicine, Hershey, Pennsylvania

Countries: USA

Study dates: April 2010 and June 2011

Participants

Active EoE or inactive EoE at beginning of study: active EoE

EoE definition/diagnostic criteria: ≥ 15 eos/hpf at enrollment (not followed for 2 participants)

Inclusion criteria:

- Male or female
- 18 to 65 years of age
- With a biopsy proven diagnosis of EoE
- Able and willing to provide consent for repeated endoscopies with esophageal biopsies and blood work as per study protocol

Exclusion criteria:

- Participants with suspected or proven inflammatory bowel disease, malignancy, or collagen-vascular disease
- Participants who had used oral, inhaled, or swallowed corticosteroids in the past 3 months
- Participants who were pregnant or breastfeeding
- Participants who were not able to swallow beclomethasone dipropionate or who were intolerant to the medication
- Participants with a history of ischemic heart disease, diabetes, or dyslipidemia, unless they had been stable in the past 6 months

Age at beginning of study per study group: not provided

Sex (m/f) per study group:

- Data were reported for participants who completed the study
- Group 1: beclomethasone dipropionate (BDP) 80 μ g 2 puffs 2x/day (4/0)
- Group 2: placebo (2/3)

Number randomized per study group: total n = 13, but

- Group 1 - BDP n = 9, group 2 - placebo n = 9

Medical treatment of eosinophilic esophagitis (Review)

Bhardwaj 2017 (Continued)

Number reaching end of study per study group (numbers of patients):

- Group 1 BDP - 4 (pre-cross-over)
- Group 2 placebo - 5 (pre-cross-over)

Interventions	Study group 1: inhaler x 8 weeks; beclomethasone dipropionate 80 µg 2 puffs 2 x/day Study group 2: inhaler (not entirely clear) x 8 weeks, drug: placebo Matched placebo swallowed 2 puffs twice-daily
Outcomes	Primary outcomes of the study: number of eosinophils (peak) in esophageal tissue measured at base-line and at the end of each treatment period (mean difference) Secondary outcomes of the study: collected non-validated score Unclear - outcomes not properly defined in the paper but defined in the protocol: "evaluate safety of montelukast in eosinophilic esophagitis"
Notes	Funding source: Foundation of Young Faculty Award of the American College of Allergy, Asthma and Immunology Conflicts of interest: none

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Author correspondence confirmed randomization was computer-generated and performed by a statistician.
Allocation concealment (selection bias)	Low risk	"A physician not involved with the study other than setting up the randomization scheme and medication packets held the randomization key, and did not play any other role in the study to prevent unblinding."
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Author correspondence confirmed participants and personnel were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Author correspondence confirmed outcome assessors were blinded.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	13 randomized, but 4 withdrew consent although unclear from which groups. Authors could not provide further information on this.
Selective reporting (reporting bias)	Unclear risk	Trial has been registered prospectively. The only outcome registered was symptom improvement at 5 months, which differed from the outcomes presented in the final paper. Adverse events not reported and authors did not provide information.
Other bias	Low risk	No obvious baseline imbalance. However, the baseline population did not meet their own inclusion criteria as 2 participants without EoE were included for unclear reasons. Authors responded that all participants had clinicopathological diagnosis at one point.

Medical treatment of eosinophilic esophagitis (Review)

Bhardwaj 2017 (Continued)

No other concerns.

Butz 2014
Study characteristics

Methods	<p>RCT design and number of study arms: RCT, 2 arms</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: USA</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: ≥ 24 eosinophils/high-power field (HPF) in the proximal/distal esophagus while being treated with a proton pump inhibitor (PPI) for at least 2 months or having a negative pH probe</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Signed informed consent for study by participant, or parent/guardian if the participant is a minor. Assent obtained from all minors 11 years of age and older. Histological findings on esophageal biopsy to include peak eosinophil density ≥ 24 per high-power field (400x) in the proximal or distal esophagus validated by a pathologist at CCHMC Allergy evaluation including skin-prick testing with multiple food antigens to ensure elimination diet is not indicated Have undergone a minimum 3 months of elimination diet as indicated by skin-prick testing without detectable resolution by repeat endoscopy with biopsies demonstrating persistent eosinophilic esophagitis OR participant/parental refusal to follow an elimination diet. If the participant/parent refuses the elimination diet, they are eligible for this study. Treatment with a proton pump inhibitor for at least 2 months (rounded to the nearest month) prior to endoscopy OR failure of histological improvement as defined by < 1 eosinophil per HPF after 2 months (rounded to the nearest month) trial of proton pump inhibitor documented by prior endoscopy. The PPI must be used prior to endoscopy to rule out the possibility of GERD. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> History of poor tolerance to fluticasone propionate defined as multiple episodes of oral candidiasis, hypothalamic-pituitary-adrenal axis suppression as evidenced by signs of Cushing syndrome, headaches, or increased respiratory infections during exposure to Flovent An inability to use a metered-dose inhaler, concurrent or recent use of systemic corticosteroids Comorbid eosinophilic disorders, or a diagnosis of or being at risk for diabetes (type I or II) <p>Age at beginning of study per study group: fluticasone 12.2 (3.54 to 26.90); placebo 13.5 (4.10 to 29.80)</p> <p>Sex (m/f) per study group: fluticasone 22/6 (79%); placebo 13/1 (93%)</p> <p>Number randomized per study group: fluticasone n = 28; placebo n = 14</p> <p>Number reaching end of study per study group: fluticasone n = 23; placebo n = 13</p>
Interventions	<p>Study group 1: placebo</p> <p>Study group 2: daily 1760 μg fluticasone propionate (2 x 880 μg)</p>
Outcomes	<p>Primary outcomes of the study: remission at 3 months, remission is considered achieved when the highest eosinophil count per high-power field (HPF) in all esophageal biopsies is ≤ 1 eosinophil/HPF after 3 months of therapy</p>

Medical treatment of eosinophilic esophagitis (Review)

Butz 2014 (Continued)

Secondary outcomes of the study:

1. The secondary objectives were to measure safety via cortisol, glucose, and adverse reaction data.
2. The relationship between fluticasone propionate responsiveness and participant age, height, weight, BMI, Z-score, race, ethnicity, atopic status, compliance, and screening eosinophil level. Atopic status was defined by a personal history of allergic disease (allergic rhinitis, hay fever, atopic dermatitis, eczema, food anaphylaxis, asthma, or positive skin prick tests).

Notes

Funding source: National Institute of Allergy and Infectious Diseases grant(U01AI088806 to M.E.R.), a GlaxoSmithKline grant (109928), and the national center for Research Resources and the National Center for Advancing Translational Sciences of the National Institutes of Health (8 UL1 TR000077-05)

Conflicts of interest: Marc Rothenberg and Ting Wen are co-inventors for a pending patent based on the Eosinophilic Esophagitis Diagnostic Panel test described in this article

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated, coded randomization method
Allocation concealment (selection bias)	Low risk	The treatment was centrally allocated.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	The allocation sequence was known only to the central pharmacist at study start. Following the first 3 months of the study, a site pharmacist and a staff member were made aware of assignments to facilitate stratification.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The authors confirmed in November 2022 that the outcome assessors were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Unclear risk	The study protocol is available. An EoE symptom score was used but scores were not reported.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Clayton 2014
Study characteristics

Methods	<p>RCT design and number of study arms: RCT</p> <p>Single-center or multi-center: dual-center, University of Utah Hospital and Primary Children's Hospital, Salt Lake City, UT, both referral centers</p> <p>Countries: USA</p> <p>Study dates: NR</p>
Participants	Active EoE or inactive EoE at beginning of study: active

Medical treatment of eosinophilic esophagitis (Review)

Clayton 2014 (Continued)

EoE definition/diagnostic criteria: did not specifically define EoE in the text. Defined in the protocol as "Eosinophilic esophagitis (EE) is an increasingly recognized condition characterized by dysphagia, food impaction or other obstructive esophageal symptoms in children and young adults."

Inclusion criteria:

- Adults (> 18 years) (3 participants in the omalizumab trial aged 15 to 17)
- Active eosinophilic esophagitis
- ≥ 15 eosinophils/hpf in esophageal biopsy specimen
- Not responsive to maximal-dose proton pump inhibitors
- Not currently being treated with steroids and had failed steroid therapy
- Serum IgE 30 to 700 IU/mL
- Acceptable medical history, physical exam, and laboratory test results, specifically no history of bleeding diathesis, significant cardio-pulmonary disease, or other contraindications to upper endoscopy
- Inclusion for controls – healthy adults > 18 years with no medical disease

Exclusion criteria:

Not mentioned in the full text but copied below from the protocol:

- Need for esophageal dilation at enrollment due to food impaction or inability to pass endoscope
- The inability of participant to provide informed consent (if ages 18 to 60), or the inability of children (ages 12 to 17) to provide assent
- History of esophagogastric surgery
- Presence of other esophageal pathology that could account for patients' symptoms including eosinophil infiltration due to gastroesophageal reflux disease (GERD)
- Incarceration
- Pregnancy
- Women of childbearing age that are not using the contraception method(s)
- Patients with elevated serum IgE levels for reasons other than atopy
- Patients taking cromolyn sodium or nedocromil sodium within 1 month of visit 1
- Patients taking oral or topical corticosteroids within one month of visit 1
- Patients taking leukotriene receptor inhibitors within one month of visit 1
- Patients with a severe medical condition(s) that in the view of the investigator prohibit participation in the study
- Patients with a history of noncompliance to medical regimens or who were considered potentially unreliable
- Use of any other investigational agent in the last 30 days
- Patients with a known hypersensitivity to any ingredient of rhuMAB-E25, study rescue medication
- Patients with Barrett's esophagus excluded if found endoscopically or pathologically at biopsy
- Currently treated with omalizumab or treated with omalizumab within the past 6 months.

Age at beginning of study per study group: mean (range; no SD given) omalizumab 32 (16 to 52), placebo 28 (15 to 39)

Sex (m/f) per study group: omalizumab m/f = 13/3, placebo m/f = 11/3

Number randomized per study group: omalizumab n = 16, placebo n = 14

Number reaching end of study per study group: omalizumab n = 16, placebo n = 14

Interventions	<p>Study group 1: placebo (saline) subcutaneous injection every 2 to 4 weeks for 16 weeks</p> <p>Study group 2: omalizumab subcutaneous injection every 2 to 4 weeks for 16 weeks, using a weight and serum IgE-based dosing protocol</p>
Outcomes	<p>Primary outcomes of the study: primary endpoint was reducing esophageal biopsy eosinophil content</p>

Medical treatment of eosinophilic esophagitis (Review)

Clayton 2014 (Continued)

Secondary outcomes of the study: a secondary endpoint was a reduction in dysphagia symptom

Notes

Funding: supported by a Castell grant (K.A.P.). Novartis funded, but did not interpret or primarily design, the omalizumab trial.

Conflict of interests: nothing to disclose

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Placebo participants were given injections of material seemingly identical to the omalizumab.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No mention in manuscript or supplemental material of all who were or were not blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"All statistical comparisons were based on blinded analysis of number-coded slides, tissue homogenates, sera, or of subjects for whom treatment status was known only by a research pharmacist."
Incomplete outcome data (attrition bias) All outcomes	Low risk	All randomized patients completed the study and their data were included in the analysis.
Selective reporting (reporting bias)	Unclear risk	The authors described their primary endpoint as a reduction in eosinophil count per high-power field. This matches the planned analysis. The authors provide P values for secondary, though not the primary outcome. The authors were contacted for clarification without response.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

De Rooij 2022
Study characteristics

Methods	<p>RCT design and number of study arms: open-label RCT; 2 arms</p> <p>Single-center or multi-center: single-center; Amsterdam UMC motility center</p> <p>Countries: Netherlands</p> <p>Study dates: December 2017 to January 2020</p>
Participants	<p>Active EoE or inactive EoE at beginning of study? (and numbers if mixed, per IG/CG): active</p> <p>EoE definition/diagnostic criteria: symptoms of esophageal dysfunction (Straumann Dysphagia Instrument score of ≥ 1) and ≥ 15 eos/HPF on baseline biopsy</p> <p>Inclusion criteria:</p>

Medical treatment of eosinophilic esophagitis (Review)

De Rooij 2022 (Continued)

- Symptoms of esophageal dysfunction (Straumann Dysphagia Instrument score of ≥ 1) and ≥ 15 eos/HPF on baseline biopsy

Exclusion criteria:

- Severe comorbidity scored as the ASA physical classification system class 4 or higher
- Recent history of GI cancer or major GI surgery
- Inability to stop anti-inflammatory drugs

Age at beginning of study per study group: NR

Sex (m/f) per study group: NR

Number randomized per study group: four food elimination diet (FFED) n = 20; : four food elimination diet + amino acid formula (FFED + AAF) n = 21

Number reaching end of study per study group (numbers of patients): four food elimination diet (FFED) n = 20; : four food elimination diet + amino acid formula (FFED + AAF) n = 20

Interventions	<p>Study group 1: four food elimination diet (FFED)</p> <p>Study group 2: four food elimination diet + amino acid formula (FFED + AAF)</p>
Outcomes	<p>Primary outcomes of the study: change in peak eos count (PEC)</p> <p>Secondary outcomes of the study: histologic remission (< 15 eos/HPF), endoscopic signs (EREFS), Straumann Dysphagia Instrument measure, EoEQoL survey, clinical and nutritional outcomes, diet feasibility, adherence, weight loss, BMI</p>
Notes	<p>Funding source: NR</p> <p>Conflicts of interest: NR</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	"Patients were randomized [...] using a blocked randomization protocol with sealed envelopes". There is no further information regarding randomization.
Allocation concealment (selection bias)	Low risk	"Patients were randomized [...] using a blocked randomization protocol with sealed envelopes".
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	The manuscript mentioned "blinded personnel" but unclear how. The authors were contacted in November 2020. No response.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	The manuscript mentioned "blinded personnel" but unclear how. The authors were contacted in November 2020. No response.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Low risk	Trial registry is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Medical treatment of eosinophilic esophagitis (Review)

Dellon 2012
Study characteristics

Methods	<p>RCT design and number of study arms: randomized, prospective, open-label, parallel-arm</p> <p>Single-center or multi-center? single-center, University of North Carolina (UNC) Center</p> <p>Countries: USA</p> <p>Dates: March 2010 and May 2011</p>
Participants	<p>Active EoE or inactive EoE at beginning of study? (and number if mixed, per IG/CG): active</p> <p>EoE definition/diagnostic criteria: included participants reported symptoms of esophageal dysfunction and had persistent esophageal eosinophilia (≥ 15 eosinophils in one high-power field) after 8 weeks of treatment with twice-daily proton pump inhibitor</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 18 years or older with an incident diagnosis of eosinophilic esophagitis • Symptoms of esophageal dysfunction • Persistent esophageal eosinophilia (≥ 15 eosinophils in one high-power field microscopy field (eos/hpf; hpf area = 0.24 mm²) after 8 weeks of treatment with a twice-daily proton pump inhibitor <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Proton pump inhibitor responsive esophageal eosinophilia • Previous diagnosis of eosinophilic esophagitis • Previous allergic reaction to steroid medications • Current use of systemic steroids • Previous treatment with topical steroids, had Barrett's esophagus or previous esophageal surgery, had an inability to read or understand English, or were pregnant <p>Age at beginning of study per study group: mean \pm SD</p> <ul style="list-style-type: none"> • Budesonide, nebulized: 34.9 \pm 7.3 • Budesonide, oral viscous: 34.4 \pm 7.5 <p>Sex (m/f) per study group:</p> <ul style="list-style-type: none"> • Budesonide, nebulized: 8/5 • Budesonide, oral viscous: 7/5 <p>Number randomized per study group: total n = 25</p> <ul style="list-style-type: none"> • Budesonide, nebulized: 13/25 • Budesonide, oral viscous: 12/25 <p>Number reaching end of study per study group:</p> <ul style="list-style-type: none"> • Budesonide, nebulized: 11/13 • Budesonide, oral viscous: 11/12
Interventions	<p>Group 1 (NEB): nebulized/swallowed budesonide solution (1 mg/2 mL) continuously swallowed for over 5 minutes until the dose was depleted</p> <p>Group 2 (OVB): viscous/swallowed budesonide solution (1 mg/2 mL) was mixed with 5 g of sucralose into a viscous slurry and swallowed</p>
Outcomes	Primary outcomes

Medical treatment of eosinophilic esophagitis (Review)

Dellon 2012 (Continued)

- Esophageal eosinophilia as measured by the eosinophil count
- Symptom of dysphagia as measured by the Mayo Dysphagia Questionnaire - 30 days

Secondary outcomes

- Mucosal contact time as measured by nuclear scintigraphy
- Levels of histologic response on esophageal biopsy
- Endoscopic findings of EoE
- Adrenal insufficiency as measured by a standard cortisol stimulation test

Notes

Funding source:

This study was sponsored by AstraZeneca

Conflicts of interest:

No potential conflicts of interest for any of the authors were reported for this study. The funding organizations had no role in the following: design and conduct of the study; collection, management, analysis, and interpretation of the data; and drafting of the manuscript.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Treatment was centrally allocated.
Blinding of participants and personnel (performance bias) All outcomes	High risk	This was an open-label study.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Biopsies were masked and provided to the study pathologists for analysis.
Incomplete outcome data (attrition bias) All outcomes	Low risk	The CONSORT flow diagram in the supplementary information states 1 patient discontinued from the NEB group (13 to 1 = 12). However, 11 completed the study with no information on n = 1. The author confirmed in November 2022 that 1 patient was lost to follow-up and 1 discontinued. There were different patients, so 11 is correct.
Selective reporting (reporting bias)	Low risk	Trial registry is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Dellon 2017
Study characteristics

Methods

RCT design and number of study arms: randomized, double-blind, placebo-controlled, parallel-group; 2 arms

Medical treatment of eosinophilic esophagitis (Review)

Dellon 2017 (Continued)

Single-center or multi-center: multi-center

Countries: USA

Study dates: July 2012 to October 2014

Participants

Active EoE or inactive EoE at beginning of study: active

EoE definition/diagnostic criteria: symptoms of esophageal dysfunction and at least 15 intra-epithelial eosinophils per high-power field (eos/hpf) (hpf area: 0.3 mm²) after an 8-week, high-dose (refers to a total daily dose, which could be administered as a once- or twice-daily dosing regimen), proton pump inhibitor (PPI) trial using any approved PPI. The PPI trial was either historical or could have been performed during the screening period of this study.

Inclusion criteria:

- Patients with EoE were required to have at least 15 eos/HPF from at least 2 esophageal levels on screening endoscopy, at least 4 days with symptoms of dysphagia over the last 2 weeks of a 4-week blinded placebo run-in period, and at least 70% compliance with a daily symptom diary
- Males and females, age 11 to 40
- Histologic evidence of EoE
- History of clinical symptoms of EoE including dysphagia
- Willing to continue with dietary, environmental, or medical therapy
- Ability to read and understand English
- Written consent

Exclusion criteria:

- Older patients who are more likely to have fibrostenotic disease and typically are not amenable to anti-inflammatory treatment alone
- Other potential causes of esophageal eosinophilia had also been excluded
- Non-EoE gastrointestinal diseases, including eosinophilic gastroenteritis/colitis, inflammatory bowel disease, celiac disease, *Helicobacter pylori* infection, esophageal candidiasis (defined based on investigator discretion), or esophageal varices; diseases causing systemic eosinophilia; or esophageal stricture on screening endoscopy that precluded passage of an adult upper endoscope
- Gastroesophageal reflux disease and erosive esophagitis were not formal exclusion criteria, but patients with esophageal eosinophilia related to gastroesophageal reflux disease were excluded, as based on PPI-responsive eosinophilia. Other exclusion criteria were: use of corticosteroids (topical or systemic) in the 4 weeks preceding the screening endoscopy; use of immunomodulatory therapy in the 8 weeks preceding the screening endoscopy; change in the dosing regimen of PPIs, allergy medications, or inhaled corticosteroids; pregnancy; and medical instability.

Age at beginning of study per study group: budesonide oral suspension (BOS) = 22.3 ± 7.9; placebo = 20.8 ± 7.5

Sex (m/f) per study group: Budesonide oral suspension = 35/16; placebo = 29/13

Number randomized per study group: Budesonide oral suspension = 51; placebo = 42

Number reaching end of study per study group: Budesonide oral suspension = 49/51; placebo = 38/42

Interventions

Study group 1: placebo suspension

Study group 2: budesonide oral suspension (BOS) 2 mg twice-daily (given as 10 mL, once in the morning after breakfast and once in the evening before bedtime to provide a total daily dose of 4 mg)

Outcomes

Primary outcomes of the study:

- Change in the DSQ
- Histologic response, defined as eos ≤ 6/hpf

Dellon 2017 (Continued)

Secondary outcomes of the study: endoscopic findings and safety

Notes

Funding source: This study was sponsored by Meritage Pharma, Inc, now a part of the Shire group of companies. Meritage Pharma, Inc contributed to the design and conduct of the study, collection and management of the data, and reviewed the manuscript for medical accuracy. Approval of the manuscript and the decision to submit the manuscript for publication were the responsibility of the authors.

Conflicts of interest: The authors disclose the following: Evan S. Dellon has received research funding from Meritage, Receptos, and Regeneron; and is a consultant for Aptalis, Banner Life Sciences, Novartis, Receptos, Regeneron, and Roche. David A. Katzka has received research funding from Meritage. Margaret H. Collins has received research funding from Meritage, Receptos, Regeneron, and Biogen Idec. Mohamed Hamdani is an employee and stockholder of Shire. Sandeep K. Gupta has received research funding from Meritage. Ikuo Hirano has received research funding from Meritage, and is a consultant for Meritage, Receptos, and Regeneron.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The randomization schedule was generated by SynteractHCR, Inc. and was verified for accuracy using strict quality-control procedures".
Allocation concealment (selection bias)	Low risk	The active study medication and placebo were dispensed in identical amber glass bottles to maintain the blind. However, the authors did not give explicit details on the similarity of both suspensions. The author was contacted in November 2020 and confirmed the allocators were not involved in any other part of the study.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Participants, investigators, the sponsor, study site personnel, and the central pathologist were blinded to patients' treatment, until after all patients had completed the treatment period and the database was locked."
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Participants, investigators, the sponsor, study site personnel, and the central pathologist were blinded to patients' treatment, until after all patients had completed the treatment period and the database was locked."
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Low risk	Trial registry is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Dellon 2019
Study characteristics

Methods

RCT design and number of study arms: RCT, 2 arms

Single-center or multi-center? single-center

Countries: USA

Medical treatment of eosinophilic esophagitis (Review)

Dellon 2019 (Continued)

Study dates: 2014 to 2018

Participants

Active EoE or inactive EoE at beginning of study: active EoE

EoE definition/diagnostic criteria: dysphagia or other symptoms of esophageal dysfunction, > or equal to 15 eos/hpf in at least one field after 8 weeks of treatment with twice-daily PPI and other causes of esophageal eosinophilia excluded

Inclusion criteria: dysphagia or other symptoms of esophageal dysfunction, > or equal to 15 eos/hpf in at least one field after 8 weeks of treatment with twice-daily PPI and other causes of esophageal eosinophilia excluded

Exclusion criteria: concomitant EG, swallowed/topical steroids for EoE or systemic steroids within 4 weeks before baseline endoscopy, inability to pass 9 mm upper endoscope due to narrowing, previous esophageal surgery, esophageal or gastric cancer, esophageal varices, inability to stop anticoagulation or active GI bleeding, medical instability precluding endoscopy, inability to read or understand English, or pregnancy

Age at beginning of study per study group: mean age \pm SD budesonide, oral viscous: 36.2 \pm 19.1 and mean age fluticasone 39.0 \pm 14.5

Sex (m/f) per study group: budesonide, oral viscous 40 males/25 females and fluticasone 44 M/20 F

Number randomized per study group: 65 to budesonide, oral viscous and 64 to fluticasone

Number reaching end of study per study group: 1 did not receive intervention in each group and 8 lost in follow-up in each group

Interventions

Study group 1: budesonide, oral viscous (1 mg/4 mL budesonide with 10 g sucralose twice-daily) + placebo inhaler

Study group 2: fluticasone (220 μ g, 4 puffs (880 μ g) twice-daily) + placebo slurry

Outcomes

Primary outcomes of the study:

 Primary: post-treatment peak eos (eos/hpf; hpf 0.24 mm²)

Co-primary: dysphagia score by the Dysphagia Symptom Questionnaire

Secondary outcomes of the study:

Secondary: EREFS, levels of histologic response (< 15 eos/hpf, < 5 eos/hpf, < 1 eos/hpf), EoE Symptom Activity Index (EEsAI)

Also: medication compliance and adverse events

Notes

Funding source:

This study was supported by National Institutes of Health (NIH) R01 DK101856, and used resources from University of North Carolina (UNC) Center for GI Biology and Disease (NIH P30 DK034987) and the UNC Translational Pathology Lab, which is supported in part by grants from the National Cancer Institute (2-P30-CA016086-40), National Institute of Environmental Health Sciences (2-P30ES010126-15A1), University Cancer Research Fund, and North Carolina Biotechnology Center (2015-IDG-1007)

Conflicts of interest:

These authors disclose the following: Dr Dellon has received research funding from Adare, Allakos, GSK, Meritage, Miraca, Nutricia, Celgene/Receptos, Regeneron, and Shire; has received consulting fees from Adare, Alivio, Allakos, AstraZeneca, Banner, Calypso, Enumeral, EsoCap, Celgene/Receptos, GSK, Regeneron, Robarts, Shire, and educational grants from Allakos, Banner, and Holoclara. The remaining authors disclose no conflicts.

Risk of bias
Medical treatment of eosinophilic esophagitis (Review)

Dellon 2019 (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Medications were premixed by the pharmacy and dispensed such that placebo and active treatments were identical.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	There was one unblinded study pharmacist who dispensed the study medications.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Subjects, investigators, endoscopists, statisticians, and study staff were all masked as to treatment allocation".
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced and explained attrition that likely has no effect on outcome. Within the CONSORT diagram, the authors do not fully explicate attrition per treatment arm. The author was contacted in November 2022 and confirmed attrition details.
Selective reporting (reporting bias)	Low risk	Trial registry is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Dellon 2021b
Study characteristics

Methods	<p>RCT design and number of study arms: RCT with 3 arms, NCT02736409 (extension of NCT02605837)</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: 60 US sites</p> <p>Study dates: 2016 to 2019</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: mixed because the beginning of this study was a continuation of an induction treatment study</p> <p>EoE definition/diagnostic criteria: based on Hirano et al (2022). Histologic EoE with > 15 eos/hpf from at least 2 levels of the esophagus and dysphagia (using DSQ) on at least 4 days in 2 consecutive weeks including the 2 weeks before randomization</p> <p>Inclusion criteria: patients who completed the induction study. For that study, patients willing and able to maintain dietary/environmental therapy and medical regimens in place at screening.</p> <p>Inclusion criteria:</p> <ol style="list-style-type: none"> Participant completed SHP621-301 induction study Participant is able to provide written informed consent (participant, parent or legal guardian and, as appropriate, participant assent) to participate in the study before completing any study-related procedures Participant is male or female aged 11 to 55 years, inclusive, at time of consent for SHP621-301 study

Medical treatment of eosinophilic esophagitis (Review)

Dellon 2021b (Continued)

4. Participant is willing and able to continue any dietary therapy, environmental therapy, and/or medical regimens (including gastric acid suppression; see exclusions below) in effect at the screening visit (Visit 0). There should be no changes to these regimens during study participation.
5. All female participants must have a negative serum pregnancy test (beta-human chorionic gonadotropin [β -hCG]) prior to enrollment into the study. Females of childbearing potential must agree to continue acceptable birth control measures (e.g. abstinence, stable oral contraceptives, or double-barrier methods) throughout study participation and for 30 days following the last dose of investigational product.
6. Participant is willing and has an understanding and ability to fully comply with study procedures including DSQ compliance (completed the DSQ on $\geq 70\%$ of days in any 2 consecutive weeks of the screening period) and restrictions defined in this protocol
7. (implicit) Patients who were defined as having undergone remission (< 7 eosinophils per high-power field (eos/hpf) and $\geq 30\%$ reduction in the Dysphagia Symptom Questionnaire score)

Exclusion criteria: patients with high-grade stricture. Also changes in diet or medications; use of immunomodulatory therapy, swallowed topical corticosteroids, or systemic corticosteroids, or P450 inhibitors; use of inhaled or nasal corticosteroids with unstable dosing; changed dosing regimen of PPI, H2 antagonists, antacids or leukotriene inhibitors; use of pure liquid or the 6-food elimination diet; and unresponsive esophageal or oropharyngeal candidiasis.

1. Participant has changes in medications that could affect the study or diet in the weeks since the final treatment evaluation visit (Visit 4) of the SHP621-301 study.
2. Participant using immunomodulatory therapy since the final treatment evaluation visit (Visit 4) of the SHP621-301 study or anticipated use of immunomodulatory therapy during the treatment period (except for any ongoing regimen of allergy shots); any temporary use (≤ 7 days) or initiation of new steroid treatment during the study should be documented and discussed with the medical monitor prospectively but cannot occur within 4 weeks of scheduled EGDs.
3. Participant using swallowed topical corticosteroid for EoE or systemic corticosteroid for any condition since the final treatment evaluation visit (Visit 4) of the SHP621-301 study or anticipated use during the treatment period; any temporary use (≤ 7 days) or initiation of new steroid treatment during the study should be documented and discussed with medical monitor prospectively but cannot occur within the 4 weeks of the scheduled EGDs.
4. Participant on inhaled or intranasal steroids and not on a stable dose between the baseline visit (Visit 1) of the SHP621-301 study and the screening EGD of this study.
5. Participant has initiated, discontinued, or changed dosage regimen of proton pump inhibitors (PPIs), H2 antagonists, antacids, antihistamines, or leukotriene inhibitors for any condition (such as gastroesophageal reflux disease, asthma or allergic rhinitis) since the final treatment evaluation visit (Visit 4) of the SHP621-301 study or anticipated changes in the use of such medications during the treatment period.
6. Participant using Cytochrome P450 3A4 inhibitors (e.g. ketoconazole, grapefruit juice) since the final treatment evaluation visit (Visit 4) of the SHP621-301 study or anticipated use of such medications during the treatment period.
7. Participant has an appearance on screening EGD of an esophageal stricture (high-grade), as defined by the presence of a lesion that does not allow passage of a diagnostic adult upper endoscope (e.g. with an insertion tube diameter of > 9 mm).
8. Participant is on a pure liquid diet or the 6-food elimination diet.
9. Participant has presence of esophageal varices at the EGD at the final treatment evaluation visit (Visit 4) of the SHP621-301 study.
10. Participant has any current disease of the gastrointestinal tract, aside from EoE, including eosinophilic gastritis, enteritis, colitis, or proctitis, inflammatory bowel disease, or celiac disease.
11. Participant has other diseases causing or associated with esophageal eosinophilia, including hypereosinophilic syndrome, collagen vascular disease, vasculitis, achalasia, or parasitic infection.

Participant has oropharyngeal or esophageal candidiasis that failed to respond to previous treatment.

Diagnosis with oropharyngeal or esophageal candidiasis at or since the final treatment evaluation visit (Visit 4) of the SHP621-301 study is not an exclusion as long as the participant received treatment for candidiasis and is expected to respond to treatment.

Dellon 2021b (Continued)

1. Participant has acute or chronic infection or immunodeficiency condition, including tuberculosis, fungal, bacterial, viral/parasite infection, ocular herpes simplex, or chicken pox/measles.
2. Participant has upper gastrointestinal bleeding identified in the EGD at the final treatment evaluation visit (Visit 4) of the SHP621-301 study or since the final treatment evaluation visit (Visit 4) of the SHP621-301 study.
3. Participant has evidence of active infection with *Helicobacter pylori*.
4. Participant has evidence of unstable asthma since the final treatment evaluation visit (Visit 4) of the SHP621-301 study.
5. Participant is female and pregnant or nursing.
6. Participant has a history of intolerance, hypersensitivity, or idiosyncratic reaction to budesonide (or any other corticosteroids), or to any other ingredients of the study medication.
7. Participant has a history or high risk of noncompliance with treatment or regular clinic visits.
8. Participant is on sucralfate or anticipates using sucralfate during the treatment period.

Age at beginning of study per study group: 1) Arm 1 budesonide oral suspension (BOS) group: 36.8 (14.1) and placebo group: 36.1 (11.7); 2) Arm 2: 33.1 (12.0); and 3) Arm 3: 33.5 (12.6)

Sex (m/f) per study groups: 1) Arm 1 BOS group 14 M/11 F and placebo group 16 M/7 F; 2) Arm 2 had 64 M/42 F; and 3) Arm 3 had 39 M/26 F

Number randomized per study group: 1) 48 induction full responders were in Arm 1 and underwent randomized withdrawal (n = 25 randomized to continue BOS and n = 23 randomized to placebo); 2) For arm 2, 106 induction partial responders and non-responders received BOS; and 3) For arm 3, 65 induction placebo patients received BOS

Number reaching end of study per study group: for safety analysis, any patient taking at least one dose and for per protocol analysis, patients without significant deviation from study protocol. For per protocol analysis (Arm 1) participants who continued BOS, n = 18 completed and of the participants who changed to placebo, n = 18 completed.

Seven patients relapsed and reinitiated BOS (from placebo group)

Interventions	<p>Study group 1 of Arm 1: full responders who continued budesonide oral suspension (continuation of BOS in previous RCT)</p> <p>Study group 2 of Arm 1: full responders who changed to placebo (change from BOS in previous RCT)</p>
Outcomes	<p>Primary outcomes of study: proportion of induction full responders (BOS or placebo) who experienced histologic and dysphagia symptom relapse</p> <p>Secondary outcomes of study:</p> <ol style="list-style-type: none"> 1. Proportions of induction partial responder and non-responders who achieved full responses at week 36 (52 weeks total of BOS) 2. Measured at weeks 12 and 36, proportions of induction full responders who achieved histologic responses (< 1, < 6, and < 15), additional dysphagia symptom responses, and maintained full responses. Percentage reductions in DSQ scores as changes from the induction and extension study baselines. 3. Also included: change in DSQ scores, peak eosinophil counts, and EREFS from the extension study baseline to weeks 12 and 36
Notes	<p>Funding source: Shire ViroPharma, Inc of Takeda Pharmaceuticals</p> <p>Conflicts of interest:</p> <p>The authors disclose the following: Evan S. Dellon has received research funding from Adare Pharmaceuticals, Allakos, AstraZeneca, GlaxoSmithKline, Meritage Pharma, Inc, Miraca Life Sciences, Nutricia, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda Company, and is a consultant for Abbott Laboratories, Adare Pharmaceuticals, Aimmune Therapeutics, Allakos, Amgen, Arena Pharmaceuticals, AstraZeneca, Biorasi, Calypso Biotech, Celldex Therapeutics, Inc, EsoCap Biotech, GlaxoSmithKline, Gossamer Bio, Lilly, Parexel/Calyx Clinical Trial Solutions, Receptos/Celgene, Regeneron Pharmaceuticals, Robarts Clinical Trials, Inc/Alimentiv, Inc, Salix Pharmaceuticals, Sanofi, and</p>

Dellon 2021b (Continued)

Shire, a Takeda Company, and has received educational grants from Allakos, Banner Life Sciences, and Holoclara; Margaret H. Collins has received research funding from MeritagePharma, Inc, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda Company, and is a consultant for Allakos, Arena Pharmaceuticals, AstraZeneca, Calypso Biotech, EsoCap Biotech, GlaxoSmithKline, Receptos/Celgene, Regeneron Pharmaceuticals, Robarts Clinical Trials, Inc/Alimentiv, Inc, and Shire, a Takeda Company; David A. Katzka has received research funding from Shire, a Takeda Company, and a consulting fee from Receptos/Celgene; Vincent A. Mukkada has received research funding from Meritage Pharma, Inc, and Shire, a Takeda Company, and is a consultant for Shire, a Takeda Company; Gary W. Falk has received research funding from Adare Pharmaceuticals, Allakos, Lucid, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda Company, and is a consultant for Adare Pharmaceuticals, Allakos, Bristol Myers Squibb, Lucid, Regeneron Pharmaceuticals, and Shire, a Takeda Company; Robin Morey, Bridgett Goodwin, Nirav K. Desai, and James Williams are employees of Takeda Development Center Americas, Inc, and stockholders of Takeda Pharmaceutical Company Limited; Jessica D. Eisner was an employee and stockholder of Takeda Pharmaceuticals USA, Inc, and a stockholder of Takeda Pharmaceutical Company Limited, at the time of the study; Lan Lan was an employee of Takeda Development Center Americas, Inc, and a stockholder of Takeda Pharmaceutical Company Limited, at the time of the study; and Ikuo Hirano has received research funding from Adare Pharmaceuticals, Allakos, Arena Pharmaceuticals, AstraZeneca, Meritage Pharma, Inc, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda company, and is a consultant for Adare Pharmaceuticals, Allakos, Arena Pharmaceuticals, AstraZeneca, EsoCap Biotech, Gossamer Bio, Lilly, MeritagePharma, Inc, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda Company.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	This was a follow-up study. For the initial induction study, randomization was performed centrally via "Interactive Web-based Response System". Full responders after induction were "randomized" to continue active treatment or to placebo. The details of the second randomization process were not stated in the paper or supplemental information.
Allocation concealment (selection bias)	Low risk	The author was contacted in November 2022 and confirmed central allocation.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Study sites and teams were blinded but no additional details were provided.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Study sites and teams were blinded. There was a separate analysis team that was not blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No major attrition imbalances or imbalances in reasons for attrition that could have impacted our outcomes, based on the flow diagram.
Selective reporting (reporting bias)	Low risk	Trial registry is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Dellon 2022
Study characteristics
Medical treatment of eosinophilic esophagitis (Review)

Dellon 2022 (Continued)

Methods	<p>RCT design and number of study arms: part A of a 3-part, randomized, placebo-controlled phase 3 study, 2 arms</p> <p>Single-center or multi-center: multi-center; 95 study locations worldwide</p> <p>Countries: US, Australia, Belgium, Canada, France, Germany, Italy, Netherlands, Spain, Sweden, Switzerland, and the UK</p> <p>Study dates: 24 September 2018 to 9 September 2021</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: "A documented diagnosis of EoE by endoscopic biopsy."</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • A documented diagnosis of EoE by endoscopic biopsy • Baseline endoscopic biopsies with a demonstration on central reading of intraepithelial eosinophilic infiltration • History (by patient report) of an average of at least 2 episodes of dysphagia (with intake of solids) per week in the 4 weeks prior to screening <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Body weight \leq 40 kg • Prior participation in a dupilumab clinical trial, or past or current treatment with dupilumab • Initiation or change of a food-elimination diet regimen or re-introduction of a previously eliminated food group in the 6 weeks prior to screening. • Other causes of esophageal eosinophilia or the following conditions: hypereosinophilic syndrome and eosinophilic granulomatosis with polyangiitis (Churg-Strauss syndrome) • Active <i>Helicobacter pylori</i> infection • History of achalasia, Crohn's disease, ulcerative colitis, celiac disease, and prior esophageal surgery • Any esophageal stricture unable to be passed with a standard, diagnostic, 9 mm to 10 mm upper endoscope or any critical esophageal stricture that requires dilation at screening • History of bleeding disorders or esophageal varices • Pregnant or breastfeeding women, or women planning to become pregnant or breastfeed during the study <p>Age at beginning of study per study group: not stated</p> <p>Sex (m/f) per study groups: not stated</p> <p>Number randomized per study group: 42 patients treated with dupilumab and 39 patients treated with placebo</p> <p>Number reaching end of study per study group:</p> <p>Dupilumab 300 mg weekly: 42 (100%)</p> <p>Placebo: 39 (100%)</p>
Interventions	<p>Study group 1: placebo</p> <p>Study group 2: dupilumab 300 mg weekly</p>
Outcomes	<p>Primary outcomes of study: this was included in another study</p> <ul style="list-style-type: none"> • Proportion of patients achieving a peak esophageal intraepithelial eosinophil (eos) count of $<$ 6 eos/hpf • Absolute change from baseline in Dysphagia Symptom Questionnaire (DSQ) score

Dellon 2022 (Continued)

Secondary outcomes of study: dupilumab's effect versus placebo on HRQoL and symptom burden

- Absolute change from baseline in EoE histologic scoring system (EoEHSS) mean grade and stage scores
- Absolute change in total EoE Endoscopic Reference Score (EREFS)
- Proportion of patients achieving a peak eos count of < 15 eos/hpf

Notes
Funding source:

Research sponsored by Sanofi and Regeneron Pharmaceuticals, Inc.

Conflicts of interest:

Dr. Dellon - Consultant: Abbott, Adare, Aimmune, Allakos, Amgen, Arena, AstraZeneca, Biorasi, Calypso, Eli Lilly, EsoCap, Gossamer Bio, GlaxoSmithKline, Parexel, Receptos/Celgene/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Robarts, Salix, Shire/Takeda; research funding: Adare, Allakos, GlaxoSmithKline, Meritage, Miraca, Nutricia, Receptos/Celgene/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Shire/Takeda; educational grant: Allakos, Banner, Holoclara.

Dr. Rothenberg - Consultant: Allakos, AstraZeneca, Bristol Myers Squibb, ClostraBio, Pulm One, Spoon Guru; equity interest: ClostraBio, Pulm One, Spoon Guru; royalties from reslizumab: Teva Pharmaceuticals; royalties from PEESV2: Mapi Research Trust; royalties: UpToDate; inventor of patents owned by Cincinnati Children's Hospital.

Dr. Collins - consultant: Allakos, Arena, AstraZeneca, Bristol Myers Squibb, Calypso, Esocap, GlaxoSmithKline, Regeneron Pharmaceuticals, Inc., Shire; research funding: Receptos/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Shire.

Dr. Hirano - consultant: Adare, Receptos/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Shire; research funding: Meritage, Receptos/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Shire.

Dr. Chehade - consultant: Adare, Allakos, AstraZeneca, Nutricia, Regeneron Pharmaceuticals, Inc., Shire; research funding: Allakos, Regeneron Pharmaceuticals Inc., Shire; honoraria for lectures: Medscape, Nutricia.

Dr. Bredenoord - consultant: Arena, AstraZeneca, Calypso, EsoCap, Falk, Gossamer Bio, Medtronic, Laborie, RB, Regeneron, Robarts; research funding: Bayer, Nutricia, SST; equity interest: SST.

Dr. Lucendo - Consultant: EsoCap, Dr. Falk Pharma; research funding: Dr. Falk Pharma, Regeneron Pharmaceuticals.

Dr. Spergel - Consultant: Regeneron, Shire, Takeda, Allakos, DBV Technology, Novartis; Grant Support: Regeneron, DBV Technology.

Q Zhao, JD Hamilton, B Beazley, S Kamat, M Ruddy, B Akinlade, N Amin, A Radin, B Shumel, J Maloney: Regeneron Pharmaceuticals, Inc. - employees and shareholders.

I Guillemin: Sanofi - prior employee, may hold stock and/or stock options in the company

L Mannent, E Laws: Sanofi - employees, may hold stock and/or stock options in the company.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The author was contacted in November 2022 and confirmed that randomization was computer-generated.
Allocation concealment (selection bias)	Low risk	The author was contacted in November 2022 and confirmed central allocation.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	The author was contacted in November 2022 and confirmed blinding of participants and personnel.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The author was contacted in November 2022 and confirmed blinding of outcome assessors.

Medical treatment of eosinophilic esophagitis (Review)

103

Dellon 2022 (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	There was no loss to follow-up in either study arm.
Selective reporting (reporting bias)	Low risk	Protocol is available and all pre-specified outcomes of interest were reported. Full data reported for all patients.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Dellon 2022a
Study characteristics

Methods	<p>RCT design and number of study arms: RCT; 5 arms (4 intervention, 1 placebo)</p> <p>Single-center or multi-center? multi-center (93 centers in 6 countries: US, Canada, Belgium, Switzerland, Spain, and Germany)</p> <p>Countries: several</p> <p>Study dates: May 2017 to August 2018</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: "defined as 3 episodes of dysphagia per week during the last 14 days of the 4-week baseline symptom assessment phase and a Global EoE Symptom Score of >3), and active esophageal eosinophilia (after evaluation of 5 biopsies from proximal and distal esophageal locations and at least 1 biopsy with a peak count of 15 eos/HPF) after documentation of failed histologic response on 8 weeks of high-dose PPI." "High-dose PPI was defined as 20 to 40 mg daily of any marketed PPI."</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> Adults (> or equal to 18 to < or equal to 75) with active EoE (with dysphagia at least 3 times weekly and global EoE symptom score > 3 as well as at least 15 eos/hpf) and failed histologic response on at least 8 weeks high-dose PPI (20 mg to 40 mg daily) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> Known esophageal mucosal disease or esophageal dysmotility unrelated to EoE History of esophageal stricture requiring dilation within 12 weeks prior or with severe stricture precluding passage of standard 8 mm to 10 mm scope was also an exclusion criterion Corticosteroids, change in diet, biologics, and immunomodulators were prohibited <p>Age at beginning of study per study group: mean age overall: 39.3 ± 12.0 years. For 3 mg twice-daily (n = 20): 36.8 ± 9.2; 3 mg every night at bedtime (n = 21): 42.9 ± 11.5; 1.5 mg twice-daily (n = 22): 41.3 ± 12.2; 1.5 mg every night at bedtime (n = 21): 36.8 ± 11.5; placebo (n = 19): 38.6 ± 14.7</p> <p>Sex (m/f) per study group: Sex: males 70/103 overall. For 3 mg twice-daily: 16/20 male; 3 mg every night at bedtime: 11/21 male; 1.5 mg twice-daily: 15/22 male; 1.5 mg every night at bedtime: 14/21 male; and placebo: 14/19 male. 106 patients randomized (details for each group shown above)</p> <p>Number randomized per study group:</p> <ul style="list-style-type: none"> 3 mg twice-daily – n = 20 3 mg every night at bedtime – n = 22 1.5 mg twice-daily – n = 22 1.5 mg every night at bedtime – n = 21

Medical treatment of eosinophilic esophagitis (Review)

Dellon 2022a (Continued)

- Placebo twice-daily – n = 21

Number reaching end of study per study group:

- 3 mg twice-daily – n = 19
- 3 mg every night at bedtime – n = 20
- 1.5 mg twice-daily – n = 20
- 1.5 mg every night at bedtime – n = 17
- Placebo twice-daily – n = 16

Interventions	Study group 1 (placebo): placebo twice-daily Study groups 2, 3, 4, 5 (fluticasone): <ul style="list-style-type: none"> • 3 mg twice-daily • 3 mg every night at bedtime • 1.5 mg twice-daily • 1.5 mg every night at bedtime
Outcomes	Primary outcomes of the study: histologic response at 12 weeks (% participants with < or equal to 6 eos/HPF) Secondary outcomes of the study: <ul style="list-style-type: none"> • Percentage of responders with sustained histologic response at weeks 26 and 52 • Endoscopic severity measured by change from baseline in EREFS at weeks 12, 26, and 52 • Percentage with < 1 and < 15 eos/HPF at weeks 12, 26, and 52 • Secondary symptomatic outcomes: <ul style="list-style-type: none"> ◦ Change from baseline in the Global EoE Symptom Score multiple times in week 52 ◦ Change in frequency of reported dysphagia episodes over a 14-day period from baseline through weeks 12, 26, and 52
Notes	Funding source: Adare/Ellodi Pharmaceuticals, Inc. Conflicts: Dellon, Lucendo, Schoepfer, Falk, and Hirano research funding from Adare/Ellodi; Dellon, Schlag, Schoepfer, Falk, and Hirano consulting fees from Adare/Ellodi; Eagle, Nezamis, Comer, Knoop employees of Ellodi

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Randomization with fixed block size of five using Interactive Web Response System"
Allocation concealment (selection bias)	Low risk	"Randomization with fixed block size of five using Interactive Web Response System"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Blinding is unclear in the manuscript. Study drug was identical to placebo. Confirmed by authors
Blinding of outcome assessment (detection bias) All outcomes	Low risk	No details provided except stating "blinded". The author was contacted in November 2022 and confirmed assessors were blinded.
Incomplete outcome data (attrition bias)	Low risk	There was no loss to follow-up for this study.

Medical treatment of eosinophilic esophagitis (Review)

105

Dellon 2022a (Continued)

All outcomes

Selective reporting (reporting bias)	Low risk	Trial registry is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Dellon 2022b
Study characteristics

Methods	<p>RCT design and number of study arms: RCT, 3 arms</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: Australia, Netherlands, United States</p> <p>Study dates: 6 July 2020 to December 2021</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: esophagus: ≥ 15 eos/high-power field (hpf) in 1 hpf and active moderate to severe symptoms – Dysphagia Symptom Questionnaire (DSQ) ≥ 12</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Dysphagia Symptom Questionnaire (DSQ) ≥ 1 • Confirmed diagnosis of EoE and esophageal intraepithelial eosinophilic infiltration of ≥ 15 eosinophils/HPF in 1 hpf from a biopsy collected during the screening EGD without any other cause for the esophageal eosinophilia • History (by patient report) of an average of ≥ 2 episodes of dysphagia with intake of solid foods per week during the 4 weeks prior to screening • Participants must have failed or not be adequately controlled on standard of care treatments for EoE symptoms, which could include PPI, systemic or topical corticosteroids, and/or diet, among others • If on an allowed treatment for EoE, stable dose for at least 4 weeks prior to screening and willingness to continue that dose for the study duration • If patient is on pre-existing dietary restrictions, willingness to maintain dietary restrictions throughout the study, as much as possible • Able and willing to comply with all study procedures • Female participants must be either post-menopausal for at least 1 year with FSH level > 30 mIU/mL at screening or surgically sterile (tubal ligation, hysterectomy, or bilateral oophorectomy) for at least 3 months, or if of childbearing potential, have a negative pregnancy test and agree to use dual methods of contraception, or abstain from sexual activity from screening until the end of the study, or for 120 days following the last dose of study drug, whichever is longer. Male participants with female partners of childbearing potential must agree to use a highly effective method of contraception from screening until the end of the study or for 120 days following the last dose of study drug, whichever is longer. All fertile men with female partners of childbearing potential should be instructed to contact the Investigator immediately if they suspect their partner might be pregnant at any time during study participation. <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Concomitant EG, EoD, or eosinophilic colitis (EC) • EG and/or EoD (≥ 30 eosinophils/hpf in 5 hpf in the stomach and/or ≥ 30 eosinophils/hpf in 3 hpf in the duodenum) as determined by central histology assessment of biopsies collected during the screening EG

Medical treatment of eosinophilic esophagitis (Review)

Dellon 2022b (Continued)

- Causes of esophageal eosinophilia other than EoE or one the following: hypereosinophilic syndrome, eosinophilic granulomatosis with polyangiitis, or peripheral blood absolute eosinophil count of > 1500 eosinophils/ μ L
- History of inflammatory bowel disease, celiac disease, achalasia, and/or esophageal surgery
- Any esophageal stricture unable to be passed with a standard diagnostic 9 mm to 10 mm upper endoscope or any critical esophageal stricture that requires dilation during screening
- History of bleeding disorders or esophageal varices
- History of malignancy; except carcinoma in situ, early stage prostate cancer, or non-melanoma skin cancers. However, cancers that have been in remission for more than 5 years and are considered cured, can be enrolled (with the exception of breast cancer). All history of malignancy (including diagnosis, dates, and compliance with cancer screening recommendations) must be documented and certified by the Investigator, along with the statement that in their clinical judgment the tissue eosinophilia is attributable to EGID, rather than recurrence of malignancy.
- Active *Helicobacter pylori* infection (as determined by central histology staining of the biopsy collected during the screening EGD), unless treated and confirmed to be negative prior to randomization and symptoms remain consistent.
- Positive Ova and Parasite (O&P) test at screening, seropositive for *Strongyloides stercoralis* at screening, and/or treatment for a clinically significant helminthic parasitic infection within 6 months of screening
- Seropositive for HIV or hepatitis at screening, except for vaccinated patients or patients with a history of hepatitis that has since resolved.
- Prior exposure to AK002 or hypersensitivity to any constituent of AK002
- Change in dose of inhaled corticosteroids, nasal corticosteroids, PPI, and/or diet therapy within 4 weeks prior to screening
- Use of oral corticosteroids (swallowed topical or systemic corticosteroids) within 8 weeks prior to screening
- Use of any biologics or medications that may interfere with the study, such as immunosuppressive or immunomodulatory drugs including azathioprine, JAK inhibitors, 6-mercaptopurine, methotrexate, cyclosporine, tacrolimus, anti-TNF, anti-IL-4 receptor, e.g. dupilumab), anti-IL-5 (e.g. mepolizumab), anti-IL-5 receptor (e.g. benralizumab), anti-IL-13 (e.g. lebrikizumab), anti-IgE (e.g. omalizumab), within 12 weeks prior to screening.
- Participation in a concurrent interventional study with the last intervention occurring within 30 days prior to administration of study drug or 90 days or 5 half-lives, whichever is longer, for biologic products
- Vaccination with live attenuated vaccines \leq 30 days prior to initiation of treatment in the study, during the treatment period, or vaccination expected \leq 5 half-lives (\leq 4 months) following study drug administration
- Treatment with chemotherapy or radiotherapy in the preceding 6 months
- The presence of abnormal laboratory values is considered by the investigator to be clinically significant
- Any disease, condition (medical or surgical), or cardiac abnormality, in the opinion of the Investigator, would place the participant at increased risk
- Known history of alcohol, drug, or other substance abuse or dependence
- Women who are pregnant, breastfeeding or planning to become pregnant while participating in the study
- Any other reason that in the opinion of the Investigator or Medical Monitor makes the patient unsuitable for enrollment

Age at beginning of study per study group

- Group 1 high-dose lircatelimab (n = 91) 29 (12 to 69)
- Group 2 low-dose lircatelimab (n = 93) 34 (12 to 67)
- Group 3 placebo (n = 92) 32 (12 to 70)

Sex (m/f) per study group

- Group 1 high-dose lircatelimab (n = 91) 29% (26)

Dellon 2022b (Continued)

- Group 2 low-dose lirentelimab (n = 93) 43% (40)
- Group 3 placebo (n = 92) 40% (37)

Number randomized per study group

- Group 1 high-dose lirentelimab (n = 91)
- Group 2 low-dose lirentelimab (n = 93)
- Group 3 placebo (n = 92)

Number reaching end of study per study group

NR

Interventions	Study group 1: high-dose lirentelimab 3 mg/kg Study group 2: low-dose lirentelimab 1 mg/kg Group 3: placebo
Outcomes	Primary outcomes of the study: 1. Histologic co-primary endpoint – proportion of tissue eosinophil responders: esophagus: ≤ 6 eos/hpf in peak hpf 2. Symptom co-primary endpoint – absolute change in Dysphagia Symptom Questionnaire (DSQ) score Secondary outcomes of the study: 1. Secondary endpoints – percent change in DSQ from baseline Achieved peak esophageal eos ≤ 1 eos/hpf at week 24
Notes	Funding source: Allakos Conflicts of interest: Evan S. Dellon, MD, MPH

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	This is not stated, and only a protocol and abstract presentation are available for review. Authors contacted for further details and confirmed in November 2022 that the schedule was computer-generated and centrally allocated.
Allocation concealment (selection bias)	Low risk	This is not stated, and only a protocol and abstract presentation are available for review. Authors contacted for further details and confirmed in November 2022 that the schedule was computer-generated and centrally allocated.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	The manuscript states "Quadruple (Participant, Care Provider, Investigator, Outcomes Assessor)" but doesn't state how this was achieved. Authors contacted for further details and confirmed in November 2022 that personnel and participants were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Quadruple masking Outcomes Assessor"; does not state how this was achieved. Authors contacted for further details and confirmed in November 2022 that personnel and participants were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	277 recruited and 276 received medication - outcome for all is recorded with no attrition.

Medical treatment of eosinophilic esophagitis (Review)

Dellon 2022b (Continued)

Selective reporting (reporting bias)	Low risk	Trial registration reviewed. Primary outcomes appropriate and match the published abstract with none missing. NCT04322708
Other bias	Low risk	No balance bias from the baseline table. No other concerns.

Dohil 2010
Study characteristics

Methods	<p>RCT design and number of study arms: RCT; 2 arms</p> <p>Single-center or multi-center: single-center; Eosinophilic Esophagitis Clinic at Rady Children's Hospital, San Diego</p> <p>Countries: USA</p> <p>Study dates: patients were recruited between February 2008 and July 2009 and the study treatment lasted 3 months</p>
Participants	<p>Active EoE or inactive EoE at beginning of the study: active</p> <p>EoE definition/diagnostic criteria: peak eos \geq 20/hpf</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Histologic evidence of eosinophilic esophagitis defined as greater than 20 eosinophils per HPF on esophageal biopsy • Ages 1 years and older • Ability to continue the same diet that the patient was on at the time of EGD with biopsy <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Adverse reaction or allergy to budesonide • Pregnancy • Chronic diseases requiring immunomodulatory therapy • Use of swallowed topical corticosteroids for eosinophilic esophagitis within the past 3 months • Use of systemic steroids 2 months prior to study entry • Upper gastrointestinal bleed within 4 months of study entry • Chronic use of medications that predispose to upper gastrointestinal bleeding including non-steroidal anti-inflammatory medications or anticoagulants • Evidence of adrenal suppression prior to study entry • Evidence of concurrent eosinophilic gastritis, enteritis, colitis, or proctitis • Recent changes in asthma or allergic rhinitis therapy for 3 months <p>Age at beginning of study per study group: NR</p> <p>Sex (m/f) per study group: NR</p> <p>Number randomized per study group:</p> <p>Budesonide, oral viscous: n = 21</p> <p>Placebo: n = 11</p> <p>Number reaching the end of study per study group:</p> <p>Budesonide, oral viscous: n = 15</p>

Medical treatment of eosinophilic esophagitis (Review)

Dohil 2010 (Continued)

Placebo n = 9

Interventions	Study group 1: sterile water Study group 2: budesonide oral viscous suspension (0.5 mg/2 mL)
Outcomes	Primary outcomes of the study: primary outcome measure was an improvement of esophageal eosinophilia. This was determined by comparing peak eos counts/hpf at baseline and after treatment. Patients were categorized into responders (0 to 6 eos/hpf), partial responders (7 to 19 eos/hpf), and non-responders (> 20 eos/hpf). Secondary outcomes of the study: secondary outcome measurements included the response of symptoms and endoscopic and histologic features to treatment
Notes	Conflicts of interest: The authors disclose the following: The University of California, San Diego has a financial interest in Meritage Pharma, the company sponsoring this research. Drs Dohil, Bastian, and Aceves and the University of California may financially benefit from this interest if the company is successful in developing and marketing its own product that is related to this research. The terms of this arrangement have been reviewed and approved by the University of California, San Diego in accordance with its conflict of interest policies. The remaining authors disclose no conflicts. Funding: support was provided by a grant from Meritage Pharma, San Diego, CA

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Suspensions of the medications were placed by the investigational pharmacist into sealed light-protective vials. Only the pharmacist had access to the randomization code.
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	The manuscript stated blinded personnel and patients. Patients were blinded due to sealed envelope but no details given on personnel blinding. Only the pharmacist had access to the randomization code.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"During the study, collected data were stored in a locked cabinet. One month after study completion, the database was locked and the randomisation code revealed".
Incomplete outcome data (attrition bias) All outcomes	Low risk	"Thirty-one patients were randomized; 24 completed the study and were included in the final analysis. Of these 24 participants, 15 received OVB and PPI and 9 received placebo and PPI." The authors explain exclusions from analysis. Eight patients (2 placebo and 6 intervention) did not have a follow-up endoscopy.
Selective reporting (reporting bias)	Low risk	Protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	There were no major baseline differences between the groups. 1 patient included in the placebo group with 15 eosinophils per high-power field, which was below the cutoff for inclusion. All patients were continued on PPI. Also diet differences from patient to patient as some patients were avoiding food groups.

Medical treatment of eosinophilic esophagitis (Review)

Gupta 2015

Study characteristics

Methods	<p>RCT design and number of study arms: RCT with 4 arms, age stratified (2 to 9 and 10 to 18 years old), parallel assignment, dose-ranging</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: United States</p> <p>Study dates: January 2009 to April 2010</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: esophageal biopsy must show ≥ 20 eos per HPF (400x, 0.3 mm² HPF) at 2 or more levels of the esophagus following 4 weeks of high dose PPI (type, actual dosage not specified)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Male and female participants between the ages of 2 to 18 years, inclusive • History of clinical symptoms of esophageal dysfunction intermittently or continuously • Histologic evidence of EoE with a peak eosinophil count of greater than or equal to 20 eosinophils per HPF, from 2 or more levels of the esophagus, within 6 weeks prior to the Baseline Visit • At the Baseline Visit, participants must have symptoms with a total EoE Clinical Symptom Score of greater than or equal to 3 • Willingness and ability to continue the dietary therapy, environmental therapy, and/or medical regimens (including gastric acid suppression, if any) in effect at the Screening Visit • Females of childbearing potential must have a negative serum pregnancy test (beta human chorionic gonadotropin) prior to randomization into the study and sexually active participants must agree to continue acceptable birth control measures throughout the duration of the study • Written informed consent (parent or legal guardian) and, as appropriate, participant assent <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Current use of immunomodulatory therapy (or anticipated use within 12 weeks following the Baseline Visit) • Diagnosis of inflammatory bowel disease • Chronic viral infection or immunodeficiency condition (current) • Use of swallowed topical corticosteroids for EoE in the 1 month prior to the biopsy required for entrance to this study or at any time between the biopsy and the Baseline Visit • Use of systemic (oral or parenteral) corticosteroid within 1 month prior to the biopsy required for entrance to this study or at any time between the biopsy and the Baseline Visit • Morning plasma cortisol level below the lower limit of normal (per central laboratory reference range) at the Screening Visit • Upper gastrointestinal bleeding within 1 month prior to the Screening Visit or between the Screening Visit and Baseline Visit • Current use of anticoagulants • Current disease of the gastrointestinal tract aside from the current EoE diagnosis • Evidence of concurrent eosinophilic gastritis, enteritis, colitis, or proctitis • Evidence of active infection with <i>Helicobacter pylori</i> • Evidence of unstable asthma or changes in asthma or allergic rhinitis therapy within 1 month prior to the biopsy required for entrance to this study • Any female who is pregnant, who is planning to become pregnant, or who is breastfeeding • Current evidence or history of hypersensitivity or idiosyncratic reaction to budesonide or any other ingredients of the study medication • Current evidence of oropharyngeal or esophageal candidiasis

Gupta 2015 (Continued)

- Receipt of an investigational drug within 30 days prior to the biopsy required for entrance to this study
- Any condition or abnormality that, in the opinion of the Principal Investigator, would compromise the safety of the participant or successful conduct of the study

Age at beginning of study per study group: placebo, 9.2 (4.36); low-dose oral budesonide suspension (OBS), 9 (5.88); medium-dose OBS, 10.2 (4.89); high-dose OBS, 8.1 (4.58)

Sex (m/f) per study group: placebo, 16/5; low-dose OBS, 17/4; medium-dose OBS, 17/4; high-dose OBS, 16/5

Number randomized per study group (numbers of patients): placebo, 21; low-dose OBS, 21; medium-dose OBS, 19; high-dose OBS, 20

Number reaching end of study per study group (numbers of patients): placebo, 18; low-dose OBS, 17; medium-dose OBS, 19; high-dose OBS, 17

Interventions

(Group 1) placebo: participants received placebo twice-daily at bedtime (hs) and after breakfast (once a day in the morning, pc) for 12 weeks with a 3 week taper period

(Group 2) low-dose OBS: participants received oral budesonide suspension (OBS) 0.05 mg/mL at bedtime (hs) and placebo after breakfast (once a day in the morning, after meals) for 12 weeks, with a total daily dose of 0.35 mg (2 to 9 years) or 0.50 mg (10 to 18 years), followed by a 3-week taper period

(Group 3) medium-dose OBS: participants received oral budesonide suspension (OBS) 0.2 mg/mL at bedtime (hs) and placebo after breakfast (once a day in the morning, after meals) for 12 weeks, with a total daily dose of 1.4 mg (2 to 9 years) or 2.0 mg (10 to 18 years), followed by a 3-week taper period

(Group 4) high-dose OBS: participants received oral budesonide suspension (OBS) 0.2 mg/mL at bedtime (hs) and after breakfast (once a day in the morning, after meals) for 12 weeks, with a total daily dose of 2.8 mg (2 to 9 years) or 4.0 mg (10 to 18 years), followed by a 3-week taper period

Outcomes

Primary outcome: percent of participants who responded to therapy following 12 weeks of treatment

The response was defined as a $\geq 50\%$ reduction from baseline in the eosinophilic esophagitis (EoE) clinical symptom score (CSS) and a reduction in peak eosinophil count to ≤ 6 /high-power field (light microscopy) from esophageal biopsies collected at the final evaluation

Secondary outcome:

- Percent of participants with histologic response following 12 weeks of treatment
 - Histologic response was defined as a maximum peak eosinophil count at the final treatment evaluation of ≤ 6 eosinophils/high-power field (light microscopy). The maximum peak was identified by examining the peak eosinophil counts obtained from the proximal, mid, and distal esophageal biopsies and selecting the maximum value.
- Percent of participants with histologic remission following 12 weeks of treatment
- Histologic remission was defined as a maximum peak eosinophil count at the final treatment evaluation of ≤ 1 eosinophils/high-power field (light microscopy). The maximum peak was identified by examining the peak eosinophil counts obtained from the proximal, mid, and distal esophageal biopsies and selecting the maximum value.
- Percent change from baseline in peak eosinophil count following 12 weeks of treatment
 - The maximum peak number of eosinophils at baseline and at the final treatment evaluation was identified by examining the peak eosinophil counts obtained from the proximal, mid, and distal esophageal biopsies and selecting the maximum value. A negative change from baseline indicates that eosinophil count has decreased.
- Change from baseline in endoscopy score following 12 weeks of treatment
- Percent of participants with clinical response following 12 weeks of treatment
 - Response was defined as a $\geq 50\%$ reduction from baseline in the eosinophilic esophagitis (EoE) clinical symptom score (CSS)
- Percent of participants with clinical remission following 12 weeks of treatment
 - Clinical remission was defined as an eosinophilic esophagitis (EoE) clinical symptom score (CSS) of zero

Gupta 2015 (Continued)

- Percent change from baseline in eosinophilic esophagitis (EoE) clinical symptom score (CSS) following 12 weeks of treatment
- Change from baseline in physician's global assessment score of disease severity following 12 weeks of treatment
- Maximum plasma concentration (C_{max}) of budesonide (time frame: week 2, 4, or 8, or at the final treatment evaluation)
- Time to maximum (T_{max}) and half maximum ($T_{1/2}$) plasma concentration of budesonide (time frame: week 2, 4, or 8, or at the final treatment evaluation)
- Area under the plasma concentration-time curve (AUC) of budesonide from time zero to time of the last measurable concentration (AUC_{0-last}) (time frame: week 2, 4, or 8, or at the final treatment evaluation)
- Percent of participants with potential corticosteroid-related treatment-emergent adverse events (TEAEs) (time frame: 15 weeks after the start of treatment)
- Mean change in blood pressure (BP) at end of treatment (time frame: baseline, 12 weeks after the start of treatment)

Notes

Sponsor: Takeda (Shire)/Meritage Pharma, Inc

Conflicts of interest: The authors disclose the following: Sandeep K. Gupta was a principal investigator for this study, is a consultant to Meritage Pharma, Inc, Abbott, and Receptos, Inc, and is a member of the Medical Advisory Panel of the American Partnership for Eosinophilic Disorders and the executive committee of the International Gastrointestinal Eosinophil Researchers. Joanne M. Vitanza was employed by Meritage Pharma, Inc, as the medical monitor during the design and conduct of this study, is a consultant to Meritage Pharma, Inc, and owns shares of Meritage Pharma, Inc, stock. Margaret H. Collins performed the central pathology slide review for this study, is a consultant to Meritage Pharma, Inc, Novartis, Receptos, Inc, and Aptalis, and is a member of the Medical Advisory Panel of the American Partnership for Eosinophilic Disorders and the executive committee of the International Gastrointestinal Eosinophil Researchers.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The study sponsor confirmed that the randomization schedule was generated using a computer program.
Allocation concealment (selection bias)	Low risk	The study sponsor has confirmed that "Labeling of study treatment bottles was performed by an independent clinical services provider according to a blinded randomization scheme; active study drug and matching placebo were dispensed in identical bottles according to a blinded randomization scheme; and study participants, investigators, and the sponsor remained blinded to the randomization scheme until the blind was formally broken after all participants completed the study and the database was locked".
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Stated as "blinded participant, care provider, and investigator," and placebo drug looked similar to the intervention.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The study sponsor confirmed that the outcome assessors were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced and explained attrition that likely has no effect on outcome.

Gupta 2015 (Continued)

Selective reporting (reporting bias)	Low risk	Protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. Noted differences in PPI and H2 use, but this appears to be random associated with the selection process.

Heine 2019
Study characteristics

Methods	<p>RCT design and number of study arms: RCT, 2 arms</p> <p>Single-center or multi-center: multi-center - Australia; The Royal Childrens Hospital - Parkville; Women's and Childrens Hospital - North Adelaide; Monash Medical Centre - Clayton campus - Clayton; The Children's Hospital at Westmead - Westmead</p> <p>Countries: USA, Australia</p> <p>Study dates: August 2012 to July 2015</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active at randomization</p> <p>EoE definition/diagnostic criteria: EoE (≥ 15 eosinophils per high-power field; HPF)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Stage 1: All patients with possible EoE (age 1 to 18 years) undergoing a diagnostic gastroscopy at any of the 4 participating hospitals (at Royal Children's Hospital Melbourne, Monash Medical Centre Melbourne, Children's Hospital at Westmead Sydney, and Women's and Children's Hospital Adelaide) will be assessed for inclusion in the study. • Patients undergoing a diagnostic gastroscopy will be assessed for upper gastrointestinal symptoms, including: nausea and vomiting, abdominal pain, diarrhea, weight loss, dysphagia, regurgitation, food bolus obstruction, food refusal • Stage 2: Patients will be included in the randomized trial if a diagnosis of EoE was confirmed based on biopsies from the first gastroscopy. Biopsies will be analyzed by the hospital's anatomical pathologist to assess whether the diagnostic criteria for EoE is confirmed. The diagnosis of EoE is based on the presence of at least 20 mucosal eosinophils/HPF (abstract says ≥ 15) in any of the upper or lower esophageal biopsies. NB: Patients who are already avoiding a food prior to the study as part of a known food allergy (e.g. egg or peanut allergy) can still be considered for the study if EoE was diagnosed while avoiding the food (which suggests that the food is not causing the EoE). <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • The patient does not fulfil the histological criteria for EoE (fewer than 20 mucosal eosinophils/HPF on histology). Recent treatment with systemic corticosteroids (prednisolone) or topical or inhaled corticosteroids (fluticasone aerosol or viscous budesonide) within 3 months of the gastroscopy. • Recent or current treatment with omeprazole • Patient already on multiple food elimination diet • Previous failure of elimination diets (as it may skew the data towards non-response) • Inability to obtain informed consent • Inability to comply with the prescribed 4-food elimination diet or PPI treatment • Contraindications to gastroscopy or general anesthesia <p>Age at beginning of study per study group: mean age total study 9.1 years</p> <p>Sex (m/f) per study group (numbers of patients): NR</p>

Medical treatment of eosinophilic esophagitis (Review)

Heine 2019 (Continued)

Number randomized per study group:

Proton pump inhibitor (PPI) plus 4-food elimination diet: 32

Proton pump inhibitor (PPI): 32

Number reaching end of study per study group (numbers of patients):

Proton pump inhibitor (PPI) plus 4-food elimination diet: 27/32

Proton pump inhibitor (PPI): 30/32

Interventions	<p>Study group 1: active treatment, oral proton pump inhibitor (PPI) plus 4-food elimination diet (strictly avoiding all foods containing cow's milk, soy, wheat or egg) + omeprazole: 7.5 kg to 9.9 kg: 5 mg morning and 10 mg night, 10.0 kg to 14.9 kg: 10 mg twice-daily, 15.0 kg to 19.9 kg: 15 mg twice-daily, > 20 kg: 20 mg twice-daily</p> <p>Study group 2: omeprazole: 7.5 kg to 9.9 kg: 5 mg morning and 10 mg night, 10.0 kg to 14.9 kg: 10 mg twice-daily, 15.0 kg to 19.9 kg: 15 mg twice-daily, > 20 kg: 20 mg twice-daily</p>
Outcomes	<p>Primary outcomes of the study: histological as number of eosinophils per high-power microscopic field</p> <p>Secondary outcomes of the study: Clinical response (symptom score) Endoscopic appearance (endoscopy score)</p>
Notes	<p>Funding source: National Health and Medical Research Council (NHMRC Project Grant #1029972) - Australia</p> <p>Conflicts of interest: NR</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Treatment was centrally allocated.
Blinding of participants and personnel (performance bias) All outcomes	High risk	This was an open-label trial.
Blinding of outcome assessment (detection bias) All outcomes	High risk	This was an open-label trial. The authors were contacted in November 2022 to determine if assessors were blinded. No response.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Attrition numbers were given per group but no reasons provided. Authors contacted in November of 2022. No response.
Selective reporting (reporting bias)	High risk	The authors did not provided their outcome of complete remission of < 5 eosinophils per high-power field. Authors contacted in November of 2022. No response.

Medical treatment of eosinophilic esophagitis (Review)

Heine 2019 (Continued)

Other bias	Low risk	No major baseline differences between groups. No other concerns.
------------	----------	--

Hirano 2019
Study characteristics

Methods	<p>RCT design and number of study arms: multi-center, double-blind trial, 3 arms</p> <p>Single-center or multi-center: multi-center (30)</p> <p>Countries: US, Canada, Switzerland</p> <p>Study dates: September 2014 through December 2015</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: symptoms of dysphagia for a minimum of 4 days over 2 weeks (within the 4-week screening period) and histologic evidence of EoE, defined as a peak count of ≥ 15 eosinophils per high-power field (eos/hpf; microscope hpf = 0.3 mm²) at any 2 of 3 levels of the esophagus (proximal, mid, distal) when off anti-inflammatory therapy for EoE.</p> <p>Inclusion criteria: Patients must have previously received an adequate trial of a proton pump inhibitor to exclude gastroesophageal reflux disease and proton pump inhibitor-responsive esophageal eosinophilia as the primary cause of their symptoms. Prior treatment of patients with steroids for EoE was recorded, with steroid refractory defined as an adequate trial of systemic or swallowed topical steroids failing to result in a meaningful reduction in symptoms, as judged by the investigator. Participants with a partial response to a proton pump inhibitor (PPI) who met all other eligibility criteria could be enrolled; prospective participants who discontinued use of a PPI had to wait at least 4 weeks before their screening endoscopy; if a prospective participant was receiving a PPI at screening, the participant must have been receiving a stable dose for at least 4 weeks before the screening endoscopy and agreed to continue on the same dose through week 16; men and women of childbearing potential had to agree to use adequate birth control measures during the trial and for 5 months after their last dose of study drug; all women of childbearing potential must have had a negative serum pregnancy test at screening and a negative urine (or serum) pregnancy test before dosing on day 1.</p> <p>Exclusion criteria: Exclusion criteria included clinical or endoscopic evidence of the presence of any other disease that may have interfered with or affected the histologic, endoscopic, and clinical symptom endpoints for this trial (e.g. erosive esophagitis grade 2 or above, Barrett's disease, upper gastrointestinal bleed, eosinophilic gastritis or gastroenteritis, duodenal or gastric eosinophilia on screening endoscopy, inflammatory bowel disease, significant hiatal hernia (> 3 cm)); presence of esophageal varices; evidence of severe endoscopic structural abnormality in esophagus (e.g. high-grade stenosis where an 8 mm to 10 mm endoscope could not pass through the stricture without dilation at the time of endoscopy); primary causes of esophageal eosinophilia other than EoE; evidence of immunosuppression or were receiving systemic immunosuppressive or immunomodulating drugs (e.g. methotrexate, cyclosporine, interferon alpha, tumor necrosis factor alpha inhibitors, antibodies to immunoglobulin E) within 5 drug half-lives before screening; were receiving systemic or swallowed topical corticosteroid medication; prospective participants with EoE treated with a corticosteroid must have not received a systemic corticosteroid within 8 weeks or swallowed topical corticosteroids within 4 weeks of the screening endoscopy or the start of the daily clinical symptom diary data collection during screening, whichever was performed first; presence of any other disease making conduct of the protocol or interpretation of the trial results difficult or that would have put the prospective participant at risk by participating in the trial (e.g. infection causing eosinophilia, gastritis, colitis, irritable bowel syndrome, and celiac disease, which have similar symptoms, neurologic or psychiatric illness that compromised the prospective participant's ability to accurately document symptoms of EoE); liver function impairment or persisting elevations of aspartate aminotransferase or alanine aminotransferase > 2 times the upper limit of normal (ULN), or direct bilirubin > 1.5 times the ULN; systemic or diarrheal illness following trav-</p>

Medical treatment of eosinophilic esophagitis (Review)

Hirano 2019 (Continued)

el or residence in endemic areas of parasitic/helminthic infections, history of clinical schistosomiasis, history of travel to endemic areas within preceding 6 months; ongoing infection (e.g. hepatitis B or C, human immunodeficiency virus, active tuberculosis); pregnancy or lactation; concurrent treatment with another investigational drug; prospective participants could not have participated in a concurrent investigational drug trial or have received an investigational drug within 5 drug half-lives before signing the informed consent form for this trial; weight less than 40 kg (88.2 pounds) or greater than 125 kg (275 pounds); history of February 2019 RPC4046 EoE Phase 2 Trial 603.e2 idiopathic anaphylaxis or a known history of a major immunologic reaction (such as anaphylactic reaction, anaphylactoid reaction, or serum sickness) to an immunoglobulin G-containing agent; history of cancer or lymphoproliferative disease, other than a successfully treated nonmetastatic cutaneous squamous cell or basal cell carcinoma or adequately treated cervical carcinoma in situ, within 10 years of screening; esophageal dilation for symptom relief during the screening period and within 4 weeks before baseline assessment of dysphagia or anticipated to be performed during the trial.

Age at beginning of study per study group:

Mean \pm SD: **placebo** 38.6 \pm 11.03; **RPC4046 (cendakimab) 180 mg** 39.1 \pm 9.87; **RPC4046 (cendakimab) 360 mg** 33.9 \pm 10.92

Median (range): **placebo** 38.5 (19, 64); **RPC4046 (cendakimab) 180 mg** 40.0 (19, 59); **RPC4046 (cendakimab) 360 mg** 31.5 (18, 63)

Sex (m/f) per study group:

Sex, n (%): male **placebo** 22 (64.7); **RPC4046 (cendakimab) 180 mg** 19 (61.3); **RPC4046 (cendakimab) 360 mg** 20 (58.8); female: **placebo** 12 (35.3); **RPC4046 (cendakimab) 180 mg** 12 (38.7); **RPC4046 (cendakimab) 360 mg** 14 (41.2)

Number randomized per study group (numbers of patients):

Placebo: 34

RPC4046 (cendakimab) 180 mg: 32

RPC4046 (cendakimab) 360 mg: 34

Number reaching end of study per study group (numbers of patients):

Placebo: 32

RPC4046 (cendakimab) 180 mg: 28

RPC4046 (cendakimab) 360 mg: 30

Interventions	<p>Study group 1 (placebo): placebo</p> <p>Study group 2: 180 mg RPC4046 (cendakimab) SC once-weekly (with initial loading dose of 5 mg/kg IV)</p> <p>Study group 3: 360 mg RPC4046 (cendakimab) SC once-weekly (with initial loading dose of 10 mg/kg IV)</p>
Outcomes	<p>Primary outcomes of the study: change in mean esophageal eosinophil count in the 5 HPF with the highest level of inflammation</p> <p>Secondary outcomes of the study: mean change in the dysphagia clinical symptom frequency and severity from baseline to week 16 as assessed by DSD completed over 2 weeks before the week 16 end-point. Other secondary outcomes included change in EEsAI PRO score, peak esophageal eosinophil count, EREFS, patient's and clinician's global assessments of disease severity, patient's global impression of change in EoE symptoms, and esophageal histologic severity (grade) and extent (stage).</p>
Notes	<p>Funding source: This study was sponsored by Celgene Corporation</p> <p>Conflicts of interest: These authors disclose the following: Ikuo Hirano has served as a consultant for Adare, Allakos, Celgene Corporation, Regeneron, and Shire, and has received grant/research support</p>

Hirano 2019 (Continued)

from Adare, Celgene Corporation, Regeneron, and Shire. Margaret H. Collins has served as a consultant for Celgene Corporation, Regeneron, and Shire and has received grant/research support from Celgene Corporation, Regeneron, and Shire. Sandeep Gupta has received grant/research support from Shire and served as a consultant for Abbott, Adare, Allakos, Celgene Corporation, and QOL. Alain M. Schoepfer has received grant/research support from Adare, Celgene Corporation, Falk, Merck Sharp & Dohme, and Regeneron, and has served as a consultant and advisor for AbbVie, Adare, Celgene Corporation, Falk, Merck Sharp & Dohme, and Regeneron. Alex Straumann has served as a consultant for Actelion, Calypso, Celgene Corporation, Falk, GlaxoSmithKline, Merck, Merck Sharp & Dohme, Novartis, Nutricia, Pfizer, Regeneron-Sanofi, Roche-Genentech, and Tillotts, and has received grant/research support from Celgene Corporation. Ekaterina Safroneeva has served as a consultant for Aptalis Pharma, Celgene Corporation, Novartis, and Regeneron. Michael Grimm, Heather Smith, Cindy-ann Tompkins, Amy Woo, Robert Peach, Paul Frohna, Sheila Gujrathi, Darryl N. Penenberg, Caiyan Li, and Richard Aranda were employees of Receptos at the time of the study; Receptos is now a wholly owned subsidiary of Celgene Corporation. Gregory J. Opitck and Allan Olson are employees of Celgene Corporation. Marc E. Rothenberg has served as a consultant for Adare, Allakos, AstraZeneca, Celgene Corporation, GlaxoSmithKline, NKT Therapeutics, Novartis, Pulm One, Shire, and Spoon Guru; has an equity interest in Immune Pharmaceuticals, NKT Therapeutics, Pulm One, and Spoon Guru; has received royalties from Teva for reslizumab; and is an inventor of patents owned by Cincinnati Children's Hospital Medical Center. Evan S. Dellon has served as a consultant for Adare, Alivio, Allakos, Banner, Celgene Corporation, Enumeral, GSK, Regeneron, Robarts, and Shire, has received grant/research support from Adare, Banner, Celgene Corporation, Meritage, Miraca, Nutricia, Regeneron, and Shire, and has received educational grants from Banner and Holoclara. The remaining authors disclose no conflicts.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Treatment was allocated centrally. "The study drug, RPC4046, and placebo solutions were identical in physical appearance."
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"The treatment each patient received was not disclosed to the investigator, trial center personnel, patient, sponsor, or their representatives. Each patient's treatment group assignment blind was not broken until all patients completed the doubleblind treatment period".
Blinding of outcome assessment (detection bias) All outcomes	Low risk	A central pathologist blinded to treatment allocation determined histologic changes.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Low risk	Trial protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	Most baseline features balanced across both groups. IgE and eosinophils higher in intervention groups than in placebo group.

Hirano 2020
Study characteristics
Medical treatment of eosinophilic esophagitis (Review)

Hirano 2020 (Continued)

Methods

RCT design and number of study arms: RCT (2 arms)

Single-center or multi-center: multi-center

Countries: USA

Study dates: May 2015 to July 2017

Participants

Active EoE or inactive EoE at beginning of study: active

EoE definition/diagnostic criteria:

Active esophageal inflammation was to be evident at screening (i.e. peak cell count 15 eosinophils per high-power field (eos/HPF): 400 magnification of a 0.3 mm² field) as indicated by esophageal pinch biopsy specimens from at least 2 of 3 esophageal sites from endoscopy performed no more than 2 weeks after at least 8 weeks of treatment with high-dose (or twice-daily dosed) PPIs

Inclusion criteria:

1. Male or female, 18 to 65 years old, with a documented diagnosis of EoE by endoscopy before or at screening. Note: Must include a demonstration of intraepithelial eosinophilic infiltration (peak cell count 15 eos/HPF (400, 0.3 mm²)) from esophageal biopsy specimens from endoscopy performed no more than 2 weeks after at least 8 weeks of treatment with high-dose (or twice-daily dosing) PPIs.
2. History (by patient report) of, on average, at least 2 episodes of dysphagia (with intake of solids off anti-inflammatory therapy) per week in the 4 weeks before screening and, on average, at least 2 episodes of documented dysphagia per week in the weeks between screening and baseline. Dysphagia is defined as trouble swallowing solid food, or having solid food stick, by patient report.
3. Must remain on a stabilized diet for at least 6 weeks before screening and during the course of the study; stable diet is defined as no initiation of single or multiple elimination diets or reintroduction of previously eliminated food groups.
4. SDI PRO score 5 at screening and baseline.
5. Documented history of or presence of 1 or more of any of the following: allergic disease (e.g. allergic asthma, allergic rhinitis, atopic dermatitis, or food allergies); blood eosinophil count 0.25 GI/L; serum total IgE 100 kU/L.
6. Willing and able to comply with all clinic visits and study-related procedures.
7. Able to understand and complete study-related questionnaires.
8. Provide signed informed consent.
9. Endoscopy with photographs performed at screening, with a demonstration of intraepithelial eosinophilic infiltration (peak cell count 15 eos/HPF) in at least 2 of the 3 biopsied esophageal regions (proximal, mid, or distal).

Exclusion criteria:

1. Prior participation in a dupilumab (anti-IL4R) clinical trial.
2. Other causes of esophageal eosinophilia or the following diseases: hypereosinophilic syndromes, Churg-Strauss vasculitis, or eosinophilic gastroenteritis.
3. History of achalasia, active *Helicobacter pylori* infection, Crohn's disease, ulcerative colitis, celiac disease, or prior esophageal surgery before screening.
4. Any esophageal stricture unable to be passed with a standard, diagnostic, adult (9 mm to 10 mm) upper endoscope, or any critical esophageal stricture that required dilation at screening.
5. History of bleeding disorders or esophageal varices.
6. Use of chronic aspirin, nonsteroidal agents, or anticoagulants within 2 weeks before screening; patients should not stop these agents solely to become eligible for entry into this study.
7. Treatment with an investigational drug within 2 months or within 5 half-lives (if known), whichever is longer, before screening.
8. Use of systemic glucocorticoids within 3 months or swallowed topical glucocorticoids within 6 weeks before screening.
9. Use of inhaled or nasal glucocorticoids within 3 months before screening and during the study, except stable dose for at least 3 months before screening biopsy (which cannot be changed during the study).

Medical treatment of eosinophilic esophagitis (Review)

119

Hirano 2020 (Continued)

10. Treatment with oral immunotherapy within 6 months before screening.
11. Allergen immunotherapy (sublingual immunotherapy and/or subcutaneous immunotherapy), unless receiving stable dose for at least 1 year before screening.
12. The following treatments within 3 months before the screening visit, or any condition that, in the opinion of the investigator, is likely to require such treatment(s) during the 3 months of study treatment: systemic immunosuppressive/immunomodulating drugs (e.g. omalizumab, cyclosporine, mycophenolate mofetil, interferon gamma, Janus kinase inhibitors, azathioprine, methotrexate, and leukotriene inhibitors (except stable dose for at least 3 months before screening)).
13. Diagnosis of active parasitic infection or having suspected parasitic infection, unless clinical and (if necessary) laboratory assessments have ruled out active infection before randomization.
14. Chronic or acute infection requiring treatment with systemic antibiotics, antivirals, or antifungals within 1 month before screening.
15. Use of oral antibiotics/anti-infectives within 2 weeks before screening.
16. Known or suspected immunosuppression, including history of invasive opportunistic infections (e.g. tuberculosis, nontuberculous mycobacterial infections, histoplasmosis, listeriosis, coccidioidomycosis, pneumocystosis, aspergillosis) despite infection resolution, otherwise recurrent infections of abnormal frequency, or prolonged infections suggesting an immunocompromised status, as judged by the investigator.
17. Known history of human immunodeficiency virus infection.
18. Positive or indeterminate hepatitis B surface antigen or hepatitis C antibody at screening.
19. Elevated transaminases (alanine aminotransferase and/or aspartate aminotransferase) more than 3 times the upper limit of normal at screening.
20. History of malignancy within 5 years before screening, except completely treated in situ carcinoma of the cervix and completely treated and resolved nonmetastatic squamous or basal cell carcinoma of the skin.
21. History of patient-reported alcohol or drug abuse within 6 months before screening.
22. Any other medical or psychological condition, including relevant laboratory result abnormalities at screening, that, in the opinion of the investigator, suggest a new and/or insufficiently understood disease, may present an unreasonable risk to the study patient as a result of his/her participation in this clinical trial, may make the patient's participation unreliable, or may interfere with study assessments. The specific justification for patients excluded under this criterion was noted in study documents (chart notes, case report form, etc).
23. Severe concomitant illness(es) that, in the investigator's judgment, would adversely affect the patient's participation in the study.
24. Planned or anticipated use of any prohibited medications or procedures during study treatment.
25. Treatment with a live (attenuated) vaccine within 3 months before screening.
26. Patient or his/her immediate family is a member of the investigational team.
27. Pregnant or breastfeeding women, or women planning to become pregnant or breastfeed during the study.
28. Women unwilling to use adequate birth control, if of reproductive potential*, and sexually active. Adequate birth control is defined as agreement to consistently practice an effective and accepted method of contraception for the duration of the study and for 120 days after the last dose of study drug; these include hormonal contraceptives, an intrauterine device, double barrier contraception (i.e. condom þ diaphragm), or male partner with documented vasectomy.

*For female participants, menopause is defined as at least 12 consecutive months without menses (if in question, follicle-stimulating hormone of 25 U/mL must be documented). Hysterectomy, bilateral oophorectomy, or bilateral tubal ligation must be documented, as applicable, and women with these documented conditions are not required to use additional contraception.

Age at beginning of study per study group:

Age, y, mean (SD): placebo 36.1 (12.75); dupilumab 33.1 (8.70)

Sex (m/f) per study group: placebo: 10/14; dupilumab: 13/10

Number randomized per study group (numbers of patients): placebo: 24; dupilumab: 23

Hirano 2020 (Continued)

Number reaching end of study per study group (numbers of patients): placebo: 20; dupilumab: 22

Interventions	<p>Study group 1: placebo</p> <p>Study group 2: weekly subcutaneous dupilumab 300 mg (loading dose, 600 mg on day 1)</p>
Outcomes	<p>Primary outcomes of the study:</p> <p>The primary efficacy endpoint was the change in SDI PRO dysphagia score from baseline to week 10</p> <p>Secondary outcomes of the study:</p> <p>Secondary SDI PRO endpoints included percent change in SDI PRO score from baseline to week 10 and percentage of patients with an SDI PRO score decrease of 3 points relative to baseline at week 10, which was proposed by Straumann et al as evidence of a clinical response. Other secondary endpoints, primarily evaluated at week 12, included histologic measures of type 2 inflammation in the esophagus (as measured by esophageal intraepithelial eosinophilia), endoscopically anatomic measures of esophageal disease (i.e. exudate, rings, edema, furrows, and strictures), distensibility measures of esophageal function, and additional PROs. These endpoints were assessed by measuring percent change in peak esophageal intraepithelial eos/HPF from baseline to week 12 and change in EoE Endoscopic Reference Scoring System (EREFS) score from baseline to week 12. Other secondary efficacy endpoints were percentage of patients requiring rescue medication or a procedure (e.g. esophageal dilation) through week 12 and the PRO and quality of life endpoints of absolute and percent change in weekly Eosinophilic Esophagitis Activity Index (EESAI) PRO score from baseline to week 10, percentage of patients with 40% improvement or > 15- or > 30-point improvement in EESAI PRO score from baseline to week 10, and change in Adult Eosinophilic Esophagitis Quality of Life (EoE-QOL-A) score, version 3.0 from baseline to week 12. Symptomatic remission of EoE, defined as an EESAI score of 20 at weeks 10 and 12, was also assessed in a post hoc analysis, as were the proportions of patients who achieved both histologic (< 6 eos/hpf at week 12) and symptomatic remission (SDI score reduction of 3 points relative to baseline at week 10) and both histologic and endoscopic remission. Safety was evaluated by incidence of treatment-emergent adverse events (TEAEs) and serious adverse events from baseline to week > 28.</p>
Notes	<p>Funding source: This research was sponsored by Sanofi and Regeneron Pharmaceuticals, Inc.</p> <p>Conflicts of interest:</p> <p>These authors disclose the following: Ikuo Hirano has been a consultant for Adare, Allakos, Receptos/Celgene, Regeneron Pharmaceuticals, Shire, Gossamer, Esocap and has received research funding from Adare, Allakos, Meritage, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire. Evan S. Dellon has been a consultant for Alivio, Adare, Allakos, Banner, Calypso, Enumeral, EsoCap, GlaxoSmithKline, Receptos/Celgene, Regeneron Pharmaceuticals, Robarts, and Shire; has received research funding from Adare, Allakos, Meritage, Miraca, Nutricia, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire; and received educational grants from Banner and Holoclara. Jennifer D. Hamilton, Qiong Zhao, Zhen Chen, Neil N. M. Graham, Bolanle Akinlade, and Allen Radin are employees and shareholders of Regeneron Pharmaceuticals. Margaret H. Collins has been a consultant for Allakos, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire and has received research funding from Receptos/Celgene, Regeneron Pharmaceuticals, and Shire. Kathryn Peterson has received research funding from Janssen, Receptos/Celgene, and Regeneron Pharmaceuticals. Mirna Chehade has been a consultant for Actelion, Allakos, and Shire and received research funding from Nutricia, Regeneron Pharmaceuticals, and Shire. Alain M. Schoepfer has been a consultant for Adare, Aptalis, Dr Falk Pharma, and Regeneron Pharmaceuticals and has received research funding from AstraZeneca, Aptalis, Dr Falk Pharma, GlaxoSmithKline, Nestlé, Novartis, Receptos/Celgene, and Regeneron Pharmaceuticals. Ekaterina Safroneeva has been a consultant for Aptalis, Novartis, Receptos/Celgene, and Regeneron Pharmaceuticals. Marc E. Rothenberg has been a consultant for AstraZeneca, Celgene, GlaxoSmithKline, NKT Therapeutics, Novartis, PulmOne, Shire, and Spoon Guru; holds equity interest in Immune Pharmaceuticals, NKT Therapeutics, PulmOne, and Spoon Guru; receives royalties from reslizumab from Teva Pharmaceutical; and is the inventor of patents owned by Cincinnati Children's Hospital Medical Center. Gary W. Falk has received research funding from Allakos; has been a consultant for Adare and Banner; and has received research funding from Adare, Meritage, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire. Gianluca Pirozzi and Leda Mannent L are employees of Sanofi and may hold stock and/or stock options</p>

Hirano 2020 (Continued)

in the company. Brian N. Swanson is a former employee of Sanofi and may hold stock and/or stock options in the company. The remaining author discloses no conflicts.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"Using a central interactive voice/web response system randomization".
Allocation concealment (selection bias)	Low risk	"Using a central interactive voice/web response system randomization".
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Blinded study drug kits coded with a medication numbering system were used, and everyone involved was blinded to all randomization assignments. Study patients and study site personnel remained blinded via the use of matching placebo.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome assessors were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Low risk	The trial registry is available and all pre-specified outcomes of interest were reported.
Other bias	Low risk	Baseline characteristics evenly distributed between treatment arms except for total IgE level.

Hirano 2020f
Study characteristics

Methods	<p>RCT design and number of study arms: randomized 3 arms</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: USA</p> <p>Study dates: October 2011 and October 2012</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: histologically confirmed EoE (esophageal mucosal peak eosinophil count ≥ 24 per high-power field (HPF) (HPF; radius = 0.275 mm; 400\times) in at least one biopsied site, within 30 days prior to and 21 days after the screening visit)</p> <p>Inclusion criteria:</p> <ol style="list-style-type: none"> 1. Eligible patients between 12 and 55 years age had histologically confirmed EoE (esophageal mucosal peak eosinophil count ≥ 24 per high-power field (HPF) (HPF; radius = 0.275 mm; 400\times) in at least one biopsied site (proximal/mid and/or distal), within 30 days prior to and 21 days after the screening visit) 2. Histologically confirmed prior treatment failure of a high-dose proton pump inhibitor (PPI), defined as peak eosinophil counts ≥ 24 per HPF after 8 weeks of 2\times standard PPI dose per investigator, and at least

Medical treatment of eosinophilic esophagitis (Review)

Hirano 2020f (Continued)

one of the following symptoms: chest pain or discomfort, dysphagia or food impaction continuously or intermittently present within 30 days prior to the screening visit. Specific symptom severity was not required for this study.

3. Females of child-bearing potential must have agreed to use adequate contraception during the study and could not be pregnant or lactating at time of enrollment
4. Written informed consent was obtained from each participant or caregiver/parent/guardian at screening

Exclusion criteria: Patients were excluded from participating in the study if they met any of the following criteria:

1. Presence of any condition, other than EoE, that affected the esophageal mucosa or motility
2. Any contraindication to completing esophagogastroduodenoscopy, including stricture that blocked the passage of a standard endoscope
3. History or presence of Crohn's disease, celiac disease, or other gastrointestinal inflammatory disease
4. Use of systemic, inhaled, intranasal or high-potency dermal topical corticosteroids during the 30 days prior to enrollment
5. Morning serum cortisol level $\leq 5 \mu\text{g/dL}$ or use of anti-inflammatory or immunosuppressant drugs

Also per the NCT registry:

1. Known contraindication, hypersensitivity or intolerance to corticosteroids
2. Any physical, mental, or social condition, history of illness or laboratory abnormality that, in the investigator's judgment, might interfere with the study procedures or the ability of the participant to adhere to and complete the study
3. Oral or esophageal mucosal infection of any type
4. Any medical condition in which the use of anti-inflammatory or immunosuppressant drugs are required or may be anticipated to be required during the study
5. History of esophageal or gastric surgery
6. Gastrointestinal bleeding
7. Current chronic infection, immunosuppression, immunodeficiency
8. Alcohol or drug abuse
9. Participation in a clinical study involving an investigations drug within 30 days of the screening visit

Age at beginning of study per study group:

Age (y), mean (SD):

Placebo 29.8 (13.9);

APT-1105 (fluticasone propionate orally disintegrating tablet) 1.5 mg twice-daily: 23.4 (11.3)

APT-1105 (fluticasone propionate orally disintegrating tablet) 3.0 mg once-daily: 24.6 (10.6)

Sex (m/f) per study group:

Placebo: 5/3

APT-1105 (fluticasone propionate orally disintegrating tablet) 1.5 mg twice-daily: 4/4

APT-1105 (fluticasone propionate orally disintegrating tablet) 3.0 mg once-daily: 6/2

Number randomized per study group:

Placebo: 8

APT-1105 (fluticasone propionate orally disintegrating tablet) 1.5 mg twice-daily: 8

APT-1105 (fluticasone propionate orally disintegrating tablet) 3.0 mg once-daily: 8

Number reaching end of study per study group:

Hirano 2020f (Continued)

Placebo: 6

APT-1105 (fluticasone propionate orally disintegrating tablet) 1.5 mg twice-daily: 8

APT-1105 (fluticasone propionate orally disintegrating tablet) 3.0 mg once-daily: 8

Interventions

Study group 1: placebo

Study group 2: APT-1105 (fluticasone propionate orally disintegrating tablet) 1.5 mg twice-daily

Study Group 3: APT-1105 (fluticasone propionate orally disintegrating tablet) 3.0 mg once-daily

Outcomes

Primary outcomes of the study:

1. Morning serum cortisol (change from baseline measure) (time frame: screening visit (up to 21 days), week 4, week 8 and follow-up)
2. Standard safety laboratory tests (time frame: screening visit (up to 21 days), week 4, week 8 and follow-up)
3. Treatment-emergent adverse events collection (time frame: screening visit (up to 21 days), randomization day, week 2 and week 6 (phone visit), week 4 and week 8 (office visit), follow-up (office visit, up to 11 days after week 8 visit))
4. Physical examination and vital signs collection (time frame: screening (up to 21 days), week 4, week 8 and follow-up)

Secondary outcomes of the study:

1. Esophagoduodenoscopy with multiple biopsies (time frame: screening (up to 21 days) and week 8)
2. Patient-reported outcomes and physician global assessment (time frame: screening (up to 21 days), week 4 and 8)

Notes

Funding source: This study was funded by Adare Pharmaceuticals, Inc. The study sponsor had a role in the study design, collection, analysis and interpretation of the data, as well as in the writing of the report.

Conflicts of interest:

IH has received research funding from Adare, Allakos, Meritage, Celgene/Receptos, Regeneron, Shire/Takeda; and consulting fees from Adare, Allakos, Arena, AstraZeneca, Biorasi, Celgene/Receptos, Eli Lilly, EsoCap, Gossamer Bio, Regeneron, Shire/Takeda. GMC is a consultant for Adare. ES has served as a consultant for Adare, Aptalis, Novartis, Receptos and Regeneron. MCR has no conflicts of interest to report. AS has served as consultant for AbbVie, Adare, Falk Pharma GmbH, MSD, Receptos, Regeneron, Novartis, Pfizer, Takeda and Vifor, and has received research funding from Adare, Falk Pharma GmbH, Receptos and Regeneron. GWF has received research support from Allakos, Receptos/Celgene and Regeneron, and has received research support and has served as a consultant for Adare and Shire/Takeda. GE has served as a consultant for and is currently employed by Adare.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Unclear risk	Allocation concealment is not reported. "Participants were given two bottles – one for the morning dose and one for the evening dose".
Blinding of participants and personnel (performance bias)	Low risk	All personnel, participants, caregivers, and the sponsor were blinded.

Medical treatment of eosinophilic esophagitis (Review)

Hirano 2020f (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	Low risk	The outcome assessors were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Minimal and explained attrition that likely has no effect on the outcome.
Selective reporting (reporting bias)	Unclear risk	In protocol outcomes not clearly defined, i.e. specific measures.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Hirano 2021
Study characteristics

Methods	<p>RCT design and number of study arms: randomized, double-blind, placebo-controlled trial 2 arms</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: United States</p> <p>Study dates: 2015 to 2019</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: (15 eosinophils/high-power field (eos/hpf) from at least 2 levels of the esophagus) during screening, with dysphagia on at least 4 days in any 2 consecutive weeks during screening and in the 2 weeks before randomization measured using the DSQ</p> <p>Inclusion criteria: Patients enrolled in this trial were 11 to 55 years of age. The lower limit of the age range chosen was considered to be the minimum age at which patients with eosinophilic esophagitis (EoE) could self-report symptoms (PRO) using the DSQ and was also the lower-bound age for which the DSQ had been validated. In addition, pediatric patients can present with different signs and symptoms than older patients with EoE. The upper limit of the age range was chosen as 55 years because patients older than this may present with fibrostenotic disease, so they are less likely to respond to anti-inflammatory therapy alone. Other key inclusion criteria included a history of clinical symptoms of esophageal dysfunction intermittently or continuously at screening as previously described; an absence of histologic response to 6 to 8 weeks of high-dose proton pump inhibitor (PPI) therapy, as per consensus guidelines in effect at the time of study onset (high-dose therapy refers to the total daily dose, which may have been administered as once- or twice-daily dosing); and a stable diet for at least 3 months before screening. A PPI trial may have occurred at the time of the qualifying esophagogastroduodenoscopy (EGD); in which case, the same PPI regimen was to continue, or this may have been done previously (in which case, PPI therapy may have been stopped if there was no response to therapy based on esophageal biopsy results). If PPI responsiveness was excluded by a previous EGD and biopsy, the historical EGD and biopsy must have been performed after the patient had been on a minimum of 6 weeks of high-dose PPI therapy.</p> <p>Exclusion criteria: Key exclusion criteria included the following: immunomodulatory therapy use 8 weeks before the qualifying EGD or anticipated use during the study; use of swallowed topical corticosteroids for EoE or systemic corticosteroids for any condition 4 weeks before the qualifying EGD and baseline or anticipated use during the study; presence of a high-grade esophageal stricture (defined as the presence of a lesion not allowing passage of a diagnostic adult upper endoscope (insertion tube diameter > 9 mm)); or following either a pure liquid diet or a 6-food elimination diet. In addition to having a stable (i.e. no changes) diet 3 months before screening, dosing with inhaled or nasal corticosteroids</p>

Medical treatment of eosinophilic esophagitis (Review)

Hirano 2021 (Continued)

and PPIs was to be stable for a specified period of time (inhaled corticosteroids for 3 months before screening; nasal corticosteroids and PPIs for 4 weeks prior to qualifying EGD).

Age at beginning of study per study group:
Mean age, y (SD) :

Placebo: 33.9 (12.1)

Budesonide oral suspension (BOS) 2.0 mg twice-daily: 33.8 (11.9)

Sex (m/f):

Placebo: 62/43

Budesonide oral suspension (BOS) 2.0 mg twice-daily: 129/84

Number randomized per study group:

Placebo: 107

Budesonide oral suspension (BOS) 2.0 mg twice-daily: 215

Number reaching end of study per study group:

Placebo: 94

Budesonide oral suspension (BOS) 2.0 mg twice-daily:202

Interventions	Study group 1: placebo Study group 2: budesonide oral suspension (BOS) 2.0 mg twice-daily
Outcomes	Primary outcomes of the study: The co-primary efficacy endpoints were (1) the proportion of stringent histologic responders (6 eos/hpf across all available esophageal levels (proximal, middle, or distal)) and (2) the proportion of dysphagia symptom responders (30% reduction in DSQ score) from baseline to week 12 of therapy Secondary outcomes of the study: The key secondary efficacy endpoint was the change in DSQ score from baseline to week 12 of treatment. Other secondary efficacy endpoints included the proportion of full responders, defined as a combined 2 stringent histologic response and dysphagia symptom response (6 eos/hpf and 30% reduction in DSQ score); the mean change in EoE Endoscopic Reference Score (EREFS) and maximum peak eosinophil count; the proportion of patients achieving a deep histologic response or histologic response (deep histologic response, 1 eos/hpf; histologic response, < 15 eos/hpf); and the mean change in the EoE Histology Scoring System (EoEHSS) total score ratios from baseline to week 12 of therapy
Notes	Funding source: This study was funded by Shire ViroPharma, Inc, a member of the Takeda group of companies. Conflicts of interest: "The authors disclose the following: Ikuo Hirano has received research funding from Adare Pharmaceuticals, Allakos, Arena Pharmaceuticals, AstraZeneca, Meritage Pharma, Inc, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda company; and served as a consultant for Adare Pharmaceuticals, Allakos, Arena Pharmaceuticals, AstraZeneca, EsoCap Biotech, Gossamer Bio, Lilly, Meritage Pharma, Inc, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda company. Margaret H. Collins has received research funding from Meritage Pharma, Inc, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda company; and served as a consultant for Allakos, Arena Pharmaceuticals, AstraZeneca, Calypso Biotech, EsoCap Biotech, GlaxoSmithKline, Receptos/Celgene, Regeneron Pharmaceuticals, Robarts Clinical Trials, Inc./Alimentiv, Inc, and Shire, a Takeda company. David A. Katzka has received research funding from Shire, a Takeda company; and served as a consul-

Medical treatment of eosinophilic esophagitis (Review)

Hirano 2021 (Continued)

tant for Receptos/Celgene. Vincent A. Mukkada has received research funding from Meritage Pharma, Inc, and Shire, a Takeda company; and served as a consultant for Shire, a Takeda company. Gary W. Falk has received research funding from Adare Pharmaceuticals, Allakos, Lucid, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda company; and served as a consultant for Adare Pharmaceuticals, Allakos, Bristol Myers Squibb, Lucid, Regeneron Pharmaceuticals, and Shire, a - 2021 Budesonide Oral Suspension for EoE 9 Downloaded for Anonymous User (n/a) at Nemours Children's Hospital Delaware from ClinicalKey.com by Elsevier on February 06, 2022. For personal use only. No other uses without permission. Copyright ©2022. Elsevier Inc. All rights reserved." Takeda company. Robin Morey, Nirav K. Desai, and James Williams are employees of Takeda Development Center Americas, Inc, and stockholders of Takeda Pharmaceutical Company Limited. Lan Lan was an employee of Takeda Development Center Americas, Inc, and a stockholder of Takeda Pharmaceutical Company Limited at the time of the study. Evan S. Dellon has received research funding from Adare Pharmaceuticals, Allakos, AstraZeneca, GlaxoSmithKline, Meritage Pharma, Inc, Miraca Life Sciences, Nutricia, Receptos/Celgene, Regeneron Pharmaceuticals, and Shire, a Takeda company; and served as consultant for Abbott Laboratories, Adare Pharmaceuticals, Aimmune Therapeutics, Allakos, Amgen, Arena Pharmaceuticals, AstraZeneca, Biorasi, Calypso Biotech, Celldex Therapeutics, Inc, EsoCap Biotech, GlaxoSmithKline, Gossamer Bio, Lilly, Parexel, Receptos/Celgene, Regeneron Pharmaceuticals, Roberts Clinical Trials, Inc/ Alimentiv, Inc, Salix Pharmaceuticals, Sanofi, and Shire, a Takeda company; and received educational grants from Allakos, Banner Life Sciences, and Holoclara.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated coded randomization method.
Allocation concealment (selection bias)	Unclear risk	Allocation concealment is not described.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	The study team, study sites, and patients were blinded. An unblinded and independent data team handled data processing, review and validation.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Outcome assessors were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Balanced and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Low risk	Trial protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Kliewer 2019
Study characteristics

Methods **RCT design and number of study arms:** RCT; 2 groups for 12 weeks - A and B. Non-responders at 12 weeks go from A to B; and B go to C (1-food, 4-food, steroids are ABC)

Single-center or multi-center: multi-center

Medical treatment of eosinophilic esophagitis (Review)

Kliewer 2019 (Continued)

Countries: USA

Study dates: March 2016 to May 2018

Participants

Active EoE or inactive EoE at beginning of study: active EoE (> 15 eos/hpf)

EoE definition/diagnostic criteria: > 15 eos/hpf

Inclusion criteria:

1. Have diagnosis of EoE (based on consensus criteria)
2. Aged 6 to 17 years
3. Have histologically confirmed active disease > 15 eosinophils/hpf in either distal or proximal esophagus within 12 weeks of screening visit
4. PPI confirmation
5. Symptomatic (have experienced symptoms within the last month prior to enrollment)
6. Has a negative urine pregnancy test at screening if of childbearing potential. Females of childbearing potential must have a negative urine pregnancy test (β -hCG) prior to enrollment into the study (i.e. at screening). Subsequently, these participants must agree to use adequate birth control measures (e.g. condom, oral/injectable/subcutaneous contraceptives, intrauterine device, or sexual abstinence) during the study and for at least 1 month after the last dose of study drug which will be documented in the source documents.

Exclusion criteria:

1. Have been treated with topical swallowed steroids within the last 2 months or systemic steroids within the past 3 months
2. Have eosinophilia in segments of the GI tract other than the esophagus
3. Have been diagnosed with a GI malabsorption disorder (i.e. inflammatory bowel disease, Crohn's disease) or celiac disease
4. Are currently on dietary therapy avoiding milk
5. Have concurrent *H. pylori* gastritis or parasitic infection
6. Are unable to obtain esophagogastroduodenoscopy with esophageal biopsies at Cincinnati Children's Hospital Medical Center (CCHMC) or other participating institution within 4 weeks of study completion
7. Have previously failed (in a clinical trial setting) dietary therapy with one of these regimens or topical steroid treatment with fluticasone at a total dose of 1760 μ g per day
8. Have definitely responded (in a clinical trial setting) to either dietary therapy avoiding these antigens or to swallowed fluticasone at a total dose of 1760 μ g per day
9. Are concurrently receiving any of the prohibited medications listed in Table 2
10. On immunotherapy for pollen (if not on maintenance therapy) or immunoglobulin E (IgE)-mediated food allergy

Age at beginning of study per study group: 6 to 17 years

Sex (m/f) per study group: NR

Number randomized per study group: group A and 25 group B

Number reaching end of study per study group: 33 group A; 16 group B. Data NA in published literature on A to B and B to C.

Interventions

 Study group 1 (control or placebo): no control group
 Group A got 1 food, B got 4 foods and C was steroids

Outcomes

Primary outcome measures:

1. Change from baseline in Pediatric EoE Symptom Score Version 2.0 (PEESS V2.0) at 12 weeks (time frame: baseline and 12 weeks)
2. Within-group comparisons (baseline vs week 12) of PEESS V2.0 scores (time frame: baseline and 12 weeks)

Kliewer 2019 (Continued)

Secondary outcome measures:

1. Percent of participants in histologic remission (< 15 eosinophils per high-power field) at 12 weeks (time frame: 12 weeks)
2. Percent of participants on swallowed glucocorticoids (SGC) in histologic remission (< 15 eos/hpf) at 12 weeks in phase 2 (time frame: 12 weeks)
3. Percent of 1FED non-responders on 4FED in histologic remission (< 15 eos/hpf) at 12 weeks in phase 2 (time frame: 12 weeks)
4. Change from baseline in Pediatric Quality of Life Inventory Version 3.0 EoE Module (PedsQL 3.0EoE) at 12 weeks (time frame: baseline and 12 weeks)
5. Change from baseline in Pediatric Quality of Life Inventory Version 4.0 (PedsQL 4.0) Generic Core Scales at 12 weeks (time frame: baseline and 12 weeks)
6. Change from baseline in Endoscopic Reference Score at 12 weeks (time frame: baseline and 12 weeks)
7. Percent of participants with positive and negative milk skin prick tests responding to 1FED (time frame: baseline and 12 weeks)

Notes
Funding source: Patient-Centered Outcomes Research Institute

Conflicts of interest: NR, several of the authors are conflicted

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The author confirmed "a computer-generated randomization schedule was used (SAS random number generator)".
Allocation concealment (selection bias)	Low risk	The author confirmed that "the Data Management Coordinating Center who was not involved in the study allocated the participants to a treatment arm. The DMCC generated an electronic notification (email) to the study coordinator at randomization".
Blinding of participants and personnel (performance bias) All outcomes	High risk	This was an open-label study.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The author confirmed that "the pathologists were blinded to the treatment allocation. The endoscopist was not actively blinded to treatment allocation".
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	Slightly more people withdrew their participation from the 4FED group (n = 6) than the 1FED group (n = 2) during phase 1. In total 34 people completed 4FED compared to 17 completing 1FED. This might have impacted our outcomes.
Selective reporting (reporting bias)	Low risk	The results have been posted on the trial registry website and have been appropriately reported for the pre-cross-over phase 1 of the trial.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Kliewer 2021
Study characteristics
Methods
RCT design and number of study arms: RCT, 2 arms

Medical treatment of eosinophilic esophagitis (Review)

Kliewer 2021 (Continued)

Single-center or multi-center: multi-center, 10 sites

Countries: USA

Study dates: May 2016 to May 2019

Participants

Active EoE or inactive EoE at beginning of study: active EoE

EoE definition/diagnostic criteria: ≥ 15 eos/hpf + symptoms, and lack of PPI response

Inclusion criteria:

- Have diagnosis of EoE (based on consensus criteria)
- Have histologically confirmed active disease > 15 eosinophils/hpf in either distal or proximal esophagus within 12 weeks of screening visit
- Symptomatic (have experienced symptoms within the last month prior to enrollment)
- Proton pump inhibitor (PPI) confirmation
- Have a negative urine pregnancy test at screening if of childbearing potential

Exclusion criteria:

- Have been treated with topical swallowed steroids within the last 2 months or systemic steroids within the last 3 months
- Have pathological eosinophilia in segments of the GI tract other than the esophagus determined by local review
- Have been diagnosed with a GI malabsorption disorder (i.e. inflammatory bowel disease, Crohn's disease) or celiac disease
- Are currently on dietary therapy strictly avoiding milk or on a 6FED
- Have concurrent *H. pylori* gastritis or parasitic infection
- Have history of anaphylaxis to milk (with current avoidance of milk)
- Have previously failed strict dietary therapy clearly documented with one of these regimens or topical steroid treatment (i.e. have achieved histological remission of < 15 eos/hpf after having been on fluticasone or > 1 mg budesonide per day)
- Use of investigational drugs within 4 weeks (one month) prior to enrollment
- Are concurrently receiving any of the prohibited medications for the study
- On immunotherapy for pollen (if not on maintenance therapy) or immunoglobulin-E (IgE)-mediated food allergy
- Past or current medical problems or findings from physical examination or laboratory testing that are not listed above, which, in the opinion of the investigator, may pose additional risks from participation in the study, may interfere with the participant's ability to comply with study requirements or that may impact the quality or interpretation of the data obtained from the study.

Age at beginning of study per study group:

1-food elimination: 36.4 (10.2)

6-food elimination: 37.8 (10.4)

Sex (m/f) per study group:

1-food elimination: 55%, 37/67, m/f

6-food elimination: 53%, 33/62, m/f

Number randomized per study group:

1-food elimination: 67

6-food elimination: 62

Number reaching end of study per study group: NR

Kliewer 2021 (Continued)

Interventions	Study groups: <ul style="list-style-type: none"> • 1-food elimination: animal milk • 6-food elimination: animal milk, wheat, egg, soy, tree nuts/peanuts, seafood 	
Outcomes	Primary outcome measures: <ol style="list-style-type: none"> 1. Percent of participants in histologic remission (< 15 eos/hpf) at 6 weeks Secondary outcome measures: <ol style="list-style-type: none"> 1. Percent of participants in complete (≤ 1 eos/hpf) and partial histologic remission (2 to 14 eos/hpf) at 6 weeks 2. Percent of participants in histologic remission following SGC in Phase 2 at 6 weeks 3. Percent of participants in histologic remission following 6FED in Phase 2 at 6 weeks 4. Change from baseline in peak eosinophil count at 6 weeks 5. Change from baseline in EoE histology scoring system (EoEHSS) at 6 weeks 6. Change from baseline in endoscopic reference score (EREFS) at 6 weeks 7. Change from baseline in EoE Symptom Activity Index (EESAI) at 6 weeks 8. Change from baseline in quality of life (EoE-QoL-A) at 6 weeks 	
Notes	Funding source: NIDDK NIAID NCATS, Office of Rare Diseases Conflict of Interest: NR	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The author confirmed "a computer-generated randomization schedule was used (SAS random number generator)".
Allocation concealment (selection bias)	Low risk	The author confirmed that "the Data Management Coordinating Center who was <u>not</u> involved in the study allocated the participants to a treatment arm. The DMCC generated an electronic notification (email) to the study coordinator at randomization".
Blinding of participants and personnel (performance bias) All outcomes	High risk	This was an open-label study.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The primary outcome was histologic response determined by blinded central pathology review.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All participants completed the study.
Selective reporting (reporting bias)	Low risk	Protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way, however 2 additional secondary outcomes were added to the study that were not included in the initial protocol, one of which was our secondary outcome of quality of life (using EoE-QoL-A). However, we did not think this could potentially bias our outcome results.
Other bias	Unclear risk	Chest pain is significantly higher in the 1-food group relative to the 6-food group at baseline. 85% of the participants in the 1-food group and 74% of the

Medical treatment of eosinophilic esophagitis (Review)

131

Kliewer 2021 (Continued)

participants in 6-food received an endoscopy. Peak eos/hpf are 1.4-fold higher in 1-food group relative to the 6-food group at baseline.

Konikoff 2006
Study characteristics

Methods	<p>RCT design and number of study arms: double-blind RCT, 2 arms</p> <p>Single-center or multi-center: Cincinnati Children's Hospital Medical Center; 2 patients were also enrolled at Children's Hospital, San Diego</p> <p>Countries: USA</p> <p>Study dates: 10 January 2003 and 16 August 2005</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: eosinophilic esophagitis was defined as the presence of ≥ 24 eosinophils in any 400x HPF in at least one biopsy specimen from either the proximal or distal esophagus and the presence of epithelial hyperplasia after careful examination of all microscopic fields</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • 3 and 30 years • the presence of ≥ 24 eosinophils in any 400x HPF in at least one biopsy specimen from either the proximal or distal esophagus and the presence of epithelial hyperplasia after careful examination of all microscopic fields <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Patients were excluded from the study if they had a history of poor tolerance to fluticasone propionate, were unable to co-operate with the use of a metered-dose inhaler, were pregnant, or had taken any corticosteroid (including inhaled, nasal, or systemic) within 3 months <p>Age at beginning of study per study group:</p> <p>Group: age mean (SE, range)</p> <p>Fluticasone propionate: 8.5 (0.8, 3 to 16)</p> <p>Placebo: 11.2 (1.3, 3 to 18)</p> <p>Sex (m/f) per study group:</p> <p>Fluticasone propionate: M/F (M%) = 17/4</p> <p>Placebo: M/F (M%) = 9/6</p> <p>Number randomized per study group:</p> <p>Fluticasone propionate: n = 21</p> <p>Placebo: n = 15</p> <p>Number reaching end of study per study group:</p> <p>Fluticasone propionate: 20/21 (95%)</p> <p>Placebo: 11/15 (73%)</p>
Interventions	<p>Study group 1: placebo twice-daily for 3 months via metered dose inhaler</p>

Medical treatment of eosinophilic esophagitis (Review)

Konikoff 2006 (Continued)

Study group 2: fluticasone propionate (400 µg twice-daily for 3 months) via metered dose inhaler and swallowed

Outcomes	<p>Primary outcomes of the study: histological remission (≤ 1 eosinophil per hpf)</p> <p>Secondary outcomes of the study: adverse effects</p>
Notes	<p>Funding: supported by the Burroughs Wellcome Fund, the CURED Foundation, the Buckeye Foundation, an American Academy of Allergy Asthma & Immunology/Sanofi-Aventis Women Physician in Allergy grant, and a grant from the US Public Health Service (NIH T32 DK007727)</p> <p>Conflicts of interest: NR</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	"A clinical research coordinator dispensed metered-dose inhalers containing either active drug or placebo to each patient according to a computer-generated randomization list".
Blinding of participants and personnel (performance bias) All outcomes	Low risk	All participants and study personnel (with the exception of the above clinical research co-ordinator) were blinded to treatment assignment for the duration of the study. "Only the study statisticians had access to the unblinded data, but they did not have contact with study participants".
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All participants and study personnel (with the exception of the above clinical research co-ordinator) were blinded to treatment assignment for the duration of the study. "Only the study statisticians had access to the unblinded data, but they did not have contact with study participants".
Incomplete outcome data (attrition bias) All outcomes	Low risk	Minimal and explained attrition that likely has no effect on outcomes.
Selective reporting (reporting bias)	Low risk	"The primary outcome measure [...] was complete histologic response to treatment. Secondary outcome measures included presence of endoscopic furrowing, presence of epithelial hyperplasia, and presence of clinical symptoms". The outcomes were all reported.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Lieberman 2018
Study characteristics

Methods	<p>RCT design and number of study arms: RCT, 2 arms</p> <p>Single-center or multi-center: single-center</p> <p>Countries: USA</p> <p>Study dates: December 2014 to December 2017</p>
Participants	Active EoE or inactive EoE at beginning of study: active

Medical treatment of eosinophilic esophagitis (Review)

Lieberman 2018 (Continued)

EoE definition/diagnostic criteria: ≥ 15 eosinophils per high-power field (eos/hpf) following at least 8 weeks of high-dose PPI therapy and a normal esophageal pH probe

Inclusion criteria:

- Diagnosis of eosinophilic esophagitis (previously PPI non-responsive)
- Age 2 to 17 years of age

Exclusion criteria:

- Concomitant treatment with swallowed corticosteroids (any prior use of swallowed corticosteroids required an 8-week washout period)
- Pregnancy
- Evidence of pathologic eosinophilia in other locations in the gastrointestinal tract
- Active participation in another research protocol
- Renal or hepatic insufficiency

Age at beginning of study per study group: (mean (SD))

- Cromolyn: 10.4 (4.1)
- Placebo: 12.8 (3.7)

Sex (m/f) per study group:

- Cromolyn: 4/5 (m/f)
- Placebo: 4/7 (m/f)

Number randomized per study group:

- Cromolyn: 9
- Placebo: 7

Number reaching end of study per study group:

- Cromolyn: 9/9
- Placebo: 6/7

Interventions	<p>Study group 1: saline ampules, participants 2 to 12 years of age - 1 ampule mixed with 1 teaspoon of sugar 4 times daily, participants 13 to 18 years of age - 2 ampules mixed with 2 teaspoons of sugar 4 times daily</p> <p>Study group 2: participants 2 to 12 years of age - 100 mg of cromolyn - 1 ampule mixed with 1 teaspoon of sugar 4 times daily, participants 13 to 18 years of age - 200 mg cromolyn - 2 ampules mixed with 2 teaspoon of sugar 4 times daily</p>
Outcomes	<p>Primary outcomes of the study:</p> <ul style="list-style-type: none"> • Change in peak esophageal eosinophilia after 8 weeks of therapy <p>Secondary outcomes of the study:</p> <ul style="list-style-type: none"> • Change in PEES score at 4 and 8 weeks of therapy • Adverse events from therapy
Notes	<p>Funding source:</p> <ul style="list-style-type: none"> • American College of Allergy, Asthma, and Immunology Young Investigator Award (to Dr Lieberman) • Drug provided for free by MEda Pharmaceuticals (Somerset, NJ) <p>Conflicts of interest:</p> <ul style="list-style-type: none"> • Dr Lieberman is an associate editor for Annals of Allergy, Asthma, & Immunology

Lieberman 2018 (Continued)

- He has served on the advisory board for Aimmune Therapeutics
- He is a principal investigator for clinical trials sponsored by Aimmune Therapeutics and Biotest Pharma
- Drs Zhang, Whitworth, and Cavender have no conflicts of interest to report

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The author confirmed that they performed a computer-generated randomization.
Allocation concealment (selection bias)	Low risk	The author confirmed that "a research pharmacist had the blinded randomization schedule. She was not involved in any other part of study. She was contacted when someone was enrolled and allocated treatment based on the blinded computer generated sequence that only she had".
Blinding of participants and personnel (performance bias) All outcomes	Low risk	The author confirmed that "study drug was matched to placebo in terms of the liquid vials as much as possible. Investigator never saw study drug subject received. Assignment was blinded to subject and investigators during the trial until study completion".
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The author confirmed that outcome assessors were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Low and explained attrition that likely has no effect on outcomes.
Selective reporting (reporting bias)	Low risk	Trial protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Lucendo 2019
Study characteristics

Methods	<p>RCT design and number of study arms: double-blind, randomized, placebo-controlled, 2 study arms</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: 26 centers in 6 countries (Belgium, Germany, Netherlands, Spain, Switzerland, UK)</p> <p>Study dates: November 2015 to October 2016</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: adults with active EoE</p> <p>EoE definition/diagnostic criteria: clinico-histologic active EoE:</p> <p>Patients had to have a severity of 4 points on a 0 to 10 numerical rating scale (NRS) for either dysphagia or odynophagia for 1 day in the week before randomization. Additionally, Patient's Global Assessment (PatGA) of EoE activity was to be 4 points on a 0 to 10 NRS. Histologic activity with peak eos $\geq 65/\text{mm}^2$ hpf in at least 1 hpf (corresponding to 20 eos/hpf), as measured in a total of 6 hpf derived from 6 biopsies, 2 each from the proximal, mid, and distal segments of the esophagus</p>

Lucendo 2019 (Continued)

Inclusion criteria: aged 18 to 75 years with clinico-histologic active EoE and refractory to treatment with a PPI for a 4-week period severity of ≥ 4 points on NRS for dysphagia or odynophagia PatGA was ≥ 4 points. Histologic activity with peak EOS $\geq 65/\text{mm}^2$ hpf in at least 1 hpf.

Exclusion criteria:

1. Clinical and endoscopic suspicion for GERD
2. Achalasia/scleroderma
3. Evidence of causes other than EoE for esophageal eosinophilia
4. Pathologic eosinophilic infiltration in gastric and duodenal biopsies
5. History of esophageal surgery at any time/esophageal dilation procedures within the last 8 weeks before screening
6. Any relevant systemic disease
7. Systemic glucocorticosteroids, immunosuppressants, biologic drugs within 4 weeks before screening/topical glucocorticosteroids within 2 weeks before screening and onset of dietary restrictions within 4 weeks before screening

Age at beginning of study per study group: y, mean (SD): budesonide orodispersible tablet (BOT): 37 (11.5); placebo: 37 (9.2)

Sex (m/f) per study group: male, n (%): BOT: 48/59 (81%); placebo: 25/29 (86%)

Number randomized per study group: 59 intervention group; 29 placebo group

Number reaching end of study per study group: 56 intervention group; 25 placebo group

Interventions	<p>Study group 1: matching placebo</p> <p>Study group 2: budesonide orodispersible tablet 1 mg twice-daily x 6 weeks, if no remission by 6 weeks were offered 6 weeks of open-label treatment with BOT</p>
Outcomes	<p>Primary outcomes of the study:</p> <p>Primary outcomes: clinical remission (including dysphagia and odynophagia severity + histologic remission)</p> <p>Secondary outcomes of the study:</p> <p>Secondary outcomes: histologic remission, change in peak eosinophil count, resolution of symptoms on each day in the week before the EoT and rate of clinical remission (EEsAI-PRO ≤ 20 at EoT)</p> <p>Further secondary efficacy variables</p> <p>Clinical weekly sum of daily 0 to 10 NRS dysphagia (range: 0 to 70)</p> <p>Rate of patients with overall symptoms resolution defined as PatGA 2 at week 6 (LOCF)</p> <p>Change from baseline to week 6 (LOCF) in blood eosinophil counts (eos/mm^3)</p> <p>Change from baseline to week 6 (LOCF) in total modified EREFS endoscopic score</p> <p>Change from baseline to week 6 (LOCF) in modified EREFS inflammatory signs subscore (0 to 4)</p> <p>Change from baseline to week 6 (LOCF) in modified EREFS fibrotic signs subscore (0 to 4)</p> <p>Rate of patients with histologic remission (i.e. peak eos $< 48/\text{mm}^2$ hpf; equivalent to < 15 eos/hpf) at week 6 (LOCF)</p> <p>Change from baseline to EoT DB phase in modified SHS symptom burden</p> <p>Change from baseline to EoT DB phase in modified SHS social function</p> <p>Change from baseline to EoT DB phase in modified SHS disease-related worry</p>

Lucendo 2019 (Continued)

- Change from baseline to EoT DB phase in modified SHS general well-being
- Change from baseline to EoT DB phase in EoE-QoL-A 30 items (weighted average)
- Change from baseline to EoT DB phase in EoE-QoL-A 24 items (weighted average)
- Change from baseline to EoT DB phase in EoE-QoL-A eating/diet impact 10 items (weighted average)
- Change from baseline to EoT DB phase in EoE-QoL-A eating/diet impact 4 items (weighted average)
- Change from baseline to EoT DB phase in EoE-QoL-A social impact (weighted average) (weighted average)
- Change from baseline to EoT DB phase in EoE-QoL-A emotional impact (weighted average)
- Change from baseline to EoT DB phase in EoE-QoL-A disease anxiety (weighted average)
- Change from baseline to EoT DB phase in EoE-QoL-A swallowing anxiety (weighted average)

Notes
Funding source:

Dr Falk Pharma GmbH, Freiburg, Germany funded this study and contributed to the design and conduct of the study; collection, management, analysis and scientific interpretation of the data; supported the manuscript preparation and reviewed the manuscript for medical and scientific accuracy. Approval of the manuscript, and the decision to submit the manuscript for publication was the responsibility of the authors.

Conflicts of interest:

These authors disclose the following: Alfredo J Lucendo has received research funding from Dr Falk Pharma; Stephan Miehle is a member of advisory boards for Celgene and EsoCap and has received speaker's fee from Dr Falk Pharma GmbH and Falk Foundation; Christoph Schlag has received consultant fees from EsoCap and speaker fees, travel and research funding from Dr Falk Pharma GmbH; Michael Vieth has received speaker and consultant fees from Dr Falk Pharma GmbH; Ulrike von Arnim is a member of MSD national advisory board, has received speaker fees from AbbVie, MSD, Falk Foundation, Pfizer, Takeda, and Vifor; Javier Molina-Infante has received speaker and consultant fees from Dr Falk Pharma GmbH; Dirk Hartmann has no conflicts of interest to declare; Albert Jan Bredenoord has received research funding from Nutricia, Norgine, and Bayer and received speaker and/or consulting fees from Laborie, EsoCap, Diversatek, Medtronic, Dr Falk Pharma GmbH, Calypso, Thelial, Regeneron, Celgene, Bayer, Norgine, AstraZeneca, Almirall, and Allergan; Constanza Ciriza de los Rios has received speaker fees from Casen Recordati; Ahmed Madisch has received speaker fees from Dr Falk Pharma GmbH and Falk Foundation; Jamal Hayat has received speaker fees from Dr Falk Pharma GmbH; Stephen Attwood has received speaker and consulting fees from Dr Falk Pharma GmbH; Ralph Mueller and Roland Greinwald are employees of Dr Falk Pharma GmbH; Alain Schoepfer is a member of an advisory board for Dr Falk Pharma GmbH, Adare Pharmaceuticals, Celgene Pharmaceuticals, and Regeneron Pharmaceuticals. He has received research funding from Dr Falk Pharma GmbH, Adare Pharmaceuticals, Celgene Pharmaceuticals, and Regeneron Pharmaceuticals. He has received speaker's fees from Dr Falk Pharma GmbH and Celgene Pharmaceuticals; Alex Straumann is a consultant of Calypso, EsoCap, Dr Falk Pharma GmbH, GSK, Receptos-Celgene, Regeneron-Sanofi, Shire and Tillotts, and has received speaker fees and research funding from Dr Falk Pharma GmbH. The remaining authors disclose no conflicts.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method
Allocation concealment (selection bias)	Low risk	"Allocation concealment was ensured as patients, investigators and their study team, the sponsor, monitoring staff, central laboratory, and central pathologist, were all kept blinded to the randomization sequence, the block

Medical treatment of eosinophilic esophagitis (Review)

137

Lucendo 2019 (Continued)

		size, and patient's treatment, until all patients had completed the study and the database was clean and locked. No individual unblinding was needed or performed".
Blinding of participants and personnel (performance bias) All outcomes	Low risk	All members of the study team and the participants were blinded for the study. BOT and corresponding placebo were identical in physical appearance and were administered twice-daily
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The study team including the central pathologist were blinded for the study.
Incomplete outcome data (attrition bias) All outcomes	Low risk	There was no attrition imbalance, and the attrition was explained.
Selective reporting (reporting bias)	Low risk	Trial protocol is available and all pre-specified outcomes of interest were reported.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Miehlke 2016
Study characteristics

Methods	<p>RCT design and number of study arms: RCT design, double-blind, double-dummy, 4 arms</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: Switzerland, Germany, and Belgium</p> <p>Study dates: June 2011 to April 2013</p>
Participants	<p>Active EoE or inactive EoE at beginning of the study: active EoE</p> <p>EoE definition/diagnostic criteria: clinical symptoms of esophageal dysfunction (dysphagia score ≥ 3), peak eosinophils (eos) $\geq 65/\text{mm}^2$ high-power fields (hpf) in at least 1 hpf (corresponding to ≥ 20 eos/hpf), and eosinophilic tissue infiltration with a mean cell density ≥ 16 eos/mm^2, as measured in a total of 30 hpf derived from 6 biopsies, 2 each from the proximal, mid, and distal segments of the esophagus</p> <p>Inclusion criteria:</p> <p>Patients between 18 and 75 years of age and confirmed clinicopathological diagnosis of EoE according to the above criteria</p> <p>Exclusion criteria:</p> <ol style="list-style-type: none"> 1. Clinical and endoscopic suspicion for GERD, achalasia, or scleroderma 2. History of abnormal pH monitoring of the distal esophagus or clinicopathological response to a treatment with proton pump inhibitors (PPIs) at a standard dose with a treatment duration of at least 2 weeks 3. Other clinical evidence of causes other than EoE for esophageal eosinophilia 4. Any concomitant esophageal disease and relevant GI disease 5. History of esophageal surgery at any time or of esophageal dilation procedures within the last 8 weeks prior to screening 6. Any relevant systemic disease if careful medical monitoring was not ensured 7. Abnormal hepatic function, liver cirrhosis, or portal hypertension

Medical treatment of eosinophilic esophagitis (Review)

Miehlke 2016 (Continued)

8. Abnormal renal function
9. History of cancer in the last 5 years
10. Upper GI bleeding within 8 weeks prior to screening
11. Systemic therapies for any reason that may have affected assessment of primary and secondary end points (i.e. systemic glucocorticoids, histamine antagonists, mast cell stabilizers, leukotriene receptor antagonists, biologics, immunosuppressants) concomitantly or within 4 weeks prior to screening
12. Treatment with topical therapies for any reason that may affect assessment of primary and secondary end points (i.e. topical glucocorticoids, inhaled sodium cromoglycate) concomitant or within 2 weeks prior to screening; concomitant therapy for more than 3 days with drugs, which might influence hepatic biotransformation (CYP3A inducers/inhibitors)
13. Installation of dietary restrictions within 4 weeks prior to screening or during treatment
14. Intake of grapefruit-containing food or beverages during the study treatment phase
15. Known intolerance/hypersensitivity to study drug; lack of patient's co-operation
16. Existing or intended pregnancy or breastfeeding and positive pregnancy test at screening in women with childbearing potential

Age at beginning of study per study group: mean (SD)

Placebo: 36.3 (9.9)

BET (budesonide effervescent tablet) 2 x 1 mg/day: 38.9 (12.6)

BET (budesonide effervescent tablet) 2 x 2 mg/day: 37.2 (13.9)

BVS (budesonide viscous suspension): 2 x 2 mg/day: 46.5 (14.1)

Sex (m/f) per study group:

Placebo: 16/3

BET (budesonide effervescent tablet) 2 x 1 mg/day: 17/2

BET (budesonide effervescent tablet) 2 x 2 mg/day: 16/3

BVS (budesonide viscous suspension): 2 x 2 mg/day: 14/5

Number randomized per study group:

Placebo: 19

BET (budesonide effervescent tablet) 2 x 1 mg/day: 19

BET (budesonide effervescent tablet) 2 x 2 mg/day: 19

BVS (budesonide viscous suspension): 2 x 2 mg/day: 19

Number reaching end of study per study group:

Placebo: 19

BET (budesonide effervescent tablet) 2 x 1 mg/day: 19

BET (budesonide effervescent tablet) 2 x 2 mg/day: 19

BVS (budesonide viscous suspension): 2 x 2 mg/day: 18

Interventions

Group 1: Placebo for 2 weeks

Group 2: BET (budesonide effervescent tablet) 2 x 1 mg/day for 2 weeks

Group 3: BET (budesonide effervescent tablet) 2 x 2 mg/day for 2 weeks

Group 4: BVS (budesonide viscous suspension): 2 x 2 mg/day for 2 weeks

Miehlke 2016 (Continued)

Outcomes

Primary outcomes of the study:

- Co-primary outcome: rate of histological remission (mean of < 16 eos/mm² hpf)
- Co-primary outcome: change in the mean numbers of eos/mm² HPF (eosinophil load) from baseline

Secondary outcomes of the study:

Endoscopic abnormality score, endoscopic intensity score and its subscores, endoscopic VAS score, dysphagia score, patient's acceptance and preference of study drugs, adverse events, morning serum cortisol, and assessment of tolerability by investigator and patient

Notes

Funding source: Dr Falk Pharma GmbH, Freiburg, Germany

Conflicts of interest: SM has received speaker's honoraria from Dr Falk Pharma GmbH. MV, MB, AM, HW, HDA and MR have received speaker's honoraria from the Falk Foundation. SS has received speaker's honoraria from Abbvie, the Falk Foundation and MSD. RM, KD and RG are employees of Dr Falk Pharma GmbH. AS is a consultant to Dr Falk Pharma GmbH and has received consulting fees and/or speaker fees and/or research grants from Actelion, AG, Switzerland, AstraZeneca, AG, Switzerland, Aptalis Pharma, Glaxo SmithKline, AG, Nestlé S. A., Switzerland, Novartis, AG, Switzerland, Pfizer, AG, and Regeneron Pharmaceuticals

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	There was central allocation of treatment.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Double-dummy strategy to maintain blinding when using different pharmaceutical preparations.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Double-dummy strategy to maintain blinding when using different pharmaceutical preparations. As the personnel used a double-dummy strategy, outcome assessment should be blind. "The biopsies were immediately placed into separate tubes with neutral-pH-buffered 4% paraformaldehyde solution and sent to the primary central pathologist". Histology seems to have been blinded to the pathologies but endoscopy was assessed for endoscopic appearance by a gastroenterologist. Details whether the gastroenterologists performing endoscopy was involved or not were not provided.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Attrition was minimal and explained.
Selective reporting (reporting bias)	Unclear risk	The authors did not report clinical response as defined in outcomes. They also stated significance in clinical outcome without numerical results.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Moawad 2013

Study characteristics

Methods	<p>RCT design and number of study arms: single (investigator)-blinded RCT, 2 arms</p> <p>Single-center or multi-center: single-center, Esophageal Clinic at Mayo Clinic in Rochester Minnesota</p> <p>Countries: USA</p> <p>Study dates: April 2008 to October 2010</p>
Participants	<p>Active EoE or inactive EoE at beginning of the study: active</p> <p>EoE definition/diagnostic criteria: one clinical symptom of esophageal dysfunction (dysphagia, food impaction, heartburn) with ≥ 15 eosinophils/hpf</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Adult patients (age ≥ 18 years) • All patients had at least one clinical symptom of esophageal dysfunction (dysphagia, food impaction, heartburn) with ≥ 15 eosinophils/HPF on index endoscopy <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Secondary hypereosinophilic disorders • Severe coagulopathy • Pregnancy <p>Age at beginning of study per study group (mean (SD), years):</p> <p>Esomeprazole: 37.0 ± 11.1</p> <p>Fluticasone: 38.0 ± 8.8</p> <p>Sex (m/f) per study group:</p> <p>Esomeprazole: 19/2</p> <p>Fluticasone: 19/2</p> <p>Number randomized per study group:</p> <p>Esomeprazole: 21</p> <p>Fluticasone: 21</p> <p>Number reaching end of study per study group:</p> <p>Esomeprazole: 21</p> <p>Fluticasone: 21</p>
Interventions	<p>Study group 1: esomeprazole 40 mg once daily for 8 weeks</p> <p>Study group 2: fluticasone proportionate 440 μg twice-daily for 8 weeks (metered dose inhaler)</p>
Outcomes	<p>Primary outcome: histological response defined as achieving < 7 eos/hpf in both proximal and distal esophageal biopsies following 8 weeks of treatment</p> <p>Secondary outcomes: symptomatic change in dysphagia (score from the Mayo Dysphagia Questionnaire); change in endoscopic and other histological findings</p>
Notes	<p>Funding source: none</p>

Moawad 2013 (Continued)

Conflicts of interest: none

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Sealed, opaque envelopes containing data on the sequence of randomization were maintained by a research pharmacist.
Blinding of participants and personnel (performance bias) All outcomes	High risk	A group of patients taking an oral tablet and the other taking an inhaler.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The author confirmed that endoscopists and pathologists were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Minimal and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Low risk	Trial protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Oliva 2018
Study characteristics

Methods	RCT design and number of study arms: RCT Single-center or multi-center: single-center Countries: Italy Study dates: during 2 years (not specified)
Participants	Active EoE or inactive EoE at beginning of study: active EoE EoE definition/diagnostic criteria: not specified Inclusion criteria: <ul style="list-style-type: none"> • Active • Pediatric Exclusion criteria: <ul style="list-style-type: none"> • NR Age at beginning of study per study group: not reported (abstract) Sex (m/f) per study group: not reported

Medical treatment of eosinophilic esophagitis (Review)

Oliva 2018 (Continued)

Number randomized per study group: not reported (of 74 patients evaluated, 64 were enrolled, but 50 completed the study and were considered for the final analysis)

Number reaching end of study per study group: NR

Interventions	<p>Study group 1: 6-food elimination diet</p> <p>Study group 2: swallowed fluticasone</p> <p>Study group 3: swallowed budesonide</p> <p>Study group 4: oral viscous budesonide</p>
Outcomes	<p>Primary outcomes of the study: percentage of histological responders (defined as < 15 eos/hpf)</p> <p>Secondary outcomes of the study: clinical symptom score, endoscopic score (not specified)</p>
Notes	<p>Funding source: NR</p> <p>Conflicts of interest: NR</p>

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Not reported.
Allocation concealment (selection bias)	Unclear risk	Not reported.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not reported but different treatments (including elimination diet) administered.
Blinding of outcome assessment (detection bias) All outcomes	High risk	Not reported but different treatments (including elimination diet) administered.
Incomplete outcome data (attrition bias) All outcomes	Unclear risk	64 patients randomized and 50 patients included (dropouts for the entire study, no report per treatment arm).
Selective reporting (reporting bias)	Unclear risk	Only histologic response rate reported.
Other bias	Unclear risk	No possible assessment of baseline imbalance.

Peterson 2010
Study characteristics

Methods	<p>RCT design and number of study arms: RCT</p> <p>Single-center or multi-center: single-center, University of Utah Health Sciences Center</p> <p>Countries: USA</p>
---------	---

Medical treatment of eosinophilic esophagitis (Review)

Peterson 2010 (Continued)

	Study dates: NR				
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: ≥ 15 eosinophils averaged over 5 high-power fields on esophageal biopsy in participants with symptoms of dysphagia, food impaction or chest pain</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Patients aged 18 to 80 with eosinophilic esophagitis, are defined as a) dysphagia, food impaction or other upper gastrointestinal symptoms (chest pain, heartburn, regurgitation); b) multiple esophageal rings or furrows; c) the presence of > 20 eosinophils/high-power field in the squamous epithelium or deeper tissues of the esophagus • Ability to undergo esophageal manometry and ambulatory pH monitoring • No history of bleeding diathesis, significant cardiopulmonary disease, or other contraindication to upper endoscopy • Those who have had a 1-month holiday from either esomeprazole therapy or fluticasone if they have been prescribed this prior to enrollment <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Patients were excluded if they had to be dilated on the first exam (due to the inability to pass an upper endoscope) • Had a history of a contraindication or intolerance of either fluticasone or PPIs • Were pregnant or incarcerated • Had a history of prior upper gastrointestinal surgery • Had other potential alternative mechanisms for dysphagia demonstrated at endoscopy • Current use of PPIs or glucocorticoids • Symptoms of food impaction to food bolus size < 1 cm <p>Age at beginning of study per study group: esomeprazole = 38.8 years (26 to 79), fluticasone = 34.6 years (18 to 58)</p> <p>Sex (m/f) per study group (numbers of patients): 12/M 3/F, 11/M 4/F</p> <p>Number randomized per study group: esomeprazole (n = 15), fluticasone (n = 15)</p> <p>Number reaching end of study per study group: NR (6 overall)</p>				
Interventions	<p>Study group 1 (control or placebo): esomeprazole 40 mg a day for 8 weeks</p> <p>Study group 2: swallowed aerosolized fluticasone (440 μg twice-daily)</p>				
Outcomes	<p>Primary outcomes of the study: "We anticipated an 80% response rate (defined as a decrease in dysphagia score of at least two points) in the fluticasone arm as compared to a 33% response rate in the esomeprazole arm as the primary endpoint".</p> <p>Secondary outcomes of the study: Secondary endpoints included changes in eosinophilic infiltration in esophageal biopsies. "We arbitrarily defined partial resolution as B15 eos/HPF and complete resolution as B5 eos/HPF".</p>				
Notes	<p>Funding source: funding support was provided in part by an ASGE Research grant</p> <p>Conflicts of interest: NR</p>				
Risk of bias					
Bias	<table border="1"> <thead> <tr> <th>Authors' judgement</th> <th>Support for judgement</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> </tr> </tbody> </table>	Authors' judgement	Support for judgement		
Authors' judgement	Support for judgement				

Peterson 2010 (Continued)

Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	The research staff assigned (central allocation) the participants to the treatment group after the investigator who performed the endoscopy enrolled the participant.
Blinding of participants and personnel (performance bias) All outcomes	High risk	Neither the participants nor the investigators were blinded to treatment.
Blinding of outcome assessment (detection bias) All outcomes	High risk	The pathologist who evaluated histology was blinded, however no other outcome assessment was blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	There was low attrition that was explained by the personnel.
Selective reporting (reporting bias)	High risk	The authors swapped the primary and secondary outcomes between the protocol and manuscript.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Rothenberg 2015
Study characteristics

Methods	<p>RCT design and number of study arms: RCT with 2 arms</p> <p>Single-center or multi-center: multi-center</p> <p>Countries: United States</p> <p>Study dates: December 2009 to February 2012</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: esophageal biopsy must show ≥ 24 eos per HPF (400x) in the proximal or distal esophagus</p> <p>Inclusion criteria:</p> <ol style="list-style-type: none"> 1. Male and female participants with symptomatic eosinophilic esophagitis aged 18 to 50 years. 2. Female participants were required to be surgically sterilized at least 6 months before study participation or postmenopausal (no regular bleeding for at least 2 years) confirmed by a plasma FSH level of > 40 IU/L at screening or baseline. 3. Esophageal biopsy must show ≥ 24 eos per HPF (400x) in the proximal or distal esophagus, validated by a central laboratory pathologist. 4. Elimination diet must either: not be indicated following allergy evaluation including skin prick testing with multiple food antigens, or have been refused to be followed by participants, or have undergone a minimum of 3 months of elimination diet as indicated by skin prick testing without detectable resolution by repeat endoscopy with biopsies, or participant refusal to follow elimination diet. 5. Participants treated with a proton pump inhibitor (PPI) must have been on treatment for at least 2 months before enrollment. The PPI must be used before endoscopy to rule out the possibility of GERD in the proximal or distal esophagus.

Medical treatment of eosinophilic esophagitis (Review)

Rothenberg 2015 (Continued)

6. Failure of histological improvement is defined as eosinophil density ≥ 24 per HPF (400x) after 2 months treatment with a PPI or documented by prior endoscopy or lack of complete disappearance of symptoms.
7. Participants had to be able to communicate well with the investigator, to understand and comply with the requirements of the study and also to understand and sign the written informed consent.

Exclusion criteria:

1. Participants have received systemic corticosteroid therapy, by any route of administration, within 3 months before dosing or have received topical corticosteroids (swallowed aerosolized fluticasone or budesonide) within 2 months before dosing.
2. Comorbid eosinophilic disorders (other than atopic dermatitis not requiring chronic steroid therapy).
3. History of exposure to human therapeutic antibody, immunoglobulin, or other plasma product within 6 months of dosing (e.g. Xolair).
4. History of clinical schistosomiasis, or stool examination positive for ova or parasites or travel within the preceding 6 months to an area with endemic schistosomiasis, including but not limited to South-east and Southwest Asia, South America, and Africa. Travel to these areas was not to be planned for at least 6 months after the last dose.
5. Participants who have not had a trial of PPI or prior allergy testing/elimination diet and who did not fulfill entry criteria after PPI or initiation of elimination diet therapy.
6. Participation in any clinical intervention with any drug administration within 4 weeks before initial dosing or longer if required by local regulations, and for any other limitation of participation based on local regulations.
7. Donation or loss of 400 mL or more of blood within 8 weeks before initial dosing, or longer if required by local regulation.
8. A past medical history of clinically significant ECG abnormalities. An abnormal ECG is defined as PR > 220 ms, QRS complex > 120 ms, QTcB > 430 ms, or any significant morphological changes, other than non-specific T-wave changes.
9. Any surgical or medical condition that might significantly alter the absorption, distribution, metabolism, or excretion of drugs, or which may jeopardize the participant in case of participation in the study. The investigator was to make this determination in consideration of the participant's medical history and/or clinical or laboratory evidence of any of the following: major gastrointestinal tract surgery such as gastrectomy, gastroenterostomy, or bowel resection.
10. History of immunodeficiency diseases, including a positive HIV (ELISA and Western blot) test result.
11. Positive hepatitis B surface antigen (HBsAg) or hepatitis C test result.
12. History of drug or alcohol abuse within the 12 months before dosing, or evidence of such abuse as indicated by the laboratory assays conducted during the screening.
13. Participants that took acetaminophen (paracetamol) chronically, that is, more than 1 g/day for more than 3 out of 7 days, or more than 2 g/day for more than 1 out of 7 days (added as part of Amendment 5).

Age at beginning of study per study group:

QAX576 (dectrekumab), 6 mg/kg: mean (SD), 30.7 (9.58)

Placebo: mean (SD), 29.5 (11.22)

Sex (m/f) per study group:

QAX576 (dectrekumab), 6 mg/kg: m/f n, 16/1

Placebo: m/f n, 8/0

Number randomized per study group:

QAX576 (dectrekumab), 6 mg/kg: 17

Placebo: 8

Number reaching end of study per study group:

Rothenberg 2015 (Continued)

QAX576 (dectrekumab), 6 mg/kg: 13

Placebo: 5

Interventions	Study group 1: placebo Study group 2: QAX576 (dectrekumab), 6 mg/kg	
Outcomes	Primary outcomes of the study: 1. The primary endpoint in this study is the number of patients (responder) with a reduction of 75% or more in eosinophils per HPF (distal or proximal esophagus) from baseline to week 13 Secondary outcomes of the study: 1. Efficacy: symptoms. Mayo Dysphagia Questionnaire (MDQ) with a 2-week recall period (Grudell AB, Alexander JA, Enders FB, Pacifico R, Fredericksen M, Wise JL, et al. Validation of the Mayo Dysphagia Questionnaire. <i>Dis Esophagus</i> 2007;20:202-5). 2. Efficacy: molecular. The EoE transcriptome before and after study medication (QAX576 and placebo) was compared by using the NanoString nCounter technology (NanoString Technologies, Seattle, Wash) for direct multiplexed measurement of gene expression. Circulating protein biomarkers and histochemistry. Quantitative determination of periostin was performed by means of liquid chromatography mass spectrometry. Histochemical staining for collagen, periostin, and eotaxin was performed.	
Notes	Funding source: Novartis Pharmaceuticals Conflicts of interest: M. E. Rothenberg has received consultancy fees from Immune Pharmaceuticals, Receptos, Pluristem Pharmaceuticals, Regeneron, and Novartis; has an equity interest in Immune Pharmaceuticals and Receptos and can receive royalty fees from Teva for reslizumab, which is under development; and is a coinventor on patent applications owned by Cincinnati Children's, concerning the eosinophilic esophagitis transcriptome. T. Wen is a coinventor of a patent application, owned by Cincinnati Children's Hospital, concerning the eosinophilic esophagitis transcriptome. B. Enav reports personal fees from QOL Medical outside the submitted work. I. Hirano reports personal fees from Novartis, Meritage, Aptalis, and Receptos outside the submitted work. S. Kaiser, T. Peters, I. Jones, J. P. Arm, and K. A. Gunawardena are employees of Novartis. R. Strieter is the Global Head for Translational Medicine for Respiratory disorders at Novartis Institutes of Biomedical Research and has stock equity in Novartis. R. Sabo and A. Perez are consultants for Novartis. The rest of the authors declare that they have no relevant conflicts of interest. Paper states "Sample management, sample analysis, and technical assistance were provided by Martin Letzkus, Urs Affentranger, Aurelie Seguin, Tiziana Valensise, Stephan Bek, Junli Yu, and Shenglin Ma of the Biomarker Development Group of Novartis Institutes for Biomedical Research. support was provided by Saurabh Aggarwal, Senior Scientific Writer, Medical Communications, Novartis", suggesting that author and editorial input came from an interested pharmaceutical company they were unable to provide how much input they had.	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The authors responded that "this study was sponsored by Novartis and we received a randomization ID via email. Presumably the allocations were generated using a random sequence generator". On balance, we have judged this likely to be random.
Allocation concealment (selection bias)	Low risk	The randomization was performed by using a "Request & Response" exchange system by an email between the study site's unblinded pharmacist and Novartis.

Medical treatment of eosinophilic esophagitis (Review)

Rothenberg 2015 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	Low risk	Patients, investigators, and study personnel were blinded to treatment assignment.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	A blinded central pathologist assessed the primary outcome. Blinded personnel assessed clinical symptoms. Molecular outcomes were analyzed by the Biomarker Development group at Novartis.
Incomplete outcome data (attrition bias) All outcomes	Low risk	Minimal and explained attrition that likely has no effect on outcome.
Selective reporting (reporting bias)	Low risk	Protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Rothenberg 2022
Study characteristics

Methods	<p>RCT design and number of study arms: part B of a 3-part, randomized, placebo-controlled phase 3 study, 2 arms</p> <p>Single-center or multi-center: multi-center; 95 study locations worldwide</p> <p>Countries: US, Australia, Belgium, Canada, France, Germany, Italy, Netherlands, Spain, Sweden, Switzerland, and the UK</p> <p>Study dates: 24 September 2018 to 9 September 2021</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: "A documented diagnosis of EoE by endoscopic biopsy."</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • A documented diagnosis of EoE by endoscopic biopsy • Baseline endoscopic biopsies with a demonstration on central reading of intraepithelial eosinophilic infiltration • History (by patient report) of an average of at least 2 episodes of dysphagia (with intake of solids) per week in the 4 weeks prior to screening <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Body weight \leq 40 kg • Prior participation in a dupilumab clinical trial, or past or current treatment with dupilumab • Initiation or change of a food-elimination diet regimen or re-introduction of a previously eliminated food group in the 6 weeks prior to screening • Other causes of esophageal eosinophilia or the following conditions: hypereosinophilic syndrome and eosinophilic granulomatosis with polyangiitis (Churg-Strauss syndrome) • Active <i>Helicobacter pylori</i> infection • History of achalasia, Crohn's disease, ulcerative colitis, celiac disease, and prior esophageal surgery • Any esophageal stricture unable to be passed with a standard, diagnostic, 9 mm to 10 mm upper endoscope or any critical esophageal stricture that requires dilation at screening • History of bleeding disorders or esophageal varices

Medical treatment of eosinophilic esophagitis (Review)

Rothenberg 2022 (Continued)

- Pregnant or breastfeeding women, or women planning to become pregnant or breastfeed during the study

Age at beginning of study per study group: NR, "adolescents and adults"

Sex (m/f) per study group: NR

Number randomized per study group:

Dupilumab: 80

Placebo: 79

Number reaching end of study per study group: NR

Interventions	<p>Study group 1: placebo, type not specified</p> <p>Study group 2: dupilumab 300 mg weekly</p>
Outcomes	<p>Primary outcomes of the study:</p> <ul style="list-style-type: none"> • Proportion of patients achieving a peak esophageal intraepithelial eosinophil (eos) count of < 6 eos/hpf • Absolute change from baseline in Dysphagia Symptom Questionnaire (DSQ) score <p>Secondary outcomes of the study:</p> <ul style="list-style-type: none"> • Safety
Notes	<p>Funding source:</p> <p>Research sponsored by Sanofi and Regeneron Pharmaceuticals, Inc.</p> <p>Conflicts of interest:</p> <p>Dr. Dellon - Consultant: Abbott, Adare, Aimmune, Allakos, Amgen, Arena, AstraZeneca, Biorasi, Calypso, Eli Lilly, EsoCap, Gossamer Bio, GlaxoSmithKline, Parexel, Receptos/Celgene/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Robarts, Salix, Shire/Takeda; research funding: Adare, Allakos, GlaxoSmithKline, Meritage, Miraca, Nutricia, Receptos/Celgene/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Shire/Takeda; educational grant: Allakos, Banner, Holoclara.</p> <p>Dr. Rothenberg - Consultant: Allakos, AstraZeneca, Bristol Myers Squibb, ClostraBio, Pulm One, Spoon Guru; equity interest: ClostraBio, Pulm One, Spoon Guru; royalties from reslizumab: Teva Pharmaceuticals; royalties from PEESv2: Mapi Research Trust; royalties: UpToDate; inventor of patents owned by Cincinnati Children's Hospital.</p> <p>Dr. Collins - consultant: Allakos, Arena, AstraZeneca, Bristol Myers Squibb, Calypso, Escap, GlaxoSmithKline, Regeneron Pharmaceuticals, Inc., Shire; research funding: Receptos/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Shire.</p> <p>Dr. Hirano - consultant: Adare, Receptos/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Shire; research funding: Meritage, Receptos/Bristol Myers Squibb, Regeneron Pharmaceuticals, Inc., Shire.</p> <p>Dr. Chehade - consultant: Adare, Allakos, AstraZeneca, Nutricia, Regeneron Pharmaceuticals, Inc., Shire; research funding: Allakos, Regeneron Pharmaceuticals Inc., Shire; honoraria for lectures: Medscape, Nutricia.</p> <p>Dr. Bredenoord - consultant: Arena, AstraZeneca, Calypso, EsoCap, Falk, Gossamer Bio, Medtronic, Laborie, RB, Regeneron, Robarts; research funding: Bayer, Nutricia, SST; equity interest: SST.</p> <p>Dr. Lucendo - Consultant: EsoCap, Dr. Falk Pharma; Research funding: Dr. Falk Pharma, Regeneron Pharmaceuticals.</p> <p>Dr. Spergel - Consultant: Regeneron, Shire, Takeda, Allakos, DBV Technology, Novartis; Grant Support: Regeneron, DBV Technology.</p> <p>Q Zhao, JD Hamilton, B Beazley, S Kamat, M Ruddy, B Akinlade, N Amin, A Radin, B Shumel, J Maloney: Regeneron Pharmaceuticals, Inc. - Employees and Shareholders.</p> <p>I Guillemin: Sanofi - Prior employee, may hold stock and/or stock options in the company.</p> <p>L Mannent, E Laws: Sanofi - Employees, may hold stock and/or stock options in the company.</p>

Rothenberg 2022 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	The author confirmed the randomization was computer-generated.
Allocation concealment (selection bias)	Low risk	The author confirmed that central allocation was completed by the sponsor.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	The author confirmed the study was appropriately blinded.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The author confirmed the pathologist and endoscopist were blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No patients were lost to follow-up. Imputation was used for continuous outcomes.
Selective reporting (reporting bias)	Unclear risk	Not all of the many registered outcomes have been reported as per the trial registration in this abstract publication.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Schaefer 2008
Study characteristics

Methods	<p>RCT design and number of study arms: RCT with 2 arms</p> <p>Single-center or multi-center: single-center</p> <p>Countries: United States</p> <p>Study dates: February 2000 to November 2004</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: esophageal mucosal biopsy specimens showing ≥ 15 eos/hpf with negative pH probe studies.</p> <p>Inclusion criteria:</p> <ol style="list-style-type: none"> Age range 1 to 18 years Active eosinophilic esophagitis with ≥ 15 eosinophils per hpf and negative pH probe study <p>Exclusion criteria:</p> <ol style="list-style-type: none"> Co-existing esophageal conditions (e.g. stricture, Barrett's esophagus, caustic injury), <i>Helicobacter pylori</i> infection, inflammatory bowel disease, and inability to tolerate corticosteroids <p>Age at beginning of study per study group:</p> <p>Prednisone: mean (SD), 7.0 (4.3)</p>

Medical treatment of eosinophilic esophagitis (Review)

Schaefer 2008 (Continued)

Fluticasone: mean (SD), 7.2 (4.1)

Sex (m/f) per study group:

Prednisone: m/f, 31/9

Fluticasone: m/f, 28/12

Number randomized per study group:

Prednisone: 40

Fluticasone: 40

Number reaching end of study per study group:

Prednisone: 32 through follow-up EGD after 4 weeks of full strength therapy; 27 through 8-week wean; 25 through week 18 follow-up; 17 through week 24 follow-up

Fluticasone: 36 through follow-up EGD after 4 weeks of full strength therapy; 27 through 8-week wean; 22 through week 18 follow-up; 19 through week 24 follow-up

Interventions	Study group 1: Prednisone: oral prednisone suspension/tablet (1 mg/kg/dose twice a day; maximum 30 mg twice a day) Study group 2: Fluticasone: swallowed fluticasone by metered dose inhaler (110 g per puff for ages 1 to 10 years and 220 g per puff for ages 11 years or older, 2 puffs 4 times/day)	
Outcomes	Primary outcomes of the study: 1. A histologic response by an improvement in biopsy grade after 4 weeks of corticosteroid therapy. Points were assigned based on (1) basal cell zone thickness as a percentage of the epithelial thickness, and (2) the maximum number of eos/hpf (at 400 power using an eyepiece grid covering an area of 0.4 mm ²). Points were summed and the totals were translated into histologic grades (normal, mild, moderate, and severe). Grades were assigned a numeric value for statistical analysis. Secondary outcomes of the study: 1. Clinical response to corticosteroids based on the presence or absence of the presenting symptom by patient/guardian report and by physician assessment at predetermined intervals. Reported symptoms included vomiting, abdominal pain, epigastric pain, heartburn, dysphagia, feeding problems, foreign body/food impaction, and weight loss.	
Notes	Funding source: Clarian Values Grant, Clarian Health Partners, Inc, Indianapolis, IN Conflicts of interest: NR	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Random number assignment was used to generate a concealed allocation schedule that was maintained by a research co-ordinator.
Allocation concealment (selection bias)	Low risk	Concealed allocation schedule that was maintained by a research co-ordinator.

Schaefer 2008 (Continued)

Blinding of participants and personnel (performance bias) All outcomes	High risk	This was an open-label trial.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	The authors confirmed in November 2022 that the outcome assessors were all blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No significant imbalance in withdrawals, which were all explained.
Selective reporting (reporting bias)	Low risk	Protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Spergel 2012
Study characteristics

Methods	<p>RCT design and number of study arms: RCT, 4 arms</p> <p>Single-center or multi-center: multi-center, 34 sites in the United States and 2 sites in Canada</p> <p>Countries: USA and Canada</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active</p> <p>EoE definition/diagnostic criteria: defined as ≥ 24 eosinophils in ≥ 1 high-power field (hpf)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> At least 1 active symptom (i.e. vomiting, regurgitation, abdominal pain, chest pain/heartburn, or dysphagia) of moderate severity or worse (as assessed by the patient) within the week before randomization An OGD with biopsy documenting active eosinophilic esophagitis (defined as > 24 eosinophils in > 1 high-power field (hpf)) A history of eosinophilic esophagitis symptoms, and treatment with proton pump inhibitors with or without histamine H2 receptor antagonists for at least 4 weeks without symptom resolution or a normal pH probe (regardless of whether the patient had undergone a failed course of proton pump inhibitors) <p>Exclusion criteria:</p> <ul style="list-style-type: none"> If they had another disorder that could cause esophageal eosinophilia (e.g. hypereosinophilic syndrome, Churg-Strauss vasculitis, eosinophilic gastroenteritis, or parasitic infection) Had a history of abnormal gastric or duodenal biopsy results or documented gastrointestinal disorders (e.g. celiac disease, Crohn disease, recurrent acute or chronic diarrhea, ulcerative colitis, malrotation, or active <i>Helicobacter pylori</i> infection) Had a history of gastrointestinal procedures (i.e. esophageal surgery (not including esophageal dilation), fundoplication, gastric surgery, or surgery for intestinal atresia) Had used systemic immunosuppressive or immunomodulating agents (e.g. oral corticosteroids, antibodies to IgE, methotrexate, cyclosporine, IFN-α, or TNF-α inhibitors) within 6 months before study entry Had received live attenuated vaccines within 3 months before study entry

Medical treatment of eosinophilic esophagitis (Review)

Spergel 2012 (Continued)

- Had swallowed corticosteroids formulated for inhalation for the treatment of eosinophilic esophagitis within 1 month before study entry
- Had a stricture that would have prevented the passage of the endoscope
- Had an infection that might have interfered with study assessments; or had a concurrent immunodeficiency
- Patients were also excluded if they had previously participated in any investigational drug or device study within 30 days before study entry or any investigational study of a biologic therapy within 3 months before study entry or received mepolizumab within 4 months or reslizumab any time before study entry

Age at beginning of study per study group:

1 mg/kg reslizumab: 12.3 (3.83)

2 mg/kg reslizumab: 11.8 (3.82)

3 mg/kg reslizumab: 11.5 (4.04)

Placebo group: 11.9 (4.17)

Sex (m/f) per study group: NR

Number randomized per study group:

1 mg/kg reslizumab = 56

2 mg/kg reslizumab = 57

3 mg/kg reslizumab = 57

Placebo = 57

Number reaching end of study per study group:

1 mg/kg reslizumab = 48

2 mg/kg reslizumab = 47

3 mg/kg reslizumab = 50

Placebo = 51

Interventions
Placebo group: placebo/saline infusion only

Intervention group 2, 3, 4:

Intervention group - arm 1: 1 mg/kg reslizumab + saline infusion

Intervention group - arm 2: 2 mg/kg reslizumab + saline infusion

Intervention group - arm 3: 3 mg/kg reslizumab + saline infusion

Outcomes
Primary outcomes of the study:

1. Co-primary efficacy measures were the percentage change from baseline to the end of therapy (or early withdrawal) in the peak esophageal eosinophil count
2. The change from baseline to the end of therapy (or early withdrawal) in the physician's eosinophilic esophagitis global assessment score

Secondary outcomes of the study: no secondary outcome in FT - secondary outcome in protocol as follows:

1. Mean change from baseline to end of treatment in eosinophilic esophagitis predominant symptom assessment (time frame: baseline (day 1, pre-treatment), end of treatment (week 15, 3 weeks (\pm 4 days) after the last dose of study drug, or at early withdrawal))

Spergel 2012 (Continued)

2. Mean percent change from baseline to end of treatment in the Child Health Questionnaire (CHQ) (time frame: baseline, end of treatment (up to 15 weeks \pm 4 days))

Notes

Funding: Sponsored by Ception Therapeutics, Inc, which has since been acquired by Cephalon, Inc.

Conflict of interest: Sponsored by Ception Therapeutics, Inc, which has since been acquired by Cephalon, Inc. Disclosure of potential conflict of interest: J. M. Spergel is a consultant for DBV; has received research support from the Department of Defense (DOD), Cephalon, and the National Institutes of Health (NIH); is a member of the American Academy of Allergy, Asthma & Immunology; and is on the American Partnership for Eosinophilic Disorders (APFED) Medical Advisory Board. M. E. Rothenberg has equity interest in reslizumab through Cephalon; is consultant and chief scientific officer of Immune Pharmaceuticals; has received research support from the NIH, the Food Allergy & Anaphylaxis Network, and the DOD; is on the APFED Medical Advisory Board; and is on the International Eosinophil Society Executive Council. M. H. Collins is a central review pathologist for Cephalon, GlaxoSmithKline, and Meritage Pharma; is a consultant for Sunovion; and is president of the APFED Medical Advisory Board. G. T. Furuta is a consultant for Nutricia and Meritage and has received research support from the NIH, AstraZeneca, and the Thrasher Foundation. G. Fuchs III has received research support from Shire and Cephalon. J. P. Abonia has received research support from the NIH, Ception Therapeutics, and the Children's Digestive Health and Nutrition Foundation. T. Henkel is a consultant for Cephalon and a shareholder in Ception Therapeutics. C. A. Liacouras is a speaker for Nutricia and is on the American Partnership for Eosinophilic Disorders Physician Board. The rest of the authors declare that they have no relevant conflicts of interest. Received for publication 21 September 2011.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Central allocation was confirmed by the author.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Patients, investigators, and study personnel were blinded to treatment assignment throughout the study. The study site's pharmacist was unblinded and was responsible for preparing and dispensing study medication.
Blinding of outcome assessment (detection bias) All outcomes	Low risk	All staff were blinded except the study site's pharmacist was unblinded and was responsible for preparing and dispensing study medication.
Incomplete outcome data (attrition bias) All outcomes	Low risk	1 patient was discontinued due to an adverse event. The adverse event was not reported in the paper. The author responded that "for the one discontinuation due to mild abdominal pain and upper respiratory tract congestion. It was thought be unrelated to study medication". We thought this was unlikely to have influenced our outcomes of interest.
Selective reporting (reporting bias)	Low risk	Trial protocol is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No information on participant sex.

Spergel 2020

Study characteristics

Methods	<p>RCT design and number of study arms: parallel, double-blind RCT, 2 arms</p> <p>Single-center or multi-center: single-center</p> <p>Countries: USA</p> <p>Study dates: screening 11 November 2015 to 20 December 2016. After enrollment, 9 months of treatment (milk out of the diet) and then 2 months with milk in the diet before another scope. After that scope 11 months open-label extension.</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: inactive EoE at beginning of study. Patients had to have < 10 eos/hpf after milk-free diet for 2 months to be eligible to participate.</p> <p>EoE definition/diagnostic criteria: the diagnosis of EoE was confirmed with an esophagogastroduodenoscopy (EGD) and biopsy showing 15 eos/HPF after at least a 2-month period of high-dose proton pump inhibitor (PPI) (1 to 2 mg/kg dose twice-daily)</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • EGD on a milk-containing diet showed 15 eos/HPF • EGD with biopsy on milk-free diet for 2 months showed < 10 eos/HPF • Age 4 to 17 years <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • IgE-mediated milk allergy (positive skin test or specific IgE to milk using standard techniques) • Were pregnant or breastfeeding • Were currently taking systemic corticosteroids, tricyclic antidepressants, beta-blockers, or other medications for EoE outside of a PPI • Patients with a history consistent with poorly controlled persistent asthma; comorbid cardiac, autoimmune, infectious gastrointestinal (GI) or pulmonary conditions; or diagnosis of eosinophilic colitis or gastritis <p>Age at beginning of study per study group:</p> <p>For patients enrolled in the study there were 5 in the placebo group (median age 12.54 (range 9.55 to 14.55)) and 15 in the treatment group (median age 10.83 (range 5.86 to 15.37))</p> <p>Sex (m/f) per study group:</p> <p>Study group 1 (placebo) included 5 males and 0 females and Study group 2 (treatment) included 10 males and 5 females</p> <p>Number randomized per study group:</p> <p>N = 5 for Study group 1 (placebo) and N = 15 for Study group 2 (treatment)</p> <p>Number reaching end of study per study group:</p> <p>2 reached the end of the study (were analyzed) for Study group 1 and 7 reached the end of the study for Study group 2</p>
Interventions	<p>Study group 1: Viaskin placebo participants epicutaneously administered daily (up to 24 hours application per day) with a patch containing a matching placebo formulation</p> <p>Study group 2: Viaskin milk 500 µg participants epicutaneously administered daily (up to 24 hours application per day) with a patch containing 500 µg cow's milk proteins</p> <p>For both groups the same dose of PPI was continued (as well as the same dose of medications for asthma and allergic rhinitis)</p>

Spergel 2020 (Continued)

Outcomes	<p>Primary outcomes of the study: the primary efficacy endpoint is each patient's maximum esophageal eosinophil count on all specimens obtained from the biopsy at the end of double-blind treatment, after milk reintroduction</p> <p>Secondary outcomes of the study: 3 different symptom assessments (PEESS-parent, PEESS-patients and investigator assessment of symptoms; EREFS score)</p>
----------	---

Notes	<p>Funding source: DBV technologies and the Children's Hospital of Philadelphia Eosinophilic Esophagitis Family Fund</p> <p>Conflicts of interest: consulting agreements and clinical trial grants with DBV Technologies and the first author has stock equity with DBV Technologies</p>
-------	--

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	"The randomization list and treatment allocation were done and computer generated by an independent party, eXSTAT (Paris, France), and they had no other role in the study."
Allocation concealment (selection bias)	Low risk	"The randomization list and treatment allocation were done and computer generated by an independent party, eXSTAT (Paris, France), and they had no other role in the study."
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Individuals providing care, assessing outcomes, and participants were masked to group assignment." "Masking was done with identical-looking Viaskin patches"
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Individuals providing care, assessing outcomes, and participants were masked to group assignment." "Masking was done with identical-looking Viaskin patches"
Incomplete outcome data (attrition bias) All outcomes	Low risk	No attrition imbalances, or imbalances in reasons for attrition, based on the flow diagram of the study.
Selective reporting (reporting bias)	Low risk	Outcomes reported as per published protocol.
Other bias	Low risk	Low imbalance. No female patients in the placebo group compared to 5 (33%) of the Viaskin milk group.

Straumann 2010a
Study characteristics

Methods	<p>RCT design and a number of study arms: double-blind RCT; 2 arms</p> <p>Single-center or multi-center: single-center (Olten, Switzerland)</p> <p>Countries: Switzerland</p> <p>Dates: December 2005 and May 2006</p>
---------	--

Participants	Active EoE or inactive EoE at beginning of the study: active
--------------	---

Medical treatment of eosinophilic esophagitis (Review)

Straumann 2010a (Continued)

EoE definition/diagnostic criteria: EoE with a history of at least one episode of dysphagia per week in the 4 weeks prior to the start of study medication and a peak esophageal eosinophilia of > 20 eosinophils per hpf (peak eosinophil density)

Inclusion criteria:

- Document evidence/presence of esophagitis prior to commencing trial drug
- Histological evidence of esophagitis: greater than 20x eosinophils per high-power field (x400) on histology of esophageal mucosal biopsy
- At least one episode of dysphagia per week
- Inadequate response to routine eosinophilic esophagitis treatment (topical or systemic steroids)
- No other known causes of esophagitis, or esophageal or generalized eosinophilia
- Not pregnant or nursing

Exclusion criteria:

- History of seasonal worsening of eosinophilic esophagitis symptoms or requirement of esophageal dilation
- Churg-Strauss syndrome
- Wegener's granulomatosis
- Lymphoma, hematological malignancy, advanced and metastatic solid tumors
- Active *H. pylori* infection
- Any previous treatment with anti-IL-5, anti-IgE monoclonal antibody, or other biological agents
- Other causes of esophagitis (hypereosinophilic syndromes, eosinophilic gastroenteritis, and parasitic infection). GERD was excluded in all patients by pretreatment with PPIs in standard dosages plus negative endoscopy for signs of reflux disease, and by pH monitoring (optional)
- Any condition with the risk of requiring esophageal dilatation during the course of the study
- A history of seasonal exacerbation of EoE symptoms is expected to coincide with the period of investigation
- Active *Helicobacter pylori* infection and any unstable medical conditions
- Patients using mast cell stabilizers, leukotriene receptor antagonists, or immunosuppressive/immunomodulatory agents, and those with a history of allergic reactions to previous antibody treatment
- Any previous treatment with an anti-IL-5 antibody or any other biopharmaceutical agent
- Female patients were excluded if pregnant or breastfeeding, or if they were not taking adequate contraceptive measures

Age at beginning of study per study group: intervention = 32.4; placebo = 34.0 (mean, no SD reported)

Sex (m/f) per study group: M/F = 4/1; M/F = 3/3

Number randomized per study group: intervention = 5; placebo n = 6

Number reaching the end of study per study group: intervention = 5; placebo n = 6

Interventions	<p>Study group 1: placebo</p> <p>Study group 2: mepolizumab (GlaxoSmithKline, Greenford, UK) was administered by intravenous infusion at a dose of 750 mg diluted in 150 mL of 0.9% sodium chloride solution 2 doses day 0 and day 7</p>
Outcomes	<p>Primary outcomes of the study: to reduce peak esophageal eosinophilia to ≤ 5 eos/hpf</p> <p>Secondary outcomes of the study: effect of treatment on symptoms, eosinophil levels, and inflammation biomarkers in esophagus tissue and blood</p>
Notes	<p>Funding: This study was supported by GlaxoSmithKline (GSK), Greenford, UK. The trial was conducted under GSK protocol number MEE103226.</p> <p>Conflicts of interest: declared here.</p>

Straumann 2010a (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Central allocation via a telephonic randomization system. "To keep the treatment blinded, the infusions were made up by an independent pharmacist who obtained the treatment allocation via a telephonic randomisation system"
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"Of the study personnel, only the pharmacist, responsible for the preparation of infusions, had access to the treatment assignments." "Patients allocated to the placebo arm received the corresponding infusions of saline only. To keep the treatment blinded, the infusions were made up by an independent pharmacist".
Blinding of outcome assessment (detection bias) All outcomes	Low risk	"Of the study personnel, only the pharmacist, responsible for the preparation of infusions, had access to the treatment assignments".
Incomplete outcome data (attrition bias) All outcomes	Low risk	There were no dropouts in any group.
Selective reporting (reporting bias)	Low risk	Outcomes were pre-specified and were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Straumann 2010b
Study characteristics

Methods	<p>RCT design: randomized, placebo-controlled, parallel-group clinical trial</p> <p>Single-center or multi-center: single-center</p> <p>Countries: Switzerland</p> <p>Study dates: December 2005 to December 2008</p>
Participants	<p>Active EoE or inactive EoE: active eosinophilic esophagitis</p> <p>EoE definition/diagnostic criteria: clinicopathologic definition of esophageal symptoms in combination with a dense esophageal eosinophilia, both being refractory to proton pump inhibition</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Patients older than 14 years old • Isolated esophageal eosinophilia • Dysphagia almost always occurring with intake of solids when off anti-inflammatory therapy or dietary restriction • Eosinophilic tissue infiltration (mean cell density of ≥ 20 eosinophils per high-power field) on esophageal histology biopsy specimens

Medical treatment of eosinophilic esophagitis (Review)

Straumann 2010b (Continued)

- Exclusion of other causes of esophageal or systemic eosinophilia
- Exclusion of gastroesophageal reflux disease by proton pump inhibitor pretreatment plus negative endoscopy for signs of reflux disease and pH monitoring (optional)

Exclusion criteria:

- Current use of specific treatments for eosinophilic esophagitis
- Secondary causes of esophageal eosinophilia
- Intolerance to budesonide
- Concomitant therapies for any reason that may affect the assessment
- Use of an investigational drug within 30 days of entering the study
- Recent history of suspicion of current drug abuse and alcohol abuse
- A positive serum pregnancy test at the screening visit
- Any unstable serious co-existing medical condition

Age:

- Budesonide – mean (SD): 33.1 ± 13.1
- Placebo – mean (SD): 38.2 ± 12.4

Sex:

- Budesonide – 18 total patients: (m/f): 17/1
- Placebo – 18 total patients: (m/f): 14/4

Number randomized:

- Budesonide n = 18
- Placebo n = 18

Number reaching the end:

- Budesonide: 18/18 (100%)
- Placebo: 18/18 (100%)

Interventions

Study group 1: placebo

- 0.9% saline
- Administered via nebulizer
- 4 mL twice per day at bedtime and in the morning after breakfast
- Administered into the oropharynx, patient instructed to swallow continuously the accumulated liquid
- Nothing to eat or drink for 30 minutes after administration

Study group 2: budesonide suspension 2.0 mg

- 0.25 mg/mL suspension
- Administered via nebulizer
- 4 mL twice per day at bedtime and in the morning after breakfast
- Administered into the oropharynx, patient instructed to swallow continuously the accumulated liquid
- Nothing to eat or drink for 30 minutes after administration

Outcomes

Outcomes assessed on day 15

Primary outcomes:

- Reduction in the esophageal eosinophil load achieved with budesonide
 - Eosinophil load – mean eosinophil number measured in a total of 40 higher-power fields taken from 4 biopsy specimens each of the proximal and mid esophagus
- Also compared post-treatment mean eosinophils per high-power field < 5 cells, 5 to 20 cells, > 20 cells

Straumann 2010b (Continued)

Secondary outcomes: the effects of budesonide on reducing eosinophilic esophagitis-associated symptoms and on eosinophilic esophagitis relevant biomarkers in the esophagus and peripheral blood

Notes

Funding source:

- Swiss National Science Foundation
- AstraZeneca, Switzerland

Other: included patients who completed a 4-week run-in period after stopping eosinophilic esophagitis-relevant therapies (steroids, leukotriene antagonists, histamine blockers, mast cell stabilizers) besides proton pump inhibition

Conflicts of interest: Dr. Christoph Beglinger and Dr. Simon received research support from AstraZeneca

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated, coded randomization method.
Allocation concealment (selection bias)	Unclear risk	How allocation was concealed is not addressed in the manuscript or trial registry.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Patients, study center personnel, laboratory personnel, and the sponsor were blinded to treatment allocation.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Neither the manuscript nor the trial registry specifically address whether the interpreting pathologist(s) were blinded to treatment arm.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All randomized patients completed the study and were included in the analysis.
Selective reporting (reporting bias)	Low risk	All pre-specified outcomes including histologic, endoscopic, and symptoms were reported.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Straumann 2011
Study characteristics

Methods

RCT design: randomized, placebo-controlled, single-center clinical trial

Single-center or multi-center: single-center

Countries: Switzerland

Study dates: December 2005 to December 2008

Participants

Active EoE or inactive EoE at beginning of study: inactive eosinophilic esophagitis at beginning of the study

Medical treatment of eosinophilic esophagitis (Review)

Straumann 2011 (Continued)

EoE definition/diagnostic criteria: clinically, endoscopically, and histologically confirmed eosinophilic esophagitis after proton pump inhibitor trial

Inclusion criteria:

- Patients older than 14 years with clinically, endoscopically, and histologically confirmed eosinophilic esophagitis, previously brought successfully into remission with short-term (= 15 days), high-dose (= 2 mg/day) budesonide were eligible
- Remission was defined as follows: (1) a mean cell density for eosinophilic tissue infiltration of fewer than 5 eosinophils/high-power field on esophageal histology biopsies, and (2) symptom score of 2 points or fewer

Exclusion criteria:

- Current use of specific treatments for eosinophilic esophagitis
- Secondary causes of esophageal eosinophilia
- Intolerance to budesonide
- Concomitant therapies for any reason that may affect assessment
- Use of an investigational drug with 30 days of entering the study
- Recent history or suspicion of current drug abuse and alcohol abuse
- Positive serum pregnancy test at the screening visit
- Any unstable serious co-existing medical condition

Age:

- Budesonide: mean (SD): 38.0 ± 11.7
- Placebo: mean (SD): 34.0 ± 13.9

Sex:

- Budesonide: m/f: 13/1
- Placebo: m/f: 11/3

Number randomized: 14 intervention; 14 placebo

Number reaching end of study

- Budesonide: 9/14
- Placebo: 5/14

Interventions

Study group 1: placebo: 0.9% saline 1 mL via an inhalation system consisting of a PARI UNI light compressor and PARI TIA nebulizer, twice per day at bedtime and in the morning after breakfast; patients instructed to nebulize the suspension into the oral cavity and to swallow continuously the accumulated liquid

Study group 2: treatment arm: 0.5 mg/day budesonide as 0.25 mg/mL suspension formulation applied using an inhalation system consisting of a PARI UNI light compressor and PARI TIA nebulizer, twice per day at bedtime and in the morning after breakfast; patients instructed to nebulize the suspension into the oral cavity and to swallow continuously the accumulated liquid

Outcomes

Primary outcomes:

The primary endpoint was to determine the ability of a long-term budesonide therapy in maintaining eosinophilic esophagitis in histologic remission, defined as an esophageal eosinophil load of fewer than 5 eosinophils per high-power field. Eosinophil load is defined as the mean eosinophil number measured in a total of 40 high-power fields from 2 x 4 biopsy specimens each, taken from the proximal and distal esophagus

Secondary outcomes:

- To determine the course of the disease under placebo (natural course) after successful short-term therapy

Straumann 2011 (Continued)

- To assess the ability of budesonide in controlling eosinophilic esophagitis symptoms
- To determine the effect of long-term corticosteroid therapy on markers of inflammation and tissue damage, as well as on signs of esophageal remodeling
- To evaluate the safety of long-term topical esophageal corticosteroid administration

Notes

Funding source: supported by grants from the Swiss National Science Foundation

AstraZeneca

Conflicts of interest: Christoph Beglinger and Hans-Uew Simon received research support for the clinical trial from AstraZeneca

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Unclear risk	The method of allocation concealment is not fully explained.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Patients, study center, laboratory staff, and sponsor were blinded to treatment allocation.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	Patients, study center, laboratory staff, and sponsor were blinded to treatment allocation. The manuscript does not specifically address whether the interpreting pathologist was blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	There were data on all patients randomized at either study end or following clinical relapse.
Selective reporting (reporting bias)	Low risk	Per NCT00271349 and the manuscript, all pre-specified outcomes were reported.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Straumann 2013
Study characteristics

Methods

RCT design and number of study arms: randomized, placebo-controlled, parallel-group, 2 arms

Single-center or multi-center: single-center

Countries: Switzerland

Study dates: August 2010 to June 2011

Participants

Active EoE of inactive EoE at beginning of study: active EoE

EoE definition/diagnostic criteria: patient aged 18 to 75 with previously clinically, endoscopically, and histologically confirmed EoE (according to Liacouras 2011 definition)

Age at beginning of study per group:

Medical treatment of eosinophilic esophagitis (Review)

Straumann 2013 (Continued)

Placebo: 38.83 ± 14.48 years

OC000459 (timapiprant): 43.71 ± 13.49 years

Sex (m/f) per study group:

Placebo: 8/4

OC000459 (timapiprant): 14/0

Number randomized per study group:

Placebo: 12

OC000459 (timapiprant): 14

Number reaching end of study per study group:

Placebo: 12

OC000459 (timapiprant): 14

Inclusion criteria:

1. Aged 18 to 75
2. Previously diagnosed and symptomatic isolated EoE
3. Able to swallow placebo medication successfully under supervision in the clinic
4. Free of all medications for EoE (including topical steroids) for at least 2 weeks prior to baseline and free of systemic steroids for at least 90 days before screening. A PPI is allowed if required for treatment of secondary reflux.
5. Inadequate response to topical and/or systemic corticosteroid therapy (e.g. corticosteroid refractory, corticosteroid dependency or necessity of high doses to control the inflammation)
6. Dysphagia with almost every intake of solids
7. Relevant eosinophilic infiltration of the epithelial layer of the esophagus (mean cell density of ≥ 20 eos/hpf in 8 biopsies)
8. Exclusion of other causes of esophageal or systemic eosinophilia
9. Gastroesophageal reflux disease was excluded by proton pump inhibitor (PPI) pretreatment plus negative endoscopy for signs of reflux disease and pH monitoring (optional)

Exclusion criteria:

1. Other causes of esophagitis (GERD, peptic ulceration, infection, etc.)
2. Other causes of esophageal or generalized eosinophilia (i.e. hypereosinophilic syndromes, parasitic infection, GERD)
3. The patient's EoE is dependent on the level of seasonal allergies and the patient's participation in the study will occur during the allergy season
4. History of abnormal gastric or duodenal eosinophilia (e.g. HES, Churg Strauss vasculitis, EG or a parasitic infection)
5. Receipt of forbidden prescribed or over-the-counter medication with the 4 weeks prior to the baseline visit and for the duration of the trial, including vitamins and herbal remedies

Interventions

Study group 1: placebo; identical-appearing placebo tablets

Study group 2: OC000459 (timapiprant) monotherapy; 100 mg tablets, twice-daily after meals for 8 weeks

Outcomes

Primary outcomes of the study:

Reduction in the esophageal eosinophil load, defined as the mean eosinophil number measured in a total of 40 hpf from 2 x 4 biopsies taken from the proximal and distal esophagus

Secondary outcomes:

Medical treatment of eosinophilic esophagitis (Review)

Straumann 2013 (Continued)

The effects of OC000459 (timapiprant) on patient-reported outcomes (PROs), on endoscopic alterations, and on EoE-relevant biomarkers in the esophagus and peripheral blood

Safety and tolerability

Notes

Funding source:

Swiss National Science Foundation

Oxagen LTD

Conflicts of interest:

Alex Straumann and Christian Bussmann received research support for the clinical trial from the sponsor. Mike Perkins, Lisa Pearce Collins, Roy Pettipher, Michael Hunter, and Jan Steiner are employed by Oxagen Ltd.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Unclear risk	Allocation concealment was not explicitly addressed in the manuscript or registry.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	Patients, study center personnel, and the sponsor were blinded.
Blinding of outcome assessment (detection bias) All outcomes	Unclear risk	It is not explicitly mentioned that the interpreting pathologist was blinded to allocation.
Incomplete outcome data (attrition bias) All outcomes	Low risk	All randomized patients completed the study and were included in the analysis.
Selective reporting (reporting bias)	Low risk	Trial registry is available and all pre-specified outcomes of interest were reported in the pre-specified way.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

Straumann 2020
Study characteristics

Methods

RCT design and number of study arms: phase 3, randomized, double-blind, placebo-controlled trial

Single-center or multi-center: multi-center

Countries: multiple

Study dates: 4 August 2016 to 4 October 2016

Participants

Active EoE or inactive EoE at beginning of study? inactive

Medical treatment of eosinophilic esophagitis (Review)

Straumann 2020 (Continued)

EoE definition/diagnostic criteria: previously confirmed diagnosis of PPI-refractory EoE according to consensus guidelines (Dellon et al. Gastroenterology 2018; Lucendo AJ et al. United European Gastroenterol J. 2017)

Inclusion criteria:

- Confirmed clinico-histologic remission at baseline after achieving study goals of a double-blind, controlled induction treatment study (EOS-1) with BOT 1.0 mg twice-daily, or be receiving open-label induction with budesonide orodispersible tablet (BOT) 1.0 mg twice-daily for 6 weeks
- Clinical remission defined as a severity of ≤ 2 points on 1- to 10-point numerical rating scale (NRS) for dysphagia and a severity of ≤ 2 points on a 0- to 10-point NRS for odynophagia on each day in the last week of induction treatment
- Histologic remission defined as peak eosinophil count < 16 eos/mm² hpf (corresponding to < 5 eos/hpf via prior method) at baseline, measured in hpf derived from 6 biopsies, 2 of each esophageal third

Exclusion criteria:

- Clinical and endoscopic suspicion for gastroesophageal reflux disease; achalasia or scleroderma; evidence of reasons for esophageal eosinophilia other than EoE; pathologic eosinophilic infiltration in gastric and duodenal biopsies
- History of esophageal surgery at any time or of esophageal dilation procedures within the last 8 weeks before induction treatment
- Any relevant systemic disease

Age at beginning of study per study group:

- Placebo: 36 ± 9.9
- BOT 0.5 mg twice-daily: 36 ± 10.9
- BOT 1.0 mg twice-daily: 37 ± 11.1

Sex (m/f) per study group:

- Placebo: 55/13
- BOT 0.5 mg twice-daily: 57/11
- BOT 1.0 mg twice-daily: 57/11

Number randomized per study group:

- Placebo: 68/204
- BOT 0.5 mg twice-daily: 68/204
- BOT 1.0 mg twice-daily: 68/204

Number reaching end of study per study group:

- Placebo: 23/68
- BOT 0.5 mg: 59/68
- BOT 1.0 mg: 59/68

Interventions

Study group 1: placebo

Study group 2: BOT 0.5 mg twice-daily

Study group 3: BOT 1.0 mg twice-daily

Outcomes

Primary:

Remission at week 48

Secondary:

- Rate of histologic relapse (defined in primary outcome) at 48 weeks

Straumann 2020 (Continued)

- Change in the peak eos/mm² hpf from baseline to EoT
- Rate of clinical relapse (defined in primary outcome)
- Food impaction requiring endoscopic intervention or endoscopic dilation
- Rate of clinical remission (eosinophilic esophagitis activity index-patient reported outcome score ≤ 20)

Notes

Funding source: Dr Falk Pharma GmbH, Freiburg, Germany

Conflicts of interest: Alex Straumann reports receiving consulting fees from Allakos, AstraZeneca, EsoCap, Dr Falk Pharma, Gossamer, GSK, Receptos-Celgene, and Regeneron-Sanofi; receiving lecture fees from Dr Falk Pharma and Vifor; receiving payment from Dr Falk Pharma for the development of educational presentations; receiving payment from AstraZeneca for serving as member independent data monitor committee; and serving as a board member for European Society of Eosinophilic Oesophagitis (EUREOS) and The International Gastrointestinal Eosinophil Researchers (TIGERS). Alfredo Lucendo reports receiving consulting fees from EsoCap, and Dr Falk Pharma; receiving lecture fees from Dr Falk Pharma; and serving as a board member for EUREOS. Stephan Miehle reports receiving consulting fees from Celgene, Dr Falk Pharma, and EsoCap; receiving lecture fees from Dr Falk Pharma and Vifor; receiving payment for the development of educational presentations from Dr Falk Pharma; and serving as a board member for EUREOS; Michael Vieth reports receiving lecture fees from Dr Falk Pharma, Janssen-Cilag, Malesci, Menarini, Olympus, and Shire. Christoph Schlag reports receiving consulting fees from Adare, Celgene, EsoCap, and Dr Falk Pharma; receiving lecture fees from Dr Falk Pharma; and serving as a board member for EUREOS. Luc Biedermann reports receiving consulting fees from Calypso Biotech SA, Switzerland; Esocap AG, Switzerland; Vifor AG, Switzerland; receiving lecture fees from Dr Falk Pharma, Germany; Sanofi-Aventis AG, Switzerland; and serving as a board member for EUREOS. Cecilio Santander Vaquero reports receiving lecture fees from Allergan and receiving payment for the development of educational presentations from Laborie. Constanza Ciriza de los Rios reports receiving consulting and/or lecture fees from Allergan and Casen Recordati. Ahmed Madisch reports receiving lecture fees from Dr Falk Pharma. Jamal Hayat reports receiving consulting fees from Dr Falk Pharma; and receiving lecture fees from Dr Falk Pharma. Ulrike von Arnim reports receiving consulting fees from Abbvie, Amgen, Eso Cap, Janssen, MSD, and Takeda; receiving lecture fees from Abbvie, Falk Foundation, Janssen, MSD, Reckitt Benckiser, Takeda, and Vifor; and serving as a board member for EUREOS. Albert Jan Bredenoord reports receiving research funding from Nutricia, Norgine, SideSleep Technologies, and Bayer; receiving lecture and/or consulting fees from Laborie, Arena, EsoCap, Diversatek, Medtronic, Dr Falk Pharma, Calypso Biotech, Thelial, Robarts, Reckitt Benckiser, Regeneron, Celgene, Bayer, Norgine, AstraZeneca, Almirall, Arena, and Allergan. Stefan Schubert reports receiving consulting fees from Abbvie, Takeda, Biogen, Amgen, and Janssen; receiving lecture fees from Abbvie, Dr. Falk Pharma, Takeda, Biogen, Amgen and Janssen; and serving as a board member for MSD, Takeda, and Janssen. Ralph Mueller reports being an employee of Dr Falk Pharma GmbH. Roland Greinwald reports being an employee of Dr Falk Pharma GmbH. Alain Schoepfer reports receiving consulting fees from Abbvie, Adare, Celgene, Dr Falk Pharma, Janssen-Cilag, MSD, Pfizer, Receptos, Regeneron, and Vifor; receiving lecture fees from Abbvie, Celgene, Dr Falk Pharma, Pfizer, Receptos, Regeneron, and Vifor; and serving as a board member for TIGERS. Stephen Attwood reports receiving consulting fees from Dr Falk Pharma, EsoCap, AstraZeneca, and Reckitt Benckiser; receiving lecture fees from Dr Falk Pharma, Medtronic; receiving payment for the development of educational presentations from Dr Falk Pharma.

The remaining authors disclose no conflicts.

Risk of bias

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Interactive Web Response System using randomly generated sequence. Patients received identical-appearing medications.
Blinding of participants and personnel (performance bias)	Low risk	Patients, investigators, and their study team, the sponsor, monitoring staff, central laboratory, and central pathologist were all kept blinded.

Medical treatment of eosinophilic esophagitis (Review)

166

Straumann 2020 (Continued)

All outcomes

Blinding of outcome assessment (detection bias) All outcomes	Low risk	Patients, investigators, and their study team, the sponsor, monitoring staff, central laboratory, and central pathologist were all kept blinded.
Incomplete outcome data (attrition bias) All outcomes	Low risk	There was comparable attrition between the treatment arms with expected attrition in the placebo arm (per design, suspect clinical). Suspected clinical failures were prematurely withdrawn from the double-blind phase of the study; exactly how this was handled statistically somewhat unclear. All attrition explained in supplementary material.
Selective reporting (reporting bias)	Low risk	All outcomes of interest were reported and qualified as a priori.
Other bias	Low risk	There were no consistent imbalances between treatment arms, though this was subjective and P values were not provided.

Tytor 2021
Study characteristics

Methods	<p>RCT design and number of study arms: RCT; 2 arms</p> <p>Single-center or multi-center: multi-center (2 centers: NU-Hospital Group, Trollhattan, and The Central Hospital, Skovde in Sweden)</p> <p>Countries: Sweden</p> <p>Study dates: April 2012 to August 2018</p>
Participants	<p>Active EoE or inactive EoE at beginning of study: active EoE</p> <p>EoE definition/diagnostic criteria: at least 15 eosinophils per high-power field (magnification 10 times 40 = x400) in any field of view in any esophageal biopsy and concurrent symptoms of esophageal dysfunction, mainly dysphagia</p> <p>Inclusion criteria:</p> <ul style="list-style-type: none"> • Age 18 years • Newly diagnosed EoE with a peak eosinophil count of at least 15 cells per HPF in any area in any of at least 6 esophageal biopsies including at least 3 biopsies from the upper- respective lower-third part of the esophagus • Total WDS score 5 <p>Exclusion criteria:</p> <ul style="list-style-type: none"> • Ongoing infection – locally or general • Glaucoma • Planned elective surgery during the treatment period • Systemic or topical corticosteroid treatment during the last 4 months • Pregnancy, ongoing or planned. Other known cause of dysphagia. • Other treatment that might affect dysphagia or motility during the trial (e.g. cisapride, erythromycin) • Allergy or intolerance to any component in the drug • Contraindication to steroid treatment (immune deficiency or immune suppression, gastroduodenal ulcer, diabetes mellitus) • PPIs were not allowed from 2 weeks before the start and during the treatment period

Medical treatment of eosinophilic esophagitis (Review)

Tytor 2021 (Continued)

- Other cause of dysphagia (cancer, autoimmune disease, neurologic disease)
- Inability to understand and provide autonomous informed consent

Age at beginning of study per study group:

Median (IQR)

- Placebo = 49.0 (37.0 to 63.0)
- Mometasone furoate = 42.0 (34.0 to 50.0)

Sex (m/f) per study group:

- Placebo m/f = 18/1
- Mometasone furoate m/f = 15/2

Number randomized per study group:

- Placebo = 19
- Mometasone furoate = 17

Number reaching end of study per study group:

- Placebo = 17
- Mometasone furoate = 16

Interventions	Study group 1: placebo Study group 2: mometasone furoate 4 spray doses at 50 µg by mouth to be swallowed 4 times daily after meals with no eating or drinking allowed 30 minutes after intake Duration of treatment is 8 weeks	
Outcomes	Primary outcomes of the study: <ol style="list-style-type: none"> 1. Watson Dysphagia Scale Score (WDS) (time frame: 2 months) 2. Difference in WDS score during treatment in active as compared to placebo group Secondary outcomes of the study: <ol style="list-style-type: none"> 1. The EORTC QLQ-OES18 dysphagia scale, the eating scale and choking item (time frame: 2 months) 2. "global health and social functioning dimensions" of SF-36 (time frame: 2 months) 	
Notes	Funding source: the study was supported by funding provided by the The Health & Medical Care Committee of the Regional Executive Board, Region Vastra Gotaland (Project 109901) Conflicts of interest: no potential conflict of interest was reported by the author(s)	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Computer-generated randomization method.
Allocation concealment (selection bias)	Low risk	Treatment was allocated centrally. Intervention and placebo were delivered in similar packages.
Blinding of participants and personnel (performance bias) All outcomes	Low risk	"After providing informed consent, the patient was asked to follow the study nurse who separately registered the patient, noted the randomization number of the treatment drug package and handed it over to the study participant without knowledge of its contents".

Medical treatment of eosinophilic esophagitis (Review)

Tytor 2021 (Continued)

Blinding of outcome assessment (detection bias) All outcomes	Low risk	The author confirmed in November 2022 that the random code was not broken for assessors. Patient questionnaires were returned anonymously and those analyzing did not have group data.
Incomplete outcome data (attrition bias) All outcomes	Low risk	No imbalance between placebo and intervention arm. All attrition was explained.
Selective reporting (reporting bias)	Low risk	Did not numerically present the secondary endpoints, although the authors mentioned that was due to lack of significance. We emailed the authors who sent the full data set.
Other bias	Low risk	No major baseline differences between groups. No other concerns.

AAF: amino acid formula; ASA: American Society of Anesthesiologists; BMI: body mass index; CG: control group; DB: double-blind; DSD: daily symptom diary; DSQ: Dysphagia Symptom Questionnaire; EEsAI: Eosinophilic Esophagitis Activity Index; EGD: esophagogastroduodenoscopy; EG: eosinophilic gastroenteritis; EoE: eosinophilic esophagitis; EoEHSS: EoE histology scoring system; eos: eosinophils; EoT: end of treatment; EREFS: EoE Endoscopic Reference Score; f: female; FFED: four-food elimination diet; GERD: gastroesophageal reflux disease; GI: gastrointestinal; H2: histamine H2-receptor; HES: hypereosinophilic syndrome; hpf: high-power field; HRQoL: health-related quality of life; IG: intervention group; IQR: interquartile range; IV: intravenous; LOCF: last observation carried forward; m: male; NR: not reported; NRS: numerical rating scale; OVB: oral viscous budesonide; PEES: Pediatric Eosinophilic Esophagitis Symptom Severity; PPI: proton pump inhibitor; PRO: patient-reported outcome; RCT: randomized controlled trial; SD: standard deviation; SDI: Straumann Dysphagia Instrument; SGC: swallowed glucocorticoids; VAS: visual analogue scale; WDS: Watson Dysphagia Scale

Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Braathen 2006	Wrong population - not EoE at baseline
Ceves 2005	Wrong study type - not an RCT
Comer 2017	Wrong population - not EoE
Della 2017	Wrong study type
Dellon 2020c	Wrong population
Eluri 2017	Wrong study type
Eluri 2017a	Wrong study type
EUCTR2014-002465-30-IT 2014	Wrong study type
Francis 2012	Wrong study type
Hefner 2016	Wrong population
Helou 2008	Wrong study type
Hudgens 2017	Wrong study type
JPRN-UMIN000021041 2016	Wrong study type - not randomized
JPRN-UMIN000026704 2017	Wrong study type

Medical treatment of eosinophilic esophagitis (Review)

Study	Reason for exclusion
Kagalwalla 2006	Wrong study type - not an RCT
Kavitt 2016	Wrong intervention (dilation)
Kruszewski 2016	Wrong study type
Kuzumoto 2021	Wrong study type
Molina-Infante 2017	Wrong study type
NCT01458418 2011	Abandoned RCT - no results
NCT01498497 2011	Wrong study type - not an RCT
NCT01702701 2012	Abandoned RCT - no results
NCT01821898 2013	Abandoned RCT - no results
NTR4892 2014	Wrong study type - not an RCT
Safroneeva 2015	Wrong study type - not an RCT
Safroneeva 2018	Wrong study type - not an RCT
Safroneeva 2018a	Wrong study type - not an RCT
Savarino 2015	Wrong study type - not an RCT
Song 2020	Wrong study type - not an RCT
Spergel 2002	Wrong study type - not an RCT
Spergel 2005	Wrong study type - not an RCT
Syverson 2020	Wrong study type - not an RCT
Tripp 2017	Wrong population – not EoE
Vazquez-Elizondo 2013	Wrong study type - not an RCT
Wang 2017	Wrong study type - not an RCT
Warners 2016	Wrong study type - not an RCT
Wechsler 2017	Wrong study type - not an RCT
Wright 2020	Wrong population – not EoE
Wright 2021	Wrong population

EoE: eosinophilic esophagitis; RCT: randomized controlled trial

Characteristics of studies awaiting classification *[ordered by study ID]*

Medical treatment of eosinophilic esophagitis (Review)

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Amini 2022

Methods	RCT
Participants	30 children
Interventions	Both groups received the same treatment (elimination diet, topical steroid, and proton pump inhibitor). A synbiotic (KidiLact) was added to the medication regimen of 15 patients (case), while the next 15 patients received a placebo (control).
Outcomes	Severity and frequency of symptoms were assessed with a checklist derived from a validated scoring tool in both groups before and after 8 weeks of treatment
Notes	This study was identified during our update search and will be fully included when this review is updated

NCT01846962 2012

Methods	RCT
Participants	64
Interventions	6-food elimination diet Fluticasone Budesonide Oral viscous budesonide (OVB)
Outcomes	Primary: Efficacy clinical severity score Secondary: Efficacy severity score for endoscopy and histology
Notes	Author contacted on 22 March 2022 for update - no response received

RCT: randomized controlled trial

Characteristics of ongoing studies [ordered by study ID]

ACTRN12619000141145 2019

Study name	'Two food group elimination diet versus swallowed fluticasone for the management of adult eosinophilic esophagitis, a single-centred randomised prospective study'
Methods	RCT
Participants	194
Interventions	Arm 1 (swallowed topical steroids): fluticasone 500 µg swallowed twice-daily using a metered dose inhaler for 8 weeks Arm 2 (2-food group elimination diet): eliminating cow's milk and wheat under the guidance of a gastrointestinal dietitian for 8 weeks

Medical treatment of eosinophilic esophagitis (Review)

ACTRN12619000141145 2019 (Continued)

Outcomes	<p>Primary outcomes:</p> <p>Proportion of patients that have responded to each intervention as defined by histological remission. Histological remission will be defined as reduction of eosinophils to < 15 per HPF on both the distal and proximal esophageal biopsies.</p> <p>Secondary outcomes:</p> <p>Comparison of the Dysphagia Symptom Questionnaire between proton pump inhibitor responsive eosinophilia and patients with eosinophilic esophagitis</p> <p>Comparing the differences in eosinophil counts (using histology, esophageal biopsies) between PPI responsive eosinophilia and eosinophilic esophagitis</p> <p>Comparing the differences in serum anti-TTG antibodies (using standard laboratory blood tests) between PPI responsive eosinophilia and eosinophilic esophagitis</p> <p>Comparing the differences in Dysphagia Symptom Score between PPI responsive eosinophilia and eosinophilic esophagitis</p> <p>Comparing the differences in Adult Eosinophilic Esophagitis Quality of Life Questionnaire between PPI responsive eosinophilia and eosinophilic esophagitis</p> <p>Comparing the differences between swallowed topical fluticasone 500 µg twice-daily for 8 weeks (Arm 1) and the 2-food (cow's milk and wheat) elimination diet for 8 weeks in Dysphagia Symptom Score</p> <p>Comparing the differences in serum IgE levels (using standard laboratory blood tests) between the PPI responsive eosinophilia and eosinophilic esophagitis</p>
Starting date	4 February 2019
Contact information	Dr Abdunnasser Lafta nasserhawas@gmail.com
Notes	—

ACTRN12621001406897

Study name	'Efficacy of fucoidan for eosinophilic oesophagitis: a phase 2 pilot study'
Methods	RCT phase 2
Participants	Adults
Interventions	<p>Patients will be treated with steroids and proton pump inhibitor (PPI) therapy for a minimum of 6 weeks as part of the routine clinical management (prescribed by Gastroenterologist and/or in consultation with participants own GP). Only the study medication will be provided as part of the study. Participants will take the study medication in combination with routine clinical management for an initial period of 6 weeks, and then continue taking the study medication for a further 6 weeks (total fucoidan/placebo supplementation will be 12 weeks). Participants will be randomized to one of 2 treatment groups.</p> <p>(1) A formulation containing 100 mg daily of 85% Maritech (Marinova, Tasmania, Australia)</p> <p>(2) Placebo: same formulation as the fucoidan supplement but without the active ingredient</p> <p>The active treatment will be administered via an oral gel, once per day. Participants will be asked to drink the gel slowly, allowing it to coat the esophagus thoroughly, and asked to refrain from eating for 30 minutes afterwards.</p>

Medical treatment of eosinophilic esophagitis (Review)

ACTRN12621001406897 (Continued)

Outcomes	<p>Primary outcomes:</p> <p>Change in EoE disease activity based on esophageal eosinophil count (13 weeks post treatment initiation)</p> <p>Change in EoE disease activity based on change in Dysphagia Symptom Questionnaire score of > 30% (13 weeks post treatment initiation)</p> <p>Secondary outcomes:</p> <p>Esophageal microbiome: diversity score (count of identified operational taxonomic units) (13 weeks post treatment initiation)</p> <p>Fecal microbiome: diversity score (count of identified operational taxonomic units) (13 weeks post treatment initiation)</p> <p>Esophageal biopsy immune gene expression profiling: the ratio of Th/Th2 inflammatory pathways and cells (13 weeks post treatment initiation)</p> <p>Fecal microbiome: microbial composition (13 weeks post intervention initiation)</p> <p>Esophageal microbiome: microbial composition (13 weeks post treatment initiation)</p>
Starting date	13 February 2023
Contact information	n.west@griffith.edu.au and r.ramsey@griffith.edu.au
Notes	—

EUCTR2017-003516-39-ES 2021

Study name	'A phase III clinical study in adult and adolescent patients with eosinophilic inflammation of the gullet to prove superiority compared to placebo of an episodic and/or a continuous 48-week treatment with budesonide orodispersible tablets for maintaining remission'
Methods	RCT
Participants	110
Interventions	Budesonide 1 mg orodispersible tablets Budesonide 0.5 mg orodispersible tablets
Outcomes	<p>Primary outcome:</p> <p>Proportion of patients free of treatment failure after a 48 weeks DB treatment phase</p> <p>Secondary outcomes:</p> <p>Proportion of patients with histological relapse at DB week 48 Proportion of patients with clinical relapse, or who have experienced a food impaction, who need endoscopic intervention during the DB treatment phase Proportion of patients in clinico-histological remission at DB week 48</p>
Starting date	—
Contact information	—

Medical treatment of eosinophilic esophagitis (Review)

EUCTR2017-003516-39-ES 2021 (Continued)

Notes —

EUCTR2017-003737-29-ES 2019

Study name	'Double-blind (neither physician nor patient knows of the actual treatment which can be with or without active substance), randomized (patient will be allocated to a certain treatment group by chance), placebo-controlled (one of the treatment groups receives medication without active substance), phase II/III study on the efficacy and tolerability of oral budesonide suspension in comparison with placebo in children and adolescents with eosinophilic esophagitis'
Methods	RCT
Participants	75
Interventions	Budesonide oral suspension (0.2 mg/mL)
Outcomes	<p>Primary outcome:</p> <p>Double-blind phase: rate of patients with pathological remission and clinical response at DB week 12</p> <p>Secondary outcomes:</p> <p>Double-blind phase: to further assess efficacy of budesonide oral suspension in children and adolescents with eosinophilic esophagitis (EoE)</p>
Starting date	—
Contact information	—
Notes	—

EUCTR2019-002871-32-ES 2019

Study name	'A study of benralizumab in patients with eosinophilic esophagitis'
Methods	RCT
Participants	170
Interventions	Benralizumab
Outcomes	<p>Primary outcomes:</p> <p>Dual-primary endpoints:</p> <ol style="list-style-type: none"> 1. Proportion of patients with a histologic response at week 24, defined as a peak esophageal intraepithelial eosinophil count = 6 eos/hpf 2. Changes from baseline in DSQ score at week 24 <p>Secondary outcomes:</p> <p>DB treatment period: To evaluate the effect of benralizumab dose regimen 1 on:</p>

Medical treatment of eosinophilic esophagitis (Review)

EUCTR2019-002871-32-ES 2019 (Continued)

1. Clinical features of EoE and disease activity
2. Patient-reported QOL measures
3. Healthcare resource utilization due to EoE
4. Patient-reported measures of disease severity and health status

To assess the PK and immunogenicity of benralizumab dose regimen 1 in patients with EoE
 To assess the safety and tolerability of benralizumab dose regimen 1 in patients with EoE

Starting date	—
Contact information	—
Notes	—

EUCTR2019-004391-19-NL 2020

Study name	'A study of benralizumab in patients with eosinophilic esophagitis'
Methods	RCT
Participants	170
Interventions	Benralizumab
Outcomes	<p>Primary outcomes:</p> <p>Dual-primary endpoints:</p> <ol style="list-style-type: none"> 1. Proportion of patients with a histologic response at week 24, defined as a peak esophageal intraepithelial eosinophil count = 6 eos/hpf 2. Changes from baseline in DSQ score at week 24 <p>Secondary outcomes:</p> <p>DB treatment period: To evaluate the effect of benralizumab dose regimen 1 on:</p> <ol style="list-style-type: none"> 1. Clinical features of EoE and disease activity 2. Patient-reported QOL measures 3. Healthcare resource utilization due to EoE 4. Patient-reported measures of disease severity and health status <p>To assess the PK and immunogenicity of benralizumab dose regimen 1 in patients with EoE To assess the safety and tolerability of benralizumab dose regimen 1 in patients with EoE</p>
Starting date	30 July 2020
Contact information	cpaterson@allakos.com
Notes	Sponsored by Allakos Inc.

EUCTR2020-000082-16-DE 2020

Study name	'A study to investigate the efficacy and tolerability of the drug ESO-101 in adult patients with inflammation of the esophagus'
Methods	RCT
Participants	42
Interventions	ESO-101
Outcomes	<p>Primary outcome:</p> <p>Absolute change in peak eosinophil count from baseline to EOT</p> <p>Secondary outcomes:</p> <p>Histological response and clinical symptoms Clinical response assessed by patient-reported outcome Endoscopic response Safety and tolerability Patient-reported treatment satisfaction</p>
Starting date	—
Contact information	—
Notes	—

EUCTR2020-001314-37-DE 2020

Study name	'Clinical study to show equal clinical efficacy of two dosing regimen of budesonid orodispersible tablets (twice daily vs. once daily) for treatment of inflammation of the esophagus'
Methods	RCT
Participants	242
Interventions	<p>Budesonide 2 mg orodispersible tablets</p> <p>Budesonide 1 mg orodispersible tablets</p>
Outcomes	<p>Primary outcomes:</p> <p>Proportion of patients with histological remission</p> <p>Secondary outcomes:</p> <p>EoE-associated clinical, endoscopic, and histological findings after 6? weeks treatment with budesonide orodispersible tablets Safety and tolerability as assessed by adverse events and laboratory parameters Patients' quality of life</p>
Starting date	—
Contact information	—
Notes	—

Medical treatment of eosinophilic esophagitis (Review)

EUCTR2020-003226-23-BE 2020

Study name	'A study to assess the safety and efficacy of oral etrasimod in adult participants with eosinophilic esophagitis VOYAGE'
Methods	RCT
Participants	96
Interventions	Etrasimod
Outcomes	<p>Percent change from baseline in esophageal peak eosinophil count (PEC)</p> <p>Absolute change from baseline in Dysphagia Symptom Questionnaire (DSQ) score (time frame: baseline to week 16)</p> <p>Absolute change from baseline in esophageal PEC (time frame: baseline to week 16)</p> <p>Number and severity of adverse events (double-blind treatment period and extension treatment period) (time frame: up to approximately 56 weeks (24 weeks of double-blind treatment period, 28 weeks of extension treatment period, and 4 weeks of safety follow-up period))</p> <p>Proportion of participants with esophageal PEC < 15 eosinophils per high-powered field (eos/hpf) (time frame: baseline to week 16)</p> <p>Proportion of participants with esophageal PEC = 6 eos/hpf (time frame: baseline to week 16)</p>
Starting date	—
Contact information	—
Notes	—

EUCTR2020-004336-16-DE 2021

Study name	'A study to evaluate the efficacy and safety of CC-93538 in adult and adolescent patients who have eosinophilic esophagitis'
Methods	RCT
Participants	399
Interventions	Cendakimab
Outcomes	<p>Primary outcomes:</p> <ol style="list-style-type: none"> 1. Mean change in dysphagia days (DD), evaluated over the prior 14-day period using the modified Daily Symptom Diary (mDSD), from baseline to week 24 2. Proportion of participants with eosinophilic histologic response defined as a peak esophageal eosinophil count < 6/high-power field (hpf) at week 24 <p>Secondary outcomes:</p> <p>To assess the efficacy of CC-93538 versus placebo at 24 weeks in improving:</p> <ol style="list-style-type: none"> 1. Endoscopic features of eosinophilic esophagitis (EoE) 2. Histologic features of EoE

EUCTR2020-004336-16-DE 2021 (Continued)

To assess the persistence of effect of CC-93538 at 48 weeks in reducing:

1. Dysphagia symptoms
2. Esophageal eosinophil counts

To assess the persistence of effect of CC-93538 through administration of a less frequent dosing regimen at 48 weeks in reducing:

1. Dysphagia symptoms
2. Esophageal eosinophil counts

To assess the persistence of effect of CC-93538 at 48 weeks in improving:

1. Endoscopic features of EoE
2. Histologic features of EoE

To evaluate the time to and frequency of EoE flare events and use of rescue therapy during the study

To evaluate the safety and tolerability of CC-93538 including characterization of the immunogenicity profile

To assess trough concentrations of CC-93538 in participants with EoE

Starting date	—
Contact information	—
Notes	—

Henry 2019

Study name	'A novel food additive removal diet for eosinophilic esophagitis (free study): conceptual design and clinical trial methods'
Methods	RCT (NCT03657771)
Participants	72 participants
Interventions	<p>Study group 1: dairy-free diet</p> <p>Study group 2: dairy-free diet with food additive-free diet</p>
Outcomes	<p>Primary outcomes:</p> <p>Eosinophils per high-power field (eos/hpf) (time frame: 12 weeks)</p> <p>Secondary outcomes:</p> <p>Eosinophilic Esophagitis Endoscopic Reference Score (EREFS) (time frame: 12 weeks)</p> <p>Other outcomes:</p> <ol style="list-style-type: none"> 1. Pediatric Eosinophilic Esophagitis Symptom Severity Module (PEESS) (time frame: baseline, 4, 8, and 12 weeks) 2. Pediatric Eosinophilic Esophagitis Symptom Severity Module (PEESS): parent report (time frame: baseline, 4, 8, and 12 weeks) 3. Pediatric Quality of Life Inventory: Eosinophilic Esophagitis Module (PedsQL-EoE) (time frame: baseline, 4, 8, and 12 weeks)

Henry 2019 (Continued)

4. Pediatric Quality of Life Inventory: Eosinophilic Esophagitis Module (PedsQL-EoE): parent report (time frame: baseline, 4, 8, and 12 weeks)

Starting date	September 2018
Contact information	James Franciosi (review author): james.franciosi@nemours.org
Notes	Funding source: anonymous donor to the Nemours Foundation Conflicts of interest: not available

IRCT20171230038142N14 2020

Study name	'Evaluation of efficacy of synbiotics in children with eosinophilic esophagitis'
Methods	RCT
Participants	40
Interventions	Omeprazole Diet Topical steroids Cidetabic synthetic intervention
Outcomes	Clinical symptoms (time point: before and 8 weeks after intervention); method of measurement: examination by a pediatrician Endoscopic findings (time point: before and 8 weeks after intervention); method of measurement: endoscopy in hospital by pediatric gastroenterologist Pathologic findings (time point: before and 8 weeks after intervention); method of measurement: gastric biopsy and eosinophil count by microscope
Starting date	—
Contact information	—
Notes	—

IRCT20191211045703N1 2020

Study name	'Comparison of the efficacy and side effects of nebulized oral Pulmicort and inhaler budesonide in patients with eosinophilic esophagitis'
Methods	RCT
Participants	60
Interventions	Nebulized oral Pulmicort Inhaler budesonide

Medical treatment of eosinophilic esophagitis (Review)

IRCT20191211045703N1 2020 (Continued)

Outcomes	<p>The effect of oral budesonide nebulizer on improving eosinophilic esophagitis. Time point: patients in the oral Budesonide nebulizer group will receive 1 puff twice a day. If the effectiveness of the drug is not observed after 8 weeks, patients will receive 2 puffs twice a day. Then during weeks 4, 12, 8, and 16 patients will be followed up and examined.</p> <p>Method of measurement: at the beginning of the study and after the 16th week, patients will be re-examined for histological and tissue pathology, as well as blood eosinophil counts and cortisol examination by enzyme-linked immunosorbent test at 8 am. There will also be a routine laboratory test, such as hematology and biochemistry.</p>
Starting date	—
Contact information	—
Notes	—

JPRN-jRCT2051200140

Study name	'A phase 3, multicenter, multinational, randomized, double-blind, placebo-controlled induction and maintenance study to evaluate the efficacy and safety of CC-93538 in adult and adolescent subjects with eosinophilic esophagitis'
Methods	RCT
Participants	33 estimated
Interventions	CC-93538 is administered subcutaneously at a dose of 360 mg weekly in adults and adolescent 12 years of age or older. 24 weeks after the initial dose, a dose of 360 mg is administered subcutaneously weekly or biweekly.
Outcomes	<p>Primary outcomes:</p> <p>Induction phase endpoints at week 24:</p> <ol style="list-style-type: none"> 1. Change in DD clinical response: the mean change in dysphagia days (DD), evaluated over the prior 14-day period using the modified Daily Symptom Diary (mDSD), from baseline to week 24 2. Eosinophil histologic response (≤ 6/hpf): the proportion of participants with eosinophilic histologic response defined as a peak esophageal eosinophil count ≤ 6/high-power field (hpf) at week 24 <p>Secondary outcomes:</p> <p>Induction phase key secondary endpoints at week 24:</p> <ol style="list-style-type: none"> 1. Eosinophil histologic response (< 15/hpf): proportion of participants with eosinophilic histologic response defined as a peak esophageal eosinophil count < 15/hpf at week 24 2. EREFS: mean change in the endoscopic features of EoE as measured by the EoE Endoscopic Reference Score (EREFs) from baseline to week 24 3. EoEHSS grade score: mean change in the mean adjusted histology grade score as measured by the EoE histology scoring system (EoEHSS) from baseline to week 24 4. EoEHSS stage score: mean change in the mean adjusted histology stage score as measured by the EoE histology scoring system (EoEHSS) from baseline to week 24 5. mDSD composite score: mean change in the modified Daily Symptom Diary (mDSD) composite score from baseline to week 24
Starting date	1 July 2021
Contact information	Name: Changliang Zhang

Medical treatment of eosinophilic esophagitis (Review)

180

JPRN-jRCT2051200140 (Continued)

Email: MG-JP-RCO-JRCT@bms.com

Affiliation: Bristol-Myers Squibb

Notes

—

NCT02873468

Study name 'Efficacy and safety of three doses of Florence oral suspension in adults with eosinophilic esophagitis'

Methods RCT

Participants 116 estimated participants

Interventions Florence 30 µg/mL

Florence 60 µg/mL

Florence 90 µg/mL

Placebo

 Outcomes **Primary outcome:** proportion of participants presenting a histological response, defined as the presence of ≤ 6 eosinophils/high-power field, at the end of treatment (time frame: 100 days)

Secondary outcomes: incidence and severity of adverse events recorded during the study (time frame: 170 days)

Starting date 19 April 2021

Contact information Arthur M Kummer, MD+551938879851

pesquisa.clinica@ems.com.br

Notes

—

NCT03656380 2019

Study name 'Mepo for EoE Study'

Methods RCT

Participants 72

Interventions Mepolizumab 100 mg

Mepolizumab 300 mg

Outcomes Mean change in dysphagia from baseline to 3 months post-treatment (time frame: baseline, month 3 post-treatment)

Absolute peak eosinophil count (measured in eos/hpf) after 3 months of treatment (time frame: after 3 months of treatment)

Histologic response levels after 3 treatment months (time frame: after 3 months of treatment)

Medical treatment of eosinophilic esophagitis (Review)

NCT03656380 2019 (Continued)

Mean change in EoE Endoscopic Reference Score (EREFS) from baseline to 3 months post-treatment (time frame: baseline, 3 months post-treatment)

Mean change in the Straumann Dysphagia Instrument (SDI) score from baseline to 3 months post-treatment (time frame: baseline, 3 months post-treatment)

Proportion of participants with a clinical remission (EEsAI score of = 20 points) after 3 months of treatment (time frame: after 3 months of treatment)

Proportion of participants with a clinical response (EEsAI score decrease of = 20 points) after 3 months of treatment (time frame: after 3 months of treatment)

Starting date	—
Contact information	—
Notes	—

NCT03657771 2018

Study name	'A food additive removal diet for pediatric eosinophilic esophagitis FREE'
Methods	RCT
Participants	72
Interventions	DED FREE
Outcomes	Primary outcome: Eosinophils per high-power field (eos/hpf) (time frame: 12 weeks) Secondary outcomes: Eosinophilic Esophagitis Endoscopic Reference Score (EREFS) (time frame: 12 weeks)
Starting date	—
Contact information	—
Notes	—

NCT03781596

Study name	'EoE RCT fluticasone and omeprazole vs fluticasone alone'
Methods	RCT
Participants	100 estimated participants
Interventions	Fluticasone and omeprazole Fluticasone and placebo

Medical treatment of eosinophilic esophagitis (Review)

NCT03781596 (Continued)

Outcomes	<p>Primary outcome:</p> <p>Change in esophageal eosinophilia (time frame: week 0 and week 8)</p> <p>Secondary outcomes:</p> <p>Change in endoscopic reference score (time frame: week 0 and week 8)</p> <p>Change in symptom scoring (time frame: week 0 and week 8)</p>
Starting date	2 October 2018
Contact information	Claire P Daniels, M.D. 443-904-3353 claire.p.daniels.mil@mail.mil
Notes	—

NCT04248712 2020

Study name	'Antihistamines in eosinophilic esophagitis ATEE'
Methods	RCT
Participants	50
Interventions	Famotidine Loratadine
Outcomes	<p>Primary outcomes:</p> <p>Adverse Events (time frame: 12 weeks)</p> <p>Change in maximum eosinophil count (time frame: 12 weeks)</p> <p>Secondary outcomes:</p> <p>Change in endoscopic response, as measured by the endoscopic reference score (time frame: 12 weeks)</p> <p>Change in histologic response (time frame: 12 weeks)</p> <p>Change in symptoms of eosinophilic esophagitis, as measured by Dysphagia Symptom Questionnaire (time frame: 12 weeks)</p>
Starting date	—
Contact information	—
Notes	—

NCT04281108 2020

Study name	'Efficacy and safety APT-1011 in adult subjects with eosinophilic esophagitis (EoE) (FLUTE-2) FLUTE-2'
------------	--

Medical treatment of eosinophilic esophagitis (Review)

NCT04281108 2020 (Continued)

Methods	RCT
Participants	143
Interventions	APT-1011
Outcomes	<p>Primary outcomes:</p> <p>Histologic responder rates at the end of the randomized withdrawal phase (RWS) (time frame: week 12 to week 52)</p> <p>Mean change in number of dysphagia episodes (time frame: week 0 to week 12)</p> <p>Percentage of APT-1011 responders in Part A who remain symptomatic responders at the end of the RWS (time frame: week 0 to week 52)</p> <p>Week 12 histologic responder rates (time frame: week 12)</p> <p>Secondary outcomes:</p> <p>Change in EREFS from week 0 to week 12 (time frame: week 0 to week 12)</p> <p>Histological change from baseline to week 12 (time frame: week 0 to week 12)</p> <p>Mean change in dysphagia episodes (time frame: week 0 to week 52)</p> <p>Mean change in dysphagia-free days (time frame: week 0 to week 52)</p> <p>Mean change in EREFS from week 0 to week 52 (time frame: week 0 to week 52)</p> <p>Mean change in PROSE day-level difficulty swallowing (time frame: week 0 to week 12)</p> <p>Mean change in PROSE day-level difficulty swallowing (time frame: week 0 to week 52)</p> <p>Mean change in PROSE day-level symptom burden (time frame: week 0 to week 52)</p> <p>Mean change in PROSE symptom burden score (time frame: week 0 to week 12)</p> <p>Mean histologic change (time frame: week 0 to week 52)</p> <p>Mean number of dysphagia-free days (time frame: week 0 to week 12)</p> <p>Percentage of participants with < 1 peak eos/HPF at week 12 (time frame: week 12)</p> <p>Percentage of participants with < 15 peak eos/HPF (time frame: week 12)</p>
Starting date	—
Contact information	—
Notes	—

NCT04394351 2020

Study name	'Study to investigate the efficacy and safety of dupilumab in pediatric patients with active eosinophilic esophagitis (EoE) EoE KIDS'
Methods	RCT
Participants	90

Medical treatment of eosinophilic esophagitis (Review)

NCT04394351 2020 (Continued)

Interventions

Dupilumab

Outcomes

Primary outcome:

Proportion of patients achieving peak esophageal intraepithelial eosinophil count = 6 eos/hpf (400×) (time frame: week 16)

Secondary outcomes:

Absolute change in Eosinophilic Esophagitis-Endoscopic Reference (EoE EREFS) (time frame: week 16)

Absolute change in EoE EREFS (time frame: week 52)

Absolute change in mean eosinophilic esophagitis (EoE) Histology Scoring System (EoE-HSS) (time frame: week 16)

Absolute change in mean EoE-HSS (time frame: week 52)

Change in the proportion of days with 1 or more EoE signs as measured by the Pediatric EoE Sign/Symptom Questionnaire - caregiver version (PESQ-C) (time frame: week 16)

Change in the proportion of days with 1 or more EoE signs as measured by the PESQ-C (time frame: week 52)

Change in the proportion of days with 1 or more EoE symptoms as measured by PESQ-P (time frame: week 52)

Change in the proportion of days with 1 or more EoE symptoms as measured by the Pediatric EoE Sign/Symptom Questionnaire - patient version (PESQ-P) (time frame: week 16)

Change in the proportion of total segments within a day with 1 or more EoE signs as measured by PESQ-C (time frame: week 16)

Change in the proportion of total segments within a day with 1 or more EoE signs as measured by PESQ-C (time frame: week 52)

Change in the proportion of total segments within a day with 1 or more EoE symptoms as measured by PESQ-P (time frame: week 16)

Change in the proportion of total segments within a day with 1 or more EoE symptoms as measured by PESQ-P (time frame: week 52)

Change in the type 2 inflammation transcriptional signature (time frame: week 16)

Change in the type 2 inflammation transcriptional signature (time frame: week 52)

Change in total score as measured by the PEESV2.0 - caregiver version questionnaire (time frame: week 16)

Concentration of functional dupilumab in serum (time frame: week 52)

Incidence of treatment-emergent adverse events (TEAEs) (time frame: week 52)

Incidence of TEAEs leading to permanent discontinuation of study treatment (time frame: week 16)

Incidence of TEAEs leading to permanent discontinuation of study treatment (time frame: week 52)

Incidence of treatment-emergent anti-drug antibody (ADA) responses and titer (time frame: week 52)

Incidence of TEAEs (time frame: week 16)

Incidence of treatment-emergent adverse events of special interest (AESIs) (time frame: week 16)

NCT04394351 2020 (Continued)

- Incidence of treatment-emergent AESIs (time frame: week 52)
- Incidence of treatment-emergent ADA responses and titer (time frame: week 16)
- Incidence of treatment-emergent serious adverse events (SAEs) (time frame: week 52)
- Incidence of treatment-emergent SAEs (time frame: week 16)
- Normalized Enrichment Scores (NES) for the relative change in the EoE diagnostic panel (EDP) transcriptome signature (time frame: week 52)
- NES for the relative change in the type 2 inflammation transcriptome signature (time frame: week 16)
- NES for the relative change in the type 2 inflammation transcriptome signature (time frame: week 52)
- NES for the relative change in the EDP transcriptome signature (time frame: week 16)
- Number of sign-free days during the 14-day period preceding week 16 as measured by the PESQ-C (time frame: week 16)
- Number of sign-free days during the 14-day period preceding week 52 as measured by the PESQ-C (time frame: week 52)
- Number of symptom-free days during the 14-day period preceding week 16 as measured by the PESQ-P (patient version) (time frame: week 16)
- Number of symptom-free days during the 14-day period preceding week 52 as measured by the PESQ-P (patient version) (time frame: week 52)
- Percent change in peak esophageal intraepithelial eosinophil count (eos/hpf) (time frame: week 16)
- Percent change in peak esophageal intraepithelial eosinophil count (eos/hpf) (time frame: week 52)
- Proportion of patients achieving peak esophageal intraepithelial eosinophil count = 6 eos/hpf (400×) (time frame: week 52)
- Proportion of patients achieving peak esophageal intraepithelial eosinophil count of < 15 eos/hpf (time frame: week 16)
- Proportion of patients achieving peak esophageal intraepithelial eosinophil count of < 15 eos/hpf (time frame: week 52)

Starting date	—
Contact information	—
Notes	—

NCT04543409

Study name	'A study of benralizumab in patients with eosinophilic esophagitis (MESSINA)'
Methods	RCT
Participants	211 estimated participants
Interventions	Benralizumab

Medical treatment of eosinophilic esophagitis (Review)

NCT04543409 (Continued)

Placebo

Outcomes

Primary outcomes:

1. Proportion of patients with a histologic response at week 24, defined as a peak esophageal intraepithelial eosinophil count ≤ 6 eos/hpf (time frame: week 24)
2. Changes from baseline in Dysphagia Symptom Questionnaire (DSQ) scores (time frame: week 24)

Secondary outcomes:

1. Percent change from baseline in tissue eosinophils (time frame: week 24)
2. Change from baseline in Eosinophilic Esophagitis-Histology Scoring System (EoE-HSS) grade score (time frame: week 24)
3. Change from baseline in Eosinophilic Esophagitis-Histology Scoring System (EoE-HSS) stage score (time frame: week 24)
4. Changes from baseline in centrally read Endoscopic Reference Score (EREFs) (time frame: week 24)
5. Treatment responder rate at week 24, defined as a composite of histological response (≤ 6 eos/hpf) and clinically meaningful improvement from baseline in Dysphagia Symptom Questionnaire (DSQ) scores (30% improvement) (time frame: week 24)
6. Centrally read biopsies for additional histopathology including tissue eosinophil counts (time frame: week 24)
7. Dysphagia-free days as captured by the Dysphagia Symptom Questionnaire (DSQ) (time frame: week 24)
8. Frequency of dysphagia episodes as captured by the Eosinophilic Esophagitis Daily Dysphagia Diary (EoE-3D) (time frame: week 24)
9. Changes from baseline in dysphagia-associated pain, discomfort, and overall severity as captured by the EoE-3D (time frame: week 24)
10. Changes from baseline in abdominal pain and nausea as captured by the daily diary (time frame: week 24)
11. Changes from baseline in PEES (time frame: week 24)
12. Changes from baseline in Adult Eosinophilic Esophagitis Quality of Life Questionnaire (EOE-QoL-A) (time frame: week 24)
13. Change from baseline in Short Form 36-item health survey (version 2, acute recall) (SF-36v2) (time frame: week 24)
14. Percent of patients with relevant concomitant procedures and healthcare resource utilization during the study through week 24 (time frame: week 24)
15. Patient-reported overall severity of disease as measured by Patient Global Impression of Severity (PGI-S) (time frame: week 24)
16. Patient-reported change in health status since baseline as measured by Patient Global Impression of Change (PGI-C) (time frame: week 24)

Other outcome measures:

1. Benralizumab pharmacokinetics (time frame: minimum of 52 weeks)
2. Immunogenicity of benralizumab (time frame: minimum of 52 weeks)
3. Safety and tolerability (time frame: minimum of 52 weeks)

Starting date 22 September 2020

Contact information Marc E. Rothenberg, MD, PhD

Notes —

NCT04593251 2020

Study name	'Dose escalation study to evaluate an experimental new treatment (CALY-002) in healthy subjects and subjects with celiac disease and eosinophilic esophagitis'
Methods	RCT
Participants	95
Interventions	CALY-002
Outcomes	Primary outcome: Incidence of treatment-emergent adverse events (time frame: through study completion, an average of 3 months post last dose)
Starting date	—
Contact information	—
Notes	—

NCT04835168 2022

Study name	'Safety and pharmacokinetics of orodispersible BT-11 in active eosinophilic esophagitis'
Methods	RCT
Participants	Withdrawn (the closure of the study was driven by the decision to redesign the study protocol for future studies)
Interventions	BT-11 500 mg BT-11 1000 mg
Outcomes	Primary outcome: Incidence and severity of adverse events (time frame: 12 weeks)
Starting date	—
Contact information	—
Notes	—

NCT05084963 2021

Study name	'A study to assess the efficacy, safety and tolerability of IRL201104 in adults with active eosinophilic esophagitis'
Methods	RCT
Participants	36 participants
Interventions	Arm 1: IRL201104 Dose A

Medical treatment of eosinophilic esophagitis (Review)

NCT05084963 2021 (Continued)

Arm 2: IRL201104 Dose B

Outcomes	<p>Primary outcome:</p> <p>Change from baseline in the peak esophageal intraepithelial eosinophil count at week 4 (time frame: 4 weeks)</p> <p>The change from baseline in histologic eosinophil count in each treatment group will be summarized as the mean, standard deviation, median, minimum, and maximum</p>
Starting date	—
Contact information	—
Notes	—

NCT05214599

Study name	'Pharmacokinetics, efficacy, tolerability and safety of different budesonide oral gel doses in adults' subjects of both genders with eosinophilic esophagitis (EoE) (BESIDE)'
Methods	RCT
Participants	36 estimated
Interventions	Budesonide gel low-, medium- and high-dose
Outcomes	<p>Primary outcomes:</p> <ol style="list-style-type: none"> 1. Peak plasma concentration (C_{max}) (time frame: first 24 hours after a single drug dose administration) 2. Area under the plasma concentration versus time curve (AUC) (time frame: first 24 hours after a single drug dose administration) 3. Half-life (T_{1/2}) (time frame: first 24 hours after a single drug dose administration) 4. Oral clearance (CL/F) (time frame: first 24 hours after a single drug dose administration) 5. ≤ 6 eosinophils per high-power field (time frame: 8 weeks of treatment) 6. Improvement in dysphagia symptoms consistent with the disease EAT (Eating Assessment Tool)-10 questionnaire (time frame: 8 weeks of treatment) <p>Secondary outcomes:</p> <ol style="list-style-type: none"> 1. Assessment of non-serious and serious adverse events rate (time frame: through 8 weeks) 2. Quality of life assessment of participants (time frame: through 8 weeks)
Starting date	2 September 2023
Contact information	<p>No contacts or locations provided</p> <p>Bazell Pharma AG</p>
Notes	—

NCT05543512

Study name	'The Immune Directed Individualized Elimination Therapy (iDIET) Study (iDIET)'
Methods	RCT pilot/feasibility
Participants	100 estimated
Interventions	Participants will be randomized in a 1:1 fashion to follow an allergen-specific immune signature-directed diet or sham diet during the 8-week treatment period
Outcomes	<p>Primary outcomes:</p> <ol style="list-style-type: none"> 1. Post-treatment peak eosinophil count (time frame: 8 weeks) <p>Secondary outcomes:</p> <ol style="list-style-type: none"> 1. Dysphagia symptom score (time frame: 8 weeks) 2. Endoscopic severity (time frame: 8 weeks) 3. Percentage of histologic responders (time frame: 8 weeks) 4. Change in peak eosinophil count (time frame: baseline and week 8)
Starting date	14 October 2022
Contact information	Evan S Dellon, MD, MPH
Notes	—

NCT05583227

Study name	'Efficacy and safety of tezepelumab in patients with eosinophilic esophagitis (CROSSING)'
Methods	RCT
Participants	360 estimated
Interventions	Subjects will be randomized in a 1:1:1 ratio to receive either a low dose of tezepelumab, a high dose of tezepelumab, or placebo
Outcomes	<p>Primary outcomes:</p> <ol style="list-style-type: none"> 1. Histologic response of peak esophageal eosinophil per HPF count of ≤ 6 across all available esophageal levels (time frame: week 24) 2. Change from baseline in DSQ (Dysphagia Symptom Questionnaire) score (time frame: week 24) <p>Secondary outcomes:</p> <ol style="list-style-type: none"> 1. Change from baseline in EoE EREFS (endoscopic reference score) (time frame: week 24, week 52) 2. Change from baseline in EoE-HSS (histologic scoring system) grade score (time frame: week 24) 3. Change from baseline in EoE-HSS (histologic scoring system) stage score (time frame: week 24) 4. Histologic response of peak esophageal eosinophil per HPF count of ≤ 6 across all available esophageal levels (time frame: week 52) 5. Change from baseline in DSQ (Dysphagia Symptom Questionnaire) score (time frame: week 52) 6. Response of achieving clinico-histological remission (time frame: week 24, week 52) <p>Other outcomes:</p>

NCT05583227 (Continued)

1. Change from baseline in peak esophageal eosinophil count (EOS/HPF) (time frame: week 24, week 52)
2. Changes from baseline in PEES module at week 24 (adolescents only) (time frame: week 24, week 52)
3. Change from baseline in EoE-HSS (histologic scoring system) stage score (time frame: week 52)
4. Change from baseline in EoE-HSS (histologic scoring system) grade score (time frame: week 52)
5. Serum tezepelumab concentration (time frame: weeks 0, 4, 12, 24, and 52)
6. Anti-drug antibody (time frame: weeks 0, 12, 24, and 52)

Starting date	10 November 2022
Contact information	AstraZeneca Clinical Study Information Center 1-877-240-9479 information.center@astrazeneca.com
Notes	—

NCT05634746

Study name	'24-week induction study of APT-1011 in adult subjects with eosinophilic esophagitis (EoE) (FLUTE 3)'
Methods	RCT
Participants	200 participants estimated
Interventions	APT-1011 3 mg Placebo
Outcomes	<p>Primary outcomes:</p> <ol style="list-style-type: none"> 1. Histological remission (co-primary outcome measure) (time frame: week 24) 2. Complete symptomatic response (co-primary outcome measure) (time frame: week 24) <p>Secondary outcomes:</p> <ol style="list-style-type: none"> 1. Clinicopathologic responder rate (time frame: week 24) 2. Percentage of participants with $\geq 70\%$ reduction in dysphagia frequency (time frame: week 24) 3. Mean change in dysphagia frequency (time frame: week 24) 4. Mean change in PROSE difficulty swallowing (time frame: week 24) 5. Mean change in PROSE pain with swallowing (time frame: week 24) 6. Mean number of dysphagia-free days (time frame: week 24) 7. Percentage of responders (strictures and \geq grade 2 rings) (time frame: week 24) 8. Percentage of responders (strictures) (time frame: week 24) 9. Percentage of responders (\geq grade 2 rings) (time frame: week 24) 10. Mean change in EREFS (time frame: week 24) 11. Time to first complete symptom response (time frame: week 24)
Starting date	29 December 2022
Contact information	ClinicalTrials@ellodipharma.com

Medical treatment of eosinophilic esophagitis (Review)

NCT05634746 (Continued)

Notes

—

NCT05695456

Study name	'Targeted elimination diet in EoE patients following identification of trigger nutrients using confocal laser endomicroscopy (CLE-EoE)'
Methods	RCT cross-over
Participants	Estimated 25 participants
Interventions	Patients with a positive CLE reaction to one or two specific nutrients will then be randomized to a blinded exclusion diet for 6 weeks of those nutrients or to exclusion of another tested nutrient that yielded no change in CLE (= sham diet), in a cross-over fashion. Patients with no CLE reaction will undergo an empirical exclusion diet of gluten-containing grains for 6 weeks. To mirror the cross-over character of the intervention, CLE negative patients will then undergo a milk exclusion diet for 6 weeks (order is interchangeable).
Outcomes	<p>Primary outcome:</p> <ol style="list-style-type: none"> 1. Difference in response (time frame: 2 years) <p>Secondary outcomes:</p> <ol style="list-style-type: none"> 1. Further histological outcomes (time frame: 2 years) 2. Further histological outcomes (time frame: 2 years) 3. Symptomatic changes in patients undergoing duodenal CLE-targeted elimination diet compared to the sham diet (time frame: 2 years) 4. Endoscopic changes in patients undergoing duodenal CLE-targeted elimination diet compared to the sham diet (time frame: 2 years) 5. Change in esophageal and duodenal permeability parameters 1 (time frame: 2 years) 6. Change in esophageal and duodenal permeability parameters 2 (time frame: 2 years) 7. Baseline and post-exposure duodenal mast cell and eosinophil counts on histology (time frame: 2 years) 8. Differences in duodenal permeability measures in CLE positive, CLE negative EoE patients and healthy controls before and after nutrient application 1 (time frame: 2 years) 9. Differences in duodenal permeability measures in CLE positive, CLE negative EoE patients and healthy controls before and after nutrient application 2 (time frame: 2 years)
Starting date	16 February 2022 (retrospectively registered)
Contact information	Jan Tack, MD PhD +3216345514 jan.tack@kuleuven.be
Notes	—

CLE: confocal laser endomicroscopy; DB: double-blind; DD: dysphagia days; DED: dairy elimination diet; DSD: daily symptom diary; DSQ: Dysphagia Symptom Questionnaire; EoE-HSS: EoE histology scoring system; EoE: eosinophilic esophagitis; EOT: end of treatment; EREFS: Eosinophilic Esophagitis Endoscopic Reference Score; FREE: dairy elimination plus food additive elimination; GP: general practitioner; HPF/hpf: high-power field; PEC: peak eosinophil count; PEES: Pediatric Eosinophilic Esophagitis Symptom Severity; PK: pharmacokinetic; PPI: proton pump inhibitor; QOL: quality of life; RCT: randomized controlled trial; TTG: tissue transglutaminase

DATA AND ANALYSES

Comparison 1. Corticosteroids vs placebo for induction of remission

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.1 Clinical improvement at study endpoint (dichotomous)	6	583	Risk Ratio (M-H, Random, 95% CI)	1.74 [1.08, 2.80]
1.2 Clinical improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	6	583	Risk Ratio (M-H, Fixed, 95% CI)	1.54 [1.25, 1.89]
1.3 Clinical improvement at study endpoint (dichotomous), sensitivity analysis, validated instruments	2	360	Risk Ratio (M-H, Random, 95% CI)	1.39 [1.08, 1.79]
1.4 Clinical improvement at study endpoint (dichotomous), subgrouped by age	6	583	Risk Ratio (M-H, Random, 95% CI)	1.74 [1.08, 2.80]
1.4.1 Children (18 years and younger)	1	81	Risk Ratio (M-H, Random, 95% CI)	0.88 [0.60, 1.27]
1.4.2 Mixed children and adults (18 years and older)	5	502	Risk Ratio (M-H, Random, 95% CI)	2.13 [1.27, 3.57]
1.5 Clinical improvement at study endpoint (dichotomous), subgrouped by type of steroid	6	583	Risk Ratio (M-H, Random, 95% CI)	1.74 [1.08, 2.80]
1.5.1 Beclomethasone	1	18	Risk Ratio (M-H, Random, 95% CI)	3.00 [0.38, 23.68]
1.5.2 Budesonide	4	523	Risk Ratio (M-H, Random, 95% CI)	1.74 [0.95, 3.16]
1.5.3 Fluticasone	1	42	Risk Ratio (M-H, Random, 95% CI)	1.71 [0.84, 3.48]
1.6 Clinical improvement at study endpoint (dichotomous), subgrouped by delivery method	6	583	Risk Ratio (M-H, Random, 95% CI)	1.74 [1.08, 2.80]
1.6.1 Adapted asthma	3	96	Risk Ratio (M-H, Random, 95% CI)	2.23 [1.30, 3.83]
1.6.2 Esophageal-specific	3	487	Risk Ratio (M-H, Random, 95% CI)	1.48 [0.79, 2.77]
1.7 Clinical improvement at study endpoint (continuous)	5	475	Std. Mean Difference (IV, Random, 95% CI)	0.51 [0.17, 0.85]
1.8 Clinical improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	5	475	Std. Mean Difference (IV, Fixed, 95% CI)	0.37 [0.18, 0.56]

Medical treatment of eosinophilic esophagitis (Review)

193

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.9 Clinical improvement at study endpoint (continuous), sensitivity analysis, validated instruments	3	407	Std. Mean Difference (IV, Random, 95% CI)	0.35 [0.07, 0.64]
1.10 Clinical improvement at study endpoint (continuous), subgrouped by age	5	475	Std. Mean Difference (IV, Random, 95% CI)	0.51 [0.17, 0.85]
1.10.1 Children (18 years and younger)	1	32	Std. Mean Difference (IV, Random, 95% CI)	0.34 [-0.39, 1.08]
1.10.2 Mixed children and adults (18 years and older)	4	443	Std. Mean Difference (IV, Random, 95% CI)	0.55 [0.14, 0.97]
1.11 Clinical improvement at study endpoint (continuous), subgrouped by type of steroid	5	475	Std. Mean Difference (IV, Random, 95% CI)	0.51 [0.17, 0.85]
1.11.1 Budesonide	4	442	Std. Mean Difference (IV, Random, 95% CI)	0.51 [0.10, 0.91]
1.11.2 Mometasone	1	33	Std. Mean Difference (IV, Random, 95% CI)	0.59 [-0.11, 1.29]
1.12 Clinical improvement at study endpoint (continuous), subgrouped by delivery method	5	475	Std. Mean Difference (IV, Random, 95% CI)	0.51 [0.17, 0.85]
1.12.1 Adapted asthma	1	36	Std. Mean Difference (IV, Random, 95% CI)	1.23 [0.51, 1.95]
1.12.2 Esophageal-specific	4	439	Std. Mean Difference (IV, Random, 95% CI)	0.31 [0.11, 0.51]
1.13 Histological improvement at study endpoint (dichotomous)	12	978	Risk Ratio (M-H, Random, 95% CI)	11.94 [6.56, 21.75]
1.14 Histological improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	12	978	Risk Ratio (M-H, Fixed, 95% CI)	18.87 [10.57, 33.71]
1.15 Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold < 15 eos/hpf	4	476	Risk Ratio (M-H, Random, 95% CI)	18.47 [4.45, 76.72]
1.16 Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 6 eos/hpf	10	912	Risk Ratio (M-H, Random, 95% CI)	14.03 [6.73, 29.26]
1.17 Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 1 eos/hpf	4	424	Risk Ratio (M-H, Random, 95% CI)	10.97 [3.12, 38.55]
1.18 Histological improvement at study endpoint (dichotomous), subgrouped by age	12	978	Risk Ratio (M-H, Random, 95% CI)	11.94 [6.56, 21.75]

Medical treatment of eosinophilic esophagitis (Review)

194

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.18.1 Children (18 years and younger)	3	149	Risk Ratio (M-H, Random, 95% CI)	5.76 [2.07, 16.02]
1.18.2 Mixed children and adults (18 years and older)	9	829	Risk Ratio (M-H, Random, 95% CI)	15.17 [7.88, 29.23]
1.19 Histological improvement at study endpoint (dichotomous), subgrouped by type of steroid	12	978	Risk Ratio (M-H, Random, 95% CI)	11.94 [6.56, 21.75]
1.19.1 Budesonide	7	728	Risk Ratio (M-H, Random, 95% CI)	16.70 [7.60, 36.70]
1.19.2 Fluticasone	5	250	Risk Ratio (M-H, Random, 95% CI)	7.57 [3.36, 17.08]
1.20 Histological improvement at study endpoint (dichotomous), subgrouped by delivery method	12	978	Risk Ratio (M-H, Random, 95% CI)	11.94 [6.56, 21.75]
1.20.1 Esophageal-specific	8	822	Risk Ratio (M-H, Random, 95% CI)	18.20 [8.29, 39.95]
1.20.2 Adapted asthma, allergy	4	156	Risk Ratio (M-H, Random, 95% CI)	7.30 [3.37, 15.84]
1.21 Histological improvement at study endpoint (continuous)	5	449	Std. Mean Difference (IV, Random, 95% CI)	1.42 [1.02, 1.82]
1.22 Histological improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	5	449	Std. Mean Difference (IV, Fixed, 95% CI)	1.33 [1.12, 1.55]
1.23 Histological improvement at study endpoint (continuous), subgrouped by age	5	492	Std. Mean Difference (IV, Random, 95% CI)	1.46 [1.03, 1.89]
1.23.1 Children (18 years and younger)	1	32	Std. Mean Difference (IV, Random, 95% CI)	2.31 [1.36, 3.26]
1.23.2 Mixed children and adults (18 years and older)	4	460	Std. Mean Difference (IV, Random, 95% CI)	1.31 [0.94, 1.67]
1.24 Histological improvement at study endpoint (continuous), subgrouped by type of steroid	5	449	Std. Mean Difference (IV, Random, 95% CI)	1.42 [1.02, 1.82]
1.24.1 Beclomethasone	1	9	Std. Mean Difference (IV, Random, 95% CI)	1.02 [-0.44, 2.47]
1.24.2 Budesonide	4	440	Std. Mean Difference (IV, Random, 95% CI)	1.46 [1.02, 1.91]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.25 Histological improvement at study endpoint (continuous), subgrouped by delivery method	5	449	Std. Mean Difference (IV, Random, 95% CI)	1.42 [1.02, 1.82]
1.25.1 Adapted asthma	2	45	Std. Mean Difference (IV, Random, 95% CI)	1.72 [0.76, 2.67]
1.25.2 Esophageal-specific	3	404	Std. Mean Difference (IV, Random, 95% CI)	1.32 [0.88, 1.75]
1.26 Endoscopic improvement at study endpoint (dichotomous)	3	102	Risk Ratio (M-H, Random, 95% CI)	2.60 [0.82, 8.19]
1.27 Endoscopic improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	3	102	Risk Ratio (M-H, Fixed, 95% CI)	2.73 [1.27, 5.86]
1.28 Endoscopic improvement at study endpoint (dichotomous), sensitivity analysis, validated instruments	2	66	Risk Ratio (M-H, Random, 95% CI)	5.87 [1.11, 31.02]
1.29 Endoscopic improvement at study endpoint (dichotomous), subgrouped by age	3	102	Risk Ratio (M-H, Random, 95% CI)	2.60 [0.82, 8.19]
1.29.1 Children (18 years and younger)	1	36	Risk Ratio (M-H, Random, 95% CI)	1.57 [0.69, 3.58]
1.29.2 Mixed children and adults (18 years and older)	2	66	Risk Ratio (M-H, Random, 95% CI)	5.87 [1.11, 31.02]
1.30 Endoscopic improvement at study endpoint (dichotomous), subgrouped by delivery method	3	102	Risk Ratio (M-H, Random, 95% CI)	2.60 [0.82, 8.19]
1.30.1 Adapted asthma	2	78	Risk Ratio (M-H, Random, 95% CI)	1.78 [0.83, 3.83]
1.30.2 Esophageal-specific	1	24	Risk Ratio (M-H, Random, 95% CI)	11.12 [0.73, 168.69]
1.31 Endoscopic improvement at study endpoint (continuous)	5	596	Std. Mean Difference (IV, Random, 95% CI)	1.33 [0.59, 2.08]
1.32 Endoscopic improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	5	596	Std. Mean Difference (IV, Fixed, 95% CI)	0.93 [0.74, 1.11]
1.33 Endoscopic improvement at study endpoint (continuous), sensitivity analysis, validated instruments	4	572	Std. Mean Difference (IV, Random, 95% CI)	1.31 [0.46, 2.17]
1.34 Endoscopic improvement at study endpoint (continuous), subgrouped by age	5	596	Std. Mean Difference (IV, Random, 95% CI)	1.33 [0.59, 2.08]


Medical treatment of eosinophilic esophagitis (Review)

196

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.34.1 Children (18 years and younger)	1	24	Std. Mean Difference (IV, Random, 95% CI)	1.44 [0.50, 2.38]
1.34.2 Mixed children and adults (18 years and older)	4	572	Std. Mean Difference (IV, Random, 95% CI)	1.31 [0.46, 2.17]
1.35 Endoscopic improvement at study endpoint (continuous), subgrouped by type of steroid	5	596	Std. Mean Difference (IV, Random, 95% CI)	1.33 [0.59, 2.08]
1.35.1 Budesonide	4	493	Std. Mean Difference (IV, Random, 95% CI)	1.41 [0.44, 2.37]
1.35.2 Fluticasone	1	103	Std. Mean Difference (IV, Random, 95% CI)	1.10 [0.58, 1.62]
1.36 Withdrawals due to adverse events	14	1032	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.43, 0.96]
1.37 Withdrawals due to adverse events, sensitivity analysis, fixed-effect	14	1032	Risk Ratio (M-H, Fixed, 95% CI)	0.65 [0.44, 0.94]
1.38 Withdrawals due to adverse events, subgrouped by age	14	1032	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.43, 0.96]
1.38.1 Children (18 years and younger)	3	149	Risk Ratio (M-H, Random, 95% CI)	0.88 [0.28, 2.72]
1.38.2 Mixed children and adults (18 years and older)	11	883	Risk Ratio (M-H, Random, 95% CI)	0.57 [0.36, 0.90]
1.39 Withdrawals due to adverse events, subgrouped by type of steroid	14	1032	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.43, 0.96]
1.39.1 Beclomethasone	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
1.39.2 Budesonide	7	728	Risk Ratio (M-H, Random, 95% CI)	0.80 [0.49, 1.31]
1.39.3 Fluticasone	5	250	Risk Ratio (M-H, Random, 95% CI)	0.42 [0.20, 0.91]
1.39.4 Mometasone	1	36	Risk Ratio (M-H, Random, 95% CI)	0.56 [0.06, 5.63]
1.40 Withdrawals due to adverse events, subgrouped by delivery method	14	1032	Risk Ratio (M-H, Random, 95% CI)	0.64 [0.43, 0.96]
1.40.1 Adapted asthma	5	174	Risk Ratio (M-H, Random, 95% CI)	0.51 [0.12, 2.12]
1.40.2 Esophageal-specific	9	858	Risk Ratio (M-H, Random, 95% CI)	0.68 [0.44, 1.04]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.41 Serious adverse events	14	1032	Risk Ratio (M-H, Random, 95% CI)	0.35 [0.17, 0.73]
1.42 Total adverse events	13	1014	Risk Ratio (M-H, Random, 95% CI)	1.14 [0.94, 1.40]
1.43 Quality of life at study endpoint (continuous)	1	88	Mean Difference (IV, Random, 95% CI)	0.20 [-0.14, 0.54]


Analysis 1.1. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 1: Clinical improvement at study endpoint (dichotomous)



Risk of bias legend


- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 1.2. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 2: Clinical improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect




Risk of bias legend
 (A) Random sequence generation (selection bias)
 (B) Allocation concealment (selection bias)
 (C) Blinding of participants and personnel (performance bias)
 (D) Blinding of outcome assessment (detection bias)
 (E) Incomplete outcome data (attrition bias)
 (F) Selective reporting (reporting bias)
 (G) Other bias

Analysis 1.3. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 3: Clinical improvement at study endpoint (dichotomous), sensitivity analysis, validated instruments



Risk of bias legend
 (A) Random sequence generation (selection bias)
 (B) Allocation concealment (selection bias)
 (C) Blinding of participants and personnel (performance bias)
 (D) Blinding of outcome assessment (detection bias)
 (E) Incomplete outcome data (attrition bias)
 (F) Selective reporting (reporting bias)
 (G) Other bias


Analysis 1.4. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 4: Clinical improvement at study endpoint (dichotomous), subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.5. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 5: Clinical improvement at study endpoint (dichotomous), subgrouped by type of steroid



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.6. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 6: Clinical improvement at study endpoint (dichotomous), subgrouped by delivery method



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.7. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 7: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.8. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 8: Clinical improvement at study endpoint (continuous), sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.9. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 9: Clinical improvement at study endpoint (continuous), sensitivity analysis, validated instruments



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.10. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 10: Clinical improvement at study endpoint (continuous), subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.11. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 11: Clinical improvement at study endpoint (continuous), subgrouped by type of steroid



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.12. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 12: Clinical improvement at study endpoint (continuous), subgrouped by delivery method



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.13. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 13: Histological improvement at study endpoint (dichotomous)



Risk of bias legend


- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 1.14. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 14: Histological improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect




Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Blinding of participants and personnel (performance bias)
(D) Blinding of outcome assessment (detection bias)
(E) Incomplete outcome data (attrition bias)
(F) Selective reporting (reporting bias)
(G) Other bias

Analysis 1.15. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 15: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold < 15 eos/hpf



Risk of bias legend
(A) Random sequence generation (selection bias)
(B) Allocation concealment (selection bias)
(C) Blinding of participants and personnel (performance bias)
(D) Blinding of outcome assessment (detection bias)
(E) Incomplete outcome data (attrition bias)
(F) Selective reporting (reporting bias)
(G) Other bias


Analysis 1.16. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 16: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 6 eos/hpf



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.17. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 17: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 1 eos/hpf



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.18. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 18: Histological improvement at study endpoint (dichotomous), subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.19. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 19: Histological improvement at study endpoint (dichotomous), subgrouped by type of steroid



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.20. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 20: Histological improvement at study endpoint (dichotomous), subgrouped by delivery method



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.21. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 21: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.22. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 22: Histological improvement at study endpoint (continuous), sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.23. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 23: Histological improvement at study endpoint (continuous), subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.24. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 24: Histological improvement at study endpoint (continuous), subgrouped by type of steroid



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.25. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 25: Histological improvement at study endpoint (continuous), subgrouped by delivery method



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.26. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 26: Endoscopic improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.27. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 27: Endoscopic improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.28. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 28: Endoscopic improvement at study endpoint (dichotomous), sensitivity analysis, validated instruments



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.29. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 29: Endoscopic improvement at study endpoint (dichotomous), subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.30. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 30: Endoscopic improvement at study endpoint (dichotomous), subgrouped by delivery method



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.31. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 31: Endoscopic improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.32. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 32: Endoscopic improvement at study endpoint (continuous), sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.33. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 33: Endoscopic improvement at study endpoint (continuous), sensitivity analysis, validated instruments



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.34. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 34: Endoscopic improvement at study endpoint (continuous), subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.35. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 35: Endoscopic improvement at study endpoint (continuous), subgrouped by type of steroid



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.36. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 36: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.37. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 37: Withdrawals due to adverse events, sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.38. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 38: Withdrawals due to adverse events, subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.39. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 39: Withdrawals due to adverse events, subgrouped by type of steroid



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.40. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 40: Withdrawals due to adverse events, subgrouped by delivery method



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.41. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 41: Serious adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 1.42. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 42: Total adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 1.43. Comparison 1: Corticosteroids vs placebo for induction of remission, Outcome 43: Quality of life at study endpoint (continuous)



Risk of bias legend


- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 2. Corticosteroids vs placebo for maintenance of remission


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.1 Clinical improvement at study endpoint (dichotomous)	2	252	Risk Ratio (M-H, Random, 95% CI)	2.17 [0.75, 6.27]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
2.2 Clinical improvement at study endpoint (continuous)	3	269	Std. Mean Difference (IV, Random, 95% CI)	0.51 [-0.49, 1.52]
2.3 Histological improvement at study endpoint (dichotomous)	3	280	Risk Ratio (M-H, Random, 95% CI)	4.58 [1.66, 12.62]
2.4 Histological improvement at study endpoint (continuous)	3	269	Std. Mean Difference (IV, Random, 95% CI)	1.26 [0.74, 1.78]
2.5 Endoscopic improvement at study endpoint (continuous)	2	240	Std. Mean Difference (IV, Random, 95% CI)	1.34 [-0.27, 2.95]
2.6 Withdrawals due to adverse events	3	280	Risk Ratio (M-H, Random, 95% CI)	0.37 [0.16, 0.87]
2.7 Serious adverse events	3	280	Risk Ratio (M-H, Random, 95% CI)	1.27 [0.09, 18.03]
2.8 Total adverse events	3	280	Risk Ratio (M-H, Random, 95% CI)	1.10 [0.75, 1.62]
2.9 Quality of life at study endpoint (continuous)	1	204	Mean Difference (IV, Random, 95% CI)	0.60 [0.40, 0.80]

Analysis 2.1. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 1: Clinical improvement at study endpoint (dichotomous)




Analysis 2.2. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 2: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 2.3. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 3: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 2.4. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 4: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 2.5. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 5: Endoscopic improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 2.6. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 6: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 2.7. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 7: Serious adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 2.8. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 8: Total adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 2.9. Comparison 2: Corticosteroids vs placebo for maintenance of remission, Outcome 9: Quality of life at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 3. Biologics vs placebo for induction of remission

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
3.1 Clinical improvement at study endpoint (dichotomous)	5	410	Risk Ratio (M-H, Random, 95% CI)	1.14 [0.85, 1.52]
3.2 Clinical improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	5	410	Risk Ratio (M-H, Fixed, 95% CI)	1.10 [0.92, 1.32]
3.3 Clinical improvement at study endpoint (dichotomous), sensitivity analysis, validated instruments	3	172	Risk Ratio (M-H, Random, 95% CI)	1.37 [1.02, 1.85]


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
3.4 Clinical improvement at study endpoint (dichotomous), subgrouped by age	5	410	Risk Ratio (M-H, Random, 95% CI)	1.14 [0.85, 1.52]
3.4.1 Children (18 years and younger)	1	227	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.72, 1.14]
3.4.2 Mixed children and adults (18 years and older)	4	183	Risk Ratio (M-H, Random, 95% CI)	1.36 [1.02, 1.81]
3.5 Clinical improvement at study endpoint (dichotomous), subgrouped by mechanism	5	410	Risk Ratio (M-H, Random, 95% CI)	1.14 [0.85, 1.52]
3.5.1 Anti-IL-13 (RPC4046, alias cendakimab; QAX576, alias dectrekumab) and anti-IL-4r (dupilumab)	3	172	Risk Ratio (M-H, Random, 95% CI)	1.37 [1.02, 1.85]
3.5.2 Anti-IL-5 (mepolizumab, reslizumab)	2	238	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.73, 1.15]
3.6 Clinical improvement at study endpoint (continuous)	7	387	Std. Mean Difference (IV, Random, 95% CI)	0.50 [0.22, 0.78]
3.7 Clinical improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	7	387	Std. Mean Difference (IV, Fixed, 95% CI)	0.48 [0.28, 0.69]
3.8 Clinical improvement at study endpoint (continuous), sensitivity analysis, peer-reviewed manuscripts	5	162	Std. Mean Difference (IV, Random, 95% CI)	0.36 [-0.09, 0.81]
3.9 Clinical improvement at study endpoint (continuous), sensitivity analysis, less than high risk of bias	6	357	Std. Mean Difference (IV, Random, 95% CI)	0.61 [0.40, 0.82]
3.10 Clinical improvement at study endpoint (continuous), sensitivity analysis, validated instruments	3	247	Std. Mean Difference (IV, Random, 95% CI)	0.62 [0.37, 0.88]
3.11 Clinical improvement at study endpoint (continuous), subgrouped by mechanism	7	387	Std. Mean Difference (IV, Random, 95% CI)	0.40 [0.05, 0.74]
3.11.1 Anti-IgE (omalizumab)	1	30	Std. Mean Difference (IV, Random, 95% CI)	-0.47 [-1.20, 0.26]
3.11.2 Anti-IL-13 (RPC4046, alias cendakimab; QAX576, alias dectrekumab) and anti-IL-4r (dupilumab)	5	346	Std. Mean Difference (IV, Random, 95% CI)	0.60 [0.39, 0.82]
3.11.3 Anti-IL-5 (mepolizumab, reslizumab)	1	11	Std. Mean Difference (IV, Random, 95% CI)	-0.79 [-2.05, 0.46]
3.12 Histological improvement at study endpoint (dichotomous)	8	925	Risk Ratio (M-H, Random, 95% CI)	6.73 [2.58, 17.52]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
3.13 Histological improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect	8	925	Risk Ratio (M-H, Fixed, 95% CI)	5.12 [3.86, 6.78]
3.14 Histological improvement at study endpoint (dichotomous), sensitivity analysis, peer-reviewed manuscripts	6	685	Risk Ratio (M-H, Random, 95% CI)	6.13 [1.67, 22.51]
3.15 Histological improvement at study endpoint (dichotomous), sensitivity analysis, less than high risk of bias	8	925	Risk Ratio (M-H, Random, 95% CI)	6.73 [2.58, 17.52]
3.16 Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold < 15 eos/hpf	4	418	Risk Ratio (M-H, Random, 95% CI)	5.61 [1.00, 31.32]
3.17 Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 6 eos/hpf	6	674	Risk Ratio (M-H, Random, 95% CI)	8.85 [5.73, 13.67]
3.18 Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 1 eos/hpf	2	323	Risk Ratio (M-H, Random, 95% CI)	18.01 [7.24, 44.83]
3.19 Histological improvement at study endpoint (dichotomous), subgrouped by age	8	925	Risk Ratio (M-H, Random, 95% CI)	7.18 [2.93, 17.56]
3.19.1 Children (18 years and younger)	2	277	Risk Ratio (M-H, Random, 95% CI)	4.51 [0.40, 50.36]
3.19.2 Mixed children and adults (18 years and older)	6	423	Risk Ratio (M-H, Random, 95% CI)	9.01 [4.88, 16.62]
3.19.3 Adults only (over 18 years)	1	225	Risk Ratio (M-H, Random, 95% CI)	7.44 [4.02, 13.77]
3.20 Histological improvement at study endpoint (dichotomous), subgrouped by mechanism	8	925	Risk Ratio (M-H, Random, 95% CI)	6.73 [2.58, 17.52]
3.20.1 Anti-sialic acid-binding Ig-like lectin 8 (lirentelimab)	1	276	Risk Ratio (M-H, Random, 95% CI)	8.30 [4.61, 14.93]
3.20.2 Anti-IL-13 (RPC4046, alias cendakimab; QAX576, alias dectrekumab) and anti-IL-4r (dupilumab)	5	412	Risk Ratio (M-H, Random, 95% CI)	9.01 [4.88, 16.62]
3.20.3 Anti-IL-5 (mepolizumab, reslizumab)	2	237	Risk Ratio (M-H, Random, 95% CI)	1.75 [1.21, 2.55]
3.21 Histological improvement at study endpoint (continuous)	6	370	Std. Mean Difference (IV, Random, 95% CI)	1.01 [0.36, 1.66]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
3.22 Histological improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	6	370	Std. Mean Difference (IV, Fixed, 95% CI)	1.25 [1.01, 1.49]
3.23 Histological improvement at study endpoint (continuous), sensitivity analysis, less than high risk of bias	5	340	Std. Mean Difference (IV, Random, 95% CI)	1.39 [1.01, 1.77]
3.24 Histological improvement at study endpoint (continuous), subgrouped by age	6	370	Std. Mean Difference (IV, Random, 95% CI)	1.01 [0.36, 1.66]
3.24.1 Children (18 years and younger)	1	169	Std. Mean Difference (IV, Random, 95% CI)	1.64 [1.25, 2.02]
3.24.2 Mixed children and adults (18 years and older)	5	201	Std. Mean Difference (IV, Random, 95% CI)	0.85 [0.04, 1.66]
3.25 Histological improvement at study endpoint (continuous), subgrouped by mechanism	6	370	Std. Mean Difference (IV, Random, 95% CI)	1.01 [0.36, 1.66]
3.25.1 Anti-IgE (omalizumab)	1	30	Std. Mean Difference (IV, Random, 95% CI)	-0.43 [-1.15, 0.30]
3.25.2 Anti-IL-13 (RPC4046, alias cendakimab; QAX576, alias dectrekumab) and anti-IL-4r (dupilumab)	3	160	Std. Mean Difference (IV, Random, 95% CI)	1.32 [0.72, 1.91]
3.25.3 Anti-IL-5 (mepolizumab, reslizumab)	2	180	Std. Mean Difference (IV, Random, 95% CI)	1.42 [0.69, 2.15]
3.26 Endoscopic improvement at study endpoint (dichotomous)	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
3.27 Endoscopic improvement at study endpoint (continuous)	3	197	Std. Mean Difference (IV, Random, 95% CI)	2.79 [0.36, 5.22]
3.28 Endoscopic improvement at study endpoint (continuous), sensitivity analysis, fixed-effect	3	197	Std. Mean Difference (IV, Fixed, 95% CI)	1.20 [0.86, 1.55]
3.29 Endoscopic improvement at study endpoint (continuous), sensitivity analysis, less than high risk of bias	2	136	Std. Mean Difference (IV, Random, 95% CI)	0.82 [0.42, 1.21]
3.30 Withdrawals due to adverse events	8	792	Risk Ratio (M-H, Random, 95% CI)	1.55 [0.88, 2.74]
3.31 Withdrawals due to adverse events, sensitivity analysis, fixed-effect	8	792	Risk Ratio (M-H, Fixed, 95% CI)	1.53 [0.89, 2.64]
3.32 Withdrawals due to adverse events, sensitivity analysis, less than high risk of bias	6	681	Risk Ratio (M-H, Random, 95% CI)	1.55 [0.88, 2.74]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
3.33 Withdrawals due to adverse events, sensitivity analysis, peer reviewed manuscripts	7	711	Risk Ratio (M-H, Random, 95% CI)	1.55 [0.88, 2.74]
3.34 Withdrawals due to adverse events, subgrouped by age	8	792	Risk Ratio (M-H, Random, 95% CI)	1.55 [0.88, 2.74]
3.34.1 Children (18 years and younger)	1	227	Risk Ratio (M-H, Random, 95% CI)	1.45 [0.63, 3.35]
3.34.2 Mixed children and adults (18 years and older)	7	565	Risk Ratio (M-H, Random, 95% CI)	1.59 [0.68, 3.74]
3.35 Withdrawals due to adverse events, subgrouped by mechanism	8	792	Risk Ratio (M-H, Random, 95% CI)	1.31 [0.66, 2.59]
3.35.1 Anti-IgE (omalizumab)	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
3.35.2 Anti-sialic acid-binding Ig-like lectin 8 (lirentelimab)	1	276	Risk Ratio (M-H, Random, 95% CI)	2.12 [0.74, 6.13]
3.35.3 Anti-IL-13 (RPC4046, alias cendakimab; QAX576, alias dectrekumab) and anti-IL-4r (dupilumab)	4	248	Risk Ratio (M-H, Random, 95% CI)	0.91 [0.36, 2.33]
3.35.4 Anti-IL-5 (mepolizumab, reslizumab)	2	238	Risk Ratio (M-H, Random, 95% CI)	1.02 [0.04, 24.63]
3.36 Serious adverse events	6	685	Risk Ratio (M-H, Random, 95% CI)	0.70 [0.25, 1.97]
3.37 Total adverse events	6	685	Risk Ratio (M-H, Random, 95% CI)	1.07 [0.94, 1.23]
3.38 Quality of life at study endpoint (continuous)	1	47	Mean Difference (IV, Random, 95% CI)	0.33 [-0.06, 0.72]


Analysis 3.1. Comparison 3: Biologics vs placebo for induction of remission, Outcome 1: Clinical improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.2. Comparison 3: Biologics vs placebo for induction of remission, Outcome 2: Clinical improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.3. Comparison 3: Biologics vs placebo for induction of remission, Outcome 3: Clinical improvement at study endpoint (dichotomous), sensitivity analysis, validated instruments



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.4. Comparison 3: Biologics vs placebo for induction of remission, Outcome 4: Clinical improvement at study endpoint (dichotomous), subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.5. Comparison 3: Biologics vs placebo for induction of remission, Outcome 5: Clinical improvement at study endpoint (dichotomous), subgrouped by mechanism



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.6. Comparison 3: Biologics vs placebo for induction of remission, Outcome 6: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.7. Comparison 3: Biologics vs placebo for induction of remission, Outcome 7: Clinical improvement at study endpoint (continuous), sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.8. Comparison 3: Biologics vs placebo for induction of remission, Outcome 8: Clinical improvement at study endpoint (continuous), sensitivity analysis, peer-reviewed manuscripts



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.9. Comparison 3: Biologics vs placebo for induction of remission, Outcome 9: Clinical improvement at study endpoint (continuous), sensitivity analysis, less than high risk of bias



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.10. Comparison 3: Biologics vs placebo for induction of remission, Outcome 10: Clinical improvement at study endpoint (continuous), sensitivity analysis, validated instruments



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.11. Comparison 3: Biologics vs placebo for induction of remission, Outcome 11: Clinical improvement at study endpoint (continuous), subgrouped by mechanism



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.12. Comparison 3: Biologics vs placebo for induction of remission, Outcome 12: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.13. Comparison 3: Biologics vs placebo for induction of remission, Outcome 13: Histological improvement at study endpoint (dichotomous), sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.14. Comparison 3: Biologics vs placebo for induction of remission, Outcome 14: Histological improvement at study endpoint (dichotomous), sensitivity analysis, peer-reviewed manuscripts



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.15. Comparison 3: Biologics vs placebo for induction of remission, Outcome 15: Histological improvement at study endpoint (dichotomous), sensitivity analysis, less than high risk of bias



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.16. Comparison 3: Biologics vs placebo for induction of remission, Outcome 16: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold < 15 eos/hpf



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.17. Comparison 3: Biologics vs placebo for induction of remission, Outcome 17: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 6 eos/hpf



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.18. Comparison 3: Biologics vs placebo for induction of remission, Outcome 18: Histological improvement at study endpoint (dichotomous), sensitivity analysis, threshold ≤ 1 eos/hpf



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.19. Comparison 3: Biologics vs placebo for induction of remission, Outcome 19: Histological improvement at study endpoint (dichotomous), subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.20. Comparison 3: Biologics vs placebo for induction of remission, Outcome 20: Histological improvement at study endpoint (dichotomous), subgrouped by mechanism



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.21. Comparison 3: Biologics vs placebo for induction of remission, Outcome 21: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.22. Comparison 3: Biologics vs placebo for induction of remission, Outcome 22: Histological improvement at study endpoint (continuous), sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.23. Comparison 3: Biologics vs placebo for induction of remission, Outcome 23: Histological improvement at study endpoint (continuous), sensitivity analysis, less than high risk of bias



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.24. Comparison 3: Biologics vs placebo for induction of remission, Outcome 24: Histological improvement at study endpoint (continuous), subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.25. Comparison 3: Biologics vs placebo for induction of remission, Outcome 25: Histological improvement at study endpoint (continuous), subgrouped by mechanism



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.26. Comparison 3: Biologics vs placebo for induction of remission, Outcome 26: Endoscopic improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.27. Comparison 3: Biologics vs placebo for induction of remission, Outcome 27: Endoscopic improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.28. Comparison 3: Biologics vs placebo for induction of remission, Outcome 28: Endoscopic improvement at study endpoint (continuous), sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.29. Comparison 3: Biologics vs placebo for induction of remission, Outcome 29: Endoscopic improvement at study endpoint (continuous), sensitivity analysis, less than high risk of bias



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.30. Comparison 3: Biologics vs placebo for induction of remission, Outcome 30: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.31. Comparison 3: Biologics vs placebo for induction of remission, Outcome 31: Withdrawals due to adverse events, sensitivity analysis, fixed-effect



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.32. Comparison 3: Biologics vs placebo for induction of remission, Outcome 32: Withdrawals due to adverse events, sensitivity analysis, less than high risk of bias



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.33. Comparison 3: Biologics vs placebo for induction of remission, Outcome 33: Withdrawals due to adverse events, sensitivity analysis, peer reviewed manuscripts



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.34. Comparison 3: Biologics vs placebo for induction of remission, Outcome 34: Withdrawals due to adverse events, subgrouped by age



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.35. Comparison 3: Biologics vs placebo for induction of remission, Outcome 35: Withdrawals due to adverse events, subgrouped by mechanism



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.36. Comparison 3: Biologics vs placebo for induction of remission, Outcome 36: Serious adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 3.37. Comparison 3: Biologics vs placebo for induction of remission, Outcome 37: Total adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 3.38. Comparison 3: Biologics vs placebo for induction of remission, Outcome 38: Quality of life at study endpoint (continuous)




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 4. Cromolyn sodium vs placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
4.1 Clinical improvement at study endpoint (continuous)	1	14	Mean Difference (IV, Random, 95% CI)	4.70 [-12.09, 21.49]
4.2 Histological improvement at study endpoint (continuous)	1	15	Mean Difference (IV, Random, 95% CI)	14.20 [-36.90, 65.30]
4.3 Withdrawals due to adverse events	1	16	Risk Ratio (M-H, Random, 95% CI)	0.27 [0.01, 5.70]
4.4 Serious adverse events	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable


Analysis 4.1. Comparison 4: Cromolyn sodium vs placebo, Outcome 1: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 4.2. Comparison 4: Cromolyn sodium vs placebo, Outcome 2: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 4.3. Comparison 4: Cromolyn sodium vs placebo, Outcome 3: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 4.4. Comparison 4: Cromolyn sodium vs placebo, Outcome 4: Serious adverse events




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 5. PGD2R antagonist OC000459 vs placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
5.1 Clinical improvement at study endpoint (continuous)	1	26	Mean Difference (IV, Random, 95% CI)	-1.06 [-6.80, 4.68]
5.2 Histological improvement at study endpoint (continuous)	1	26	Mean Difference (IV, Random, 95% CI)	26.21 [-23.78, 76.20]
5.3 Endoscopic improvement at study endpoint (continuous)	1	26	Mean Difference (IV, Random, 95% CI)	-0.49 [-2.05, 1.07]
5.4 Withdrawals due to adverse events	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
5.5 Serious adverse events	1	26	Risk Ratio (M-H, Random, 95% CI)	2.60 [0.12, 58.48]


Analysis 5.1. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 1: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 5.2. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 2: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 5.3. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 3: Endoscopic improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 5.4. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 4: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 5.5. Comparison 5: PGD2R antagonist OC000459 vs placebo, Outcome 5: Serious adverse events




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 6. Swallowed fluticasone vs oral prednisone

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
6.1 Clinical improvement at study endpoint (dichotomous)	1	80	Risk Ratio (M-H, Random, 95% CI)	1.09 [0.90, 1.33]
6.2 Histological improvement at study endpoint (dichotomous)	1	80	Risk Ratio (M-H, Random, 95% CI)	1.10 [0.87, 1.38]
6.3 Histological improvement at study endpoint (continuous)	1	68	Mean Difference (IV, Random, 95% CI)	-4.45 [-9.08, 0.18]
6.4 Endoscopic improvement at study endpoint (dichotomous)	1	80	Risk Ratio (M-H, Random, 95% CI)	1.13 [0.91, 1.41]
6.5 Withdrawals due to adverse events	1	80	Risk Ratio (M-H, Random, 95% CI)	0.50 [0.16, 1.53]
6.6 Serious adverse events	1	80	Risk Ratio (M-H, Random, 95% CI)	0.14 [0.01, 2.68]
6.7 Total adverse events	1	80	Risk Ratio (M-H, Random, 95% CI)	0.38 [0.16, 0.86]


Analysis 6.1. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 1: Clinical improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 6.2. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 2: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 6.3. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 3: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 6.4. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 4: Endoscopic improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 6.5. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 5: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 6.6. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 6: Serious adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 6.7. Comparison 6: Swallowed fluticasone vs oral prednisone, Outcome 7: Total adverse events




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 7. Oral viscous budesonide vs swallowed fluticasone

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
7.1 Clinical improvement at study endpoint (continuous)	1	84	Mean Difference (IV, Random, 95% CI)	-0.60 [-3.78, 2.58]
7.2 Histological improvement at study endpoint (dichotomous)	1	129	Risk Ratio (M-H, Random, 95% CI)	1.13 [0.84, 1.51]
7.3 Histological improvement at study endpoint (continuous)	1	111	Mean Difference (IV, Random, 95% CI)	6.20 [-5.63, 18.03]
7.4 Endoscopic improvement at study endpoint (continuous)	1	111	Mean Difference (IV, Random, 95% CI)	0.70 [-0.03, 1.43]
7.5 Withdrawals due to adverse events	1	129	Risk Ratio (M-H, Random, 95% CI)	0.98 [0.42, 2.32]
7.6 Serious adverse events	1	129	Risk Ratio (M-H, Random, 95% CI)	0.33 [0.01, 7.91]
7.7 Total adverse events	1	129	Risk Ratio (M-H, Random, 95% CI)	0.66 [0.32, 1.35]


Analysis 7.1. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 1: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 7.2. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 2: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 7.3. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 3: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 7.4. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 4: Endoscopic improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 7.5. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 5: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 7.6. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 6: Serious adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 7.7. Comparison 7: Oral viscous budesonide vs swallowed fluticasone, Outcome 7: Total adverse events




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 8. Esomeprazole vs fluticasone

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
8.1 Clinical improvement at study endpoint (continuous)	2	63	Std. Mean Difference (IV, Random, 95% CI)	0.32 [-0.88, 1.52]
8.2 Histological improvement at study endpoint (dichotomous)	2	72	Risk Ratio (M-H, Random, 95% CI)	1.62 [0.77, 3.41]
8.3 Histological improvement at study endpoint (continuous)	2	67	Std. Mean Difference (IV, Random, 95% CI)	0.28 [-0.20, 0.76]
8.4 Withdrawals due to adverse events	2	72	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.07, 13.38]
8.5 Serious adverse events	2	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
8.6 Total adverse events	2	72	Risk Ratio (M-H, Random, 95% CI)	0.14 [0.01, 2.61]


Analysis 8.1. Comparison 8: Esomeprazole vs fluticasone, Outcome 1: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 8.2. Comparison 8: Esomeprazole vs fluticasone, Outcome 2: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 8.3. Comparison 8: Esomeprazole vs fluticasone, Outcome 3: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 8.4. Comparison 8: Esomeprazole vs fluticasone, Outcome 4: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 8.5. Comparison 8: Esomeprazole vs fluticasone, Outcome 5: Serious adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 8.6. Comparison 8: Esomeprazole vs fluticasone, Outcome 6: Total adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Comparison 9. One-food elimination diet vs four-food elimination diet

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
9.1 Clinical improvement at study endpoint (continuous)	1	50	Mean Difference (IV, Random, 95% CI)	-7.50 [-16.28, 1.28]
9.2 Histological improvement at study endpoint (dichotomous)	1	63	Risk Ratio (M-H, Random, 95% CI)	2.26 [1.15, 4.43]
9.3 Endoscopic improvement at study endpoint (continuous)	1	34	Mean Difference (IV, Random, 95% CI)	-0.60 [-2.15, 0.95]
9.4 Withdrawals due to adverse events	1	63	Risk Ratio (M-H, Random, 95% CI)	0.33 [0.11, 0.98]

Medical treatment of eosinophilic esophagitis (Review)

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
9.5 Serious adverse events	1	63	Risk Ratio (M-H, Random, 95% CI)	0.66 [0.04, 10.04]
9.6 Total adverse events	1	63	Risk Ratio (M-H, Random, 95% CI)	0.41 [0.15, 1.11]
9.7 Quality of life at study endpoint (continuous)	1	63	Mean Difference (IV, Random, 95% CI)	0.10 [-6.49, 6.69]


Analysis 9.1. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 1: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 9.2. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 2: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 9.3. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 3: Endoscopic improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 9.4. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 4: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 9.5. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 5: Serious adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 9.6. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 6: Total adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 9.7. Comparison 9: One-food elimination diet vs four-food elimination diet, Outcome 7: Quality of life at study endpoint (continuous)




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 10. One-food elimination diet vs six-food elimination diet

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
10.1 Clinical improvement at study endpoint (continuous)	1	129	Mean Difference (IV, Random, 95% CI)	-5.20 [-11.06, 0.66]
10.2 Histological improvement at study endpoint (dichotomous)	1	129	Risk Ratio (M-H, Random, 95% CI)	0.85 [0.54, 1.33]
10.3 Histological improvement at study endpoint (continuous)	1	129	Mean Difference (IV, Random, 95% CI)	6.80 [-10.40, 24.00]
10.4 Endoscopic improvement at study endpoint (continuous)	1	129	Mean Difference (IV, Random, 95% CI)	-0.42 [-1.67, 0.83]
10.5 Withdrawals due to adverse events	1	129	Risk Ratio (M-H, Random, 95% CI)	0.62 [0.11, 3.57]
10.6 Serious adverse events	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
10.7 Total adverse events	1	129	Risk Ratio (M-H, Random, 95% CI)	0.46 [0.04, 4.98]
10.8 Quality of life at study endpoint (continuous)	1	129	Mean Difference (IV, Random, 95% CI)	0.57 [-3.25, 4.39]


Analysis 10.1. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 1: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 10.2. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 2: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 10.3. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 3: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 10.4. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 4: Endoscopic improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 10.5. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 5: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 10.6. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 6: Serious adverse events

Study or Subgroup	1-FED		6-FED		Weight	Risk Ratio		Risk Ratio		Risk of Bias						
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI	A	B	C	D	E	F	G		
Kliewer 2021	0	67	0	62		Not estimable				+	+	-	+	+	+	?
Total (95% CI)		0		0		Not estimable										
Total events:	0		0													
Heterogeneity: Not applicable																
Test for overall effect: Not applicable																
Test for subgroup differences: Not applicable																

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 10.7. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 7: Total adverse events

Study or Subgroup	1-FED		6-FED		Weight	Risk Ratio		Risk Ratio		Risk of Bias						
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI	A	B	C	D	E	F	G		
Kliewer 2021	1	67	2	62	100.0%	0.46 [0.04 , 4.98]				+	+	-	+	+	+	?
Total (95% CI)		67		62	100.0%	0.46 [0.04 , 4.98]										
Total events:	1		2													
Heterogeneity: Not applicable																
Test for overall effect: Z = 0.64 (P = 0.52)																
Test for subgroup differences: Not applicable																

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 10.8. Comparison 10: One-food elimination diet vs six-food elimination diet, Outcome 8: Quality of life at study endpoint (continuous)




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 11. Four-food elimination diet with omeprazole vs omeprazole

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
11.1 Histological improvement at study endpoint (dichotomous)	1	64	Risk Ratio (M-H, Random, 95% CI)	1.57 [0.99, 2.48]
11.2 Histological improvement at study endpoint (continuous)	1	58	Mean Difference (IV, Random, 95% CI)	9.50 [-11.18, 30.18]
11.3 Withdrawals due to adverse events	1	64	Risk Ratio (M-H, Random, 95% CI)	5.00 [0.62, 40.44]


Analysis 11.1. Comparison 11: Four-food elimination diet with omeprazole vs omeprazole, Outcome 1: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 11.2. Comparison 11: Four-food elimination diet with omeprazole vs omeprazole, Outcome 2: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 11.3. Comparison 11: Four-food elimination diet with omeprazole vs omeprazole, Outcome 3: Withdrawals due to adverse events



Risk of bias legend


- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 12. Four-food elimination diet with amino acid formula vs four-food elimination diet

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
12.1 Clinical improvement at study endpoint (continuous)	1	41	Mean Difference (IV, Random, 95% CI)	-0.50 [-2.41, 1.41]
12.2 Histological improvement at study endpoint (dichotomous)	1	41	Risk Ratio (M-H, Random, 95% CI)	1.90 [0.79, 4.60]
12.3 Histological improvement at study endpoint (continuous)	1	41	Mean Difference (IV, Random, 95% CI)	13.80 [-9.50, 37.10]
12.4 Endoscopic improvement at study endpoint (continuous)	1	41	Mean Difference (IV, Random, 95% CI)	-1.00 [-2.83, 0.83]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
12.5 Withdrawals due to adverse events	1	41	Risk Ratio (M-H, Random, 95% CI)	0.95 [0.06, 14.22]
12.6 Serious adverse events	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
12.7 Total adverse events	1	41	Risk Ratio (M-H, Random, 95% CI)	2.86 [0.12, 66.44]


Analysis 12.1. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 1: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 12.2. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 2: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 12.3. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 3: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 12.4. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 4: Endoscopic improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 12.5. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 5: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 12.6. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 6: Serious adverse events

Study or Subgroup	4-FED + AAF		4-FED		Weight	Risk Ratio	Risk Ratio	Risk of Bias						
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI	A	B	C	D	E	F	G
De Rooij 2022	0	21	0	20		Not estimable		?	+	?	?	+	+	+
Total (95% CI)		0		0		Not estimable								
Total events:	0		0											
Heterogeneity: Not applicable														
Test for overall effect: Not applicable														
Test for subgroup differences: Not applicable														

Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 12.7. Comparison 12: Four-food elimination diet with amino acid formula vs four-food elimination diet, Outcome 7: Total adverse events

Study or Subgroup	4-FED + AAF		4-FED		Weight	Risk Ratio	Risk Ratio	Risk of Bias						
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI	A	B	C	D	E	F	G
De Rooij 2022	1	21	0	20	100.0%	2.86 [0.12, 66.44]		?	+	?	?	+	+	+
Total (95% CI)		21		20	100.0%	2.86 [0.12, 66.44]								
Total events:	1		0											
Heterogeneity: Not applicable														
Test for overall effect: Z = 0.66 (P = 0.51)														
Test for subgroup differences: Not applicable														

Risk of bias legend


- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 13. Nebulized budesonide vs viscous budesonide

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
13.1 Clinical improvement at study endpoint (continuous)	1	22	Mean Difference (IV, Random, 95% CI)	-6.00 [-18.30, 6.30]
13.2 Histological improvement at study endpoint (continuous)	1	22	Mean Difference (IV, Random, 95% CI)	78.00 [20.81, 135.19]
13.3 Serious adverse events	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
13.4 Total adverse events	1	25	Risk Ratio (M-H, Random, 95% CI)	0.92 [0.15, 5.56]


Analysis 13.1. Comparison 13: Nebulized budesonide vs viscous budesonide, Outcome 1: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 13.2. Comparison 13: Nebulized budesonide vs viscous budesonide, Outcome 2: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 13.3. Comparison 13: Nebulized budesonide vs viscous budesonide, Outcome 3: Serious adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 13.4. Comparison 13: Nebulized budesonide vs viscous budesonide, Outcome 4: Total adverse events



Risk of bias legend


- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 14. Viaskin milk patch vs placebo

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
14.1 Clinical improvement at study endpoint (continuous)	1	9	Mean Difference (IV, Random, 95% CI)	1.29 [-0.83, 3.41]
14.2 Histological improvement at study endpoint (continuous)	1	9	Mean Difference (IV, Random, 95% CI)	69.43 [-21.75, 160.61]
14.3 Endoscopic improvement at study endpoint (continuous)	1	20	Mean Difference (IV, Random, 95% CI)	-0.33 [-2.00, 1.34]
14.4 Withdrawals due to adverse events	1	20	Risk Ratio (M-H, Random, 95% CI)	1.12 [0.05, 23.99]
14.5 Serious adverse events	1	20	Risk Ratio (M-H, Random, 95% CI)	0.13 [0.01, 2.67]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
14.6 Total adverse events	1	20	Risk Ratio (M-H, Random, 95% CI)	1.00 [0.77, 1.29]
14.7 Quality of life at study endpoint (continuous)	1	9	Mean Difference (IV, Random, 95% CI)	13.60 [-16.12, 43.32]


Analysis 14.1. Comparison 14: Viaskin milk patch vs placebo, Outcome 1: Clinical improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 14.2. Comparison 14: Viaskin milk patch vs placebo, Outcome 2: Histological improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 14.3. Comparison 14: Viaskin milk patch vs placebo, Outcome 3: Endoscopic improvement at study endpoint (continuous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 14.4. Comparison 14: Viaskin milk patch vs placebo, Outcome 4: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 14.5. Comparison 14: Viaskin milk patch vs placebo, Outcome 5: Serious adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 14.6. Comparison 14: Viaskin milk patch vs placebo, Outcome 6: Total adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 14.7. Comparison 14: Viaskin milk patch vs placebo, Outcome 7: Quality of life at study endpoint (continuous)




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 15. Leukotriene receptor antagonist vs placebo for maintenance of remission

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
15.1 Clinical improvement at study endpoint (dichotomous)	1	41	Risk Ratio (M-H, Random, 95% CI)	1.68 [0.66, 4.28]
15.2 Withdrawals due to adverse events	1	41	Risk Ratio (M-H, Random, 95% CI)	2.10 [0.21, 21.39]
15.3 Serious adverse events	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable
15.4 Total adverse events	1	0	Risk Ratio (M-H, Random, 95% CI)	Not estimable


Analysis 15.1. Comparison 15: Leukotriene receptor antagonist vs placebo for maintenance of remission, Outcome 1: Clinical improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias


Analysis 15.2. Comparison 15: Leukotriene receptor antagonist vs placebo for maintenance of remission, Outcome 2: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 15.3. Comparison 15: Leukotriene receptor antagonist vs placebo for maintenance of remission, Outcome 3: Serious adverse events




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 15.4. Comparison 15: Leukotriene receptor antagonist vs placebo for maintenance of remission, Outcome 4: Total adverse events

Study or Subgroup	Leukotriene receptor antagonist		Placebo		Weight	Risk Ratio	Risk Ratio	Risk of Bias						
	Events	Total	Events	Total		M-H, Random, 95% CI	M-H, Random, 95% CI	A	B	C	D	E	F	G
Alexander 2017	0	20	0	21		Not estimable		+	+	+	+	+	?	+
Total (95% CI)		0		0		Not estimable								
Total events: 0														
Heterogeneity: Not applicable														
Test for overall effect: Not applicable														
Test for subgroup differences: Not applicable														




Risk of bias legend
 (A) Random sequence generation (selection bias)
 (B) Allocation concealment (selection bias)
 (C) Blinding of participants and personnel (performance bias)
 (D) Blinding of outcome assessment (detection bias)
 (E) Incomplete outcome data (attrition bias)
 (F) Selective reporting (reporting bias)
 (G) Other bias

Comparison 16. Mepolizumab 10 mg/kg vs mepolizumab 0.55 mg/kg

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
16.1 Histological improvement at study endpoint (dichotomous)	1	39	Risk Ratio (IV, Random, 95% CI)	1.19 [0.37, 3.77]
16.2 Withdrawals due to adverse events	1	39	Risk Ratio (IV, Random, 95% CI)	0.63 [0.12, 3.38]


Analysis 16.1. Comparison 16: Mepolizumab 10 mg/kg vs mepolizumab 0.55 mg/kg, Outcome 1: Histological improvement at study endpoint (dichotomous)

Study or Subgroup	Mepolizumab 10 mg/kg		Mepolizumab 0.55 mg/kg		Weight	Risk Ratio	Risk Ratio	Risk of Bias						
	Events	Total	Events	Total		IV, Random, 95% CI	IV, Random, 95% CI	A	B	C	D	E	F	G
Assa'ad 2011	5	20	4	19	100.0%	1.19 [0.37, 3.77]		?	+	+	+	?	+	+
Total (95% CI)		20		19	100.0%	1.19 [0.37, 3.77]								
Total events: 5														
Heterogeneity: Not applicable														
Test for overall effect: Z = 0.29 (P = 0.77)														
Test for subgroup differences: Not applicable														



Risk of bias legend
 (A) Random sequence generation (selection bias)
 (B) Allocation concealment (selection bias)
 (C) Blinding of participants and personnel (performance bias)
 (D) Blinding of outcome assessment (detection bias)
 (E) Incomplete outcome data (attrition bias)
 (F) Selective reporting (reporting bias)
 (G) Other bias

Analysis 16.2. Comparison 16: Mepolizumab 10 mg/kg vs mepolizumab 0.55 mg/kg, Outcome 2: Withdrawals due to adverse events




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 17. Mepolizumab 2.5 mg/kg vs mepolizumab 0.55 mg/kg

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
17.1 Histological improvement at study endpoint (dichotomous)	1	39	Risk Ratio (IV, Random, 95% CI)	2.14 [0.79, 5.79]
17.2 Withdrawals due to adverse events	1	39	Risk Ratio (IV, Random, 95% CI)	0.32 [0.04, 2.79]


Analysis 17.1. Comparison 17: Mepolizumab 2.5 mg/kg vs mepolizumab 0.55 mg/kg, Outcome 1: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 17.2. Comparison 17: Mepolizumab 2.5 mg/kg vs mepolizumab 0.55 mg/kg, Outcome 2: Withdrawals due to adverse events




Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Comparison 18. Mepolizumab 10 mg/kg vs mepolizumab 2.5 mg/kg

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
18.1 Histological improvement at study endpoint (dichotomous)	1	40	Risk Ratio (IV, Random, 95% CI)	0.56 [0.23, 1.37]
18.2 Withdrawals due to adverse events	1	40	Risk Ratio (IV, Random, 95% CI)	2.00 [0.20, 20.33]


Analysis 18.1. Comparison 18: Mepolizumab 10 mg/kg vs mepolizumab 2.5 mg/kg, Outcome 1: Histological improvement at study endpoint (dichotomous)



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

Analysis 18.2. Comparison 18: Mepolizumab 10 mg/kg vs mepolizumab 2.5 mg/kg, Outcome 2: Withdrawals due to adverse events



Risk of bias legend

- (A) Random sequence generation (selection bias)
- (B) Allocation concealment (selection bias)
- (C) Blinding of participants and personnel (performance bias)
- (D) Blinding of outcome assessment (detection bias)
- (E) Incomplete outcome data (attrition bias)
- (F) Selective reporting (reporting bias)
- (G) Other bias

ADDITIONAL TABLES

Table 1. Primary outcome - endoscopic improvement

Study ID	Endoscopic improvement system used	Continuous or dichotomous	Outcome data - endoscopic improvement at study endpoint
Alexander 2012	Any endoscopic findings yes/no	Dichotomous; endoscopic findings not seen	Resolution of all endoscopic findings of EoE was seen in 8.3% (1 of 12) of placebo-treated patients who completed the study and who had an abnormal baseline esophagogastroduodenoscopy. In the fluticasone-treated patients who completed the trial, resolution of pretreatment abnormal endoscopic findings was seen in 26.7% (4 of 15). Dichotomous (used for endoscopic dichotomous analysis): Fluticasone: 4/21 Placebo: 1/21
Alexander 2012	Not validated		
Alexander 2017	Endoscopic findings described by the gastroenterologist	Not reported	
Alexander 2017	No specific score was used		No quantitative data were reported – "no differences in endoscopic findings of EoE"
Assa'ad 2011	Post hoc	Not reported	Not reported
Bhardwaj 2017	Not reported	Not reported	Not reported
Butz 2014	Not reported	Not reported	Not reported
Clayton 2014	Not reported	Not reported	Not reported
Dellon 2012	Morphological endoscopic findings	Dichotomous	No prespecified aggregate score, cannot use

Table 1. Primary outcome - endoscopic improvement (Continued)

		described by the gastroenterologist	All budesonide, nebulized vs budesonide, oral viscous at end of trial:
		No specific score was used	Rings: 10/11 vs 4/11
			Narrowing: 6/11 vs 2/11
			Stricture: 3/11 vs 2/11
			Furrows: 6/11 vs 4/11
			White plaques/exudates: 3/11 vs 3/11
			Pallor/decreased vascularity: 2/11 vs 0/11
			Crepe-paper: 0/11 vs 0/11
			Erosive esophagitis: 0/11 vs 0/11
Dellon 2017	EREFS	Continuous	Change in EREFS from baseline at end of trial, mean (SD) (used for endoscopic continuous analysis):
	Hirano 2013	Edema (0 to 2)	Budesonide: -3.8 (3.9)/49
	Validated	Rings (0 to 3)	Placebo: 0.4 (6.7)/38
		Exudates (0 to 2)	
		Furrows (0 to 2)	
		Strictures (0 to 1)	
Dellon 2019	EREFS	Continuous	EREFS at end of trial, mean (SD) (used for endoscopic continuous analysis):
	Hirano 2013		Budesonide: 2.1 (1.7)/56
	Validated		Fluticasone: 2.8 (2.2)/55
Dellon 2021b	EREFS	Continuous	From digitized figure 3C, change in EREFS at end of trial, mean (SD) (used for endoscopic continuous analysis):
	Hirano 2013		Budesonide: -0.99 (-2.93)/24
	Validated		Placebo: 0.60 (3.30)/21
Dellon 2022	EREFS	Continuous	Change in EREFS from baseline at end of trial, mean (SD) (used for endoscopic continuous analysis):
	Hirano 2013		Dupilumab: -3.2 (0.41) n = 35/7 imputed
	Validated		Placebo: -0.3 (0.41) n = 26/13 imputed
Dellon 2022a	EREFS	Continuous	Change in EREFS from baseline at end of trial, mean (SD) (used for endoscopic continuous analysis):
	Hirano 2013	Endoscopic severity measured by the change from baseline in the EREFS (edema/rings/exudates/furrows/strictures (EoE Endoscopic Reference Score)) at week 12	APT-1011 3 mg twice-daily: -2.2 (1.84)/20
	Validated		APT-1011 3 mg at bedtime: -3.2 (2.28)/21
			APT-1011 1.5 mg twice-daily: -2.9 (1.92)/22
			APT-1011 1.5 mg at bedtime: -2.4 (1.85)/21
			All treatment arms:
			APT-1011: -2.68 (1.91)/84
			Placebo: -0.7 (1.31)/19

Table 1. Primary outcome - endoscopic improvement (Continued)

Dellon 2022b	Presentation, not publications	Not reported	Not reported
De Rooij 2022	EREFS	Endoscopic features are scored according to the EREFS classification and sub-classified as (i) inflammatory signs including white exudates, edema, and linear furrows (ii) fibrostenotic signs including rings and strictures	Median (IQR) reported, cannot use
Hirano 2013	Validated	The following scores were reported as median (IQR)	EREFS, median (IQR) (used for analysis)
	Inflammatory score	EREFS - post-treatment	<ul style="list-style-type: none"> Four-food elimination diet = 4 (1 to 4), SD = 3.20503 Four-food elimination diet + amino acid formula = 3 (1.5 to 4), SD = 2.746079
	Fibrostenotic score	Inflammatory score - post-treatment	Inflammatory score, median (IQR)
		Fibrostenotic score - post-treatment	<ul style="list-style-type: none"> Four-food elimination diet: 2 (1 to 2) Four-food elimination diet + amino acid formula: 2 (1 to 2)
			Fibrostenotic score, median (IQR)
			<ul style="list-style-type: none"> Four-food elimination diet: 1 (1 to 2) Four-food elimination diet + amino acid formula: 1 (1 to 2)
Dohil 2010	Endoscopy scoring tool	Continuous	Endoscopy score at end of trial, mean (SD) (used for endoscopic continuous analysis):
Aceves 2009	Not validated	Pre- and post-scores	Budesonide + PPI: 1.5 (2.5)/15
		Mucosal pallor/reduced vasculature	Placebo + PPI: 5.4 (2.8)/9
		Linear furrows/mucosal thickening, white plaques, concentric rings/stricture Friability/"tissue-paper" mucosa	
		Histology scoring tools	
		Epithelial histology score	
		Peak eosinophil count	
		Absent = 0	
		Present = 1	
Gupta 2015	Not reported	Not reported	Not reported
Heine 2019	Not reported	Not reported	Not reported
Hirano 2019	EREFS	EREFS: Continuous, mean difference	EREFS at end of trial, mean (SD) (used for endoscopic continuous analysis):
Hirano 2013	Validated		RPC4046 180 mg: 5.3 (4.2)/27
			RPC4046 360 mg: 4.8 (3.4)/30
			RPC4046 = 5.04 (3.71)/57

Table 1. Primary outcome - endoscopic improvement (Continued)

			Placebo: 7.9 (5.1)/32
Hirano 2020	EREFS Hirano 2013 Validated Change in esophageal distensibility plateau as measured by functional lumen imaging	Continuous	Change in EREFS from baseline at end of trial, LS mean change from baseline (SD) (used for endoscopic continuous analysis), N/imputed n: Dupilumab: -1.9 (1.4)/23/0 Placebo: -0.3 (1.5)/24/2
Hirano 2020f	EREFS Hirano 2013 Validated	Dichotomous Change from baseline to week 8/end of trial improvement/no change/worsening	Supplementary Table 3, data are from pre-specified analyses (used for endoscopic dichotomous analysis): APT-1011 at 1.5 mg twice-daily: 5/8 APT-1011 at 3.0 mg daily: 5/8 APT-1011: 10/16 Placebo: 0/8 Placebo: improvement 0; no change 7; worsening 1 APT-1011 1.5 mg: improvement 5; no change 2; worsening 1 APT-1011 3 mg: improvement 5; no change 3; worsening 0 Continuous outcomes, data are from post hoc analyses (cannot use) APT-1011 at 1.5 mg twice-daily: -2.92 (95% CI -4.68 to -0.88) APT-1011 at 3.0 mg daily: -2.74 (95% CI -4.5 to -0.88) APT-1011: -2.83 (1.72) n = 16 Placebo: 0 (1.72) n = 8
Hirano 2021	EREFS Hirano 2013 Validated	Continuous (mean (SD))	EREFS at end of trial, mean (SD) (used for endoscopic continuous analysis): Budesonide: 4.2 (3.3)/202 Placebo: 6.2 (3.7)/93
Kliewer 2019	EREFS Hirano 2013 Validated	Continuous (change in mean (SD))	From NCT02610816, change in EREFS from baseline at end of trial, mean (SD) (used for endoscopic continuous analysis): 1-food elimination diet change from baseline: -0.7 (2.2)/22 4-food elimination diet change from baseline: -1.3 (2.2)/12
Kliewer 2021	EREFS Hirano 2013 Validated	Continuous EREFS change from baseline mean (SD)	Change in EREFS from baseline at end of trial, mean (SD) (used for endoscopic continuous analysis): • 1-food elimination at 6 weeks: -1.32 (3.28)/67 • 6-food elimination at 6 weeks: -1.74 (3.91)/62
Konikoff 2006	No scoring system used Konikoff 2006	Dichotomous (number of patients with esophageal furrowing, epithelial hyperplasia, and esophageal mastocytosis)	Dichotomous, endoscopic: lack of furrows in the esophagus at end of trial (used for endoscopic dichotomous analysis): Fluticasone: 11/21 (52.4%) Placebo: 5/15 (33.3%)

Table 1. Primary outcome - endoscopic improvement (Continued)

After treatment, significantly fewer individuals in the fluticasone propionate (FP) group had endoscopic distal esophageal furrowing compared with the placebo group (50% vs 91%). Endoscopic distal esophageal furrowing was not present in any FP responders (0/10) after treatment, while all FP non-responders (10/10) had persistent furrowing in the distal esophagus.

Treatment with FP significantly reduced epithelial hyperplasia in both the proximal and distal esophagus, as assessed by histologic examination of H&E-stained sections. Placebo had no effect.

In the FP group, mast cell counts were significantly decreased by treatment (17.1 ± 3.5 pre-treatment vs 7.3 ± 2.2 post-treatment mast cells/hpf in the proximal esophagus and 17.9 ± 3.1 pre-treatment vs 9.8 ± 2.2 post-treatment mast cells/hpf in the distal esophagus) and post-treatment mast cell counts were significantly lower in the FP group than in the placebo group. FP responders had significantly lower post-treatment mast cell counts than FP non-responders (1.8 ± 0.5 vs 13.3 ± 3.6 mast cells/hpf in the proximal esophagus and 2.9 ± 1.0 vs 17.5 ± 2.5 mast cells/hpf in the distal esophagus).

Lieberman 2018	Not reported	Not reported	Not reported
Lucendo 2019 Hirano 2013	EREFS Validated	Continuous mean (SD) at end of trial	EREFS at end of trial, mean (SD) (used for endoscopic continuous analysis). Calculated from supplementary Table 5: Budesonide: 1.3(1.04)/59 Placebo: 4.6(1.26)/28
Miehlke 2016	Endoscopic score Global assessment of endoscopic appearance was determined using a 100 mm visual analogue scale (VAS) Not validated	Continuous Mean change in total endoscopic intensity score Endoscopic abnormalities: absent (0), mild (1), moderate (2), or severe (3): white exudates, furrows, edema, fixed rings, crêpe paper sign, short-segment stenosis, long-distance stenosis. Total endoscopic intensity score ranged from 0 to 21. Mean change in VAS endoscopic score (No SD reported)	No SD reported, cannot use data AT 2 weeks Endoscopic intensity score: Budesonide effervescent tablet 2 x 1 mg: -4.1 Budesonide effervescent tablet 2 x 2 mg: -3.4 Budesonide viscous suspension 2 x 2mg: -3.6 Placebo: -0.7 VAS endoscopic score: Budesonide effervescent tablet 2 x 1 mg: -37.4 Budesonide effervescent tablet 2 x 2 mg: -31.7 Budesonide viscous suspension 2 x 2mg: -25.2 Placebo: -9.6
Moawad 2013	Endoscopic assessment	Dichotomous Improvement of endoscopic findings	No aggregate outcome reported, cannot use data Stenosis on index endoscopy • Fluticasone: (4/5) 80% • ESO: (4/5) 80%

Table 1. Primary outcome - endoscopic improvement (Continued)

			Concentric rings <ul style="list-style-type: none"> • Fluticasone: (2/16) 13% • ESO: (7/16) 44% Longitudinal furrows <ul style="list-style-type: none"> • Fluticasone: (3/17) 18% • ESO: (7/17) 41% White plaques <ul style="list-style-type: none"> • Fluticasone: (2/4) 24% • ESO: (5/5) 100%
Oliva 2018	EREFS Hirano 2013 Validated	Not reported	Not reported
Peterson 2010	No scoring system, but morphological assessment Not validated	Continuous No threshold of success defined	No aggregate outcome reported, cannot use data Note: it is unclear if these were the findings at baseline or at 8 weeks (end of the study) Rings, n (%) <ul style="list-style-type: none"> • Fluticasone 14/15 • Esomeprazole 15/15 Furrows, n(%) <ul style="list-style-type: none"> • Fluticasone 1/15 • Esomeprazole 1/15 Abscesses, n(%) <ul style="list-style-type: none"> • Fluticasone 3/15 • Esomeprazole 2/15
Rothenberg 2015	Not reported	Not reported	Not reported
Rothenberg 2022	Not reported	Not reported	Not reported
Schaefer 2008	Endoscopy score Schaefer 2008	Dichotomous, improvement of one or more histological grades	Calculated from Table 5, improvement of one or more histological grades at end of trial (used for endoscopic dichotomous analysis): Fluticasone: 34/40 Prednisone: 30/40
Spiegel 2012	Not reported	Not reported	Not reported
Spiegel 2020	EREFS Hirano 2013 Validated	Continuous No threshold of success was identified but a mean difference of change was calculated	EREFs at end of trial, mean (SD) (used for analysis) Viaskin milk = 1.93 (1.58)/15 Placebo = 1.60 (1.67)/5 Change in EREFS from baseline at end of trial, mean (SD) (used for endoscopic continuous analysis):

Table 1. Primary outcome - endoscopic improvement (Continued)

			Mean ± SD
			Viaskin milk = -0.07 (1.49)/15
			Placebo = -0.80 (1.30)/5
Straumann 2010a	Endoscopic eosinophilic esophagitis abnormalities	Dichotomous	Absence of esophageal abnormalities at end of trial (used for endoscopic dichotomous analysis):
Straumann 2003		Endoscopic findings were graded by means of a simple overall score: absent, minor (fine nodules, fine whitish reticular structures, furrows), moderate (bright white scale- or plaque-like structures, corrugated rings) or severe (mucosal lesions, fixed stenosis)	Absent:
			Mepolizumab: 0/5
			Placebo: 0/6
			Minor n = 0/5, n = 1/6
			Moderate n = 3/5, n = 3/6
			Severe n = 2/5, n = 2/6
		Absence of features is defined as the primary outcome	
Straumann 2010b	Macroscopic assessment during endoscopy	Dichotomous	No aggregate score reported, cannot use data
		Threshold of success was not established. However, more disappearing endoscopic features counts as success.	Among the 10 patients with complete histologic remission:
		Roughly classified as absent, minimal, moderate, or severe... []. Additionally, the presence of 6 major signs of EoE (white exudates, red furrows, corrugated rings, solitary rings, crêpe paper sign, and severe stenosis impossible to pass with the standard endoscope) as well as signs of fungal infection were recorded.	<ul style="list-style-type: none"> • 10/10 had white exudates disappear • 8/9 had red furrows disappear • 8/9 had corrugated rings persist
Straumann 2011	Not reported	Not reported	Not reported
Straumann 2013	The global appearance of endoscopic abnormalities was assessed using a 10 cm visual analogue scale	Continuous; compared means; no pre-specified treatment response threshold	Change in global assessment of endoscopic appearance from baseline at end of trial, mean (SD) (used for endoscopic continuous analysis):
			OC004549: 6.06 (1.79)/14
			Placebo: -5.57 (2.20)/12
Straumann 2020	EREFS	Continuous	EREFS at end of trial, mean (SD) (used for endoscopic continuous analysis):
Hirano 2013			

Table 1. Primary outcome - endoscopic improvement *(Continued)*

Validated			Budesonide 0.5 mg twice-daily: 1 (1.2)/65
There were many endoscopic outcomes reported; however, all were exploratory			Budesonide 1.0 mg twice-daily: 1 (1.1)/65
			Budesonide: 1 (1.14)/130
			Placebo: 4 (1.8)/65
Tytor 2021	Not reported	Not reported	Not reported

AAF: amino acid-based formula; BET: budesonide effervescent tablet; BOV: budesonide, oral viscous; CG: control group; CI: confidence interval; EoE: eosinophilic esophagitis; EoT: end of treatment; EREFS: EoE Endoscopic Reference Score; FFED: four food elimination diet; IG: intervention group; ESO: esomeprazole; IQR: interquartile range; LS: least squares; NEB: nebulized/swallowed budesonide solution; OVB: viscous/swallowed budesonide solution; PPI: proton pump inhibitor; SD: standard deviation; VAS: visual analogue scale

Table 2. Included studies' characteristics

Study ID	Adults/children or both	Interventions	Control	Induction or maintenance at the time of randomization	Disease activity for induction studies/Definition of remission for maintenance studies	RCT duration and measurement time points	Concomitant medications and diet modifications (mandatory and/or allowed)
Alexander 2012	Children and adults (18 to 65)	Fluticasone 880 µg twice-daily, aerosolized/swallowed, 6 weeks	Placebo twice-daily, aerosolized/swallowed, 6 weeks	Induction	Peak eosinophil level of 20 or more eosinophils (eos)/hpf on esophageal biopsy	<p>Duration: 6 weeks</p> <p>Measurement points: 2 weeks, phone interview, med compliance, MDQ-2 week, side effects questionnaire</p> <p>Measurement points: 4 weeks, phone interview, med compliance, MDQ-2 week, side effects questionnaire</p> <p>Measurement points: 6 weeks, EDG, MDQ-2 week, side effects questionnaire, 24-hour urine</p>	<p>PPI</p> <p>All patients enrolled after the establishment of the consensus definition of EoE in 2007 had at least 1 month of twice-daily PPI therapy without resolution of dysphagia (fluticasone: 52.4% (11 of 21); placebo: 57.1% (12 of 21)). Repeat endoscopy post-PPI therapy was not performed routinely before study initiation. The baseline MDQ-30 documenting dysphagia was completed after PPI treatment. Patients on PPI medications with symptomatic relief of heartburn or regurgitation and with persistent dysphagia were allowed to continue their PPI medications at the same dose during the study (fluticasone: 26.3% (5 of 19); placebo: 0% (0 of 15)).</p> <p>Diet</p> <p>Four treatment patients avoided fibrous foods: 2 patients had a partial symptom response and complete histologic response, 1 patient had a complete symptom response and partial histologic response, and 1 patient had a partial symptom response and no histologic response. Two placebo-treated patients avoided fibrous foods: 1 patient had a complete symptom response and no histologic response, and 1 patient had a partial symptom response and no histologic response. Fibrous food avoidance remained unchanged throughout the study in 5 of the 6 patients. One treat-</p>

Table 2. Included studies' characteristics (Continued)

<p>Alexander 2017</p>	<p>Children and adults (18 to 65)</p>	<p>Montelukast 2 x 10 mg/day, orally at bedtime, 26 weeks</p>	<p>Placebo tablets 2/day, orally at bedtime, 26 weeks</p>	<p>Maintenance (after steroid induction successful on endoscopic screening)</p>	<p>Remission was defined as the absence of dysphagia as defined as an answer of yes to the question of "Have you had trouble swallowing unrelated to a sore throat or cold?", a severity of at least moderate, and a frequency of at least 1 or more times per week</p>	<p>Duration: 26 weeks Measurement points: side effects: 2, 4, 8, 12, 16, 20, and 24 weeks Symptoms: 2, 4, 8, 12, 20, and 26 weeks</p>	<p>ment patient, who had a complete symptomatic response, advanced to an unrestricted diet for the last 2 weeks of the study.</p> <p>Steroids Not reported</p> <p>PPI Patients on PPI medications with symptomatic relief of heartburn or regurgitation and with persistent dysphagia before topical steroid treatment were allowed to continue their PPI medications at the same dose during the study.</p> <p>Diet No restrictions applied.</p> <p>Steroids A.Mandatory prior to randomization – Patients were given topical steroids in the form of swallowed aerosolized fluticasone at 880 µg twice-daily OR swallowed budesonide - Rincinolgel 3 mg twice-daily for at least 6 weeks. Patients kept compliance logs that were reviewed at telephone interviews at 2- to 4-week intervals during the study; 90% compliance was required for continued study inclusion. B.Allowed during study – Patients on nasal/inhaled steroids for rhinitis and/or asthma were allowed to continue on the same dose. C. No new topical steroid medication was initiated during the study or during the pre study swallowed steroid treatment period.</p>
-----------------------	---------------------------------------	---	---	---	---	---	--

Table 2. Included studies' characteristics (Continued)

Assa'ad 2011	Children (2 to 17)	Mepolizumab 3 x 0.55 mg/kg, intravenous infusion, 3 monthly doses Mepolizumab 3 x 2.5 mg/kg, intravenous infusion, 3 monthly doses Mepolizumab 3 x 10 mg/kg, intravenous infusion, 3 monthly doses	Only comparator arms	Induction	Peak eosinophil level of 20 or more eosinophils (eos)/hpf on esophageal biopsy	Duration: 12 weeks Measurement points: histologic, safety, tolerability, mean intraepithelial eosinophil counts, improvement of histopathologic and endoscopic findings, blood eosinophil counts, and frequency and severity of EoE symptoms at 9 to 12 weeks	PPI Mepolizumab 3 x 0.55 mg/kg: 6/19 (31.6%) Mepolizumab 3 x 2.5 mg/kg: 6/20 (30.0%) Mepolizumab 3 x 10 mg/kg: 6/20 (30.0%) Diet Mepolizumab 3 x 0.55 mg/kg: 4/19 (21.0%) Mepolizumab 3 x 2.5 mg/kg: 6/20 (30.0%) Mepolizumab 3 x 10 mg/kg: 8/20 (40%) Steroids Required to terminate steroid therapy.
Bhardwaj 2017	Children and adults (18 to 65)	Beclomethasone diphosphate 80 µg twice-daily, aerosolized/swallowed, 8 weeks	Placebo twice-daily, aerosolized/swallowed, 8 weeks	Induction	Peak eosinophil level of 15 or more eosinophils (eos)/hpf on esophageal biopsy	Duration: 8 weeks Measurement points: histologic, symptoms, peripheral blood eosinophil counts, the tissue MCT level, tissue IL-13, CCL2, CCL-5, IL-17F, IL-10, IL-25, and thymic stromal lymphopoietin (TSLP) expression all at 8 weeks	PPI All patient continued PPI (except n = 1 in the placebo group) Diet No diet elimination. During the screening period all patients were asked to discontinue dietary restrictions, if any. Steroids During the screening period of 12 weeks before the treatment periods, the enrolled patients were asked to discontinue all previous topical corticosteroids for EoE

Table 2. Included studies' characteristics (Continued)

Butz 2014	Children and adults (3 to 30)	Fluticasone propionate 880 µg twice-daily, aerosolized/swallowed, 12 weeks	Placebo twice-daily, aerosolized/swallowed, 12 weeks	Induction	24 or more eosinophils/hpf in the proximal or distal esophagus while being treated with a PPI for at least 2 months or having a negative pH probe	Duration: 12 weeks Measurement points: histologic and EoE symptom score at 12 weeks	PPI Participants were instructed not to change PPI dosage and/or diet therapy during the study. Diet Not reported Steroids Not reported
Clayton 2014	Children and adults (≥ 15)	Omalizumab 0.016 mg/kg/IgE (IU/mL), subcutaneous every 2 to 4 weeks, 16 weeks	Placebo, subcutaneous every 2 to 4 weeks, 16 weeks	Induction	> 15 eosinophils/hpf in esophageal biopsy specimen, not responsive to maximal-dose PPI	Duration: 16 weeks Measurement points: histologic and dysphagia symptoms at 16 weeks	PPI The participants to were only on proton pump inhibitors during the trial once the consensus criteria for eosinophilic esophagitis were published in 2007. A majority of the participants (all but 5 in the treatment group and 4 in the control group) were treated with high-dose, twice-daily proton pump inhibitors for the duration of the study and for at least 8 weeks prior to the initial biopsy and the beginning of the study. Diet Not reported Steroids Not reported
Dellon 2012	Children and adults (≥ 18)	Budesonide solution (1 mg/2 mL) twice-daily, nebulized/swallowed, 8 weeks Budesonide solution (1 mg/2 mL) with 5 g of sucralose twice-	Only comparator arms	Induction	Symptoms of esophageal dysfunction and had persistent esophageal eosinophilia (≥ 15 eosinophils in one high-power field) after 8 weeks or treatment with	Duration: 8 weeks Measurement points: histologic, dysphagia symptom scores, endoscopic,	PPI Previously prescribed PPIs were discontinued as patients included in this study did not have either a symptomatic or histologic response to a high-dose PPI trial. Diet

Table 2. Included studies' characteristics (Continued)

		daily, swallowed, 8 weeks			twice-daily proton pump inhibitor	and safety at 8 weeks	"No dietary elimination therapy was allowed in either group during the study period, and no other concurrent therapy for eosinophilic esophagitis was allowed. " Steroids Subjects were excluded if previously treated with topical steroids.
Dellon 2017	Children and adults (11 to 40)	Budesonide oral suspension 2 mg/10 mL twice-daily, swallowed, 12 weeks	Placebo 10 mL twice-daily, swallowed, 12 weeks	Induction	Symptoms of esophageal dysfunction and at least 15 intra-epithelial eosinophils per hpf after an 8-week, high-dose PPI	Duration: 12 weeks Measurement points: histologic, dysphagia symptom score, and endoscopic score at 12 weeks	PPI PPI-responsive patients were excluded. PPI-responsive is defined as < 15 eos/hpf. Changing the PPI regimen for non-responsive patients was also a reason for exclusion. Diet Not reported Steroids The use of corticosteroids (topical or systemic) in the 4 weeks preceding the screening endoscopy was an exclusion criterion. Changes to the inhaled corticosteroid regimen were also exclusion criteria.
Dellon 2019	Children and adults (16 to 80)	Budesonide 1 mg/4 mL twice-daily with 10 g of sucralose, swallowed + placebo inhaler twice-daily, aerosolized/ swallowed, 12 weeks Placebo 4 mL twice-daily with 10 g of sucralose, swallowed + fluticasone 880	Only comparator arms	Induction	Cases had to have dysphagia or other symptoms of esophageal dysfunction, persistent esophageal eosinophilia (15 eosinophils in at least 1 high-power field (eos/hpf)) after 8 weeks of treatment with a twice-daily PPI, and other competing caus-	Duration: 8 weeks Measurement points: histologic, dysphagia symptom score, and endoscopic score at 8 weeks	PPI No changes in baseline PPI medication dose were allowed during the study period. Diet No dietary changes were allowed during the study period. Steroids Not reported

Table 2. Included studies' characteristics (Continued)

		µg twice-daily, aerosolized/swallowed, 12 weeks			es of esophageal eosinophilia excluded. A symptom threshold was not required for study entry.		
Dellon 2021b	Children and adults (11 to 55)	Budesonide oral suspension 2.0 mg twice-daily	Placebo	Maintenance	Eosinophil histology relapse was defined as an eosinophil count of greater than or equal to (\geq) 15 per high-power field (eos/hpf) from at least 2 of 3 levels of the esophagus. Dysphagia symptom relapse was defined as having at least 4 days of dysphagia (with answer 'Yes' for question 2 in DSQ (Dysphagia Symptom Questionnaire)) in the 2-week period prior to the scheduled visit, as determined by the DSQ.	Duration: 36 weeks Measurement points: histologic, dysphagia symptom score, endoscopic score at 36 weeks	PPI Budesonide oral suspension: 22 (88.0%) Placebo: 20 (87.0%) Participants were 100% prior PPI failures. Diet Steroids Budesonide oral suspension: 6 (24.0%) Placebo: 3 (13.0%)
Dellon 2022	Children and adults (\geq 12)	Dupilumab 300 mg subcutaneously weekly	Placebo	Induction	"A documented diagnosis of EoE by endoscopic biopsy"	Duration: 24 weeks Measurement points: histologic, dysphagia symptom score, endoscopic score at 24 weeks	PPI Not reported Participants were 100% prior PPI failures Diet Dupilumab: 17/42 Placebo: 16/39 Steroids

Table 2. Included studies' characteristics (Continued)

Dellon 2022a	Children and adults (18 to 75)	Active APT-1011 with 4 arms varying in dosage: 3 mg twice-daily 3 mg at bedtime 1.5 mg twice-daily 1.5 mg at bedtime	Placebo disintegrating tablet	Induction	Defined as 3 episodes of dysphagia per week during the last 14 days of the 4-week baseline symptom assessment phase and a Global EoE Symptom Score of > 3), and active esophageal eosinophilia (after evaluation of 5 biopsies from proximal and distal esophageal locations and at least 1 biopsy with a peak count of 15 eos/HPF) after documentation of failed histologic response on 8 weeks of high-dose PPI	Duration: 12 weeks Measurement points: Measurement points: histologic, dysphagia symptom score, endoscopic score at 12 weeks	Not reported PPI Patients on a PPI were required to maintain a stable regimen Diet Changes in diet were prohibited Steroids Corticosteroids are prohibited. However, randomization was stratified by current esophageal stricture(s) and a positive response to prior corticosteroid use Biologics and immunomodulator Biologics and immunomodulator were prohibited
Dellon 2022b	Children and adults (12 to 70)	Lirentelimab 1 + 3 + 3 + 3 + 3 + 3 mg/kg, intravenous infusion, 6 monthly doses Lirentelimab 1 + 1 + 1 + 1 + 1 + 1 mg/kg, intravenous infusion, 6 monthly doses	Placebo 1 + 1 + 1 + 1 + 1 matching saline, intravenous infusion, 6 monthly doses	Induction	Peak eosinophil level of 15 or more eos/hpf on esophageal biopsy	Duration: 24 weeks Measurement points: histologic and Dysphagia Symptom Score at 23 to 24 weeks	PPI Not reported Diet Not reported Steroids Not reported
De Rooij 2022	Children and adults (not reported)	Elimination of 4 foods including wheat/gluten, milk, egg, and either soy or legumes (FFED) +	Elimination of 4 foods including wheat/gluten, milk, egg, and either soy	Induction	Symptoms of esophageal dysfunction (Straumann Dysphagia Instrument (SDI) score of ≥ 1) and	Duration: 6 weeks Measurement points: histologic, endoscopic, clinical,	PPI Patients were on PPI from baseline Diet

Table 2. Included studies' characteristics (Continued)

		amino acid-based formula (AAF)	or legumes (FFED)		≥ 15 eosinophils (eos) per microscopic hpf on baseline biopsy	and nutritional outcomes were evaluated between week 1 and week 6	Dietitian specialized in allergies for extensive nutritional evaluation. To guarantee sufficient intake and to improve diet adherence, patients subsequently received personalized nutritional advice with restriction of gluten, milk, soy, and eggs (four food elimination diet). The amount of prescribed amino acid-based formula added to the four food elimination diet in the intervention group was 30% of patients' daily caloric requirements based on body mass index and weekly physical activity.
Dohil 2010	Children (1 to 17)	Budesonide suspension (0.5 mg/2 mL) + PPI	Sterile water + PPI	Induction	Peak eosinophil level of 20 or more eosinophils (eos)/hpf on esophageal biopsy	Duration: 12 weeks Measurement points: histologic, symptomatic, and endoscopic score at 12 weeks	<p>PPI</p> <p>Patients were a mixture of being on PPI and not, prior to PPI therapy as part of both arms. Once assigned a group, patients were given PPI as part of the intervention group or the placebo group.</p> <p>Diet</p> <p>Patients who were on diet restrictions as part of their treatment to EoE were allowed to continue with these restrictions. Dietary restrictions were reported in a non-specific pattern throughout the study. Restrictions were limited to: E, eggs; F, fish; milk; nuts; soy and wheat.</p> <p>Steroids</p> <p>Patients were excluded from the study of they needed a systemic corticosteroid</p>

Table 2. Included studies' characteristics (Continued)

Gupta 2015	Children (2 to 18)	<p>Low-dose OBS: oral budesonide suspension (OBS) 0.05 mg/mL at bedtime and placebo after breakfast for 12 weeks, with a total daily dose of 0.35 mg (2 to 9 years) or 0.50 mg (10 to 18 years), followed by a 3-week taper period</p>	<p>Placebo twice-daily at bedtime and after breakfast for 12 weeks with a 3-week taper period</p>	Induction	<p>Esophageal biopsy must show ≥ 20 eos per HPF (400x, 0.3 mm² HPF) at 2 or more levels of the esophagus following 4 weeks of high-dose PPI (type, actual dosage not specified)</p>	<p>Duration: 12 weeks</p> <p>Measurement points: clinical symptom scores, and safety at weeks 2, 4, 8, and 12. Histologic at 12 weeks.</p>	<p>PPI</p> <p>Four weeks of high-dose PPI therapy (type, actual dosage not specified) were required for inclusion</p> <p>Diet</p> <p>Dietary therapy, environmental therapy, and/or medical regimens (including gastric acid suppression, if any) in effect at the Screening Visit, were continued during treatment</p> <p>Steroids</p> <p>Not reported</p>
		<p>Medium-dose OBS: oral budesonide suspension (OBS) 0.2 mg/mL at bedtime and placebo after breakfast for 12 weeks, with a total daily dose of 1.4 mg (2 to 9 years) or 2.0 mg (10 to 18 years), followed by a 3-week taper period</p>					
		<p>High-dose OBS: oral budesonide suspension (OBS) 0.2 mg/mL at bedtime (hs) and after breakfast for 12 weeks, with a total daily dose of 2.8 mg (2 to 9 years) or 4.0 mg (10 to 18 years), followed</p>					

Table 2. Included studies' characteristics (Continued)
 by a 3-week taper period

Heine 2019	Children (1 to 18)	4-food elimination diet + PPI 4-food elimination diet (strictly avoiding all foods containing cow's milk, soy, wheat or egg) Omeprazole: 7.5 kg to 9.9 kg: 5 mg in the morning and 10 mg at night, 10.0 kg to 14.9 kg: 10 mg twice-daily, 15.0 kg to 19.9 kg: 15 mg twice-daily, > 20 kg: 20 mg twice-daily	PPI alone Omeprazole: 7.5 kg to 9.9 kg: 5 mg in the morning and 10 mg at night, 10.0 kg to 14.9 kg: 10 mg twice-daily, 15.0 kg to 19.9 kg: 15 mg twice-daily, > 20 kg: 20 mg twice-daily	Induction	≥ 15 eosinophils per high-power field; HPF	Duration: 8 to 12 weeks Measurement points: histologic at 8 to 12 weeks	PPI Use controlled Diet Controlled Steroids No steroids allowed
Hirano 2019	Children and adults (18 to 65)	RPC4046 180 mg (n =31), or RPC4046 360 mg (n = 34) subcutaneously once weekly	Placebo subcutaneous injections once weekly	Induction	Symptoms of dysphagia for a minimum of 4 days over 2 weeks (within the 4-week screening period) and histologic evidence of EoE, defined as a peak count of ≥ 15 eosinophils per high-power field (eos/hpf; microscope hpf = 0.3 mm ²) at any 2 of 3 levels of the esophagus (proximal, mid, distal) when off antiinflammatory therapy for EoE	Duration: 16 weeks Measurement points: histologic, dysphagia symptoms, endoscopic, and participants global assessment of disease severity score at 16 weeks	PPI If patient was screened while on PPI they had to agree to maintain the same dose over the 16-week study period Diet No mention Steroids Excluded

Table 2. Included studies' characteristics (Continued)

Hirano 2020	Children and adults (18 to 65)	Weekly subcutaneous dupilumab 300 mg (loading dose, 600 mg on day 1) for 12 weeks	Placebo subcutaneously for 12 weeks	Induction	Active esophageal inflammation was to be evident at screening (i.e. peak cell count ≥ 15 eosinophils per high-power field (eos/HPF): 400 magnification of a 0.3 mm ² field) as indicated by esophageal pinch biopsy specimens from at least 2 of 3 esophageal sites from endoscopy performed no more than 2 weeks after at least 8 weeks of treatment with high dose	Duration: 12 weeks Measurement points: Stroumann Dysphagia Instrument (SDI) patient-reported outcome (PRO) score, histologic endoscopic reference score, esophageal distensibility, and safety at 12 weeks	Patients could receive concomitant medications as needed at the investigator's discretion, except for those that were prohibited. If medically necessary, rescue medications or emergency esophageal dilation could be provided. Patients who received rescue therapy were discontinued from study treatment and considered non-responders. Prohibited concomitant medications included medications used for the treatment of EoE, allergen immunotherapy, live attenuated vaccines, and any investigational drug other than dupilumab.
PPI							
Patients using stable doses of PPIs at screening were permitted to continue the same dosing regimen until the end-of-treatment visit; those not using PPIs in the 8 weeks before screening were prohibited from starting them.							
Prior history of treatment with high-dose PPIs at baseline:							
Dupilumab: 23 (100%)							
Placebo: 24 (100%)							
PPI treatment ongoing at baseline:							
Dupilumab: 14 (60.9%)							
Placebo: 15 (62.5%)							
Diet							
Patients were instructed not to modify their diets during the study							
Steroids							

Table 2. Included studies' characteristics (Continued)

							Prohibited. If required for rescue therapy, the patient was discontinued from the study.
Hirano 2020f	Children and adults (12 to 55)	APT-1011 (fluticasone propionate tablets) at 1.5 mg once in morning and once in the evening, APT-1011 at 3.0 mg once a day APT-1011 3.0 mg once a day group had a placebo tablet in the morning bottle and a 3.0 mg fluticasone propionate tablet in the evening bottle	The placebo group had a placebo tablet in both morning and evening bottles	Induction	Esophageal mucosal peak eosinophil count \geq 24 per high-power field (HPF) (HPF; radius = 0.275 mm; 400 \times)	Duration: 8 weeks Measurement points: treatment-emergent adverse events at 2, 4, 6, and 8 weeks Exploratory outcomes: histologic and endoscopic at 8 weeks. Physician Global Assessment of the participant's overall EoE activity, the Patient Global Assessment of symptom severity and EEsAI PRO, and Mayo Dysphagia Questionnaire at weeks 4 and 8.	PPI Current: APT-1011 2 x 1.5 mg: 5 (62.5%) APT-1011 1 x 3 mg: 5 (62.5%) Placebo: 6 (75%); Prior: APT-1011 2 x 1.5 mg: 3 (37.5%) APT-1011 1 x 3 mg: 3 (37.5%) Placebo: 2 (25%) Diet Not reported Steroids Not reported
Hirano 2021	Children and adults (11 to 55)	Study group 2: budesonide oral suspension (BOS) 2.0 mg twice-daily (10 mL at a concentration of 0.2 mg/mL)	Placebo	Induction	(\geq 15 eosinophils/high-power field (eos/hpf) from at least 2 levels of the esophagus	Duration: 12 weeks Measurement points: Measurement points: histologic, dysphagia symptom score, endoscopic score at 12 weeks	PPI Budesonide oral suspension: 176 (82.6%) Placebo: 92 (87.6%) Diet Budesonide oral suspension: 11 (10.5%) Placebo: 21 (9.9%)

Table 2. Included studies' characteristics (Continued)

							Steroids Budesonide oral suspension: 40 (18.8%) Placebo: 19 (18.1%)
Kliewer 2019	Children (6 to 17)	1 - food (milk) elimination diet (1FED); 4 - food (milk, egg, wheat, soy) elimination diet (4FED)	No placebo, only comparator arms	Induction	Histologically confirmed active EoE (≥ 15 eos/hpf) and symptoms of esophageal dysfunction	Duration: 12 weeks Measurement points: histologic, symptomatic, and quality of life at 12 weeks	PPI Not reported. Failure of a PPI trial was required for inclusion. Diet Not reported Steroids Exclusionary.
Kliewer 2021	Children and adults (18 to 60)	1-food elimination: animal milk 6-food elimination: animal milk, wheat, egg, soy, tree nuts/peanuts, seafood	No placebo, only comparator arms	Induction	≥ 15 eos/hpf + symptoms and lack of PPI response	Duration: 6 weeks Measurement points: histologic, EoE Histologic Scoring System (EoE-HSS), EoE Endoscopic Reference Score (EREFs), EoE Symptom Activity Index (EoE-SAI), and quality of life (EoE-QoL-A) at 6 weeks	PPI PPI failure required for inclusion, not clear how concomitant PPI was handled Diet Elimination determined by randomization. No other restrictions specified. Steroids Excluded
Konikoff 2006	Children (1 to 18)	Swallowed fluticasone propionate (FP) (880 mg/day)	Patients were treated with swallowed FP or placebo. Patients were given identical metered-dose in-	Induction	The primary outcome measure, as specified before the study was initiated, was complete histologic response to treatment as defined by a peak	Duration: 12 weeks Measurement points: histologic, endoscopic, and	PPI Fluticasone: 8 (38%) Placebo: 5 (33%) Acid suppression (PPI or H2-RA) Fluticasone: 10 (48%)



Table 2. Included studies' characteristics (Continued)

			halers of FP or placebo.		eosinophil count of 1 eosinophil in all 400x HPFs in both the proximal and distal esophagus.	vomiting at 12 weeks	Placebo: 7 (47%) Diet Not reported Montelukast Fluticasone: 4 (19%) Placebo: 0 (0%) Steroids Not reported
Lieberman 2018	Children (2 to 17)	Participants 2 to 12 years of age - 100 mg of cromolyn - 1 ampule mixed with 1 teaspoon of sugar 4 times daily Participants 13 to 18 years of age - 200 mg cromolyn - 2 ampules mixed with 2 teaspoons of sugar 4 times daily (ClinicalTrials.gov)	Saline ampules, participants 2 to 12 years of age - 1 ampule mixed with 1 teaspoon of sugar 4 times daily Participants 13 to 18 years of age - 2 ampules mixed with 2 teaspoons of sugar 4 times daily (ClinicalTrials.gov)	Induction	≥ 15 eosinophils per high-power field (eos/hpf) following at least 8 weeks of high-dose PPI therapy and a normal esophageal pH probe	Duration: 8 weeks Measurement points: symptoms at 4 and 8 weeks. Endoscopic and symptoms at 8 weeks.	PPI Not reported Diet Not reported Steroids Patients on concomitant treatment with swallowed corticosteroids were excluded. Any prior use of swallowed corticosteroids required a 4-week washout period.
Lucendo 2019	Children and adults (18 to 75)	Budesonide orodispersible tablets (BOT; 1 mg twice-daily)	Placebo	Induction	Patients had to have a severity of 4 points on a 0 to 10 numerical rating scale (NRS) for either dysphagia or odynophagia for 1 day in the week before randomization.	Duration: 6 weeks Measurement points: histologic at 6 weeks. Dysphagia, EEsAI-Pro at 2, 4, and 6 weeks	PPI Budesonide: 7 (12%) Placebo: 3 (10%) Diet Not reported Steroids

Table 2. Included studies' characteristics (Continued)

					<p>Additionally, Patient's Global Assessment (PatGA) of EoE activity was to be 4 points on a 0 to 10 NRS. Histologic activity with peak eos $\geq 65/\text{mm}^2$ hpf in at least 1 hpf (corresponding to ≥ 20 eos/hpf), as measured in a total of 6 hpf derived from 6 biopsies, 2 each from the proximal, mid, and distal segments of the esophagus.</p>		Not reported
Miehlke 2016	Children and adults (18 to 75)	<p>Budesonide effervescent tablet (BET) 2 x 1 mg/day</p> <p>Budesonide effervescent tablet (BET) 2 x 2 mg/day</p> <p>Budesonide viscous suspension 2 x 2 mg/day</p>	Placebo	Induction	<p>Clinical symptoms of esophageal dysfunction (dysphagia score ≥ 3), peak eosinophils (eos) $\geq 65/\text{mm}^2$ high-power fields (hpf) in at least 1 hpf (corresponding to ≥ 20 eos/hpf), and eosinophilic tissue infiltration with a mean cell density ≥ 16 eos/mm^2, as measured in a total of 30 hpf derived from 6 biopsies, 2 each from the proximal, mid, and distal segments of the esophagus</p>	<p>Duration: 2 weeks</p> <p>Measurement points: histologic, symptomatic, endoscopic, and safety at 2 weeks</p>	<p>PPI</p> <p>Patients with a clinicopathological response to a treatment with proton pump inhibitors (PPIs) at a standard dose with a treatment duration of at least 2 weeks were excluded</p> <p>Diet</p> <p>Patients with dietary restrictions within 4 weeks prior to screening or during treatment were excluded. Patients who had an intake of grapefruit food/drinks were excluded</p> <p>Steroids</p> <p>Patients excluded if they received:</p> <p>Topical/systemic therapies for any reason that may have affected assessment of primary and secondary end points (i.e. systemic glucocorticoids, histamine antagonists, mast cell stabilizers, leukotriene receptor antagonists, biologics, immunosuppressants)</p>

Table 2. Included studies' characteristics (Continued)

							concomitantly or within 4 weeks prior to screening
							Topical therapy (topical steroids, inhaled sodium cromoglycate concomitant or within 2 weeks prior to screening)
Moawad 2013	Children and adults (≥ 18)	Esomeprazole 40 mg once daily 8 weeks	Fluticasone propionate 440 μg twice-daily 8 weeks	Induction	One clinical symptom of esophageal dysfunction (dysphagia, food impaction, heart-burn) with ≥ 15 eosinophils/hpf	Duration: 8 weeks Measurement points: histologic and dysphagia symptom score at 8 weeks	PPI/steroid From NCT00895817, patients had to agree to a 1-month washout of both PPI and steroids to be eligible Diet Not reported
Oliva 2018	Children (not reported)	Six-food elimination diet Swallowed fluticasone Swallowed budesonide Oral viscous budesonide	No placebo, only comparator arms	Induction	Not reported	Duration: 8-week induction, 34 week maintenance Measurement points: histologic, clinical symptoms and endoscopic scores at 8 and 42 weeks	PPI Not reported Diet Not reported Steroids Not reported
Peterson 2010	Children and adults (18 to 80)	Esomeprazole (40 mg by mouth every morning) for 8 weeks	Aerosolized, swallowed fluticasone (440 μg by mouth twice a day) for 8 weeks	Induction	≥ 15 eosinophils averaged over 5 high-power fields on esophageal biopsy in participants with symptoms of dysphagia, food impaction or chest pain	Duration: 8 weeks Measurement points: histologic and dysphagia scores at 8 weeks	PPI If patients were on esomeprazole or fluticasone prior the trial, they needed to withhold the treatment for a month prior to be included. One arm received esomeprazole (40 mg by mouth every morning) for 8 weeks. Diet Not reported Steroids

Table 2. Included studies' characteristics (Continued)

							Not reported
Rothenberg 2015	Children and adults (18 to 50)	IV QAX576 6 mg/kg at weeks 0, 4, and 8	IV placebo at weeks 0, 4, and 8	Induction	Peak eosinophil density of 24 cells or greater per high-power field (hpf; 3400 magnification) in the proximal or distal esophagus validated by a central laboratory pathology service	Duration: 12 weeks Measurement points: histologic and dysphagia symptom score at 12 weeks	<p>PPI/steroids</p> <p>Patients already on PPIs, nasal, or inhaled steroids were allowed to continue these throughout the study.</p> <p>Diet</p> <p>Patients were instructed to maintain their baseline diet throughout the study.</p>
Rothenberg 2022	Children and adults (≥ 12)	Dupilumab 300 mg subcutaneously weekly	Placebo	Induction	"A documented diagnosis of EoE by endoscopic biopsy"	Duration: 24 weeks Measurement points: histologic and dysphagia symptom scores at 24 weeks	<p>PPI / Steroids</p> <p>No information reported for concomitant PPI or steroids. Participants were 100% prior PPI failures.</p> <p>Diet</p> <p>Food elimination diet at screening</p> <p>Dupilumab: 60/161</p> <p>Placebo: 29/79</p>
Schaefer 2008	Children (1 to 18)	Fluticasone: swallowed fluticasone by metered dose inhaler (110 µg per puff for ages 1 to 10 years and 220 µg per puff for ages 11 years or older, 2 puffs 4 times/day) for 4 weeks	Prednisone: oral P suspension/tablet (1 mg/kg/dose twice a day; maximum 30 mg twice a day) for 4 weeks	Induction	Esophageal mucosal biopsy specimens showing ≥ 15 eos/hpf with negative pH probe studies	Duration: 4-week induction Measurement points: histologic, clinical symptoms and safety at 4 weeks	<p>PPI</p> <p>Not reported</p> <p>Diet</p> <p>During the entire study, patients continued a regular diet except for foods identified as possible allergens by allergy testing.</p> <p>Steroids</p> <p>None of the patients were on corticosteroids at the time of initial endoscopy or at study enrollment.</p>
Spergel 2012	Children (5 to 18)	1, 2, or 3 mg/kg reslizumab	Placebo	Induction	Defined as ≥ 24 eosinophils in ≥ 1	Duration: 15 weeks	PPI

Table 2. Included studies' characteristics (Continued)

					high-power field (hpf))	Measurement points: histologic, and the physician's global assessment score at week 15	"Patients were also allowed to take medications for acid reflux if the doses remained stable throughout the study; use of these therapies on an as-needed basis was not permitted."
							<p>Diet</p> <p>"Patients were instructed to maintain their baseline diet throughout the study."</p> <p>Steroids</p> <p>"Patients were allowed to take inhaled corticosteroids, and nasal corticosteroids for allergies if they were started before the first dose of study medication, if the patients had symptoms of eosinophilic esophagitis while taking these medications, and if the doses remained stable during the study period."</p>
Spergel 2020	Children (4 to 17)	Viaskin milk 500 µg participants epicutaneously administered daily (up to 24 hours application per day) with a patch containing 500 µg cows milk proteins	Viaskin placebo participants epicutaneously administered daily (up to 24 hours application per day) with a patch containing a matching placebo formulation	Induction (no previous agent to induce induction prior randomization)	≥ 15 eos/HPF	Duration: 44 weeks Measurement points: histologic at 44 weeks	<p>PPI</p> <p>Patients were diagnosed with EoE if EGD and biopsy showed ≥ 15 eos/hpf after at least 2-month period of being on a high-dose PPI (1 to 2 mg/kg dose twice-daily)</p> <p>Diet</p> <p>All patient had 2 biopsies. First biopsy whilst being on milk-rich diet. Second biopsy on a milk-free diet. Milk was reintroduced after 9 months of randomization, where the primary endpoint was measured at the 11th month.</p> <p>Steroid</p> <p>Participants on swallowed corticosteroids for eosinophilic esophagitis were excluded</p>

Table 2. Included studies' characteristics (Continued)

Straumann 2010a	Children and adults (≥ 18)	Mepolizumab intravenous infusion at a dose of 750 mg diluted in 150 mL of 0.9% sodium chloride solution for the first 2 infusions Day 0 and day 7	Placebo IV day 0 and day 7	Induction	At least one episode of dysphagia per week in the 4 weeks prior to the start of study medication and a peak esophageal eosinophilia of ≥ 20 eosinophils	Duration: 13 weeks Measurement points: histologic, symptomatic and endoscopic at 4 and 13 weeks	Any current anti-eosinophil treatment was discontinued and patients were directed to avoid any changes in their feeding habits during the entire study period. PPI GERD was excluded in all patients by pretreatment with PPIs in standard dosages plus negative endoscopy for signs of reflux disease, and by pH monitoring (optional). Diet 1 patient excluded in placebo group due to lack of efficacy and was on elemental diet. Steroids Selected patients unresponsive to steroids. No steroids permitted at least 6 weeks prior to starting the trial.
Straumann 2010b	Children and adults (> 14)	Budesonide administered as 0.25 mg/mL suspension twice per day at bedtime and in the morning after breakfast via a nebulizer. Patients instructed swallowing continuously the accumulated liquid.	0.9% saline administered as 4 mL twice per day at bedtime and in the morning after breakfast via a nebulizer. Patients instructed swallowing continuously the accumulated liquid.	Induction	Clinicopathologic definition of esophageal symptoms in combination with ≥ 20 eosinophils per high-power field	Duration: 2 weeks Measurement points: histologic, dysphagia scores, endoscopic at 2 weeks	PPI Previously established proton pump inhibition was continued throughout the study period. Budesonide: 13/18 Placebo: 10/18 Diet Not reported Steroids Not reported
Straumann 2011	Children and adults (> 14)	0.5 mg/day budesonide as 0.25 mg/mL suspension formula-	0.9% saline 1 mL via an inhalation system con-	Maintenance	Clinically, endoscopically, and histologically confirmed eosinophilic	Duration: 50 week maintenance	PPI

Table 2. Included studies' characteristics (Continued)

		tion applied using an inhalation system consisting of a PARI UNI light compressor and PARI TIA nebulizer, twice per day at bedtime and in the morning after breakfast; patients instructed to nebulize the suspension into the oral cavity and to swallow continuously the accumulated liquid	sisting of a PARI UNI light compressor and PARI TIA nebulizer, twice per day at bedtime and in the morning after breakfast; patients instructed to nebulize the suspension into the oral cavity and to swallow continuously the accumulated liquid		esophagitis after proton pump inhibitor trial	Measurement points: histologic, symptomatic, endoscopic, and safety at 50 weeks	Previously established proton pump inhibition was continued throughout the study period. Diet Not reported Steroids Throughout the study period, participants took no other anti-eosinophil medication.
Straumann 2013	Children and adults (18 to 75)	OC004549 100 mg tablets, twice-daily after meals for 8 weeks	Placebo	Induction	Patient with previously clinically, endoscopically, and histologically confirmed EoE (according to Liacouras 2011 definition)	Duration: 8 weeks Measurement points: histologic and physician's global assessment of disease activity at 8 weeks	PPI / Steroids Patients discontinued all specific treatments for EoE (e.g. corticosteroids, leukotriene antagonists, histamine blockers, mast cell stabilizers); medications stopped 2 weeks prior to baseline exam; previously established PPI therapies for secondary reflux were continued throughout the study in a constant dose. Diet Not reported
Straumann 2020	Children and adults (18 to 75)	Budesonide orodispersible tablet 0.5 mg twice-daily and 1.0 mg twice-daily	Placebo	Maintenance	Previously confirmed diagnosis of PPI-refractory EoE according to consensus guidelines (Dellon et al. Gastroenterology 2018; Lucendo AJ et al. United Euro-	Duration: 48 week maintenance Measurement points: histologic, EEsAI-PRO an 48 weeks	PPI Concomitant PPI treatment was to be kept stable. Diet Dietary restriction was not permitted. Steroids

Table 2. Included studies' characteristics (Continued)

					pean Gastroenterol J. 2017)		The use of other swallowed topical steroids, systemic glucocorticoids, immunosuppressants or biologic drugs was not permitted.
Tytor 2021	Children and adults (≥ 18)	Mometasone furoate 4 spray doses 50 µg by mouth to be swallowed 4 times daily after meals with no eating or drinking allowed 30 minutes after intake. Duration of treatment is 8 weeks.	Placebo For 8 weeks	Induction	Newly diagnosed EoE with a peak eosinophil count of at least 15 cells per HPF in any area in any of at least 6 esophageal biopsies including at least 3 biopsies from the upper-respective lower-third part of the esophagus, and total WDS score ≥ 5	Duration: 8 weeks Measurement points: dysphagia score at 8 weeks	<p>PPI</p> <p>PPIs were not allowed from 2 weeks before the start and during the treatment period.</p> <p>Diet</p> <p>Not reported</p> <p>Steroids</p> <p>Systemic or topical corticosteroid treatment during the last 4 months was not allowed.</p>

AAF: amino acid-based formula; CG: control group; DB: double-blind; DSQ: Dysphagia Symptom Questionnaire; EDG: esophagogastroduodenoscopy; EoE: eosinophilic esophagitis; FED: food elimination diet; FFED: four-food elimination diet; GERD: gastroesophageal reflux disease; HPF/hpf: high-power field; IG: intervention group; IV: intravenous; MCT: mast cell tryptase; MDQ: Mayo Dysphagia Questionnaire; NR: not reported; PPI: proton pump inhibitor; PRO: patient-reported outcome; SC: subcutaneous; SDI: Straumann Dysphagia Instrument; WDS: Watson Dysphagia Scale

Table 3. Primary outcome - clinical improvement

Study ID	Validated symptom scoring system	Continuous or dichotomous	Outcome data - clinical symptom treatment success at study endpoint
Alexander 2012	Mayo Dysphagia Questionnaire (MDQ)- 2 week	Dichotomous: A complete symptom response was defined as an answer of "no" to the question, "In the past 2 weeks, have you had trouble swallowing, not associated with other cold symptoms (such as strep throat or mononucleosis)?" on the Mayo Dysphagia Questionnaire 2-week version. A partial symptom response was defined as an answer of "yes" to the earlier-described question and a decrease in the severity of at least 2 levels (or to a level of "Doesn't bother me at all"), or a decrease in the frequency of at least 1 level. If there was a decrease in one variable (frequency or severity) and an increase in the other variable then this was classified as no response.	Partial or complete response at end of trial (used for clinical dichotomous analysis): Fluticasone: 12/21 Placebo: 7/21 Complete at end of trial: Fluticasone: 9/21 Placebo: 6/21
	Peloquin 2006		
	Validated		
Alexander 2017	Mayo Dysphagia Questionnaire (MDQ) - 2 weeks	Dichotomous Remission was defined as the absence of dysphagia	In remission at end of trial (used for clinical dichotomous analysis): Montelukast: 8/20 Placebo: 5/21
	Peloquin 2006 Validated		
Assa'ad 2011	Pain in stomach severity score Regurgitation bothersome score Feeling something stuck in throat bothersome score	Continuous The presence and severity of the following symptoms were assessed: abdominal and chest or throat pain, regurgitation, vomiting, solid and liquid food dysphagia (age 8 to 17 years only), difficulty drinking, and difficulty eating solid foods. Reported as mean CI. Pain in stomach severity score: 0 = none 1 = a little 2 = somewhat 3 = quite a bit 4 = a whole lot Regurgitation bothersome score: 1 = not bothered at all	No primary outcome defined, no aggregate score defined. Data not used. Pain in stomach severity score: -0.277 (-0.617, 0.062), -0.149 (-0.412, 0.115), -0.157 (-0.458, 0.144) Proportion of days with pain in stomach: -14.02 (-25.93,-2.12), -12.44 (-21.66, -3.21), -10.11 (-20.70, 0.48) Pain in chest/throat severity score: -0.419 (-0.796,-0.042), -0.063 (-0.356, 0.230), -0.049 (-0.382, 0.285) Proportion of days with pain in chest/throat: -24.37 (-39.34,-9.40), -5.09 (-16.64, 6.47), -10.16 (-23.38, 3.06) Regurgitation bothersome score: -0.307 (-0.741, 0.128), 0.017 (-0.320, 0.354), -0.047 (-0.431, 0.337) Proportion of days with regurgitation: -10.26 (-24.44, 3.91), 3.8 (-7.32, 14.93), -4.64 (-17.15, 7.88)
	Flood 2008		
	Not validated		

Table 3. Primary outcome - clinical improvement (Continued)

		2 = bothered a little		Feeling something stuck in throat bothersome score: -0.751 (-1.135,-0.368), 0.238 (-0.621, 0.145), -0.510 (-0.982, -0.038)
		3 = somewhat bothered		
		4 = bothered quite a lot		Proportion of days with feeling of something stuck in throat: -21.56 (-33.73,-9.40), -12.11 (-24.29, 0.06), -17.44 (-32.34, -2.54)
		5 = bothered a whole lot		Pain with drinking: -0.005 (-0.363, 0.354), 0.085 (-0.194, 0.364), -0.166 (-0.483, 0.151)
		Feeling something stuck in throat bothersome score: no details given apart from that it is applicable to 8 to 17 years		Difficulty with drinking: 0.008 (-0.193, 0.208), 0.046 (-0.110, 0.202), -0.152 (-0.329, 0.026)
				Pain with eating solid food: -0.493 (-0.819,-0.167), -0.123 (-0.375, 0.130), -0.137 (-0.444, 0.170)
				Difficulty with eating solid food: -0.474 (-0.794,-0.153), -0.174 (-0.422, 0.074), -0.119 (-0.418, 0.180)
				Proportion of days with vomiting: -2.40 (-6.58, 1.77), -3.54 (-6.74,-0.34), -4.56 (-8.32, -0.80)
				Vomiting frequency: -0.048 (-0.296, 0.199), 0.04 (-0.149, 0.229), -0.060 (-0.281, 0.161)
Bhardwaj 2017	Daily diary card The frequency of dysphagia ± heart-burn	Dichotomous: Improved yes/no	Dichotomous improvement, first 8 weeks only (used in clinical dichotomous analysis): Budesonide: 3/9 Placebo: 1/9	
	Bhardwaj 2017 Not validated			
Butz 2014	EoE symptom score	Continuous	Available as a figure, not possible to digitize. Data not used.	
	Pentiuk 2009 Not validated	Score asks participants about the frequency and the severity of their symptoms The threshold for success is not clearly defined		
Clayton 2014	Dysphagia symptom score (DSQ)	Continuous	Change in DSQ score from baseline, mean (SD) at end of trial (used in clinical continuous analysis): Omalizumab: -1.2 (1.22)/16 Placebo: -1.7 (0.78)/14 Dysphagia score before treatment means (SD): Omalizumab: 4.0 (0.7)/16 Placebo: 5.5 (0.5)/14 Dysphagia score after treatment means (SD): Omalizumab: 2.8 (1.39)/16 Placebo: 3.8 (0.84)/14	
	Clayton 2014 Not validated	Score as follows: 0 = no dysphagia; 1 = solid food dysphagia monthly; 2 = solid food dysphagia < weekly; 3 = solid food dysphagia > weekly, < daily; 4 = solid food dysphagia daily; 5 = solid food dysphagia with every meal; and 6 = dysphagia for solid and liquid food The threshold for success is defined as a mean difference that is significant on a statistical test reported as the mean difference and the corresponding P value.		

Table 3. Primary outcome - clinical improvement (Continued)

Dellon 2012	Mayo Dysphagia Questionnaire (MDQ) Score - 30 day	Continuous The authors compared mean MDQ pre- and post-treatment	MDQ score at end of trial, mean (SD) (used for clinical continuous analysis): Nebulized budesonide solution: 10 (12)/11 Oral viscous budesonide: 16 (17)/11
	McElhiney 2009 Validated		
Dellon 2017	Dysphagia symptom questionnaire (DSQ)	Continuous The authors compared mean DSQ pre- and post-treatment	Change in score at end of trial, means (SD) (used for clinical continuous outcomes): Budesonide oral suspension: -14.3 (13)/49 Placebo: -7.5 (10.7)/38
	Hudgens 2017 Validated		
Dellon 2019	Dysphagia symptom questionnaire (DSQ)	Continuous	DSQ score at end of trial, mean (SD) (used for clinical continuous analysis): Budesonide: 4.8 (7.3)/46 Fluticasone: 4.2 (7.5)/38
	Hudgens 2017 Validated EEsAI-PRO Schoepfer 2014 Validated		EEsAI post-treatment, mean (SD): Budesonide: 22.1 (18.9)/32 Fluticasone: 28.0 (20.4)/38
Dellon 2021b	Dysphagia symptom questionnaire (DSQ)	Dichotomous and continuous Percentage of group that did not relapse	In remission at end of trial (used for clinical dichotomous analysis): Budesonide oral suspension: 19/25 Placebo: 13/23 Digitized from Figure 3A, change in DSQ at end of trial, mean (SD) (used for clinical continuous analysis): Budesonide oral suspension: -1.50 (-10.70)/24 Placebo: -0.11 (-12.06)/21 Symptom relapse at end of trial: Budesonide oral suspension: 6/25 Placebo: 10/23
	Hudgens 2017 Validated		
Dellon 2022	Dysphagia symptom questionnaire (DSQ)	Continuous	Change in DSQ score at end of trial, LS mean (SD) (used for clinical continuous analysis): Dupilumab: -21.92 (13.39) n = 28/11 imputed Placebo: -9.60 (17.20) n = 38/4 imputed
	Hudgens 2017 Validated		DSQ score at end of trial, LS mean (SE): Dupilumab: -21.92 (2.53) n = 28/11 imputed Placebo: -9.60 (2.79) n = 38/4 imputed
Dellon 2022a	Dysphagia symptom questionnaire (DSQ)	Continuous	No SDs reported. Can not use data. DSQ at end of trial: APT-1011 3 mg twice-daily: 5.6/20
	Hudgens 2017 Validated		

Table 3. Primary outcome - clinical improvement (Continued)

	EEsAI-PRO		APT-1011 3 mg at bedtime: 3.6/21
	Schoepfer 2014		APT-1011 1.5 mg twice-daily: 11.8/22
	Validated		APT-1011 1.5 mg at bedtime: 3.8/21
			Placebo: 9.1/16
Dellon 2022b	Dysphagia symptom questionnaire (DSQ)	Continuous	No SDs reported. Cannot use data.
	Hudgens 2017	Mean absolute change in DSQ at weeks 23 to 24 (no SD reported)	DSQ score, mean change at end of trial, mean, no SD
	Validated		Lirentelimab 3 mg/kg: -17.4
			Lirentelimab 1 mg/kg: -11.9
			Placebo: -14.6
De Rooij 2022	Straumann Dysphagia Instrument (SDI)	Continuous	Change in total SDI score from baseline to week 6, median [IQR] (SD).
	Straumann 2010	Change from baseline to week 6	FFED + AAF: -2 [-4, -2] (2.34)/21
	Not validated		FFED: -2.5 [-4.25, -1] (3.70)/20
Dohil 2010	Symptom scoring tool	Continuous	Symptom scoring tool at end of trial, mean (SD) (used for clinical continuous analysis):
	Dohil 2010	Pre- and post- scores	Oral viscous budesonide + PPI: 1.2 (1.87)/21
	Not validated	Heartburn/regurgitation; abdominal pain; nausea/vomiting; anorexia/early satiety; dysphagia symptom induced nocturnal waking; gastrointestinal bleeding	Placebo + PPI: 1.85 (1.8)/11
Gupta 2015	EoE Clinical Symptom Score	Dichotomous	Dichotomous symptom response (Figure 3) (used for clinical dichotomous analysis):
	Dohil 2010	Based on a physician's assessment of the frequency and disruptiveness of multiple symptoms within 6 categories (heartburn; abdominal pain; nocturnal awakening with symptoms; nausea, regurgitation, or vomiting; anorexia or early satiety; and dysphagia, odynophagia, or food impaction) and the use and disruptiveness of coping behaviors, determined by questioning of the subject and/or caregiver	Budesonide, low-dose: 11/21 (64.7%) Budesonide, medium-dose: 15/19 (78.9%) Budesonide, high-dose: 9/20 (52.9%) All treatment groups: 35/60 (58.3%) Placebo: 14/21 (77.8%)
	Not validated		
Heine 2019	Clinical symptom score	Not reported	Not reported
	Not validated		
Hirano 2019	Daily Dysphagia Symptom Diary (DSD) scores	DSD: continuous, mean change	DSD, digitized from Figure 3D, week 16, mean reduction in the DSD composite score (SD) (used for clinical continuous analysis):
	Hirano 2019	EEsAI: continuous, mean change	RPC4046: -13.31 (15.26)/34
			Placebo: -6.41 (15.40)/34

Medical treatment of eosinophilic esophagitis (Review)

323

Table 3. Primary outcome - clinical improvement (Continued)

	Not validated	Patient's global impression of EoE symptoms: dichotomous	Patient's global impression of EoE symptoms, dichotomous, from Figure 4C (used for clinical dichotomous analysis): RPC4046: $0.444 (31) + 0.222 (31) + 0.645 (34) + 0.194 (34)/66 = 14 + 7 + 22 + 7 = 50/66$ Placebo: $0.364 (34) + 0.212 (34)/34 = 12 + 7 = 19/34$
	Eosinophilic Esophagitis Activity Index (EESAI)		
	Schoepfer 2014		
	Validated		
	Patient's global impression of EoE symptoms		
	Hirano 2019		
Hirano 2020	Straumann Dysphagia Instrument (SDI)	Continuous (primary outcome was the change in value of the PRO score at week 10. A secondary outcome was the change as a percentage)	SDI PRO mean change Mean (SE) (used for clinical continuous analysis): Dupilumab: $-3.2 (0.61)/17$ SD: 2.52
	Straumann 2010		
	No validated		
	Eosinophilic Esophagitis Activity Index (EESAI) scores	Dichotomous (one of the secondary outcomes was a PRO score change of equal to or greater than 3)	Placebo: $-1.1 (0.67)/14$ SD: 2.51
	Schoepfer 2014		EESAI PRO \geq 40% improvement from baseline (used for clinical dichotomous analysis): Dupilumab: 6/23 Placebo: 2/24
Hirano 2020f	EESAI was adapted for use in this trial	Continuous	All clinical response data are for post hoc analyses only, cannot use
	Schoepfer 2014		
	Not validated in adapted form		
	Patient eosinophilic esophagitis global assessment		
	Schoepfer 2014		
	Validated		
	Physician eosinophilic esophagitis Global Assessment		
	Schoepfer 2014		
	Validated		
	Mayo Dysphagia Questionnaire-30		
	McElhiney 2009		
	Validated		

Table 3. Primary outcome - clinical improvement (Continued)

	Gastrointestinal Symptom Rating Scale	Validated	
Hirano 2021	Dysphagia Symptom Questionnaire (DSQ)	Continuous	DSQ at end of trial, mean (SD) (used for clinical continuous analysis):
Hudgens 2017		Dichotomous $\geq 30\%$ reduction in DSQ score from baseline to week 12	Budesonide oral suspension: 19.5 (17.0)/198 Placebo: 22.6 (17.5)/89
			Dichotomous (used for clinical dichotomous analysis): Budesonide oral suspension: 112/213 Placebo: 41/105
Kliewer 2019	PEESS V2.0	Continuous	PEESS at end of trial, mean (SD) (used for clinical continuous analysis):
Franciosi 2011			1-FED: 23.5 (18.3)/33 4-FED: 16.0 (13.0)/17
	Validated		
Kliewer 2021	EoE Symptom Activity Index (EESAI)	Continuous	Change in EESAI at end of trial, mean (SD) (used for clinical continuous analysis):
Schoepfer 2014			1-FED: -3.0 (16.9)/67 6-FED: -8.2 (17.0)/62
	Validated		
Konikoff 2006	Clinical symptom assessment	Dichotomous	No prespecified aggregate value reported, cannot use data.
Konikoff 2006			FP improves vomiting. The most common clinical symptoms at the start of the study were abdominal pain (reported in 16/28 patients for whom symptom information was available (57%)), vomiting (15/28 (54%)), and dysphagia (13/29 (45%)). Only vomiting improved significantly with treatment with FP (67% pretreatment vs 27% post-treatment). All patients who responded histologically had a concurrent resolution of their vomiting (6/6), while vomiting in FP non-responders did not resolve (0/4).
	Not validated		
Lieberman 2018	PEESS V1.0	Continuous: PEES	Digitized from Figure 3 mean (SD) at end of trial (used for clinical continuous analysis):
Pentiuk 2009			Cromolyn sodium: 17.5 (19.2)/8 Placebo: 22.2 (12.8)/6
	Not validated		
Lucendo 2019	NRS for dysphagia, odynophagia	Dichotomous	Rate of patients with clinical remission (as defined in the primary end point) at end of trial (used as clinical dichotomous outcome):
Lucendo 2019		Clinical remission (symptoms severity of 2 points on each 0 to 10 NRS for dysphagia and odynophagia, respectively on each day in the week before end of trial)	Budesonide: 35/59 Placebo: 4/29
	NRS for PatGA and PGA		
	EESAI-PRO		

Table 3. Primary outcome - clinical improvement (Continued)

Schoepfer 2014			
	Validated		
	Dysphagia-free days		
Miehlke 2016	Straumann Dysphagia Instrument (SDI)	Ordinal (0 to 9)	Decrease in mean dysphagia score from baseline at end of trial. Digitized from supplementary Figure 2, mean dysphagia score (SD) (used in clinical continuous analysis): Budesonide: -2.34 (2.66)/53 Placebo: -1.99 (2.85)/17
	Straumann 2010	Dichotomous	
	Not validated	Clinical response defined as a decrease in the dysphagia score of at least 3 points compared with baseline Frequency of dysphagia ranging from none (0) to several times per day (4) and intensity of dysphagia ranging from unhindered swallowing (0) to long-lasting complete obstruction requiring endoscopic intervention (5). Total scores ranged from 0 to 9.	
Moawad 2013	Mayo Dysphagia Questionnaire (MDQ) - 2 weeks	Continuous	MDQ at end of trial, mean (SD) (used for clinical continuous analysis): Esomeprazole Pos: 1.4 (4.5)/21 Fluticasone Pos: 12 (16)/19
	Peloquin 2006		
Oliva 2018	Not reported	Not reported	Not reported
Peterson 2010	Dysphagia scale	Continuous	Dysphagia score at end of trial, mean (SD) (used for clinical continuous analysis):
	DiSario 2002	The dysphagia scale ranged from 0 to 7	Esomeprazole: 2.3 (2.0)/11
	Not validated	A score of 0 = no dysphagia; 1 = solid food dysphagia once in 3 to 12 months; 2 = solid food dysphagia once in 1 to 3 months; 3 = solid food dysphagia once every 2 to 4 weeks; 4 = solid food dysphagia once every 1 to 2 weeks; 5 = solid food dysphagia once every 1 to 7 days; 6 = solid food dysphagia with every meal; 7 = dysphagia to solid and liquid food	Fluticasone: 1.7 (1.6)/12
	Revalidated reflux disease questionnaire (RDQ)	No details were provided for the revalidated reflux disease questionnaire (RDQ) apart from a pre-treatment score	
	Aanen 2006		
	Not validated		

Table 3. Primary outcome - clinical improvement (Continued)

		No threshold of success provided for both score, but a percentage of change is noted	
Rothenberg 2015 Mayo Dysphagia Questionnaire (MDQ) - 2 weeks Peloquin 2006	Dichotomous Sum change of Mayo Dysphagia Questionnaire items 1, 2, 4, 9, 10, 13, 14, 16, 20, 21. A positive sum change was scored as an improvement, a negative sum change was scored as a worsening, and a zero-sum change was scored as unchanged. Continuous MDQ score digitized from supplement article Figure E1	Sign of change from baseline in MDQ score, positive vs no change and negative at end of trial (used for clinical dichotomous analysis): Treatment: 10/17 Placebo: 4/8 Data extracted from supplementary Figure E1, MDQ score, mean (SD) at end of trial (used for clinical continuous analysis): Treatment: 1.933333 (3.494213)/15 Placebo: 0.285714 (3.638419)/7	
Rothenberg 2022 Hudgens 2017	Continuous	Least squares mean absolute changes in DSQ score at end of trial, mean (SE) (used for clinical continuous analysis): Dupilumab: -23.78 (1.86), n not reported 80?, SD (16.64) Placebo: -13.86(1.91), n not reported 79?, SD (16.98)	
Schaefer 2008	Resolution of the presenting symptom(s) including, abdominal pain, dysphagia, epigastric pain, foreign body, feeding problems, heartburn, regurgitation, vomiting, and weight loss. Schaefer 2008 Not validated A daily symptom diary was maintained by the patient/guardian while on corticosteroid therapy. Clinical assessment was performed at weeks 4, 12, 18, and 24 to monitor for the presence or absence of the presenting esophageal symptom(s). Schaefer 2008 Not validated	Dichotomous Proportion of symptom-free patients at follow-up (4 weeks). Kaplan–Meier analysis was performed including all 80 patients based on intention-to-treat analysis. A log-rank test was used to compare survival curves between treatments.	Proportion of symptom-free patients at end of trial (used for clinical dichotomous analysis): Prednisone: 32/40 Fluticasone: 35/40

Table 3. Primary outcome - clinical improvement (Continued)

Spergel 2012	Physician's EoE global assessment	Continuous and dichotomous	Physician's EoE global assessment
	Schoepfer 2014	Taking into account physical findings, vital signs, the patient's predominant eosinophilic esophagitis symptom assessment, the patient's symptom diary, and dietary questions, physicians answered the following question: "In the review of the subject's symptoms and the physical assessment, what is your global assessment of the subject's eosinophilic esophagitis?" Physicians answered "none", "mild", "moderate", "severe", or "very severe".	Mean shift score from baseline to end of therapy. No SD reported, cannot use.
	Validated		<ul style="list-style-type: none"> • Reslizumab (1 mg) = -0.85 • Reslizumab (2 mg) = -1.02 • Reslizumab (3 mg) = -1.12 • Placebo = -1.14
	Patient's predominant EoE symptom		Patient's predominant EoE symptom
	Schoepfer 2014		Mean shift score from baseline to end of therapy. No SD reported, cannot use.
	Validated		<ul style="list-style-type: none"> • Reslizumab (1 mg) = -0.94 • Reslizumab (2 mg) = -1.20 • Reslizumab (3 mg) = -1.28 • Placebo = -1.44
		Patient's predominant EoE symptom	Dichotomous: Physician's Global Assessment at end of trial (used for clinical dichotomous analysis): Reslizumab (1 mg): 31/56 Reslizumab (2 mg): 32/57 Reslizumab (3 mg): 37/57 Reslizumab: 100/170 Placebo: 37/57
Spergel 2020	PEESS V2.0	Continuous	Change from baseline in PEESS V2.0, mean (SD) at end of trial (used for clinical continuous analysis):
	Franciosi 2011	Total score is reported with a range of 0 to 9. A lower score is better.	Viaskin milk: 0.71 (1.11)/7
	Validated		Placebo: 2.00 (1.41)/2
Straumann 2010a	Esophagus-related symptom score	Dichotomous	Improvement at end of trial (used for clinical dichotomous analysis):
	Straumann 2003	Clinical improvement in eosinophilic esophagitis	Mepolizumab: 3/5 Placebo: 3/6
	Not validated	Continuous, percent dysphagia days	Digitized from Figure 3B percent dysphagia days at end of trial, mean (SD) (used for clinical continuous analysis):
	Dysphagia days		Mepolizumab: 71.91 (17.34)/5
	Straumann 2010a		Placebo: 55.14 (20.83)/6
	Not validated		
Straumann 2010b	Clinical symptoms were assessed by frequency and intensity of dysphagia events without use of a validated PRO	Dichotomous	Clinical response (used for clinical dichotomous analysis):
		Clinical response	Budesonide: 13/18
		Continuous	Placebo: 4/18
		Post-scores (mean ± SD)	Symptom scores at end of trial, mean (SD) (used for clinical continuous analysis):
	Straumann 2010	The following non-validated scores were used to assess dysphagia. Frequency of dysphagia events; none = 0; once per	Budesonide: 2.22 (2.02)/18

Table 3. Primary outcome - clinical improvement (Continued)

		<p>week = 1; several times per week = 2; once per day = 3; and several times per day = 4. Intensity of dysphagia events: swallowing unhindered = 0; slight sensation of resistance = 1; slight retching with delayed passage = 2; short period of obstruction necessitating intervention = 3; longer-lasting period obstruction only removable by vomiting = 4; and long-lasting complete obstruction requiring endoscopic intervention = 5</p> <p>A clinical response was defined as a decrease in the dysphagia score of at least 3 points compared with baseline</p>	Placebo: 4.72 (1.96)/18
Straumann 2011	Dysphagia symptom score	Continuous, pre- and post-scores	Symptom score at end of trial, mean (SD) (used for clinical continuous analysis):
	Straumann 2010		Budesonide: 0.79 (1.37)/14
	Not validated		Placebo: 0.71 (1.20)/14
Straumann 2013	Dysphagia was assessed using a visual dysphagia questionnaire (VDQ)	Continuous (compared post-treatment means); no pre-specified response threshold	Total PRO score at end of trial, mean (SD) (used for clinical continuous analysis):
	Straumann 2013		OC000459: 10.79 (6.52)/14
	Not validated		Placebo: 9.73 (8.16)/12
	Chest pain was assessed using a "pain questionnaire"		
	Straumann 2010		
	Not validated		
	Combined the VDQ and pain questionnaire for a "total score"		
	Physician's global assessment		
	Validated		
Straumann 2020	Dysphagia assessed via numerical rating scale (1 to 10)	Dichotomous for dysphagia and odynophagia (clinical remission defined as a severity of ≤ 2 points on 1- to 10-point numerical rating scale (NRS) for dysphagia and a severity of ≤ 2 points on a 0- to 10-point NRS for odynophagia on each day in the last week of induction treatment)	Weekly EEsAI-PRO score of ≤ 20 at end of trial (used for clinical dichotomous analysis):
	Straumann 2020		Budesonide 0.5 mg twice-daily: 49/68 Budesonide 1.0 mg twice-daily: 50/68
	Not validated		Budesonide: 99/136 Placebo: 14/68

Table 3. Primary outcome - clinical improvement (Continued)

	Odynophagia assessed via numerical rating scale (1 to 10)	EEsAI-PRO score also dichotomous ≤ 20 at end of treatment	EEsAI-PRO at end of trial, mean (SD) (used for clinical continuous analysis): Budesonide 0.5 mg: 14 (18.5)/65 Budesonide 1.0 mg: 11 (18.0)/66
	Straumann 2020		
	Not validated		Budesonide: 12.49 (18.10)/131 Placebo: 39 (21.4)/65
	EEsAI		
	Schoepfer 2014		
	Validated		
Tytor 2021	Watson Dysphagia Score (WDS)	Continuous, difference in Watson Dysphagia Scale score (0 to 45) at 8 weeks from screening	Change in Watson Dysphagia Scale score at end of trial, mean (SD) (used for clinical continuous analysis): Mometasone: -6.0 (7.1)/16 Placebo: -1.8 (6.7)/17
	Dakkak 1992		
	Not validated		

BOS: budesonide oral suspension; BOT: budesonide orodispersible tablet; CG: control group; CI: confidence interval; DSD: daily symptom diary; DSQ: Dysphagia Symptom Questionnaire; EEsAI: Eosinophilic Esophagitis Activity Index; EoT: end of treatment; FP: fluticasone propionate; HD: high-dose; IG: intervention group; LD: low-dose; LS: least squares; NEB: nebulized/swallowed budesonide solution; NRS: numerical rating scale; OVB: viscous/swallowed budesonide solution; PEES: Pediatric Eosinophilic Esophagitis Symptom Severity; PRO: patient-reported outcome; SE: standard error

Table 4. Primary outcome - histological improvement

Study ID	Histological improvement system used	Continuous or dichotomous	Outcome data - histological improvement at study endpoint
Alexander 2012	Eosinophils were counted using a 40x objective, a field diameter of 0.625 mm, and a field area of 0.307 mm ² . The peak eosinophil count per high-powered field was reported. From the area of the greatest density under low-powered review, 5 random fields were chosen. Peak eosinophil counts from these 5 fields were used to calculate a mean eosinophil count.	Continuous A complete histologic response was defined as a decrease in the mean eosinophil level of more than 90% from the pre-treatment value. A partial response was defined as a decrease in more than 50% from the pre-treatment value.	Partial or complete at end of trial (used for histological dichotomous analysis): Fluticasone: 17/21 Placebo: 1/21 Complete (≤ 2) at end of trial: Fluticasone: 13/21 Placebo: 0/21
Alexander 2017	No histological scoring system used	Not reported	Not reported
Assa'ad 2011	Peak eosinophils count	Dichotomous Measuring number of patients with mean	Used for histological dichotomous analysis at end of trial (< 15): Mepolizumab 0.55 mg/kg: 4/19

Medical treatment of eosinophilic esophagitis (Review)

Table 4. Primary outcome - histological improvement (Continued)

		peak eos \leq 5 (complete responders)	Mepolizumab 2.5 mg/kg: 9/20 Mepolizumab 10 mg/kg: 5/20 Mepolizumab combined: 18/59
Bhardwaj 2017	Eosinophils/hpf	Continuous Amount of eosinophils in esophageal tissue (cells/hpf) compared to baseline	Eos/hpf at end of trial from Table 1, mean (SD) (used for histological continuous analysis): Beclomethasone: 2 (4) n = 4 Placebo: 22.2 (23.1) n = 5
Butz 2014	Peak eosinophils/hpf	Dichotomous Defined as mean peak eosinophils of \leq 1 eos/hpf in both distal and proximal	Dichotomous outcome mean peak \leq 1 at end of trial: Fluticasone: 15/28 Placebo group: 0/14 Dichotomous outcome mean peak \leq 6 at end of trial: Fluticasone: 17/28 Placebo group: 0/14 Dichotomous outcome mean peak \leq 14 at end of trial (used for histological dichotomous analysis): Fluticasone: 18/28 Placebo group: 1/14
Clayton 2014	Eosinophils/hpf	Continuous Pre- and post-treatment eosinophils/hpf	Eos/hpf at end of trial, mean (SD) (used for histological continuous analysis): Omalizumab: 39 (15)/16 Placebo: 33 (12)/14
Dellon 2012	Eosinophils/hpf 2 biopsies were procured from the distal esophagus (3 cm above the gastroesophageal junction), 1 from the mid-esophagus (8 cm above), and 2 from the proximal esophagus (13 cm above). A total of 5 hpf (hpf; hpf size = 0.24 mm ²) were examined per each of the 5 biopsy specimens from each patient and the maximum eosinophil count (eos/hpf) was defined as the count in the hpf in the area of the highest eosinophil density after review of all 25 hpf in each patient. The mean eosinophil count for each patient	Continuous Post-treatment max and mean eos count was reported Dichotomous Histological response (complete, near-complete, and partial responses) for both groups as well	Post-treatment peak eosinophil count (SD) at end of trial (used for histological continuous analysis): Budesonide, nebulized: 89 (94)/11 Budesonide, oral viscous: 11 (23)/11 Complete response (mean peak < 1 eos/hpf at end of trial): Budesonide, nebulized: 3/13 Budesonide, oral viscous: 7/12 Near-complete (mean peak < 7 eos/hpf at end of trial): Budesonide, nebulized: 4/13 Budesonide, oral viscous: 8/12 Partial (mean peak < 15 eos/hpf at end of trial) (used for histological dichotomous analysis): Budesonide, nebulized: 5/13 Budesonide, oral viscous: 8/12

Table 4. Primary outcome - histological improvement *(Continued)*

was calculated after examination of all 25 hpf.

Dellon 2017	Peak ≤ 6 eosinophils/hpf and histological responders	Dichotomous Peak ≤ 6 eos/hpf Continuous Mean difference	Mean peak difference (SD) eosinophils/hpf at end of trial (used for histological continuous analysis): Budesonide: $-117.0 (111.6)/49$ Placebo: $-17.3 (83.8)/38$ Mean peak ≤ 6 eos/hpf at end of trial (used for histological dichotomous analysis): Budesonide: $n = 19/51$ Placebo: $n = 1/42$
Dellon 2019	Eosinophils/hpf	Continuous and dichotomous Mean value and < 15 eos/hpf	Post-treatment, eos/hpf, mean peak (SD) at end of trial (used for histological continuous analysis): Budesonide: $14.7 (29.0)/56$ Fluticasone: $20.9 (34.3)/55$ Mean peak < 15 eos/hpf at end of trial (used for histological dichotomous analysis): Budesonide: $40/65$ Fluticasone: $35/64$
Dellon 2021b	Eosinophils/hpf	Continuous and dichotomous Percentage that did not relapse	Histologic response across all esophageal regions, peak mean eos/hpf at end of trial: Mean peak ≤ 1 eos/hpf at end of trial: Budesonide: $15/25$ Placebo: $0/23$ Mean peak ≤ 6 eos/hpf at end of trial: Budesonide: $19/25$ Placebo: $1/23$ Mean peak < 15 eos/hpf at end of trial (used for histological dichotomous analysis): Budesonide: $19/25$ Placebo: $3/23$ Digitized from figure 3B (used for histological continuous analysis): Budesonide: $15.2(45.8) / 24$ Placebo: $76.8 (50.5) / 21$
Dellon 2022	Peak eosinophils/hpf	Dichotomous	Mean peak ≤ 6 eos/hpf at end of trial Dupilumab: $25/42$ Placebo: $2/39$

Table 4. Primary outcome - histological improvement (Continued)

			Mean peak < 15 eos/hpf at end of trial (used for histological dichotomous analysis): Dupilumab: 27/42 Placebo: 3/39
Dellon 2022a	≤ 6 peak eosinophils/hpf	Dichotomous Histologic response at week 12, defined as the percentage of participants with ≤ 6 peak eos/hpf	≤ 6 mean peak eos/hpf at end of trial (used for histological dichotomous analysis): APT-1011 3 mg twice-daily: 16/20 APT-1011 3 mg at bedtime: 14/21 APT-1011 1.5 mg twice-daily: 19/23 APT-1011 1.5 mg at bedtime: 10/21 APT-1011: 59/85 Placebo: 0/21
Dellon 2022b	≤ 6 eosinophils/hpf in peak hpf Peak eosinophils ≤ 1 eosinophils/hpf at week 24	Dichotomous n of patients with eos ≤ 6 at 24 weeks (primary endpoint) n of patients with eos ≤ 1 at 24 weeks	≤ 6 mean peak eos/hpf at end of trial (used for histological dichotomous analysis): Lirentelimab 3 mg/kg: 80/91 Lirentelimab 1 mg/kg: 86/93 Lirentelimab: 166/184 Placebo: 10/92 Mean peak ≤ 1 eos/hpf at end of trial: Lirentelimab 3 mg/kg: 77/91 Lirentelimab 1 mg/kg: 82/93 Lirentelimab: 159/184 Placebo: 4/92
De Rooij 2022	Peak eosinophils count	Continuous Mean of absolute change in peak eosinophil count from baseline to week 6, eos/hpf, mean (SD) Dichotomous Histological remission rates at 6 weeks	Absolute change in peak eos count at end of trial (SD) (used for histological continuous analysis): FFED: -26.2 (39.9)/20 FFED + AAF: -40 (36)/21 Mean peak < 15 eos/hpf at end of trial (used for histological dichotomous analysis): FFED: 5/20 FFED + AAF: 10/21
Dohil 2010	Eosinophils count/hpf	Continuous: peak eos/hpf Dichotomous: < 20 peak eos/hpf	Continuous outcomes, mean peak eos/hpf at end of trial (SD): Budesonide: 4.8 (7.0)/15 Placebo + PPI: 65.6 (43.3)/9 Dichotomous outcomes ≤ 6 mean peak at end of trial, (used for histological dichotomous analysis): Budesonide: 14/21 Placebo: 1/11

Table 4. Primary outcome - histological improvement (Continued)

Gupta 2015	At least 2 mucosal pinch biopsies were obtained from the proximal, mid, and distal esophagus - all 3 esophageal levels Peak eosinophil count of ≤ 6 eosinophils/hpf	Dichotomous	Peak eosinophil count of mean peak ≤ 6 /hpf at end of trial (used for histological dichotomous analysis): Budesonide, low-dose: 2/21 Budesonide, medium-dose: 8/19 Budesonide, high-dose: 13/20 Budesonide: 23/60 Placebo: 0/21
Heine 2019	The peak eosinophil count (eosinophils/hpf)	Median eosinophil counts	Median mucosal eosinophil counts (IQR) at end of trial PPI + four food elimination diet: 2.5 (IQR 0.5 to 19)/27, SD: 24.52 PPI: 12 (IQR 0 to 37)/31, SD: 52.55 <10 eosinophils/HPF at end of trial PPI + four food elimination diet: 22/32 (69%) PPI: 14/32 (44%)
Hirano 2019	Eosinophils per high-power field (eos/hpf)	Eosinophils/hpf: continuous/dichotomous Response = < 15 eos/hpf Complete response = < 6 eos/hpf	Eosinophils/hpf change from baseline at end of trial (SD) (used for histological continuous analysis): RPC4046 180 mg: -94.76 (67.27)/28 RPC4046 360 mg: -99.90 (79.53)/30 RPC4046: -97.42 (72.02)/58 Placebo: -4.42 (59.94)/32 Mean peak < 15 eos/hpf at end of trial (used for histological dichotomous analysis): RPC4046 180 mg: 14/32 (44%) RPC4046 360 mg: 15/34 (44%) RPC4046: 29/66 Placebo: 0/34 (0%) Mean peak < 6 eos/hpf at end of trial: RPC4046 180 mg: 7/32 (22%) RPC4046 360 mg: 6/34 (18%) RPC4046: 13/66 Placebo: 0/34 (0%)
Hirano 2020	Intraepithelial eosinophils/hpf	Continuous (mean intraepithelial eos/hpf change) Dichotomous	LS mean change from baseline (SE) at end of trial, eos/HPF (used for histological continuous analysis): Dupilumab: -96.4 (9.44)/23, SD = 45.3 Placebo: -9.7 (9.65)/24, SD = 47.3 Patients with response mean peak < 1 eos/hpf at end of trial: Dupilumab: 3/23 (13.0%)

Table 4. Primary outcome - histological improvement (Continued)

			Placebo: 0/24 Patients with response ≤ 6 mean peak eos/hpf at end of trial (used for histological dichotomous analysis): Dupilumab: 15/23 Placebo: 0/24 Patients with response < 15 mean peak eos/HPF at end of trial post hoc: Dupilumab: 19/23 Placebo: 0/24
Hirano 2020f	Esophageal eosinophil counts per hpf in all parts of the esophagus were assessed	Continuous (mean/median)	Mean change in eos/hpf at end of trial, no SD, cannot use: APT-1011 1 mg: -63.8 APT-1011 3 mg: -34.0 Placebo: -14.8 < 15 mean peak eos/hpf at end of trial (used for histological dichotomous analysis): APT-1011 1 mg: 6/8 APT-1011 3 mg: 5/8 APT-1011: 11/16 Placebo: 1/8 0 eos/hpf at end of trial: APT-1011: 10/16 Placebo: 1/8
Hirano 2021	Maximum peak eosinophil count, eos/hpf Proportion of strict histologic responders (≤ 6 eos/hpf across all available esophageal levels (proximal, middle, or distal)) Proportion of patients achieving a deep histologic response, < 1 eos/hpf Proportion of patients achieving a histologic response (< 15 eos/hpf)	Continuous (mean) Dichotomous (see previous entry)	Mean peak eos/hpf (SD) at end of trial (used for histological continuous analysis): Budesonide: 21.9 (34.6)/201 Placebo: 69.9 (38.4)/92 Dichotomous mean peak eos/hpf < 15 at end of trial (used for histological dichotomous analysis): Budesonide: 132/215 Placebo: 1/107 Dichotomous outcome mean peak eos/hpf ≤ 6 at end of trial: Budesonide: 113/215 Placebo: 1/107 Dichotomous outcome mean peak eos/hpf ≤ 1 at end of trial: Budesonide: 69/215 Placebo: 0/107
Kliewer 2019	Dichotomous: remission is defined as clinical esophageal peak eosinophil count < 15 eosinophils per high-power field (eos/hpf) at 12 weeks	Dichotomous	From NCT02610816 at end of trial mean peak eos/hpf < 15 (used remission for histological dichotomous analysis): 1-food elimination diet: Remission: 24/38 Partial remission: 8/38

Table 4. Primary outcome - histological improvement (Continued)

	Complete remission is defined as ≤ 1 peak eos/hpf and partial remission as 2 to 14 peak eos/hpf at 12 weeks		Complete remission: 7/38 4-food elimination diet: Remission: 7/25 Partial remission: 4/25 Complete remission: 3/25
Kliewer 2021	Peak eos/hpf < 15	Dichotomous and continuous	Dichotomous mean peak eos/hpf < 15 at end of trial (used for histological dichotomous analysis): 1-food elimination: 23/67 6-food elimination: 25/62 Continuous, decrease in peak eos/hpf from baseline (SD) at end of trial: 1-food elimination diet: -24.5 (57.6)/67 6-food elimination diet: -17.7 (41.3)/62
Konikoff 2006	Peak eosinophil count eos/hpf	Dichotomous, remission defined as a peak eosinophil count of ≤ 1 eosinophil/hpf. Also reported ≤ 6 eos/hpf.	Mean peak ≤ 1 eos/hpf at end of trial: Fluticasone: 10/21 Placebo: 1/15 Mean peak ≤ 6 eos/hpf at end of trial (used for histological dichotomous analysis): Fluticasone: 11/21 Placebo: 2/15
Lieberman 2018	Change in peak eos/hpf from baseline following 8 weeks of treatment (magnification and field of view not specified; email PI)	Continuous: eos/hpf	Eosinophils/hpf (SD) at end of trial digitized from Figure 2 (used for histological continuous analysis): Cromolyn sodium: 57.3 (44.0)/9 Placebo: 71.4 (52.8)/6
Lucendo 2019	Peak eosinophils, eos/hpf	Dichotomous (rate of patients with histologic remission (i.e. peak eos < 16/mm ² hpf; equivalent to < 5 eos/hpf) at week 6)	Mean peak eos < 16/mm ² hpf at end of trial (used for histological dichotomous analysis): Budesonide: 55/59 Placebo: 0/29
Miehlke 2016	Mean eos/mm ² /hpf (On each esophageal biopsy specimen, all levels were surveyed and the eosinophils in the most densely infiltrated area were counted in 5 hpf. Total of biopsies per patient: 6. Total of hpf evaluated 30).	Dichotomous Rate of histological remission defined as mean of < 16 eos/mm ² /hpf Continuous Change in mean number of eos/mm ² /hpf from baseline to end of trial	Mean peak of < 16 eos/mm ² /hpf at end of trial (2 weeks) (used for histological dichotomous analysis): Budesonide effervescent tablet 2 x 1 mg: 19/19 Budesonide effervescent tablet 2 x 2 mg: 17/19 Budesonide oral viscous suspension 2 x 2 mg: 17/19 Budesonide: 53/57 Placebo: 0/19 Change in mean number of eos/mm ² /hpf at end of trial, no SD reported, cannot use: Budesonide effervescent tablet 2 x 1 mg: -227 Budesonide effervescent tablet 2 x 2 mg: -287

Table 4. Primary outcome - histological improvement (Continued)

			Budesonide oral viscous suspension 2 x 2 mg: -180 Placebo: -8
Moawad 2013	Change in peak eosinophils/hpf and eosinophils \leq 7/hpf	Continuous Peak eos/hpf Dichotomous \leq 7 eos/hpf	Peak mean eos/hpf (SD) at end of trial (used for histological continuous analysis): Esomeprazole: 30.5 (33.7)/21 Fluticasone: 39.2 (29.4)/21 Mean peak \leq 7 eos/hpf at end of trial (used for histological dichotomous analysis): Esomeprazole: 7/21 Fluticasone: 4/21
Oliva 2018	Peak eosinophil count/hpf	Dichotomous Histologic response < 15 eos/hpf	Only percentages, no numbers as total number of patients in each group not reported at end of trial: 6-food elimination diet: 69% Swallowed fluticasone: 67% Swallowed budesonide: 75% Oral viscous budesonide: 85%
Peterson 2010	Eosinophils/hpf	Continuous (note participant counts in Table 1 are misleading for post therapy eos/hpf; see Figure 3 and 4) Partial resolution as \leq 15 eos/hpf and complete resolution as \leq 5 eos/HPF	Post-treatment mean max (SD) at end of trial (used for histological continuous analysis): Esomeprazole: 37.2 (28.6)/12 Fluticasone: 48.1 (42)/13 Mean peak \leq 15 eos/hpf at end of trial (used for histological dichotomous analysis): Esomeprazole: 6/15 Fluticasone: 4/15 Mean peak \leq 5 eos/hpf at end of trial: Esomeprazole: 4/15 Fluticasone: 2/15
Rothenberg 2015	Eosinophils/hpf	Continuous Mean peak eos/hpf Dichotomous Responders were defined by a reduction in the peak eosinophil counts per hpf by 75% or more at day 85, compared with the baseline counts	Mean peak eos/hpf reduced by 75% at end of trial (used for histological dichotomous analysis): QAX576: 6/17 Placebo: 1/8 Mean peak eos/hpf (SD) digitized from Figure 2 at end of trial (used for histological continuous analysis): QAX576: 42.7 (52.7)/15 Placebo: 72.5 (40.8)/8
Rothenberg 2022	Eosinophils/hpf	Dichotomous mean peak \leq 6 eosinophils	Mean peak \leq 6 eos/hpf at end of trial (used for histological dichotomous analysis): Dupilumab: 47/80

Table 4. Primary outcome - histological improvement (Continued)

			Placebo: 5/79
Schaefer 2008	Basal cell zone thickness as a percentage of the epithelial thickness and the maximum number of eos/hpf	Continuous Points were assigned based on basal cell zone thickness as a percentage of the epithelial thickness, and the maximum number of eos/hpf. Points were summed and the totals were translated into histologic grades (normal, mild, moderate, and severe). Grades were assigned a numeric value for statistical analysis.	Histologic improvement by a grade of 1 or more at end of trial (used for histological dichotomous analysis): Prednisone: 75% (30/40) Fluticasone: 85% (34/40) "Complete" histologic resolution (defined as normal biopsy specimens; at end of trial): Prednisone: 65% (26/40) Fluticasone: 45% (18/40) Continuous, data from extracted Table 4 at end of trial (used for histological continuous analysis): Mean peak eos/hpf at end of trial (SD): Prednisone: 2.13 (7.75)/32 Fluticasone: 6.58 (11.55)/36
Spergel 2012	Peak esophageal eosinophil count and the change from baseline to the end of therapy	Continuous Dichotomous: mean peak < 15 eos/hpf	Results are from supplementary Figure E1 that has been digitized; mean peak < 15 eos/hpf at end of trial (used for histological dichotomous analysis): Reslizumab 1 mg/kg: 33/55 Reslizumab 2 mg/kg: 32/57 Reslizumab 3 mg/kg: 39/57 Reslizumab: 104/169 Placebo: 20/57 Eosinophils/hpf mean (SD) at end of trial (used for histological continuous analysis): Reslizumab 1 mg/kg: 42.1 (46.5)/40 Reslizumab 2 mg/kg: 23.9 (25.0)/38 Reslizumab 3 mg/kg: 35.9 (23.3)/45 Reslizumab: 37.0 (23.4)/123 Placebo: 99.6 (62.4)/46
Spergel 2020	Change in maximum esophageal eosinophil count from baseline to end of double-blind treatment	Continuous No threshold of success was identified but a mean difference of change was calculated	Mean peak eos at end of trial (SD) (used for histological continuous analysis): Viaskin milk: 25.57 (31.19)/7 Placebo: 95.00 (63.64)/2 Change from baseline mean (SD) at end of trial: Viaskin milk: -26.86 (22.53)/15 Placebo: 42.50 (31.82)/5
Straumann 2010a	Peak eosinophil count/hpf (area 0.3072 mm ²) Mean eos/hpf	Dichotomous ≤ 5 eos/hpf Continuous Percentage of reduction in mean eos/hpf	Mean peak ≤ 5 eos/hpf at end of trial (used for histological dichotomous analysis): Mepolizumab: 0/5 Placebo: 0/6 Mean (SD) eos/hpf end of trial (used for histological continuous analysis): Mepolizumab: 33.83 (22.82)/5 Placebo: 53.99 (24.51)/6
Straumann 2010b	Reduction in the esophageal eosinophil	Continuous (means)	Mean peak eos/hpf (SD) at end of trial (used for histological continuous analysis):

Table 4. Primary outcome - histological improvement (Continued)

	load; also present peak eosinophil counts and categorized by the level of response; eosinophil load defined as mean eosinophil number measured in a total of 40 hpf from 2 x 4 biopsy specimens taken from the proximal and distal esophagus	Dichotomous (by the degree of remission) At least 4 biopsy specimens were taken endoscopically from the proximal half and from the lower half of the esophagus and additionally from any lesion. In all 8 esophageal biopsy specimens, all levels were surveyed and the eosinophils in the most densely infiltrated area were counted in 5 consecutive hpf (area of field 0.3072 mm ²).	Budesonide: 17.7 (26.7)/18 Placebo: 125.6 (67.6)/18 Budesonide vs placebo ≤ 5: 13/18 vs 2/18 5 to 20: 3/18 vs 0/18 > 20: 2/18 vs 16/18 Mean peak ≤ 20 eos/hpf (used for histological dichotomous analysis): Budesonide: 16/18 Placebo: 2/18
Straumann 2011	Looked at eosinophil load; eos load defined as the mean eosinophil number measured in a total of 40 hpf from 2 x 4 biopsy specimens each, taken from the proximal and distal esophagus; all levels were surveyed and the eos in the most densely infiltrated area were counted in 5 hpf -area of microscopic field = 0.3072 mm ²	Continuous Dichotomous	Mean peak eos/hpf (SD) at end of trial (used for histological continuous analysis): Budesonide: 29.9 (30.6)/14 Placebo: 51.1 (31.1)/14 ≤ 5 eos: 5/14 vs 0/14 5 to 20 eos: 2/14 vs 4/14 > 20 eos: 7/14 vs 10/14 ≤ 20 (used for histological dichotomous analysis): Budesonide: 7/14 Placebo: 4/14
Straumann 2013	Total of 40 hpf from 2 x 4 biopsies taken from the proximal and distal esophagus In all 8 esophageal biopsies, eosinophils in the most densely infiltrated area were counted in 5 consecutive hpf. Area of microscopic field = 0.3072 mm ² .	Continuous (compared post-treatment means); no pre-specified response threshold	Mean peak eos/hpf at end of trial (SD) (used for histological continuous analysis): OC000459: 73.26 (58.29)/14 Placebo: 99.47 (69.95)/12
Straumann 2020	2 biopsies of each esophageal third Counted eosinophils in the most densely infiltrated area (HPF 0.245 mm ²)	Dichotomous Histologic relapse (i.e. peak of ≥ 48 eos/mm ² hpf (corresponding to ≥ 15 eos/hpf in prior studies))	Histologic relapse at end of trial: Budesonide 0.5 mg twice-daily: 9/68 Budesonide 1.0 mg twice-daily: 7/68 Placebo: 61/68

Medical treatment of eosinophilic esophagitis (Review)

Table 4. Primary outcome - histological improvement (Continued)

	Continuous	Mean peak < 15 eos/hpf at end of trial (used for histological dichotomous analysis):
	Comparison of change in peak eos/hpf from baseline to end of trial	Budesonide: 120/136 Placebo: 7/68
		Change in mean peak eos/mm ² from baseline to end of treatment, mean (SD) (used for histological continuous analysis):
		Budesonide 0.5 mg twice-daily: 38 (112.6)/66 Budesonide 1.0 mg twice-daily: 21 (64)/65 Budesonide: 29.56 (91.11)/131 Placebo: 262 (216.3)/65
Tytor 2021	Not reported	Not reported

AAF: amino acid-based formula; BET: budesonide effervescent tablet; BOS: budesonide oral suspension; BOV: budesonide, oral viscous; CG: control group; ED: elimination diet; eos: eosinophils; FFED: four food elimination diet; hpf: high-power field; IG: intervention group; IQR: interquartile range; LS: least squares; NEB: nebulized/swallowed budesonide solution; OBS: oral budesonide solution; OVB: viscous/swallowed budesonide solution; PPI: proton pump inhibitor; SD: standard deviation

Table 5. Primary outcome - withdrawals due to adverse events

Study ID	Withdrawals due to adverse events	Reasons for withdrawals
Alexander 2012	Fluticasone: 2/21 (9.5%) Placebo: 6/21 (28.6%)	The causes of dropout were travel in 2 patients, scheduling in 3 patients, family issues in 2 patients, and change of mind on study involvement 1 week after initiation in 1 patient
Alexander 2017	Montelukast: 2/20 Placebo: 1/21	The causes of the withdrawals were personal and travel reasons
Assa'ad 2011	Mepolizumab 0.55 mg/kg: 3/19 Mepolizumab 2.5 mg/kg: 1/20 Mepolizumab 10 mg/kg: 2/20	Mepolizumab 0.55 mg/kg: 1 withdrew consent due to an adverse event; 2 withdrew consent; 1 was lost to follow-up Mepolizumab 2.5 mg/kg: 1 withdrew consent due to lack of efficacy Mepolizumab 10 mg/kg: 1 withdrew for other reasons
Bhardwaj 2017	Beclomethasone: 0/9 Placebo: 0/9	No withdrawals reported
Butz 2014	Fluticasone: 5/28 (17.9%) Placebo: 1/14 (7.1%)	Fluticasone: 2 because of prohibited medications, 2 because of loss to follow-up evaluation, and 1 because of an adverse event (AE) (absence seizure that was deemed unlikely to be related to FP) Placebo: 1 participant from the placebo group was lost to follow-up
Clayton 2014	Omalizumab: 0/16 Placebo: 0/14	No withdrawals reported
Dellon 2012	Budesonide, nebulized: 0/13 Budesonide, oral viscous: 0/12	No withdrawals reported

Table 5. Primary outcome - withdrawals due to adverse events (Continued)

Dellon 2017	Budesonide: 2/51 (3.9%) Placebo: 3/42 (7.1%)	Budesonide: 1 because of an adverse event, 1 owing to lack of compliance Placebo: 1 because of lack of efficacy, 1 because of lack of compliance, and 1 owing to pregnancy, 1 additional patient in the placebo arm did not have an evaluable post-treatment biopsy
Dellon 2019	Budesonide: 9/65 (0%) Fluticasone: 9/64 (0%)	Fluticasone: 1 had an adverse event of a food bolus impaction necessitating an emergency department visit and study withdrawal (from: NCT02019758, SAE was in the fluticasone arm) 1 participant in each group did not receive the intervention after randomization, and 8 in each group were lost to follow-up and did not undergo the week-8 endoscopy
Dellon 2021b	Supplementary Figure 1 used for analysis: Budesonide: 3/25 Placebo: 4/23	Budesonide: 2 withdrawal by patient, 1 adverse event Placebo: 4 withdrawal by patient
Dellon 2022	Dupilumab: 0/42 Placebo: 0/39	No withdrawals reported
Dellon 2022a	Supplementary Figure 2: APT-1011 3 mg twice-daily (n = 1/20) APT-1011 3 mg at bedtime (n = 2/21) APT-1011 1.5 mg twice-daily (n = 2/22) APT-1011 1.5 mg at bedtime (n = 4/21) Placebo (n = 5/20) Total: APT-1011: 9/84 Placebo: 5/20	APT-1011 3 mg twice-daily; 1 adverse event APT-1011 3 mg at bedtime; 1 withdrawal of consent, 1 for other reasons APT-1011 1.5 mg twice-daily; 1 withdrawal of consent, 1 adverse event APT-1011 1.5 mg at bedtime; 3 withdrawal of consent, 1 for other reasons Placebo: 2 withdrawal of consent, 2 adverse events, 1 for other reasons
Dellon 2022b	Lirentelimab 3 mg/kg: 11/91 Lirentelimab 1 mg/kg: 6/93 Lirentelimab: 17/184 Placebo: 4/92	Not reported
De Rooij 2022	Four-food elimination: 1/20 Four-food elimination + amino acid formula: 1/21	Four-food elimination: 1 non-compliance Four-food elimination + amino acid formula group: 1 non-compliant to AAF for personal reasons
Dohil 2010	Budesonide: 7/21 (33%) Placebo: 2/11 (18%)	Budesonide: 2 not wanting to take Splenda, 2 acute asthma requiring systemic corticosteroids, 1 non-compliance with therapy, 1 transient rash attributed to lansoprazole, 1 lost to follow-up

Table 5. Primary outcome - withdrawals due to adverse events (Continued)

		Placebo: reasons not given
Gupta 2015	From NCT00762073: Budesonide, low-dose: 4/21 (19.0%) Budesonide, medium-dose: 2/19 (10.5%) Budesonide, high-dose: 3/20 (15%) Budesonide: 9/60 (15%) Placebo: 3/21 (14.3%)	Budesonide, low-dose: 1 lack of efficacy, 1 non-compliance, 2 withdrawal by patient Budesonide, medium-dose: 1 adverse event, 1 withdrawal by patient Budesonide, high-dose: 1 adverse event, 1 non-compliance, 1 withdrawal by patient Placebo: 1 lack of efficacy, 1 adverse event, 1 withdrawal by patient
Heine 2019	Four food elimination diet + PPI: 5/32 PPI: 1/32	Four-food elimination diet + PPI: 5 for noncompliance PPI: 1 for non-compliance
Hirano 2019	From Figure 2: RPC4046 180: 4/31 RPC4046 360: 4/30 RPC4046: 8/61 Placebo: 2/34	RPC4046 180: 3 adverse events, 1 withdrew consent RPC4046 360: 1 adverse event, 1 withdrew consent, 2 for other reasons Placebo: 2 withdrew consent
Hirano 2020	Dupilumab: 1/23 Placebo: 4/24	Dupilumab: 1 adverse event Placebo: 1 protocol non-compliance, 3 cited as "Other"
Hirano 2020f	APT-1011 1 mg: 0/8 (0%) APT-1011 3 mg: 0/8 (0%) APT-1011: 0/16 Placebo: 2/8 (25%)	Placebo: 1 protocol violation, 1 patient needed excluded medication
Hirano 2021	Budesonide: 11/215 (5.1%) Placebo: 11/107 (10.3%)	Budesonide: 8 withdrawal by patient, 1 adverse event, 1 non-compliance, 1 physician decision Placebo: 8 withdrawal by patient, 3 adverse events
Kliewer 2019	Four-food elimination diet: 8/25 (36.0) One-food elimination diet: 4/38 (15.8)	Four-food elimination diet: 3 participants withdrew because the diet was too difficult to adhere to, 2 withdrew before initiating the diet, 1 withdrew for insurance reasons, 1 was withdrawn due to an unrelated adverse event requiring a prohibited medication, and 1 was lost to follow-up One-food elimination diet: 2 withdrew because the diet was too difficult to adhere to, 2 were lost to follow-up
Kliewer 2021	From NCT02778867 at end of phase 1: Six-food elimination: 3/62 One-food elimination: 2/67	Six-food elimination: 3 withdrawal by patient; 2 unwilling to continue, 1 non-compliant One-food elimination: 2 withdrawal by patient; 1 insurance reasons, 1 unknown
Konikoff 2006	Fluticasone: 1/21 (0%) Placebo: 4/15 (20%)	Fluticasone: 1 did not meet the inclusion criteria for the diagnosis of EoE

Table 5. Primary outcome - withdrawals due to adverse events (Continued)

		Placebo: 3 for increased symptoms, 1 for non-compliance
Lieberman 2018	Cromolyn sodium: 0.0% (0/9) Placebo: 14.3% (1/7)	Placebo: 1 increased GI symptoms within 1 week of beginning study
Lucendo 2019	Budesonide: 8/59 (13.6) Placebo: 3/29 (10.3)	Budesonide: multiple reasons possible including, 3 protocol violations, 4 in/exclusion criteria violated, 1 prohibited concomitant medicine, 1 non-compliant Placebo: multiple reasons possible including, 1 protocol violations, 1 in/exclusion criteria violated, 1 prohibited concomitant medicine, 3 non-compliant
Miehlke 2016	Budesonide effervescent tablet 2 x 1 mg: 0/19 (0%) Budesonide effervescent tablet 2 x 2 mg: 2/19 (10.5%) Budesonide oral viscous suspension 2 x 2 mg: 2/19 (10.5%) Budesonide: 4/57 Placebo: 2/19 (10.5%)	Budesonide effervescent tablet 2 x 2 mg: 1 non-compliant, 1 no post-therapy biopsy Budesonide oral viscous suspension 2 x 2 mg: 1 insufficient baseline disease, 1 no post-therapy biopsy Placebo: 2 insufficient baseline disease
Moawad 2013	Fluticasone: 2/21 (14.3%) Esomeprazole: 0/21 (0%)	Fluticasone: 1 with worsening of migraine headaches, which he attributed to FP, 1 with bothersome GERD-related symptoms and discontinued the steroid, and began treatment with a PPI
Oliva 2018	Withdrawals due to adverse events were not reported	Not reported
Peterson 2010	Fluticasone: 1/15 Esomeprazole: 3/15	Reasons for dropout were unwillingness to perform the second EGD in 3 patients. One patient was withdrawn by the IRB for an interpretation of "inadequate pathology". All dropouts occurred prior to completion of the second dysphagia questionnaire. Two patients who completed the second EGD did not complete the second dysphagia questionnaire.
Rothenberg 2015	QAX576: 2/17 Placebo: 0/8	QAX576: 1 because of a positive drug screen, 1 because of a non-drug-related serious adverse event
Rothenberg 2022	None reported	None reported
Schaefer 2008	Prednisone: 8/40 Fluticasone: 4/40	Prednisone: non-compliance with medication, n = 4; patient/family decision, n = 2; lost to follow-up evaluation, n = 1; adverse effect, n = 1 (increased appetite and abdominal pain) Fluticasone: non-compliance with medication, n = 2; patient/family decision, n = 2
Spiegel 2012	Reslizumab 1 mg/ml: 8/56 Reslizumab 2 mg/ml: 11/57 Reslizumab 3 mg/ml: 7/57 Reslizumab: 26/170 Placebo: 6/57	Reslizumab 1 mg/mL: 1 adverse event, 5 lack of efficacy, 1 lost to follow-up, 1 other Reslizumab 2 mg/mL: 9 lost to follow-up, 2 for other reasons Reslizumab 3 mg/mL: 6 lack of efficacy, 1 other Placebo: 4 lack of efficacy, 1 lost to follow-up, 1 other
Spiegel 2020	Viaskin milk: 1/15 (0%) Placebo: 0/5 (0%)	Viaskin milk: withdrawal of consent

Table 5. Primary outcome - withdrawals due to adverse events (Continued)

Straumann 2010a	Mepolizumab: 0/5 Placebo: 0/6	No withdrawals reported
Straumann 2010b	Budesonide: 0/18 Placebo: 0/18	No withdrawals reported
Straumann 2011	Budesonide: 5/14 Placebo: 9/14	All withdrawals were due to clinical relapse
Straumann 2013	OC000456: 0/14 Placebo: 0/12	No withdrawals reported
Straumann 2020	Budesonide 0.5: 9/68 Budesonide 1.0: 9/68 Budesonide: 18/136 Placebo: 45/68	Budesonide 0.5: 7 due to lack of efficacy, 2 due to lack of co-operation Budesonide 1.0: 5 due to lack of efficacy, 2 due to adverse event (retinitis and allergic dermatitis), 2 due to lack of co-operation Placebo: 42 due to lack of efficacy, 3 due to lack of patient's co-operation
Tytor 2021	Mometasone: 1/17 (5.9%) Placebo: 2/19 (10.5%)	Mometasone: 1 lost to follow-up Placebo: 2 lost to follow-up

AAF: amino acid-based formula; BOS: budesonide oral suspension; BOT: budesonide orodispersible tablet; CG: control group; EGD: esophagogastroduodenoscopy; EoE: eosinophilic esophagitis; FFED: four food elimination diet; FP: fluticasone propionate; GERD: gastroesophageal reflux disease; GI: gastrointestinal; IG: intervention group; IRB: institutional review board; N/A: not applicable; NEB: nebulized/swallowed budesonide solution; OVB: viscous/swallowed budesonide solution; SAE: serious adverse event; TEAE: treatment-emergent adverse event

Table 6. Secondary outcomes

Study ID	Patients with serious adverse events	Patients with total adverse events	Quality of life
Alexander 2012	From NCT00275561: Fluticasone: 0/21 Placebo: 0/21	From NCT00275561: Fluticasone: 7/21 (36.8%) (sore throat, 2/21 (10.53%); esophageal candidiasis, 5/21 (26.32%); hoarseness, 0/21 (0.00%); 24-hour urine cortisol, 23.2 ± 2.5 g/24 hours) Placebo: 6/21 (40%) (sore throat, 3/21 (20.00%); esophageal candidiasis, 0/21 (0.00%); hoarseness, 3/21 (20.00%); 24-hour urine cortisol, 15.5 ± 2.5 µg/24 hours)	Not reported
Alexander 2017	Montelukast: 0/20 Placebo: 0/21	Montelukast: 0/20 Placebo: 0/21	Not reported
Assa'ad 2011	From NCT00358449: Mepolizumab 0.55 mg/kg: 0/19 Mepolizumab 2.5 mg/kg: 1/20; 1 x foreign body trauma	From NCT00358449: Mepolizumab 0.55 mg/kg: 18/19 Mepolizumab 2.5 mg/kg: 14/20 Mepolizumab 10 mg/kg: 18/20 Most common: vomiting (16.9%), diarrhea (13.6%), and upper abdominal pain (10.2%)	Not reported

Medical treatment of eosinophilic esophagitis (Review)

Table 6. Secondary outcomes (Continued)

	Mepolizumab 10 mg/kg: 2/20; 1 x chest discomfort, 1 x esophageal injury		
Bhardwaj 2017	n = 0 "No significant adverse effects were reported with the study drug"	Not reported	Not reported
Butz 2014	From NCT00426283: Fluticasone: 0/28 Placebo: 0/14 n = 1 SAE: absence seizure that was deemed unlikely to be related to fluticasone group	From NCT00426283: Fluticasone: 19/28 Placebo: 9/14 Fluticasone (n = 28): Eye disorders 0/14 (0.0%); gastrointestinal disorders 7/14 (25.0%); chest pain 1/14 (3.6%); immune system disorders 2/14 (7.1%); infections and infestations 3/14 (10.7%); injury, poisoning, and procedural complications: scrapes and cuts 1/14 (3.6%); investigations: abnormal laboratory values 5/14 (17.9%); nervous system disorders 4/14 (14.3%); respiratory, thoracic, and mediastinal disorders 3/14 (10.7%); skin and subcutaneous tissue disorders 0/14 (0.0%) Placebo (n = 14): Eye disorders 2/14; gastrointestinal disorders 2/14 (14.3%); chest pain 1/14 (7.1%); immune system disorders 0/14 (0.0%); infections and infestations 3/14 (21.4%); injury, poisoning, and procedural complications: scrapes and cuts 0/14 (0.0%); investigations: abnormal laboratory values 1/14 (7.1%); nervous system disorders 0 (0.0%); respiratory, thoracic, and mediastinal disorders 3/14 (21.4%); skin and subcutaneous tissue disorders 1/14 (7.1%)	Not reported
Clayton 2014	From NCT00123630: Omalizumab: 0/16 Placebo: 0/14	From NCT00123630: Omalizumab: 0/16 Placebo: 0/14	Not reported
Dellon 2012	From NCT00961233: Budesonide, nebulized: 0/13 Budesonide, oral viscous: 0/12	From NCT00961233: Budesonide, nebulized: 0/13 Budesonide, oral viscous: 0/12 Used for analysis. Budesonide, nebulized: 1/11 - candidal esophagitis, 1/11 epistaxis Budesonide, oral viscous: 2/11 candidal esophagitis	Not reported
Dellon 2017	From NCT01642212: Budesonide: 1/51 Placebo: 0/42 n = 1; budesonide group	From NCT01642212: Budesonide: 20/51 Placebo: 21/42 Reported as: budesonide; placebo All TEAEs: 24/51; 21/42	Not reported

Table 6. Secondary outcomes (Continued)

Food poisoning	TEAEs related to study drug: 5/51; 4/42 Severe TEAE: 1/51; 0/42 Serious adverse events: 1/51; 0/42 TEAEs leading to withdrawal from study: 1/51; 0/42 TEAEs related to study drug and leading to withdrawal from study: 1/51; 0/42 Infections and infestations: 13/51; 7/42 Nasopharyngitis: 3/51; 4/42 Upper respiratory tract infection: 3/51; 2/42 Sinusitis: 2/51; 1/42 <i>Clostridium difficile</i> infection: 1/51; 0/42 Oral candidiasis: 1/51; 0/42 Esophageal candidiasis: 1/51; 0/42 Gastrointestinal disorders: 3/51; 9/42 Diarrhea: 0/51; 1/42 Food poisoning: 2/51; 0/42 Vomiting: 1/51; 1/42 Abdominal pain/discomfort: 0/51; 3/42 Respiratory disorders: 6/51; 3/42 Oropharyngeal pain: 2/51; 2/42 Cough: 1/51; 0/42 Dyspnea: 1/51; 0/42 Allergic rhinitis: 1/51; 0/42 Skin disorders: 3/51; 3/42 Acne: 1/51; 0/42 Contact dermatitis: 1/51; 0/42 Eczema: 0/51; 1/42 General: 3/51; 2/42 Fever: 1/51; 1/42 Fatigue: 1/51; 0/42
----------------	---

Dellon 2019	From NCT02019758: Oral viscous budesonide: 0/65 Active fluticasone: 1/64; food impaction	From NCT02019758: Oral viscous budesonide: 10/65 Active fluticasone: 15/64 Adverse event: Esophageal candidiasis: budesonide 8 (12%); fluticasone 10 (16%)	Not reported
-------------	--	--	--------------

Medical treatment of eosinophilic esophagitis (Review)

Table 6. Secondary outcomes (Continued)

		Oral candidiasis: budesonide 2 (3%); fluticasone 1 (2%)	
		Food impaction: budesonide 0 (0%); fluticasone 1 (2%)	
		Sore throat: budesonide 0 (0%); fluticasone 2 (3%)	
		Chest pain: budesonide 0 (0%); fluticasone 1 (2%)	
		Pneumonia: budesonide 0 (0%); fluticasone 1 (2%)	
Dellon 2021b	Any severe TEAE from NCT02736409: Budesonide: 0/25 (0) Placebo: 1/23 (4.3); back pain	Any TEAE From NCT02736409: Budesonide: 21/25 (84%) Placebo: 14/23 (60.9%) TEAEs experienced by 2.5% of the total Upper respiratory tract infection Budesonide: 12 (9.2%) Placebo: 2 (3.1%) Nasopharyngitis Budesonide: 7 (5.3%) Placebo: 3 (4.6%) Sinusitis Budesonide: 5 (3.8%) Placebo: 3 (4.6%) Esophageal candidiasis Budesonide: 4 (3.1%) Placebo: 5 (7.7%) Influenza Budesonide: 5 (3.8%) Placebo: 1 (1.5%) Nausea Budesonide: 9 (6.9%) Placebo: 4 (6.2%) Vomiting Budesonide: 13 (9.9%) Placebo: 1 (1.5%) Diarrhea Budesonide: 5 (3.8%) Placebo: 4 (6.2%) Dysphagia Budesonide: 3 (2.3%) Placebo: 1 (1.5%) Gastritis Budesonide: 6 (4.6%) Placebo: 0 (0.0%) Blood cortisol decreased	Not reported

Table 6. Secondary outcomes (Continued)

		Budesonide: 4 (3.1%)	
		Placebo: 4 (6.2%)	
		Cough	
		Budesonide: 5 (3.8%)	
		Placebo: 2 (3.1%)	
		Fatigue	
		Budesonide: 4 (3.1%)	
		Placebo: 1 (1.5%)	
		Mood swings	
		Budesonide: 2 (1.5%)	
		Placebo 3 (4.6%)	
Dellon 2022	None reported	Not reported as individuals/group, cannot use	Not reported
		Injection-site reactions:	
		Dupilumab: 7/42(16.7%)	
		Placebo: 14/39 (0.3%)	
		Nasopharyngitis:	
		Dupilumab: 5/42 (11.9%)	
		Placebo: 4/39 (10.3%)	
Dellon 2022a	APT-1011 3 mg at bedtime (n = 1/22) (5%) APT-1011: 1/85 Placebo: 0/21	Total adverse events = 63/85 (74%) APT-1011 3 mg twice-daily (17/20) APT-1011 3 mg at bedtime (16/22) APT-1011 1.5 mg twice-daily (17/22) APT-1011 1.5 mg at bedtime (13/21) Placebo (13/21) APT-1011 = 63/85 Placebo: 13/21	EoE Adult Quality of Life Questionnaire (EoE-QoLA) but no data reported
Dellon 2022b	Total n = 3 Lirentelimab high-dose: 2 Lirentelimab low-dose: 0 Placebo: 1 Lirentelimab: 2/184 Placebo: 1/92 Type not reported	≥ 1 TEAE • Lirentelimab high-dose: 61/91 • Lirentelimab low-dose: 65/93 • Placebo: 53/92 Lirentelimab: 126/184 Placebo: 53/92 Infusion reaction • Lirentelimab high-dose: 35/91 • Lirentelimab low-dose: 24/91 • Placebo: 11/92	Not reported

Table 6. Secondary outcomes (Continued)

		Headache	
		<ul style="list-style-type: none"> • Lirentelimab high-dose: 6/91 • Lirentelimab low-dose: 8/93 • Placebo: 6/92 	
De Rooij 2022	Four-food elimination diet + amino acid formula: 0/20 Four-food elimination diet: 0/21	Four-food elimination diet + amino acid formula: 1/20; emergency room visit due to severe abdominal pain after eating a kiwi Four-food elimination diet: 0/21	EoEQoL score Change in total EoE-QoL score from baseline to week 6, median (IQR) (cannot use) Four-food elimination diet + amino acid formula: 0.1 (0.04 to 0.56) Four-food elimination diet: 0 (-0.08 to 0.4)
Dohil 2010	From NCT00638456: Budesonide + PPI: 0/21 PPI: 0/11	From NCT00638456: Budesonide + PPI: 3/21 PPI: 5/11 Budesonide + PPI: 3 1: emesis 1: oral Candida 1: transient headache PPI: 5 1: eczema worse 1: chest infection 1: mild abdominal pain 1: transient headache 1: transient diarrhea	Not reported
Gupta 2015	From NCT00762073: Budesonide: 1/60; diet refusal Placebo: 0/21 G1 - 1 G2 - 0 G3 - 1 G4 - 1	From NCT00762073: Budesonide: 13 + 16 + 17 = 46/60 Placebo: 10/21 Budesonide most frequent adverse events: rash, 10/60 (17%); diarrhea, 10/60 (17%); pyrexia, 10/60 (17%); cough, 9/60 (15%); sinusitis, 9/60 (15%); nasopharyngitis, 8/60 (13%); oropharyngeal pain, 8/60 (13%); headache, 7/60 (12%) Placebo most frequent adverse events: pyrexia, 3/21 (14%); headache, 2/21 (10%); vomiting, 2/21 (10%); asthma, 2/21 (10%)	Not reported
Heine 2019	Not reported	Not reported	Not reported

Table 6. Secondary outcomes (Continued)

Hirano 2019	SAE:	Total AE:	Not reported
	RPC4046 180 mg: 0/32 (0%)	RPC4046 180 mg: 20/32 (63%)	
	RPC4046 360 mg: 1/34 (2.9%); appen- dicitis	RPC4046 360 mg: 29/34 (85%)	
	RPC4046: 1/66	RPC4046: 49/66	
	Placebo: 2/34 (5.9%); 1 x umbilical hernia, 1 x appen- dicitis	Placebo: 22/34 (65%)	
		Adverse events in placebo, RPC4046 180 mg, RPC4046 360 mg:	
		Headache: 5 (14.7%), 5 (16.1%), 7 (20.6%)	
		Upper respiratory tract infection: 3 (8.8%), 5 (16.1%), 5 (14.7%)	
		Arthralgia: 0, 4 (12.9%), 2 (5.9%)	
		Nasopharyngitis: 0, 3 (9.7%), 3 (8.8%)	
		Diarrhea: 2 (5.9%), 3 (9.7%), 2 (5.9%)	
		Nausea: 4 (11.8%), 2 (6.5%), 3 (8.8%)	
		Abdominal pain: 0, 2 (6.5%), 2 (5.9%)	
		Dizziness: 2 (5.9%), 3 (9.7%), 1 (2.9%)	
		Oropharyngeal pain: 0, 1 (3.2%), 3 (8.8%)	
		Sinusitis: 0, 3 (9.7%), 1 (2.9%)	
		Vomiting: 2 (5.9%), 1 (3.2%), 3 (8.8%)	
		Contact dermatitis: 0, 1 (3.2%), 2 (5.9%)	
		Fatigue: 1 (2.9%), 2 (6.5%), 1 (2.9%)	
		Injection site erythema: 2 (5.9%), 0, 3 (8.8%)	
		Urticaria: 0, 2 (6.5%), 1 (2.9%)	
		Myalgia: 0, 1 (3.2%), 2 (5.9%)	
		Contusion: 1 (2.9%), 2 (6.5%), 0	
		Cough: 1 (2.9%), 2 (6.5%), 0	
		Gastroenteritis: 1 (2.9%), 2 (6.5%), 0	
		Hypersensitivity: 0, 0, 2 (5.9%)	
		Injection site hematoma: 0, 0, 2 (5.9%)	
		Injection site pruritus: 1 (2.9%), 0, 2 (5.9%)	
		Ligament sprain: 1 (2.9%), 0, 2 (5.9%)	
Hirano 2020	From NCT02379052: Dupilumab: 3/23; 1 x-food allergy, 1 x blood creatine phosphokinase in- creased, 1 x abor- tion spontaneous Placebo: 0/24	From NCT02379052: Dupilumab: 18/23 Placebo: 14/24 Dupilumab, major adverse events: injection site erythema, 8/23 (34.78%); injection site rash, 3/23 (13.04%); injection site urticaria, 2/23 (8.70%); injection site inflammation, 3/23 (13.04%); injection site pain, 2/23 (8.70%); nasopharyngitis, 5/23 (21.74%) Placebo, major adverse events: injection site erythema, 2/24 (8.33%); injection site pain, 2/24 (8.33%); upper respiratory tract infection, 3/24 (12.50%); abdominal pain, 2/24 (8.33%); nausea, 3/24 (12.50%); dizziness, 2/24 (8.33%)	From NCT02379052: LS mean change from baseline (SE) Dupilumab: 0.80 (0.137) 23 SD = (0.66) Placebo: 0.47 (0.141) 24 SD = (0.69) QoL was assessed using Eosinophilic Esophagitis Quali- ty of Life (EoE-QoL- A) score, version 3.0 from baseline to week 12 EoE-QoL-A total score week 12, n (%): Dupilumab: 23/0 Placebo: 21/3

Table 6. Secondary outcomes (Continued)

			LS mean change from baseline (SE): Dupilumab: 0.8 (0.1) Placebo: 0.5 (0.1) Difference vs placebo (95% CI): 0.3 (-0.1 to 0.7) Unclear if 30 items or 24 items
Hirano 2020f	APT-1011: 0/16 Placebo: 0/8	APT-1011: 12/16 Placebo: 6/8 Major adverse events, placebo, APT-1011 1.5 mg, APT-1011 3.0 mg: blood cortisol decreased: 2 (25%), 3 (37.5%), 1 (12.5%); diarrhea: 0, 0, 2 (25%); nasopharyngitis: 0, 1 (12.5%), 1 (12.5%) A total of 41 TEAEs were reported by 18 participants: 12 participants receiving APT-1011 reported 26 TEAEs and 6 participants receiving placebo reported 15 TEAEs	Not reported
Hirano 2021	From NCT02605837: Budesonide: 2/215 Placebo: 1/107	From NCT02605837: Budesonide: 63/215 Placebo: 28/107 Any non-serious TEAE Budesonide: 130 (61.0%) Placebo: 64 (61.0%) Total: 194 (61.0%) Any mild TEAE Budesonide: 69 (32.4%) Placebo: 38 (36.2%) Total: 107 (33.6%) Any moderate TEAE Budesonide: 56 (26.3%) Placebo: 24 (22.9%) Total 80 (25.2%) Any severe TEAE Budesonide: 5 (2.3%) Placebo: 2 (1.9%) Total 7 (2.2%) Any serious TEAE 2 Budesonide: (0.9%)	Not reported

Table 6. Secondary outcomes (Continued)

Placebo: 1 (1.0%)
Total 3 (0.9%)
Any life-threatening TEAE
Budesonide: 0 (0.0%)
Placebo: 0 (0.0%)
Total 0 (0.0%)
TEAE related to study treatment
Budesonide: 45 (21.1%)
Placebo: 23 (21.9%)
Total 68 (21.4%)
TEAE related to EoE
Budesonide: 11 (5.2%)
Placebo: 6 (5.7%)
Total 17 (5.3%)
TEAE leading to dose discontinuation
Budesonide: 3 (1.4%)
Placebo: 5 (4.8%)
Total 8 (2.5%)
TEAE leading to study discontinuation
Budesonide: 1 (0.5%)
Placebo: 3 (2.9%)
Total 4 (1.3%)
Infections and infestations
Nasopharyngitis
Budesonide: 11 (5.2%)
Placebo: 4 (3.8%)
Total 15 (4.7%)
Sinusitis
Budesonide: 9 (4.2%)
Placebo: 3 (2.9%)
Total 12 (3.8%)
Esophageal candidiasis
Budesonide: 8 (3.8%)
Placebo: 2 (1.9%)

Table 6. Secondary outcomes *(Continued)*

Total 10 (3.1%)
Oral candidiasis
Budesonide: 8 (3.8%)
Placebo: 0 (0.0%)
Total 8 (2.5%)
Gastrointestinal disorders
Nausea
Budesonide: 6 (2.8%)
Placebo: 3 (2.9%)
Total 9 (2.8%)
Vomiting
Budesonide: 4 (1.9%)
Placebo: 4 (3.8%)
Total 8 (2.5%)
Investigations ACTH stimulation test abnormal
Budesonide: 6 (2.8%)
Placebo: 3 (2.9%)
Total 9 (2.8%)
Respiratory, thoracic, and mediastinal disorders
Cough
Budesonide: 6 (2.8%)
Placebo: 3 (2.9%)
Total 9 (2.8%)
Skin and subcutaneous tissue disorders
Acne
Budesonide: 5 (2.3%)
Placebo: 3 (2.9%)
Total 8 (2.5%)
Nervous system disorders
Headache
Budesonide: 7 (3.3%)
Placebo: 1 (1.0%)
Total 8 (2.5%)

Table 6. Secondary outcomes (Continued)

Kliwer 2019	From NCT02610816 One-food elimination diet: 1/38; abdominal pain Four-food elimination diet: 1/25; abdominal pain	From NCT02610816 One-food elimination diet: 5/38 Four-food elimination diet: 8/25 One-food elimination diet: abdominal pain, 2/38 (5.26%); vomiting, 1/38 (2.63%); cough, 1/38 (2.63%); nasal congestion, 1/38 (2.63%) Four-food elimination diet: abdominal pain, 2/25 (8.00%); vomiting, 2/25 (8.00%); cough, 2/25 (8.00%); urticaria, 2/25 (8.00%)	Change from baseline in Pediatric Quality of Life Inventory Version 3.0 EoE Module (Ped-sQL 3.0 EoE) at 12 weeks One-food elimination diet: 9.7 (11.3)/31 Four-food elimination diet: 9.8 (14.1)/16
Kliwer 2021	From NCT02778867: One-food elimination diet: 0/67 Six-food elimination diet: 0/62 One-food elimination: at 6 weeks, 0/67 Six-food elimination: at 6 weeks, 0/62	From NCT02778867: One-food elimination diet: 1/67; Diarrhea Six-food elimination diet: 2/62; Diarrhea	EoE-QoL-A (24 items), change from baseline: One-food elimination: at 6 weeks -0.9 (10.2)/67 Six-food elimination: at 6 weeks -0.33 (11.8)/62
Konikoff 2006	Fluticasone: 0/21 Placebo: 0/15	Fluticasone: 1/21; esophageal candidiasis Placebo: 0/15	Not reported
Lieberman 2018	Cromolyn: 0/9 Placebo: 0/7	Not reported as individuals/group, cannot use Adverse events reported: Nausea: cromolyn group 55.6% (5/9); placebo group 14.3% (1/7) Abdominal pain: cromolyn group 44.4% (4/9); placebo group 28.6% (2/7) Headache: cromolyn group 44.4% (4/9); placebo group 14.3% (1/7) Vomiting: cromolyn group 22.2% (2/9); placebo group 14.3% (1/7) Upper respiratory tract infection: cromolyn group 22.2% (2/9); placebo group 14.3% (1/7) Fatigue: cromolyn group 22.2% (2/9); placebo group 0.0% (0/9) Sore throat: cromolyn group 22.2% (2/9); placebo group 0.0% (0/9) Lip swelling: cromolyn group 11.1% (1/9); placebo group 0.0% (0/9) Sinus infection: cromolyn group 11.1% (1/9); placebo group 14.3% (1/7)	Not reported

Table 6. Secondary outcomes (Continued)

		Dysphagia: cromolyn group 11.1% (1/9); placebo group 0.0% (0/9)	
		Diarrhea: cromolyn group 11.1% (1/9); placebo group 0.0% (0/9)	
		Eye pain: cromolyn group 11.1% (1/9); placebo group 0.0% (0/9)	
		Mood change: cromolyn group 11.1% (1/9); placebo group 0.0% (0/9)	
		Hypernatremia: cromolyn group 11.1% (1/9); placebo group 0.0% (0/9)	
Lucendo 2019	Budesonide: 6/59 (10.1%) Placebo: 13/29 (44.8%) Severe TEAE esophageal food impaction TEAE related to study drug	Any TEAE: Budesonide: 37/59 (62.7%) Placebo: 12/29 (41.1%) TEAEs by occurring in 2 patients in any treatment group, reported as budesonide; placebo: Gastrointestinal disorders: 10/59 (16.9%); 3/29 (10.3%) Gastroesophageal reflux disease: 3/59 (5.1%); 0/29 (0%) Nausea: 2/29 (3.4%); 0/29 (0%) Infections and infestations: 21/59 (35.6%); 6/29 (20.7%) Suspected local fungal infection, thereof: 14/59 (23.7%); 0/29 (0%) Histologically confirmed: 10/59 (16.9%); 0/29 (0%) Histologically confirmed with suspected endoscopic signs: 8/59 (13.6%); 0/29 (0%) Histologically confirmed with suspected endoscopic signs and clinical symptoms: 3/59 (5.1%); 0/29 (0%) Nasopharyngitis: 2/59 (3.4%); 1/29 (3.4%) Pharyngitis: 1/59 (1.7%); 2/29 (6.9%) Investigations: 5/59 (8.5%); 0/29 (0%) Blood cortisol decreased: 3/59 (5.1%); 0/29 (0%) Nervous system disorders: 5/59 (8.5%); 1/29 (3.4%) Headache: 4/59 (6.8%); 1/29 (3.4%) Respiratory, thoracic and mediastinal disorders: 2/59 (3.4%); 2/29 (6.9%) Asthma: 0/59 (0%); 2/29 (6.9%) Vascular disorders: 3/59 (5.1%); 0/29 (0%) Hypertension: 2/59 (3.4%); 0/29 (0%)	EoE-QoL-A 30-items (weighted average) baseline, mean (SD) Budesonide: 2.3 (0.8) Placebo: 2.3 (0.8) EoT, mean (SD): Budesonide: 2.8 (0.9)/59 Placebo: 2.6 (0.7)/29 Change from baseline to EoT, mean (95% CI) Budesonide: 0.5 (0.32 to 0.62) Placebo: 0.2 (0.06 to 0.42) BOT–placebo, mean difference (95% CI) 0.23 (–0.010 to 0.472)
Miehlke 2016	Budesonide: 0/57 Placebo: 0/19	Budesonide: 4 + 5 + 6 = 15/57 Placebo: 0/19 BET 2 x 1 mg: 5/19: 3 esophageal candidiasis, 1 increased WBC count, 1 pruritus BET 2 x 2mg: 6/19: 1 nausea, 1 blistering oral mucosa, 3 esophageal candidiasis, 1 blood cortisol decreased BOV 2 x 2 mg: 6/19: 1 bowel movement irregularity, 1 lip edema, 3 esophageal candidiasis, 1 pruritus	Not reported
Moawad 2013	From NCT00895817: Esomeprazole: 0/21 Swallowed fluticasone: 0/21	From NCT00895817: Esomeprazole: 0/21 Swallowed fluticasone: 0/21 Used for analysis:	Not reported

Table 6. Secondary outcomes (Continued)

		FP arm n = 3/21 n = 1; 1 patient had worsening of migraine headaches, which he attributed to FP (discontinued) n = 1; GERD-related symptoms and discontinued the steroid, and began treatment with a PPI (discontinued) n = 1; esophageal candidiasis PPI arm n = 0/21	
Oliva 2018	Not reported	Not reported	Not reported
Peterson 2010	Esomeprazole: 0/15 Fluticasone: 0/15	Esomeprazole: 0/15 Fluticasone: 0/15	Not reported
Rothenberg 2015	QAX576: 1/17; asymptomatic cyst-like lesion in the right calf that predated enrollment in the study, but upon subsequent investigations, it turned out to be a spindle cell sarcoma Placebo: 1/8; not reported	QAX576: 10/17 Placebo: 5/8 Adverse events, reported as QAX576; placebo: Cough: 4 (23.5%), 1 (12.5%); nasal congestion: 3 (17.6%), 2 (25.0%); oropharyngeal pain: 3 (17.6%), 2 (25.0%); gastroesophageal reflux disease: 4 (23.5%), 0 (0.0%); headache: 3 (17.6%), 1 (12.5%); nausea: 3 (17.6%), 1 (12.5%); chills: 2 (11.8%), 1 (12.5%); contusion: 2 (11.8%), 1 (12.5%); vomiting: 2 (11.8%), 1 (12.5%); arthralgia: 1 (5.9%), 1 (12.5%); back pain: 1 (5.9%), 1 (12.5%); dermatitis: 2 (11.8%), 0 (0.0%); dizziness: 1 (5.9%), 1 (12.5%); fatigue: 2 (11.8%), 0 (0.0%); ligament sprain: 1 (5.9%), 1 (12.5%); nasal mucosal discoloration: 2 (11.8%), 0 (0.0%); pyrexia: 2 (11.8%), 0 (0.0%); sinus congestion: 2 (11.8%), 0 (0.0%); sinusitis: 1 (5.9%), 1 (12.5%); tonsillolith: 0 (0.0%), 2 (25.0%)	Not reported
Rothenberg 2022	None reported	Not reported as individuals/group, cannot use Injection-site reactions: Dupilumab: 30/80 (37.5%) Placebo: 26/79 (33.3%) Fever: Dupilumab: 5/80 (6.3%) Placebo: 1/79 (1.3%)	Not assessed
Schaefer 2008	Prednisone: 7.5% (3/40) Systemic adverse effects (hyperphagia, weight gain, and/or cushingoid features) Fluticasone: 0% (0/40)	Systemic adverse effects (hyperphagia, weight gain, and/or cushingoid features) Prednisone: 40% (16/40) Fluticasone: 0% (0/40) Esophageal candidal overgrowth Prednisone: 0% (0/40) Fluticasone: 15% (6/40)	Not reported

Table 6. Secondary outcomes (Continued)

Spergel 2012	From NCT00538434: Reslizumab 1 mg/kg: 1/55 Reslizumab 2 mg/kg: 1/57 Reslizumab 3 mg/kg: 1/57 Reslizumab: 3/169 Placebo: 2/57 <ul style="list-style-type: none"> • Abdominal pain 0; 0; 0; 1 • Anaphylaxis 0; 0; 0; 1 • Gastrointestinal inflammation 1; 0; 0; 0 • Respiratory distress 0; 0; 1; 0 • Syncope 0; 0; 0; 1 • Viral gastroenteritis 0; 1; 0; 0 • Patients who discontinued because of an adverse event 1; 0; 0; 0 	From NCT00538434: Reslizumab 1 mg/kg: 38/55 Reslizumab 2 mg/kg: 29/57 Reslizumab 3 mg/kg: 39/57 Reslizumab: 106/169 Placebo: 40/57 <ul style="list-style-type: none"> • Headache: 8; 6; 12; 7 • Cough: 5; 6; 6; 6 • Nasal congestion: 7; 3; 4; 8 • Pharyngo-laryngeal pain: 6; 3; 4; 9 • Upper respiratory tract infection: 5; 4; 6; 5 • Nausea: 6; 2; 4; 3 • Pyrexia: 4; 2; 3; 6 • Sinusitis: 5; 1; 5; 4 • Upper abdominal pain: 4; 2; 4; 5 • Nasopharyngitis: 3; 2; 1; 7 • Diarrhea: 4; 3; 0; 5 	Child Health Questionnaire (CHQ) Validated Landgraf 2014 Physical summary score Mean difference: <ul style="list-style-type: none"> • Arm 1 (1 mg) = -4.75 (-17.35 to 8.03) • Arm 2 (2 mg) = -1.47 (-14.17 to 11.23) • Arm 3 (3 mg) = 1.36 (-11.19 to 13.91) Psychosocial summary score Mean difference: <ul style="list-style-type: none"> • Arm 1 (1 mg) = -6.38 (-15.01 to 2.24) • Arm 2 (2 mg) = -3.41 (-12.05 to 5.23) • Arm 3 (3 mg) = -0.43 (-9.00 to 8.15) Global health summary score Mean difference: <ul style="list-style-type: none"> • Arm 1 = -1.57 (-16.98 to 13.85) • Arm 2 = -5.54 (-20.94 to 9.87) • Arm 3 = -1.06 (-16.16 to 14.06) From NCT00538434: no SDs reported, cannot use
Spergel 2020	Viaskin milk: 0/15 Placebo: 1/5; vocal cord dysfunction in a participant with asthma leading to a hospitalization at day 2 of the study	Total (n = 20) Blood and lymphatic system disorders Ear and labyrinth disorders Eye disorders Gastrointestinal disorders	PedsQL - Quality of life (validated) Viaskin milk: 24.4 (20.68)/7 Placebo: 38.00 (18.38)/2

Table 6. Secondary outcomes (Continued)

General disorders and administration site conditions

Infections and Infestations

Injury, poisoning, and procedural complications

Metabolism and nutrition disorders

Musculoskeletal and connective tissue disorders

Nervous system disorders

Respiratory, thoracic, and mediastinal disorders

Skin and subcutaneous tissue disorders

Viaskin milk (n = 15)

Blood and lymphatic system disorders

Ear and labyrinth disorders

Eye disorders

Gastrointestinal disorders

General disorders and administration site conditions

Infections and Infestations

Injury, poisoning, and procedural complications

Metabolism and nutrition disorders

Musculoskeletal and connective tissue disorders

Nervous system disorders

Respiratory, thoracic, and mediastinal disorders

Skin and subcutaneous tissue disorders

Placebo (n = 5)

Blood and lymphatic system disorders

Ear and labyrinth disorders

Eye disorders

Gastrointestinal disorders

General disorders and administration site conditions

Infections and Infestations

Injury, poisoning, and procedural complications

Metabolism and nutrition disorders

Musculoskeletal and connective tissue disorders

Nervous system disorders

Respiratory, thoracic, and mediastinal disorders

Skin and subcutaneous tissue disorders

Table 6. Secondary outcomes (Continued)

		Viaskin milk: 15/15 Placebo: 5/5	
Straumann 2010a	Mepolizumab: 0/5 Placebo: 0/6	Mepolizumab: 2/5 Placebo: 2/6 <ul style="list-style-type: none"> • Any adverse event: 2/5 (used) • Nausea = 0/5 • Esophageal food impaction = 0/5 • Vomiting = 0/5 • Fatigue = 1/5 • Upper respiratory tract infection = 1/5 <ul style="list-style-type: none"> • ANy adverse event: 2/6 (used) • Nausea = 1/6 • Esophageal food impaction = 1/6 • Vomiting = 1/6 • Vomiting = 0/6 • Upper respiratory tract infection = 0/6 	Not reported
Straumann 2010b	Budesonide: 0/18 Placebo: 0/18	Budesonide: 4/18 Placebo: 1/18 Budesonide: 3/18 mild signs of clinically asymptomatic esophageal candidiasis on follow-up endoscopy 1/18 histologic Candida without endoscopic findings of the same Placebo: 1/18 hoarseness	Not reported
Straumann 2011	Budesonide: 0/14 Placebo: 0/14	Budesonide: 0/14 Placebo: 0/14	Not reported
Straumann 2013	OC000459: 1/14 Placebo: 0/12 1/14 patients with a serous event in the OC000459 arm – acute appendicitis in follow-up period	Not reported as individuals/group, cannot use 1/14 patients with a serous event in the OC000459 arm – acute appendicitis in follow-up period 1/12 patients in the placebo arm with an adverse event - dizziness 15 minor events; 6 x OC000459 and 9 x placebo	Not reported
Straumann 2020	Budesonide 0.5 mg: 3/68 Budesonide 1.0 mg: 1/68 Placebo: 0/68 Budesonide: 4/136 Placebo: 0/68	Budesonide 0.5 mg: 57/68 Budesonide 1.0 mg: 59/68 Budesonide: 116/ Placebo = 61/68 Cartilage injury: 1 (1.5%), 0, 0; upper limb fracture: 1 (1.5%), 0, 0; sinusitis: 1 (1.5%), 0, 0; inguinal hernia: 1 (1.5%), 0, 0; skull fracture: 0, 1 (1.5%), 0; condition aggravated (clinical relapse%): 7 (10.3%), 5 (7.4%), 41 (60.3%); food impaction needing endoscopic intervention: 0, 0, 2 (2.9%); chest pain:	Eosinophilic esophagitis quality of life scale for adults (EoE-QoL-A) questionnaire version 2.0 S. Bajaj, T. Taft, L. Keefer, et al. Validity, usability, and acceptability of the eosinophilic

Table 6. Secondary outcomes (Continued)

<p>Not clear on particulars; the 1/68 likely a skull fracture</p>	<p>0, 1 (1.5%), 0; retinitis: 0, 1 (1.5%), 0; oropharyngeal pain: 0, 1 (1.5%), 0; dermatitis allergic: 0, 1 (1.5%), 0; esophageal dilation: 0, 0, 1 (1.5%); food impaction needing endoscopic intervention: 0, 0, 2 (2.9%); food impaction without need for endoscopic intervention: 0, 3 (4.4%), 0; eye disorders: 1 (1.5%), 1 (1.5%), 1 (1.5%); cataract nuclear: 0, 0, 1 (1.5%); gastrointestinal disorders: 5 (7.4%), 5 (7.4%), 0; general disorders and administration site conditions: 2 (2.9%), 2 (2.9%), 0; infections and infestations: 12 (17.6%), 10 (14.7%), 1 (1.5%); candidiasis overall: 12 (17.6%), 9 (13.2%), 0; suspected symptomatic candidiasis: 11 (16.2%), 8 (11.8%), 0; histologic confirmed and symptomatic candidiasis: 4 (5.9%), 1 (1.5%), 0; investigations: 3 (4.4%), 2 (2.9%), 0; blood cortisol decreased: 2 (2.9%), 2 (2.9%), 0; neoplasms benign, malignant and unspecified: 0, 1 (1.5%), 0; lipoma: 0, 1 (1.5%), 0; nervous system disorders: 3 (4.4%), 3 (4.4%), 0; dysgeusia: 0, 1 (1.5%), 0; reproductive system and breast disorders: 0, 1 (1.5%), 1 (1.5%); respiratory, thoracic and mediastinal disorders: 0, 1 (1.5%), 0; skin and subcutaneous tissue disorders: 1 (1.5%), 3 (4.4%), 0; vascular disorders: 0, 1 (1.5%), 0; hypertension: 0, 1 (1.5%), 0</p>	<p>esophagitis quality of life scale for adults (EoE-QoL-A)</p> <p>T.H. Taft, E. Kern, M.A. Kwiatek, et al. The adult eosinophilic oesophagitis quality of life questionnaire: a new measure of health-related quality of life</p> <p>Aliment Pharmacol Ther, 34 (2011), pp. 790-798</p> <p>EoE-QoL-A end of treatment (mean ± SD)</p> <p>BOT 0.5: -3.3 ± 0.46/68</p> <p>BOT 1.0: -3.5 ± 0.48/68</p> <p>BOT: -3.4 (0.48)/136</p> <p>Placebo: -2.8 ± 0.75/68</p>	
<p>Tytor 2021</p>	<p>Mometasone: 0/17 Placebo: 0/19</p>	<p>Mometasone: 0/17 Placebo: 0/19</p>	<p>The organ-related QoL was evaluated using the EORTC QLQ-OES18 (originally developed and validated for patients with esophagus cancer)</p> <p>(References: Blazeby JM, Conroy T, Hammerlid E, et al. Clinical and psychometric validation of an EORTC questionnaire module, the EORTC QLQOES18, to assess quality of life in patients with oesophageal cancer. Eur J Cancer. 2003;39(10):1384-1394.</p> <p>Blazeby JM, Alderson D, Winstone K, et al. Development of an EORTC questionnaire module to be used in quality of</p>

Table 6. Secondary outcomes (Continued)

life assessment for patients with oesophageal cancer. The EORTC Quality of Life Study Group. *Eur J Cancer*. 1996;32(11):1912–1917

General QoL, Short Form-36 (SF-36)

(Sullivan M, Karlsson J, Ware JE. Jr., The Swedish SF-36 Health Survey–I. Evaluation of data quality, scaling assumptions, reliability and construct validity across general populations in Sweden. *Soc Sci Med*. 1995;41(10):1349–1358)

No difference between placebo and intervention although no numerical data provided

AAF: amino acid-based formula; AE: adverse event; BET: budesonide effervescent tablet; BOS: budesonide oral suspension; BOT: budesonide orodispersible tablet; BOV: budesonide, oral viscus; CG: control group; EoE: eosinophilic esophagitis; EoT: end of treatment; FFED: four food elimination diet; FP: fluticasone propionate; GERD: gastroesophageal reflux disease; IG: intervention group; NEB: nebulized/swallowed budesonide solution; LS: least squares; QOL: quality of life; SAE: serious adverse event; SE: standard error; TEAE: treatment-emergent adverse event; WBC: white blood cell

APPENDICES

Appendix 1. CENTRAL via Cochrane Library search strategy

Date Run: 04/03/2023 01:04:38

#1 [mh "Eosinophilic Esophagitis"] OR ([mh Esophagitis] AND [mh Eosinophilia]) OR ((Eosinophil* OR Eosinophyl*) AND (Esophag* OR Oesophag*)) with Cochrane Library publication date Between Oct 2021 and Mar 2023, in Trials 94

Appendix 2. MEDLINE via Ovid SP search strategy

Database: Ovid MEDLINE(R) ALL <1946 to March 02, 2023>

1 ((Randomized Controlled Trial or Controlled Clinical Trial).pt. or (Randomi?ed or Placebo or Randomly or Trial or Groups).ab. or Drug Therapy.fs.) not (exp Animals/ not Humans.sh.) (4899157)

2 Eosinophilic Esophagitis/ or (Esophagitis/ and Eosinophilia/) or ((Eosinophil* or Eosinophyl*) and (Esophag* or Oesophag*)).tw,kw. (4480)

3 1 and 2 (1182)

4 limit 3 to ed=20211023-20230303 (173)

5 limit 3 to dt=20211023-20230303 (121)

Medical treatment of eosinophilic esophagitis (Review)

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

6 4 or 5 (194)

Appendix 3. Embase via Ovid SP search strategy

Database: Embase <1974 to 2023 Week 08>

1 Randomized controlled trial/ or Controlled clinical study/ or randomization/ or intermethod comparison/ or double blind procedure/ or human experiment/ or (random\$ or placebo or (open adj label) or ((double or single or doubly or singly) adj (blind or blinded or blindly)) or parallel group\$1 or crossover or cross over or ((assign\$ or match or matched or allocation) adj5 (alternate or group\$1 or intervention \$1 or patient\$1 or subject\$1 or participant\$1)) or assigned or allocated or (controlled adj7 (study or design or trial)) or volunteer or volunteers).ti,ab. or (compare or compared or comparison or trial).ti. or ((evaluated or evaluate or evaluating or assessed or assess) and (compare or compared or comparing or comparison)).ab. (6208790)

2 (random\$ adj sampl\$ adj7 ("cross section\$" or questionnaire\$1 or survey\$ or database\$1)).ti,ab. not (comparative study/ or controlled study/ or randomi?ed controlled.ti,ab. or randomly assigned.ti,ab.) (9365)

3 Cross-sectional study/ not (randomized controlled trial/ or controlled clinical study/ or controlled study/ or (randomi?ed controlled or control group\$1).ti,ab.) (338867)

4 (((case adj control\$) and random\$) not randomi?ed controlled).ti,ab. (21248)

5 (Systematic review not (trial or study)).ti. (251119)

6 (nonrandom\$ not random\$).ti,ab. (18726)

7 ("Random field\$" or (random cluster adj3 sampl\$)).ti,ab. (4426)

8 (review.ab. and review.pt.) not trial.ti. (1090581)

9 "we searched".ab. and (review.ti. or review.pt.) (48360)

10 ("update review" or (databases adj4 searched)).ab. (60294)

11 (rat or rats or mouse or mice or swine or porcine or murine or sheep or lambs or pigs or piglets or rabbit or rabbits or cat or cats or dog or dogs or cattle or bovine or monkey or monkeys or trout or marmoset\$1).ti. and animal experiment/ (1214071)

12 Animal experiment/ not (human experiment/ or human/) (2549714)

13 or/2-12 (4261110)

14 1 not 13 (5483547)

15 Eosinophilic Esophagitis/ or (Esophagitis/ and (Eosinophilia/ or Eosinophilic Gastrointestinal Disorder/)) or ((Eosinophil* or Eosinophyl*) and (Esophag* or Oesophag*)).tw,kw. (10952)

16 14 and 15 (2068)

17 limit 16 to em=202142-202308 (316)

Appendix 4. ClinicalTrials.gov search strategy

Advanced Search

Condition or disease: Eosinophilic Esophagitis

Study type: Interventional Studies (Clinical Trials)

First posted from 10/14/2021 **to** 03/03/2023

24 Studies found

Appendix 5. WHO ICTRP search strategy

Advanced Search

Eosinophilic Esophagitis *in the Condition*

Recruitment status is ALL

Medical treatment of eosinophilic esophagitis (Review)

Copyright © 2023 The Authors. Cochrane Database of Systematic Reviews published by John Wiley & Sons, Ltd. on behalf of The Cochrane Collaboration.

Date of registration is between 01/01/2021 **and** 03/03/2023

59 records for 40 trials found

WHAT'S NEW

Date	Event	Description
20 July 2023	New search has been performed	<p>The number of included studies has increased from 3 to 41, and we have updated the methodology to modern standards. Consequently, the results, discussion, and conclusions have changed considerably since this review was last published.</p> <p>None of the authors of the previously published version are authors on the update (EEJ, TD, MJE). All authors of the update are first-time authors on this review.</p> <p>There have been adjustments in the primary and secondary outcomes, as well as the pre-planned subgroup and sensitivity analyses, since the previous version, reflecting current knowledge on eosinophilic esophagitis.</p> <p>We were not able to perform any pre-planned subgroup or sensitivity analyses for any of the comparisons presented in summary of findings table 2 and tables 4 to 10 due to very limited data. Any pre-planned subgroup and sensitivity analyses not listed under analyses for the comparisons corticosteroids versus placebo for induction of remission, and biologics versus placebo for induction of remission, were also not performed due to lack of data.</p> <p>Funnel plots to judge publication bias were only possible in one instance (Figure 3), as in all other cases there was not a sufficient number of studies (> 10).</p>
20 July 2023	New citation required and conclusions have changed	<p>Initial version of review</p> <p>The initial version of this review, published on 17 March 2010, identified three studies that met the inclusion criteria (Konikoff 2006; Schaefer 2008; Straumann 2010a). These studies examined the efficacy of fluticasone propionate (N = 20, 400 µg twice-daily for 3 months) via metered dose inhaler and swallowed versus placebo (N = 11), oral prednisone (N = 32, 1 mg/kg/dose twice-daily for four weeks) versus topical (swallowed) fluticasone via metered dose inhaler (N = 36, 110 µg per puff (age 1 to 10 years) and 22 µg per puff (> 11 years of age)), and a dose escalation pilot study of mepolizumab compared to placebo (N = 11), respectively. The total number of included participants was 127.</p> <p>The authors concluded the following:</p> <ul style="list-style-type: none"> Comparing topical fluticasone with placebo, fluticasone decreased vomiting more than placebo (67% versus (versus) 27%, $P < 0.05$) but did not improve dysphagia. Histological remission was reported in the fluticasone group compared with the placebo group (50% versus 9%, $P = 0.05$; RR 5.5, 95% CI 0.81 to 37.49). Comparing fluticasone with oral prednisone, symptom resolution and improvement of esophagitis were similar, with the majority of participants being symptom-free at four weeks, with no difference between groups (RR 1.03, 95% CI 0.95 to 1.11).

Date	Event	Description
		<ul style="list-style-type: none"> Comparing mepolizumab to placebo, there was no difference in symptom response with mepolizumab compared to placebo, but the decrease in esophageal eosinophil count was greater with mepolizumab than placebo (67% versus 25%). <p>Current version of review</p> <p>The current version of this review includes an additional 38 studies, which increases the number of analyses from 3 to 18 and increases the total number of participants to 3253.</p> <p>Regarding the conclusions that have changed from the initial review to the present, an additional four trials compared fluticasone versus placebo for histological remission (Alexander 2012; Butz 2014; Dellon 2022a; Hirano 2020f). These included a total of 250 participants and together confirm and reinforce the original conclusion that fluticasone is effective for treatment of EoE as measured by histological dichotomous remission (68% versus 6%, $P < 0.00001$; RR 7.57, 95% CI 3.36 to 17.08; $I^2 = 0\%$). There were no new studies that compared fluticasone with oral prednisone or mepolizumab to placebo, so the results from these analyses did not change in the current review.</p> <p>In addition to these changes in the results, the current version of the review includes 14 new classes of analyses:</p> <ul style="list-style-type: none"> Corticosteroids versus placebo for induction of remission Corticosteroids versus placebo for maintenance of remission Biologics versus placebo for induction of remission Cromolyn sodium versus placebo PGD2R antagonist OC000459 versus placebo Oral viscous budesonide versus swallowed fluticasone Esomeprazole versus fluticasone One-food elimination diet versus four-food elimination diet One-food elimination diet versus six-food elimination diet Four-food elimination diet with omeprazole versus omeprazole Four-food elimination diet with amino acid formula versus four-food elimination diet Nebulized budesonide versus viscous budesonide Viaskin milk patch versus placebo Leukotriene receptor antagonist versus placebo for maintenance of remission <p>Each class of analysis examines, as primary outcomes, clinical (continuous, dichotomous), histological (continuous, dichotomous), and endoscopic (continuous, dichotomous) improvement, and withdrawals due to adverse events (dichotomous), and as secondary outcomes, serious adverse events (dichotomous), total adverse events (dichotomous), and quality of life (continuous, dichotomous), at end of trial when prespecified by the study authors. The sensitivity analyses conducted include analyses based on fixed-effect model, eos/hpf threshold, validated instruments, peer-reviewed publications, and less than high risk of bias, as appropriate. The subgroup analyses conducted include age group, type of steroid (TCSs only), delivery method (TCSs only), and mechanism (biologics only), as appropriate.</p>

HISTORY

Protocol first published: Issue 1, 2003

Review first published: Issue 3, 2004

Date	Event	Description
27 January 2010	New search has been performed	New studies added, review updated.
27 January 2010	New citation required but conclusions have not changed	Author line changed.
30 October 2008	Amended	Converted to new review format.
8 May 2006	New search has been performed	Minor update.
2 February 2006	Amended	New studies sought but none found.
15 April 2004	New citation required and conclusions have changed	Substantive amendment.

CONTRIBUTIONS OF AUTHORS

JPF: led the update of the review, secured funding, designed and developed, screened, extracted, contributed to writing and editing, advised on, and approved the final version prior to submission, and is a guarantor of the review.

MG: led the update of the review, secured funding, designed and developed, screened, extracted, resolved conflicts, assessed certainty, contributed to writing and editing, advised on, and approved the final version prior to submission.

VS: led the writing of the results, designed and developed, screened, extracted, resolved conflicts, assessed certainty, contributed to writing and editing, advised on, and approved the final version prior to submission.

ESD: screened, extracted, contributed to writing and editing, advised on, and approved the final version prior to submission.

SKG: screened, extracted, contributed to writing and editing, advised on, and approved the final version prior to submission.

CR: screened, extracted, advised on, contributed to writing, and approved the final version prior to submission.

CGJ: screened, extracted, contributed to writing and editing, advised on, and approved the final version prior to submission.

RDV: screened, extracted, advised on, contributed to writing, and approved the final version prior to submission.

EAE: screened, extracted, advised on, contributed to writing, and approved the final version prior to submission.

AbE: screened, extracted, and approved the final version prior to submission.

AsE: screened, extracted, and approved the final version prior to submission.

EBM: led the analysis, designed and developed, screened, extracted, contributed to writing and editing, advised on, and approved the final version prior to submission.

Two of the review authors (ED, SG) are the authors of 17 of the included studies ([Assa'ad 2011](#); [Dellon 2012](#); [Dellon 2017](#); [Dellon 2019](#); [Dellon 2021b](#); [Dellon 2022](#); [Dellon 2022a](#); [Dellon 2022b](#); [Gupta 2015](#); [Hirano 2019](#); [Hirano 2020](#); [Kliewer 2019](#); [Hirano 2021](#); [Kliewer 2021](#); [Rothenberg 2022](#); [Oliva 2018](#); [Schaefer 2008](#)). These studies were screened, extracted for data, and assessed for risk of bias independently by EBM, VS, CGJ, CR, RDV, EAE, AbE, and AsE.

DECLARATIONS OF INTEREST

JPF: nothing to declare.

Medical treatment of eosinophilic esophagitis (Review)

MG: Morris Gordon is a Cochrane editor. He was not involved in the editorial process for this review.

VS: nothing to declare.

ESD: Research funding: Adare/Ellodi, Allakos, Arena, AstraZeneca, GSK, Meritage, Miraca, Nutricia, Celgene/Receptos/BMS, Regeneron, Revolo, Shire/Takeda. Consultant: Abbott, Abbvie, Adare/Ellodi, Aimmune, Akesobio, Alfasigma, ALK, Allakos, Amgen, Aqilion, Arena/Pfizer, Aslan, AstraZeneca, Avir, Biorasi, Calypso, Celgene/Receptos/BMS, Celldex, Eli Lilly, EsoCap, Eupraxia, Ferring, GSK, Gossamer Bio, Holoclara, Invea, Landos, LucidDx, Morphic, Nexstone Immunology, Nutricia, Parexel/Calyx, Phathom, Regeneron, Revolo, Robarts/Alimentiv, Salix, Sanofi, Shire/Takeda, Target RWE, Upstream Bio. Educational grant: Allakos, Holoclara, Invea.

He is one of the authors of 13 of the included studies in this review ([Dellon 2012](#); [Dellon 2017](#); [Dellon 2019](#); [Dellon 2022](#); [Dellon 2021b](#); [Dellon 2022a](#); [Dellon 2022b](#); [Hirano 2020](#); [Kliewer 2019](#); [Hirano 2019](#); [Kliewer 2021](#); [Rothenberg 2022](#); [Hirano 2021](#)). These studies were screened, extracted for data, and assessed for risk of bias independently by EBM, VS, CGJ, CR, RDV, EAE, AbE, and AsE.

SKG: Consultant/DSMB member/Author – Adare, BMS, QOL, Takeda, MedScape, PVI, ViaSkin, UpToDate; Research support - Allakos, Ellodi, AstraZeneca. He is one of the authors of seven of the included studies in this review ([Assa'ad 2011](#); [Dellon 2017](#); [Gupta 2015](#); [Hirano 2019](#); [Kliewer 2021](#); [Oliva 2018](#); [Schaefer 2008](#)). These studies were screened, extracted for data, and assessed for risk of bias independently by EBM, VS, CGJ, CR, RDV, EAE, AbE, and AsE.

CR: nothing to declare.

CGJ: Research funding: Dr. Falk Pharma GmbH.

RDV: nothing to declare.

EAE: nothing to declare.

AbE: nothing to declare.

AsE: nothing to declare.

EBM: nothing to declare.

SOURCES OF SUPPORT

Internal sources

- University of Central Lancashire, UK
 - The authors provided their time in kind
- Nemours Children's Health System, USA
 - The authors provided their time in kind

External sources

- NIHR incentive grant, UK
 - NIHR incentive grant NIHR150511

DIFFERENCES BETWEEN PROTOCOL AND REVIEW

There have been adjustments in the primary and secondary outcomes, as well as the pre-planned subgroup and sensitivity analyses since the previous version, reflecting current knowledge on eosinophilic esophagitis.

We were not able to perform any pre-planned subgroup or sensitivity analyses for any of the comparisons presented in summary of findings table 2 and tables 4 to 10 due to very limited data. Any pre-planned subgroup and sensitivity analyses not listed under analyses for the comparisons corticosteroids versus placebo for induction of remission and biologics versus placebo for induction of remission, were also not performed due to lack of data.

Funnel plots to judge publication bias were only possible in one instance ([Figure 3](#)), as in all other cases there was not a sufficient number of studies ($n \geq 10$).

INDEX TERMS**Medical Subject Headings (MeSH)**

Adrenal Cortex Hormones [therapeutic use]; *Biological Products; Chronic Disease; *Eosinophilic Esophagitis [drug therapy]; Proton Pump Inhibitors [therapeutic use]; Randomized Controlled Trials as Topic; Remission Induction

MeSH check words

Adult; Child; Humans