

Human retinal secretome: A cross-link between mesenchymal and retinal cells

Luigi Donato, Concetta Scimone, Simona Alibrandi, Sergio Zaccaria Scalinci, Domenico Mordà, Carmela Rinaldi, Rosalia D'Angelo, Antonina Sidoti

Specialty type: Cell biology

Provenance and peer review:

Invited article; Externally peer reviewed.

Peer-review model: Single blind

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C, C
Grade D (Fair): 0
Grade E (Poor): 0

P-Reviewer: Gallone A, Italy; Lei XH, China; Li SC, United States

Received: December 23, 2022

Peer-review started: December 23, 2022

First decision: March 9, 2023

Revised: March 17, 2023

Accepted: April 10, 2023

Article in press: April 10, 2023

Published online: July 26, 2023



Luigi Donato, Concetta Scimone, Simona Alibrandi, Domenico Mordà, Carmela Rinaldi, Rosalia D'Angelo, Antonina Sidoti, Department of Biomedical and Dental Sciences and Morphofunctional Imaging, University of Messina, Messina 98125, Italy

Luigi Donato, Concetta Scimone, Simona Alibrandi, Domenico Mordà, Department of Biomolecular Strategies, Genetics and Cutting-Edge Therapies, Euro-Mediterranean Institute of Science and Technology, Palermo 90139, Italy

Simona Alibrandi, Department of Chemical, Biological, Pharmaceutical and Environmental Sciences, University of Messina, Messina 98125, Italy

Sergio Zaccaria Scalinci, Department of Medical and Surgical Sciences, University of Bologna, Bologna 40121, Italy

Corresponding author: Simona Alibrandi, PhD, Department of Biomedical and Dental Sciences and Morphofunctional Imaging, University of Messina, Via Consolare Valeria 1, Messina 98125, Italy. salibrandi@unime.it

Abstract

In recent years, mesenchymal stem cells (MSC) have been considered the most effective source for regenerative medicine, especially due to released soluble paracrine bioactive components and extracellular vesicles. These factors, collectively called the secretome, play crucial roles in immunomodulation and in improving survival and regeneration capabilities of injured tissue. Recently, there has been a growing interest in the secretome released by retinal cytotypes, especially retinal pigment epithelium and Müller glia cells. The latter trophic factors represent the key to preserving morphofunctional integrity of the retina, regulating biological pathways involved in survival, function and responding to injury. Furthermore, these factors can play a pivotal role in onset and progression of retinal diseases after damage of cell secretory function. In this review, we delineated the importance of cross-talk between MSCs and retinal cells, focusing on common/induced secreted factors, during experimental therapy for retinal diseases. The cross-link between the MSC and retinal cell secretomes suggests that the MSC secretome can modulate the retinal cell secretome and vice versa. For example, the MSC secretome can protect retinal cells from degeneration by reducing oxidative stress, autophagy and programmed cell death. Conversely, the retinal cell secretome can influence the MSC secretome by inducing changes in

MSC gene expression and phenotype.

Key Words: Secretome; Mesenchymal stem cells; Retinal cells; Extracellular vesicles; Retinal diseases

©The Author(s) 2023. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Recently, the mesenchymal stem cell secretome, a solution rich with paracrine bioactive factors and extracellular vesicles, acquired a significant role in immunomodulation and survival induction of damaged tissues. A secretome is also released by retinal cells, physiologically or following pathological stimuli. One of the most promising therapeutic frontiers is represented by a possible “cross-talk” between mesenchymal stem cells and retinal cells through the secretomes in order to improve the knowledge on released factors mechanisms of action during their potentially beneficial role.

Citation: Donato L, Scimone C, Alibrandi S, Scalinci SZ, Mordà D, Rinaldi C, D'Angelo R, Sidoti A. Human retinal secretome: A cross-link between mesenchymal and retinal cells. *World J Stem Cells* 2023; 15(7): 665-686

URL: <https://www.wjgnet.com/1948-0210/full/v15/i7/665.htm>

DOI: <https://dx.doi.org/10.4252/wjsc.v15.i7.665>

INTRODUCTION

In recent years, mesenchymal stem cells (MSCs) have been indicated as the most effective source for cell-based therapy, particularly in regenerative medicine. In particular, MSCs produce major therapeutic effects releasing soluble paracrine bioactive components and extracellular vesicles (EVs) constituting the so called secretome. These secreted factors play crucial roles in modulating immunity and improving survival and regeneration capabilities of injured tissue[1].

Secreted trophic factors are also key to preserving the morphofunctional integrity of the retina, regulating biological pathways involved in survival, function and response to injury[2]. Additionally, these factors can play a fundamental role in onset and progression of retinal diseases after damage of cell secretory functions[3]. In this review, we discussed the link between the secretome of MSCs to the retinal cell secretome, in order to highlight the current knowledge of secreted factor involvement in retinal diseases.

Main features of MSCS

One of the most recent fields of therapy research concerns MSCs, multipotent non-hematopoietic stem cells that originate from the mesoderm. They can reach a pathological site following the release of different biologically active immunomodulatory and regenerative factors related to different diseases[4]. There are multiple sources of MSCs, including umbilical cord blood, placenta, adipose tissue, skin and bone marrow tissue, with the latter representing the most widely used source[5]. Isolation of different types of MSCs, such as adipose tissue-derived mesenchymal stromal/stem cells (ASCs), is a non-invasive process, and this represents a fundamental advantage from an ethical and/or legal point of view[6]. The key point for MSC use is their low immunogenicity, permitting allogeneic trans-plantation in the medical setting[7].

Moreover, recent studies have shown that MSCs can produce an immune response, mediated by T cells regulated by IFN- γ [8]. When activated, MSCs can reach the correct pathological site to exert reparative functions, triggered by a huge number of secreted factors from the injured cells, such as cytokines, chemokines and growth factors[9]. Among the latter, placental growth factor plays a pivotal role, along with VEGF, EPO, SDF-1, ANG2, G-CSF, stem cell factor, PDGF, EGF, HGF and IGF-1[10]. Regarding cytokines and chemokines, the former include TNF- α and interleukins such as IL-1b, IL-2, IL-3, IL-6 and IL-8, and the latter includes, among others, CCL5 and CCL22[11].

Various studies have confirmed that human MSCs evade allorecognition, affect T lymphocytes and dendritic cell activities and produce a local immunosuppressant microenvironment by releasing the already cited cytokines[12]. Moreover, MSCs can be easily genetically manipulated, with elevated metabolic activity and low mutation rate, and can efficiently secrete a wide number of proteins[13]. Today, preclinical and clinical trials using MSCs, especially human bone marrow-derived MSCs (hBMMSCs) and human adipose mesenchymal stem cells (hADSCs), have been performed in different kinds of pathologies with promising results, such as autoimmune disease, joint reconstruction, vascular disease, nerve injury, organ transplantation, degenerative disease and severe infection[14].

In particular, one of the fields with the highest number of ongoing clinical trials is represented by eye diseases (Table 1). The protective activity of MSCs was initially linked to their direct differentiation and replacement of injured tissues, as evidenced by human MSCs becoming hepatocyte-like cells or rat MSCs turning into neuron-like cells[15]. However, today the protective action of MSCs is well known to be primarily mediated through paracrine properties, exerted by what is defined as the MSC secretome.

MSC secretome

The secretome released by MSCs consists of a conditioned medium (CM) made up of soluble elements (cytokines and growth factors) and a vesicular part made up of exosomes and microvesicles, which are fundamental for protein and genetic material transfer towards other cells[16]. The most recent *in vitro* and *in vivo* studies on the features of the MSC

Table 1 Ongoing clinical trials on eye diseases based on the use of mesenchymal stem cells

Title	Sponsor/collaborators	URL
Safety and Efficacy of Pluripotent Stem Cell-derived Mesenchymal Stem Cell Exosome (PSC-MSC-Exo) Eye Drops Treatment for Dry Eye Diseases Post Refractive Surgery and Associated With Blepharospasm	Second Affiliated Hospital, School of Medicine, Zhejiang University Zhejiang University Hangzhou yuansheng biotechnology Co., Ltd	https://ClinicalTrials.gov/show/NCT05738629
Therapeutic Effect of Stem Cell Eye Drops on Dry Eye Disease	The Affiliated Nanjing Drum Tower Hospital of Nanjing University Medical School	https://ClinicalTrials.gov/show/NCT05784519
Safety of Cultured Allogeneic Adult Umbilical Cord Derived Mesenchymal Stem Cells for Eye Diseases	The Foundation for Orthopaedics and Regenerative Medicine	https://ClinicalTrials.gov/show/NCT05147701
The Role of Transscleral Cyclophotocoagulation in Patients Undergoing a Boston Keratoprosthesis	Centre hospitalier de l'Université de Montreal (CHUM) Fonds de recherche en ophtalmologie de l'Université de Montreal	https://ClinicalTrials.gov/show/NCT04232982
Characterization of Potential Biomarkers of Eye Disease and Vision	Association for Innovation and Biomedical Research on Light and Image	https://ClinicalTrials.gov/show/NCT02500862
Long-term Safety of UC-MSC Transplantation in Patients With Retinitis Pigmentosa	PT. Prodia Stem Cell Indonesia	https://ClinicalTrials.gov/show/NCT05786287
Clinical Evaluation of Two Daily Disposable Lenses in Sphere Design	Coopervision, Inc.	https://ClinicalTrials.gov/show/NCT05516082
Efficacy of Locally Delivered Allogeneic Mesenchymal Stromal Cells	University of Illinois at Chicago United States Department of Defense	https://ClinicalTrials.gov/show/NCT05705024
Diquafosol <i>vs</i> Hyaluronic Acid for Diabetic Dry Eye	He Eye Hospital	https://ClinicalTrials.gov/show/NCT05682547
Eye Length Signal With Myopia Control	Brien Holden Vision Institute	https://ClinicalTrials.gov/show/NCT04813640
Patient Acceptability of Autonomous Telemedicine	Ufonia Buckinghamshire Healthcare NHS Trust Innovate UK	https://ClinicalTrials.gov/show/NCT04885868
Phase 2b Pivotal Study of Izokibep in Non-infectious, Intermediate-, Posterior- or Pan-uveitis	ACELYRIN Inc.	https://ClinicalTrials.gov/show/NCT05384249
Dose Optimization for Safe and Efficient Fluorescein Angiography (DOSE Study)	Seoul National University Bundang Hospital	https://ClinicalTrials.gov/show/NCT05664555
Analysis of the Results of Intense Pulsed Light Treatment Previously to Laser Refractive Surgery	Vissum, Instituto Oftalmologico de Alicante	https://ClinicalTrials.gov/show/NCT05139511
Study of the Association Between Digital Eye Syndrome With Binocular Vision and the Ocular Surface in Higher Education Students in the Area of Health Technologies	Universidade Nova de Lisboa NOVA Medical School Faculdade de Ciências Médicas, Universidade Nova de Lisboa Escola Superior de Tecnologia da Saúde de Lisboa (ESTeSL) University of Évora CINTESIS@RISE, NOVA Medical School Faculdade de Ciências Médicas, Universidade Nova de Lisboa Comprehensive Health Research Center (CHRC), Universidade Nova de Lisboa	https://ClinicalTrials.gov/show/NCT05675475
Caffeine Consumption and Cataract Prevention	Vienna Institute for Research in Ocular Surgery	https://ClinicalTrials.gov/show/NCT05194696
Rotational Stability of the TECNIS Eyhance Toric	Vienna Institute for Research in Ocular Surgery	https://ClinicalTrials.gov/show/NCT05126368
Reading Performance in Patients With Acrysof IQ Vivify Versus Acrysof IQ	Vienna Institute for Research in Ocular Surgery	https://ClinicalTrials.gov/show/NCT05194657
Performance of Two Intraocular Lenses With Extended Depth of Vision	Vienna Institute for Research in Ocular Surgery	https://ClinicalTrials.gov/show/NCT05194150
Assessment of Retinal Vascular Changes With and Without ILM Peeling in Diabetic Vitrectomy Using OCT-A	Kasr El Aini Hospital	https://ClinicalTrials.gov/show/NCT05739539
Primary Vitrectomy With Silicone Oil or SF6 for Rhegmatogenous Retinal	Cairo University	https://ClinicalTrials.gov/show/NCT05377606

Detachment		
Intravitreal Sirolimus as Therapeutic Approach to Uveitis	Stanford University Santen Inc.	https://ClinicalTrials.gov/show/NCT01280669
Clinical Trial With Artiflex Presbyopic	Ophtec BV	https://ClinicalTrials.gov/show/NCT04632784
Post-Market Evaluation of the EVO ICL	Staar Surgical Company	https://ClinicalTrials.gov/show/NCT05538754
Clinical Evaluation of Two Multifocal Contact Lenses	Coopervision, Inc.	https://ClinicalTrials.gov/show/NCT05457608
Ologen Collagen Matrix Versus Mitomycin-C in Patients With Juvenile-onset Open Angle Glaucoma	L.V. Prasad Eye Institute	https://ClinicalTrials.gov/show/NCT03548805
A Comparative Study of Visual Outcome of Two Extended Depth of Focus Intraocular Lenses After Cataract Surgery	Cairo University	https://ClinicalTrials.gov/show/NCT05647421
Keratometric Change After XEN, Trabeculectomy and Tube Shunts	Centre hospitalier de l'Université de Montreal (CHUM) Allergan	https://ClinicalTrials.gov/show/NCT04602923
Post-market Follow Up Study on Paragon CRT 100 (Paflucof D)	Coopervision, Inc. TigerMed	https://ClinicalTrials.gov/show/NCT04187599
Evaluating Two Multifocal Daily Disposable Contact Lenses	Coopervision, Inc.	https://ClinicalTrials.gov/show/NCT05579886
A Clinical Comparison of Two Soft Multifocal Contact Lenses	Coopervision, Inc.	https://ClinicalTrials.gov/show/NCT05794126
Ocular Surface Disease and IOP Monitoring With Travoprost Without Conservatives	Democritus University of Thrace	https://ClinicalTrials.gov/show/NCT05319470
Performance of Two Hydrophobic IOLs	Vienna Institute for Research in Ocular Surgery	https://ClinicalTrials.gov/show/NCT05639049
RayOne EMV Mini-monovision - Efficiency and Safety of 3 Grades of Mini-monovision	Somich, s.r.o.	https://ClinicalTrials.gov/show/NCT05417633
A Clinical Study to Evaluate the Potential Role of ACTH Gel in Patients With Scleritis	Metropolitan Eye Research & Surgery Institute Mallinckrodt Stanford University Ocular Imaging Research and Reading Center Foresight Studies, LLC	https://ClinicalTrials.gov/show/NCT03465111
Zimbabwe Eyecare And Learning (ZEAL): Formative Research on Hyperopia and Educational Outcomes in Primary School Children	Queen's University, Belfast L.V. Prasad Eye Institute University of Zimbabwe University of Ulster New England College of Optometry Peek Vision Zimbabwe Optometric Association Clearly Christian Blind Mission	https://ClinicalTrials.gov/show/NCT05538182
Vision and Balance Changes After Bilateral Implantation of Toric IOLs	University of Plymouth Carl Zeiss Meditec AG Glasgow Caledonian University University of St Mark and St John Plymouth University Hospital Plymouth NHS Trust	https://ClinicalTrials.gov/show/NCT05629078
Multicenter Study on the Efficacy and Safety of OCS-01 in Subjects With Uveitis Related and Post Surgical Macular Edema	Quan Dong Nguyen Global Ophthalmic Research Center (GORC) Oculis Stanford University	https://ClinicalTrials.gov/show/NCT05608837
Inflammatory Biomarkers in Ocular Surface in Primary Open Angle Glaucoma or Ocular Hypertension Under Topical Prostaglandins	Instituto Universitario de Oftalmobiología Aplicada (Institute of Applied Ophthalmobiology) - IOBA Hospital Clinico Universitario de Valladolid	https://ClinicalTrials.gov/show/NCT05039684
Ranibizumab vs Bevacizumab for Type 1 Retinopathy of Prematurity	Zagazig University Cairo University	https://ClinicalTrials.gov/show/NCT05033106
Dresden Corneal Disease and Treatment Study	Technische Universität Dresden	https://ClinicalTrials.gov/show/NCT04251143
Reliability, Validity of the Turkish Version of the Primary Sjogren Syndrome Quality of Life (PSS-QoL) Questionnaire	Gazi University	https://ClinicalTrials.gov/show/NCT04858464
Mean Visual Acuity Changes Following Five Injections of Aflibercept	McMaster University	https://ClinicalTrials.gov/show/NCT02645266

Targeted Fluorescence Imaging in AMD	University Medical Center Groningen	https://ClinicalTrials.gov/show/NCT05262244
Treatment of Ligneous Conjunctivitis in Children With Plasminogen Deficiency	University of Saskatchewan Canadian Blood Services	https://ClinicalTrials.gov/show/NCT05404932
Can the Risk for AMD be Modulated?	Association for Innovation and Biomedical Research on Light and Image	https://ClinicalTrials.gov/show/NCT05735730
Study to Evaluate the Response to Supplementation With Postbiotics in Patients With Macular Degeneration	Institut de la Macula y la Retina Igen BioLab SLU	https://ClinicalTrials.gov/show/NCT05056025
Development of a Tele-Physiotherapy Tool for the Early Management of Musculoskeletal Pain in People With Visual Impairment (TeleEDxPhysio)	Escuela Universitaria de Fisioterapia de la Once Universidad de Zaragoza	https://ClinicalTrials.gov/show/NCT05478200
Clinical Trial to Evaluate Safety and Efficacy of Cell Therapy in Patients With Cicatricial Conjunctivitis	Instituto de Investigacion Sanitaria de la Fundacion Jimenez Diaz Efficce Servicios Para la Investigacion S.L	https://ClinicalTrials.gov/show/NCT05520086
Advanced Glaucoma Progression Study	University of California, Los Angeles National Eye Institute (NEI)	https://ClinicalTrials.gov/show/NCT01742819
Retrobulbar Methylprednisolone as Adjunctive Treatment in Optic Neuritis Trial	Asociacion para Evitar la Ceguera en Mexico	https://ClinicalTrials.gov/show/NCT04942002
Nystagmus Assessment for Patients Consulting in the Emergency Department for Acute Vertigo	CHU de Quebec-Universite Laval	https://ClinicalTrials.gov/show/NCT05176015
Effectiveness of Periocular Drug Injection in CATaract Surgery	Luigi Rondas European Society of Cataract and Refractive Surgeons Academisch Ziekenhuis Maastricht	https://ClinicalTrials.gov/show/NCT05158699
Clinical Trial Comparing Two Non-Surgical Treatments for Severe Blepharoptosis	Massachusetts Eye and Ear Infirmary National Eye Institute (NEI)	https://ClinicalTrials.gov/show/NCT04678115
Stem Cell Ophthalmology Treatment Study II	MD Stem Cells	https://ClinicalTrials.gov/show/NCT03011541
Effect of Intravenous Methylprednisolone and Intravenous Erythropoietin in Toxic Optic Neuropathies: Randomized Clinical Trial	Asociacion para Evitar la Ceguera en Mexico	https://ClinicalTrials.gov/show/NCT05748561
SPT Screening in Irradiated Hereditary Retinoblastoma Survivors	Amsterdam UMC, location VUmc ODAS	https://ClinicalTrials.gov/show/NCT02329002
Community Access Through Remote Eyesight (CARE) Study	New England College of Optometry National Institute on Disability, Independent Living, and Rehabilitation Research University of California, Los Angeles	https://ClinicalTrials.gov/show/NCT04926974
Electro-acupuncture and Transcorneal Electrical Stimulation (TES) for Retinitis Pigmentosa	Nova Southeastern University National Eye Institute (NEI)	https://ClinicalTrials.gov/show/NCT02086890
Evaluation of NeoRetina Artificial Intelligence Algorithm for the Screening of Diabetic Retinopathy at the CHUM	Centre hospitalier de l'Université de Montreal (CHUM) DIAGNOS Inc.	https://ClinicalTrials.gov/show/NCT04699864
Clemastine Fumarate as Remyelinating Treatment in Internuclear Ophthalmoparesis and Multiple Sclerosis	Amsterdam UMC, location VUmc	https://ClinicalTrials.gov/show/NCT05338450
The K-Map Study, Global Prevalence of KC	University Hospital, Geneva ELZA Institute	https://ClinicalTrials.gov/show/NCT03115710
Methotrexate For The Prevention and Treatment of Proliferative Vitreoretinopathy in Pediatric Patients	Stanford University	https://ClinicalTrials.gov/show/NCT04830878
A Collaborative Resource of Heidelberg Multimodal Imaging of Intermediate and Early Atrophic AMD Cases to Study Prediction of Disease Progression	Association for Innovation and Biomedical Research on Light and Image European Vision Institute Clinical Research Network	https://ClinicalTrials.gov/show/NCT05698316

Clinical Trial of Multi-Periscope Prism Glasses for Hemianopia	Massachusetts Eye and Ear Infirmary National Eye Institute (NEI)	https://ClinicalTrials.gov/show/NCT04827147
Feasibility Tests for Various Prism Configurations for Visual Field Loss	Massachusetts Eye and Ear Infirmary National Eye Institute (NEI)	https://ClinicalTrials.gov/show/NCT04424979
Quality Assurance Via Telephone Interviews After Cataract Surgery	Vienna Institute for Research in Ocular Surgery	https://ClinicalTrials.gov/show/NCT05215002
0.01% Hypochlorous Acid in the Treatment of Blepharitis	Eye & ENT Hospital of Fudan University ShuGuang Hospital	https://ClinicalTrials.gov/show/NCT05608980
Birdshot Chorioretinopathy: Prospective Follow-up and Immunogenetic Studies(CO-BIRD)	Assistance Publique - Hopitaux de Paris	https://ClinicalTrials.gov/show/NCT05153057
Characterization of Retinal Disease Progression in Eyes With Non Proliferative Diabetic Retinopathy in Diabetes Type 2 Using Non-invasive Procedures (CHART)	Association for Innovation and Biomedical Research on Light and Image European Vision Institute Clinical Research Network	https://ClinicalTrials.gov/show/NCT04636307
Swiss Pediatric Inflammatory Brain Disease Registry (Swiss-Ped-IBrainD)	University of Bern Schweizerische Multiple Sklerose Gesellschaft University Hospital Inselspital, Berne Roche Pharma (Switzerland) Ltd Novartis	https://ClinicalTrials.gov/show/NCT05017142
Topotecan Episcleral Plaque for Treatment of Retinoblastoma	Targeted Therapy Technologies, LLC	https://ClinicalTrials.gov/show/NCT04156347
Comparison of Phacoemulsification and Corneal Damage Between FLACS and Standard Phaco With Two Handpieces	Centre hospitalier de l'Université de Montreal (CHUM)	https://ClinicalTrials.gov/show/NCT05119270
Computer-based Tutorial and Automated Speech Recognition for Intravitreal Drug Injections	Prim. Prof. Dr. Oliver Findl, MBA Vienna Institute for Research in Ocular Surgery	https://ClinicalTrials.gov/show/NCT04142164
OCS-05 in Patients With Acute Optic Neuritis	Oculus Neurotrials	https://ClinicalTrials.gov/show/NCT04762017
PROgressive Supranuclear Palsy CorTico-Basal Syndrome Multiple System Atrophy Longitudinal Study UK	University College, London University of Cambridge University of Oxford University of Manchester Newcastle University University of Sussex Royal Gwent Hospital	https://ClinicalTrials.gov/show/NCT02778607
Therapeutic Recommendations For The Treatment Of Children With A Retinoblastoma	French Africa Pediatric Oncology Group	https://ClinicalTrials.gov/show/NCT04425434
Temperature on Evaporative Dry Eye	He Eye Hospital	https://ClinicalTrials.gov/show/NCT05720754
Prevalence of Visual Dysfunction in Neurological Disorders	University of Florida	https://ClinicalTrials.gov/show/NCT04836715
Intravitreal Infliximab for Proliferative Vitreoretinopathy	Cairo University	https://ClinicalTrials.gov/show/NCT04891991
A Patch Free Treatment for Young Children With Amblyopia	University of Waterloo Retina Foundation of the Southwest McGill University Queensland University of Technology	https://ClinicalTrials.gov/show/NCT04086524
Effectiveness of the Serious Game 'Broodles' for Siblings of Children With Visual Impairment and/or Intellectual Disability	VU University of Amsterdam	https://ClinicalTrials.gov/show/NCT05376007
Screening for Oculocerebral Lymphoma With the Phenotype of NK Cells in Patients With Uveitis	Hospices Civils de Lyon	https://ClinicalTrials.gov/show/NCT05388838
10-year Progression of Diabetic Retinopathy: Identification of Signs and Surrogate Outcomes	Association for Innovation and Biomedical Research on Light and Image	https://ClinicalTrials.gov/show/NCT04650165
Management of DE With IPL in Combination With DQS	He Eye Hospital	https://ClinicalTrials.gov/show/NCT05694026
Endophthalmitis Post Intravitreal Injections	Rajeev Muni Unity Health Toronto	https://ClinicalTrials.gov/show/NCT04035369
Discovering Early Biomarkers in Circulating Endothelial Cells for	Aarhus University Hospital University of Aarhus	https://ClinicalTrials.gov/show/NCT05169502

Diabetes Complications by Single Cell RNA Sequencing		
Diabetic Retinopathy Classification: ETDRS 7-fields <i>vs</i> Widefield Imaging (ClarusDR)	Association for Innovation and Biomedical Research on Light and Image	https://ClinicalTrials.gov/show/NCT05746975
Evaluation of Desensitization Therapy and Re-treatment of Eye Movement Information [EMDR] in Patients With Post-traumatic Stress Disorder [PTSD]	Centre hospitalier de Ville-Evrard, France	https://ClinicalTrials.gov/show/NCT04431765
Prediction of Progression of Retinal Ischemia in Diabetes	Association for Innovation and Biomedical Research on Light and Image	https://ClinicalTrials.gov/show/NCT05581225
Personalized Parkinson Project PSP Cohort	Radboud University Medical Center UCB Pharma Verily Life Sciences LLC	https://ClinicalTrials.gov/show/NCT05501431
Enriched Eggs for Retina Health in Type 2 Diabetes	University of Manitoba Egg Farmers of Canada	https://ClinicalTrials.gov/show/NCT04496817
Pneumatic Retinopexy Versus Vitrectomy for Retinal Detachment in Patients With Extended Criteria	Unity Health Toronto	https://ClinicalTrials.gov/show/NCT02871531
Adherence to Lifestyle Changes for Age-related Macular Degeneration	Erasmus Medical Center CORR foundation	https://ClinicalTrials.gov/show/NCT05667441
Timing of Glaucoma Drainage Device With Boston Keratoprosthesis	Centre hospitalier de l'Université de Montreal (CHUM)	https://ClinicalTrials.gov/show/NCT02084745
Patient Satisfaction and Visual Function Following Implantation of Trifocals or Extended Range of Vision Intraocular Lenses	Queen's University University of Toronto	https://ClinicalTrials.gov/show/NCT04900662
A Computerized, Adaptive Therapeutic Gaming Approach Training Visual Perceptual Skills in Children With CVI	Universitaire Ziekenhuizen KU Leuven Vrije Universiteit Brussel Fund for Scientific Research, Flanders, Belgium	https://ClinicalTrials.gov/show/NCT05014503
Re-Orchestration of Interregional Oscillatory Activity to Promote Visual Recovery	Ecole Polytechnique Federale de Lausanne	https://ClinicalTrials.gov/show/NCT05220449
Effect of Type of Head Positioning on Retinal Displacement in Vitrectomy for Retinal Detachment	Unity Health Toronto	https://ClinicalTrials.gov/show/NCT04035343
Macular Perfusion Changes After Anti-VEGF Versus Targeted Retinal Photocoagulation in Proliferative Diabetic Retinopathy	Cairo University	https://ClinicalTrials.gov/show/NCT04674254
Metabo-lipidomics of the Ocular Surface for Cataract Surgery	University Hospital, Tours	https://ClinicalTrials.gov/show/NCT05802550
Macular Involvement in Diabetic Retinopathy Evaluated With Swept-Source OCT	University of British Columbia	https://ClinicalTrials.gov/show/NCT03765112
EyeConic: Qualification for Cone-Optogenetics	University Hospital, Basel, Switzerland Institute of Molecular and Clinical Ophthalmology Basel	https://ClinicalTrials.gov/show/NCT05294978
Suprachoroidal Visco-buckling for the Treatment of Rhegmatogenous Retinal Detachment	King's College Hospital NHS Trust Norfolk and Norwich University Trust Foundation St Thomas' Hospital, London University of Sunderland Moorfields Eye Hospital NHS Foundation Trust Mid and South Essex NHS Foundation Trust Sheffield Teaching Hospitals NHS Foundation Trust	https://ClinicalTrials.gov/show/NCT04557527
S.T.O.P. Technology Contact Lenses Versus Dual-focus Contact Lenses for Slowing Down Myopia Progression in Children	nthalmic Pty Ltd Brighten Optix Corporation	https://ClinicalTrials.gov/show/NCT05243836
PMCF Study on EDOF (Isopure) <i>vs</i> Monofocal (Micropure) IOL	Beaver-Visitec International, Inc. targomedGmbH	https://ClinicalTrials.gov/show/NCT04249492
ARTFL LEFFTDS Longitudinal Frontotemporal Lobar Degeneration (ALLFTD)	Mayo Clinic University of California, San Francisco National Institute on Aging (NIA) National Institute of Neurological Disorders and Stroke (NINDS)	https://ClinicalTrials.gov/show/NCT04363684
Systematic Assessment of Laryngo-	Kliniken Beelitz GmbH University Hospital	https://ClinicalTrials.gov/show/NCT04706234

pharyngeal Function in Patients With MSA, PD, and 4 repeat Tauopathies	Muenster Medical University of Warsaw University Hospital Carl Gustav Carus University of Ulm Medical University Innsbruck Hannover Medical School University of Barcelona
--	--

secretome have highlighted its role in facilitating cell survival, proliferation, differentiation and physiological processes [17]. A huge number of secreted growth factors are well known today, including VEGF, SDF-1, TGF- β , IGF-1, fibroblast growth factor (FGF), nerve growth factor-beta (NGF- β), HGF, G-CSF and EGF[18]. With regard to MSC secreted cytokines and chemokines, the most investigated are CCL2, CCL5 and CXCL12 (SDF-1)[19].

One of the most useful aspects of the MSC secretome is the possibility to tailor or modify its composition depending on the desired cell-specific therapeutic effects. This promising possibility depends on MSC tissue sources or on the number of passages, allowing the creation of distinct secretory profiles and exosomal compositions[20]. However, several controversial studies have already been published. It was shown, for example, that the impact of MSCs extracted from adipose tissue was more noticeable on axonal growth than MSCs coming from bone marrow, while cell passaging did not influence the secretome content/activities supporting postnatal neuronal survival and axonal growth[21]. During the last few years, it has been revealed that MSCs are able to modify the microenvironment by releasing EVs, primarily distinct into apoptotic bodies, microvesicles and exosomes[22] (Table 2). The latter subtype consists of a bilayered lipid film of 30-120 nm, originating from convex membranes in late endosomes, determining the production of multi-alveolar bodies[23].

Different and various proteins are typical markers of exosomes, such as tetraspanin (CD9, CD63, CD81), annexin, heat shock proteins, caveolin and clarin as well as protein characteristics of source cells[24]. Furthermore, exosomes present specific lipids, comprising lipid raft portions, ceramides, sphingomyelin, cholesterol, GM1 ganglioside and phosphatidylserine. Additionally, they can contain nucleic acids, mRNA and ncRNA[25]. MSC-derived exosome biosynthesis and secretion are complex pathways that differ in microenvironmental stimuli, like inflammation or hypoxia[26]. The mTOR and Wnt pathways seem to play a pivotal role in exosome release[27]. Interestingly, recent studies have shown that MSC-derived exosomes may be involved in antigen presentation and immunologic response, coagulation, angiogenesis and apoptosis, as confirmed by the expression of antigens such as CD9, CD44 and CD89 on their surface [28]. Thus, the secretome obtained from the culture of MSCs would appear to promote tissue repair and modulate immune response *in vitro* and *in vivo*, showing a translational impact on regenerative medicine[29]. The use of CM could present diverse advantages if compared to the original MSC implantation, such as: (1) Removal of the inherent risks of cell transplantation; (2) Simpler storage, transport and conservation requirements; and (3) Possible application as a ready-to-go biologic product[30].

Secretome preconditioning modulated by the MSC cultural microenvironment

In recent years, it has been shown how preconditioning approaches for improving paracrine secretion, such as hypoxia, biochemical stimuli and 3D microenvironment, can increase the viability, proliferation and paracrine features of MSCs, thus expanding the therapeutic potential of these cells and their derived products[31]. In detail, dynamic culture conditions, such as 3D aggregate culture and fluid flow, could noticeably impact cellular behavior[32]. Boosted levels of growth factors and cytokines were detected in 3D MSC cultures grown on rotatory orbital or shaking platforms, in stirred systems, such as stirred tank reactors or spinner flasks, and in microgravity bioreactors[33]. Nevertheless, little is still known about the dynamic culture conditions and procedures for 3D aggregate MSC cultures as a scalable and reproducible plan for secretome production. However, the possibility of culturing cells under 3D conditions in a way to better mimic the *in vivo* environment has emerged[34].

A dynamic cross-talk between the cells could permit them to constantly modify their secretome following received stimuli, generating a microenvironment able to promote secretome enrichment for specific applications. Additionally, enhancing the manufacturing process allows MSC cell populations to be obtained that can be cryopreserved for clinical applications to expand clinical efficacy[35]. Recently, the use of matrix-conjugated hydrogel cell culture materials normalized a culture of induced pluripotent stem cell-derived MSCs (iPSC-MSCs), leading to a well-defined secretory profile able to promote enhanced neovascularization both *in vitro* and *in vivo*[36]. Using such innovative biomaterials, it was possible to stimulate reproducible secretion of proangiogenic and immunomodulatory cytokines from iPSC-MSCs that improved tubulogenesis of endothelial cells in Geltrex and neovascularization in chick chorioallantoic membranes [37].

Treatment with both IFN- γ and TNF- α permitted optimization of the MSC secretome. Recently, a unique supernatant of MSCs from human umbilical cord-derived MSCs, pretreated with TNF- α , was discovered to be more powerful in promoting macrophage migration, M2 polarization and phagocytosis, due to the induced high levels of CCL2 and IL-6 [38]. Another way to overcome the expansion limitation of MSCs is to work with MSCs derived from human-induced pluripotent MSCs, serving as a reproducible and sustainable cell source. In a way similar to hBMMSCs, human-induced pluripotent MSCs can release EVs with *in vitro* immunomodulatory properties, with an increased expression of well-known immunomodulatory genes, such as *HLA-DRA*, *IDO1* and *CXCL8/IL8* and at least another 100 regulated by NFkB signaling, known to play a pivotal role in immune response[39].

Interestingly, it has recently been shown that hypoxic preconditioning appears to induce the ASC secretome to release a secretome with enhanced anti-apoptotic effects by promoting the autophagic process of ASCs[40]. Furthermore, the specific content of EVs can be modulated by hypoxia, with their source cell responding by triggering HIF at low oxygen levels. The pleiotropic effects of HIF regulate the expression of many genes involved in pathways such as inflammation, angiogenesis, migration, differentiation, metabolism, proliferation and apoptosis. Expression of these genes is reflected in the interior of secreted EVs, which showed a greater regenerative ability than those achieved under normal oxygen conditions[41]. Moreover, the preconditioning of MSCs in an oxidative stress (OS) environment provides the release of

Table 2 Secretome and extracellular vesicle features

Type	Dimension	Origin	Collection	Content
Medium-size EVs (microvesicles)	200-1000 nm	Plasma membrane shedding	Filtration, ultracentrifugation, chromatography, precipitation, immunoaffinity	Similar to exosomes + cytosolic/plasma membrane/post-translational modified proteins
Small-size EVs (exosomes)	Up to 200 nm	Multivesicular bodies pathway	Filtration, ultracentrifugation, chromatography, precipitation, immunoaffinity	Receptors, transcription factors, enzymes, proteins, lipids, nucleic acids (DNA, mRNA and miRNA)
Soluble factors	Up to 5 nm	Protein synthesis	Protein extraction methods	Proteins, growth factors, chemokines, cytokines, enzymes
Secretome	From protein size to 1000 nm	Cell secretion/shedding	Cell culture media (concentration is possible)	The combination of the other components

EV: Extracellular vesicle; miRNA: MicroRNA.

many proteins, growth factors, cytokines and exosomes that could increase the antioxidant ability of MSCs against OS, enforcing the secretome as an encouraging, novel, cell-free tissue regeneration approach[42].

Characterization of MSC secretome EVs

A detailed analysis of the secretome structure might contribute to the improvement of secretome application for regenerative purposes and allow the discovery of novel biomarkers circulating in patient blood, improving pathology diagnosis and discovering new therapeutics targets[43]. As already anticipated, secretome fractions consist of lipids, proteins and non-coding RNAs able to impact the physiology of target cells. MSC-EVs were shown to contain a significant number of microRNAs (miRNAs), such as miR-210, miR-200b-3p and miR-4732-3p, involved in improving myocardial function[44]. BM-MSC-EVs, PD-L1-MSC-EVs and human umbilical cord-derived MSC-EVs also exhibited a healing role in autoimmune conditions[45]. miR-146a and miR-27a/b, upregulated in ASC-derived EVs, were able to induce neoangiogenesis pathways, while miR-122-5p, miR-27a, miR-206 and lncRNA MALAT1 played a relevant role in osteogenic regenerative processes[46].

Other studies identified specific factors from the secretome released by tumor cells that might be actively involved in cancer progression, thus representing optimal biomarkers[47]. Additionally, a customized secretome could be rich in proapoptotic factors that are helpful against cancer or higher levels of proangiogenic and pro-osteogenic factors suitable for regenerative applications. An interesting case is represented by human fetal MSCs, producing a secretome rich in anti-apoptotic factors as well as proangiogenic and antiangiogenic and osteogenic differentiative proteins[48]. On the contrary, the multipotent fetal dermal cell secretome is enriched in upregulated proteins involved in wound healing processes, angiogenesis and cellular metabolism[49]. Such data underlined that the fetal MSC secretome could be more beneficial for regenerative purposes if compared to the adult MSC secretome. In agreement with reports on the secretome derived from 3D cultured cells, fetal cells cultured under 3D conditions might further improve the therapeutic abilities of their secretome[50].

Unique immunomodulatory properties emerged for amniotic MSCs. Their secretome was able to reduce the polarization of T cells toward inflammatory helper T cell subgroups, inducing regulatory T cells, to decrease the proliferation of activated peripheral blood mononuclear cells, to affect monocyte polarization to antigen-presenting cells stimulating the synthesis of anti-inflammatory macrophage (M2) markers and to reduce the activation of B lymphocytes into plasma cells[51]. The most intriguing aspect of secretome EV fractions is the functional mitochondria release from human mesenchymal stromal cells[52]. Recent studies in non-orthopedic tissues proposed that MSCs can rescue damaged cells by donating mitochondria, repairing mitochondrial activity in target cells, preserving cell viability and stimulating tissue repair. To obtain this goal, MSCs might be able to package mitochondria for export into EVs, and these "mitoEVs" could provide a delivery approach for cell-free mitochondria-targeted therapy[53].

Therapeutic role of the secretome in central nervous system pathologies

The MSC secretome is a significant element of the paracrine and autocrine cell signaling mechanism, playing a crucial role in the regulation of many physiological and pathological processes. In particular, its effects on immunomodulation, neuronal survival and regeneration, due to the action of soluble and vesicular factors, are pivotal in reducing or even arresting neuronal disease evolution and in promoting repair[54]. Thus, the various MSC secreted factors and vesicles seem to be an effective tool for the protection and survival of neuronal and glial cells[55]. Traumatic brain injury (TBI) is determined by external mechanical forces able to cause physical, cognitive and emotional impairments[56]. In this case, the MSC-derived secretome may be used to control the secondary injury mechanisms of TBI, modulate the abnormal inflammatory cascade, reduce proinflammatory cytokines and stimulate neural stem cell proliferation and differentiation [57]. Moreover, EVs released by MSCs reduced neuroinflammation and supported neurogenesis and angiogenesis, rescuing spatial learning and motor damage in TBI animal models[58].

Spinal cord injury is characterized by long-term functional deficits following the loss of neurons and glial cells, inflammation and demyelination[59]. The paracrine factors secreted into the lesion site by MSCs, such as HGF, BDNF and NGF could promote immunomodulation, glial scar reduction, axonal regeneration and neurite outgrowth. Additionally, the

ASC-derived secretome reduced the production of TNF- α by M1 macrophages while it improved TGF- β 1 and IL-10 production by M2 macrophages[60]. MSC exosomes could stimulate anti-inflammatory and proangiogenic effects and axonal regeneration and suppress glial scar formation and cell apoptosis, reducing lesion size and improving functional recovery after traumatic spinal cord injury[61].

Ischemic stroke is a cerebrovascular pathology induced by blood vessel occlusion or injury, leading to a blood supply defect, determining focal tissue loss and endothelial and neuronal cell death[62]. The use of MSC secreted factors such as IGF-1 and BDNF could induce neuroprotection by impeding neuronal damage and tissue loss and reduce astrocyte injury by GFAP downregulation[63].

Parkinson's disease (PD) is a neurodegenerative pathology characterized by the progressive degeneration of dopaminergic neurons. In PD, it has already been seen that the addition of the MSC secretome can promote a partial reversion of PD histological impairments and gains in animal motor ability by the secretion of immunomodulatory, anti-inflammatory, neurogenic, neurodevelopmental, neurorescuing or antiapoptotic factors[64]. Recent evidence highlighted the particular ability of the MSC secretome to reduce one of the hallmarks of the disease, the alpha-synuclein aggregates, through an MMP-2-based mechanism[65].

MAIN FEATURES OF RETINAL CELLS

Even if peripherally localized, the retina represents an important part of the central nervous system. Though it presents the same types of functional elements and neurotransmitters sited in other portions of the central nervous system, the retina includes five classes of neurons: photoreceptors (rods and cones); bipolar cells; amacrine cells; horizontal cells; and ganglion cells. Light absorption by the photopigment in the outer segment of rods and cones, the two photoreceptors, starts a cascade of events that changes the receptor membrane potential and the quantity of neurotransmitter released by the rod and cone synapses onto the adjacent bipolar cells, in the outer plexiform layer. Then, in the inner plexiform layer, the short axonal processes of bipolar cells realize a synapse with the dendritic processes of ganglion cells whose axons form the optic nerve. Horizontal and amacrine cells, instead, present their cell bodies within the inner nuclear layer and are mainly involved in lateral interactions between already described retinal cells, impacting the sensitivity of the visual system to light contrast over a wide range of intensities. The amacrine cell processes, which ramify laterally in the inner plexiform layer, are postsynaptic to bipolar cells and presynaptic to ganglion cells, while the processes of horizontal cells instead extend in the outer plexiform layer. The existence of different subgroups of amacrine cells that play a distinct role within visual pathways is relevant. Furthermore, the neural retina and the choroid are connected by a monolayer of cells constituting the retinal pigmented epithelium (RPE). Light absorption, epithelial transport, spatial buffering of ions, visual cycle regulation and phagocytosis of rod and cone outer segment membranes represent the main functions exerted by the RPE[66].

RPE cell secretome and retinal diseases

The RPE is characterized by a polarized nature, with molecules expressed by these cells either secreted to the apical or basolateral membrane by the Na⁺/K⁺-ATPase associated channel or by the anion channel, respectively. These products, mainly growth, anti/proangiogenic and neurotrophic factors, are critical for the correct functioning of the neuroretina and choroid. Among them, the most characterized are VEGF[67,68], TGF- β [67,69], PEDF[70], MMPs[71], NGF[72], FGF-1, FGF-2, and FGF-5[73], IGF-1[74], BDNF[75], PDGF[76], CTGF[77], LEDGF[78], interleukins[79], tissue inhibitor of matrix metalloproteases[80], PIGF[81], angiogenin[82], EPO[83], somatostatin[84] and apolipoprotein A1[85]. These factors could also play a fundamental role in the etiology of several retinal diseases such as diabetic retinopathy (DR), age-related macular degeneration (AMD) and retinopathy of prematurity[86].

Deep proteomic analyses of RPE cells cultivated in these pathological condition microenvironments suggested that previously described molecules could be involved in membrane and cytoskeleton dynamics, mitochondrial trafficking, protection/induction of cellular stress, apoptosis, differential modulation of multidrug resistance-associated proteins and in other metabolic events already during the first stages of the diseases[87]. In physiological conditions, RPE cells release EVs characterized by proteins associated with biological pathways involved in AMD etiology, including drusen composition. Recently, it was shown that drusen-associated proteins are secreted as cargo of EVs produced by RPE cells in a polarized apical to basal way. Remarkably, drusen-associated proteins revealed differential regulation of polarized secretion in homeostatic conditions and in response to AMD stressors[88]. Findings suggested that a finely-tuned mechanism is pivotal to regulate directional sorting and secretion of drusen-associated proteins *via* RPE secretome EVs, supporting the influential role of vesicles as a strategic source of drusen proteins and critical elements to drusen development[89].

OS changed the release of several factors implicated in neovascularization and AMD, stimulating a proangiogenic microenvironment by increasing the secretion of VEGF, PTN and CRYAB and reducing the production of anti-PEDF and CFH. Apical secretion was influenced more than basolateral for PEDF, CRYAB and CFH, while directional secretion was impacted more for VEGF, which may have implications for choroidal neovascularization[90]. VEGF-A is an important proangiogenic factor released by different retinal cytotypes (endothelial cells, Müller cells, ganglion cells and pericytes) but primarily by the basolateral side of RPE in homeostatic conditions, shifting to apical during pathological conditions [91].

In particular, VEGF overexpression was highlighted in hypoxic and hyperglycemic conditions, by both *in vitro* and *in vivo* studies, demonstrating that in pathological conditions VEGF causes alteration of tight junction proteins and transepithelial resistance[92]. It has been confirmed that VEGF R2, placed in the apical side of RPE cells, can induce

disruption of the RPE barrier by promoting VEGF signaling[93]. Considered together, such findings led to the development of anti-VEGF therapies to treat retinal neovascularization in patients with DR and other related diseases. However, many associated complications are still present, such as repeated injection requirements, increased ocular pressure, macular edema, subconjunctival hemorrhage, pain, uveitis and the compromised viability of RPE, photoreceptors, choriocapillaris and Müller glia[94].

One of the most interesting relates to splice variants of VEGF, such as VEGF165b, expressed by RPE cells. It can appear to act as a powerful antiangiogenic isoform of VEGF with significant results in treating induced choroidal neovascularization and was decreased in DR[95]. Nevertheless, while inner retinal barrier and Müller cell association with VEGF is well known, the outer retinal barrier properties of RPE in relation to VEGF in diabetes and other ocular neovascularization-related diseases should be better investigated. One of the most significant glycoproteins of the RPE secretome is PEDF, a serine protease inhibitor with neuroprotective, antiangiogenic and anti-inflammatory features. In homeostatic conditions, PEDF is apically released from the RPE and preserves retinal and choriocapillaris integrity by preventing endothelial cell proliferation[96]. It was seen that PEDF was downregulated in human hyperglycemic RPE cells as well as in patients affected by proliferative DR (PDR), diabetic macular edema (DME), retinopathy of prematurity, retinitis pigmentosa and leber congenital amaurosis[97]. Thus, PEDF is primarily considered for its therapeutic potential, showing positive effects in photoreceptor survival, morphology and function while reducing vascular permeability in correlation with reduced levels of angiogenic factors (VEGF, VEGFR-2), cytokines and chemokines[98]. Additionally, recent animal studies have proven that in an oxygen-induced retinopathy model and in a rat model of choroidal neovascularization, PEDF upregulation blocked retinal neovascularization and inflammation[99].

Another prosurvival cytokine able to stimulate fibroblast chemotaxis/proliferation and preserve pericyte viability and physiological vascularization of the retina is PDGF. Similar to VEGF, it can promote pathologic neovascularization in PDR and DR and in a hypoxia-regulated microenvironment[100]. PDGF receptor activation suggests an autocrine mechanism in epiretinal membrane development and retinal wound repair[101]. Today, one of the most promising research fields is understanding the cross-talk of PDGF with other signaling pathways in order to identify the best molecular targets for combinatorial therapies. This idea arose from several animal studies that established that antagonism to PDGF-BB (a homodimeric form of the PDGF family), together with anti-VEGF, enhanced the arrest of retinal neovascularization[102].

An important cofactor of VEGF is PIGF. It can alter retinal fibrovascular integrity and RPE permeability by interaction with VEGF and activation of Akt and HIF-1 pathways[103]. Thus, it was found at high levels in AMD and PDR patients [104]. The use of anti-PIGF monoclonal antibody in different animal models revealed reduced inflammation and vascular leakage with no adverse effects in retinal ganglion cell (RGC) viability[105]. However, novel strategies that avoid the weaknesses observed in repeated intraocular injections should consider PIGF as a valid therapeutic target. RPE cells cultured in high glucose medium also showed an elevated expression of CTGF, one of the main fibrogenic factors involved in fibroblast proliferation and extracellular matrix synthesis, which could control the microenvironment around the distal retinal/RPE/Bruch's membrane complex and protect against neurodegenerative diseases[106]. Increased retinal CTGF levels might play an essential role in DR, probably by reducing VEGF levels[107]. Thus, the combined use of anti-CTGF and anti-VEGF in treating complications of DR could exert more beneficial effects than a monotherapy drug[108].

CTGF is corroborated in its activity by the more well-known FGF, which plays a crucial role in stimulating vascularization, angiogenesis and cell survival and acting as autocrine factors. FGF1, FGF2 and FGF5 are principally released in the RPE, reaching their highest levels in non-proliferative retinopathy, PDR with active proliferative retinopathy and diabetic conditions, respectively[109]. Recently, targeting of retinal FGFs exhibited worthy results in improving visual acuity of DME and exudative AMD patients, even if further studies are mandatory to determine long-term effects[110]. The secretome produced by human RPE cells also contained IGF-1 and IGF-2, natural proteins promoting growth and insulin-like metabolic effects, together with their receptor (IGF-R) and binding protein (IGFBP-2)[111]. Both growth factors seem to play a pivotal role for RPE autocrine/paracrine-mediated modulation of proliferation[112]. Recent evidence showed that another IGFBP family member, IGFBP-3, was able to reduce DR by considerably decreasing TNF- α levels and proapoptotic markers[113].

Among secreted factors, TGF- β represents one of the main elements that can modulate main cellular physiological processes, like growth, differentiation, proliferation and apoptosis[114]. However, there is scant information on its efficacy and potential mechanisms in relation to retinal homeostasis or pathology. In detail, comparable secretion levels of TGF- β from polarized RPE, differentiated from human embryonic stem cells and human RPE, promoting retinal homeostasis and sustaining the potential of human embryonic stem cell-RPE in replacement therapies, have recently been highlighted[115]. Human stem cell-derived RPE treated with reactive oxygen species for 1 wk or 3 wk released more than 1000 proteins, many of which showed relevant changes due to induced stress.

In particular, secreted APOE and TGF- β were decreased 4-fold, and urotensin-II, one of the most effective vasoconstrictors, doubled, similar to BMP1[116]. The glycoprotein EPO represents one of the most promising molecules found in the RPE secretome. It acts as an erythropoiesis regulator with different additive features such as vessel integrity, recruitment of endothelial progenitor cells, neuroprotection and antioxidative properties[117]. High levels of EPO were recently found in DR, PDR and DME patients[118]. Especially in hyperglycemic conditions, EPO seems to protect the RPE barrier, reducing retinal vasculogenesis, downregulating VEGF and VEGFR expression and protecting tight junctions by increasing the flow of Ca²⁺ ions in blood-brain barrier animal models[119]. However, the administration of EPO in the late stage of a hypoxia-induced murine retinopathy model worsened retinal neovascularization, suggesting that EPO might play a protective role in early DR and a pathologic one in late DR[120]. This dual nature of EPO could be related to its action mechanism, whose first step is its hypoxia-modulated binding to cell surface receptor EPOR. Thus, it can be predicted that in the first stages of DR, EPO exerts neuroprotective functions, while in the advanced stage of DR EPO acts as a neovasculogenesis inducing molecule that is regulated by hypoxia[121].

MMPs, apically secreted by RPE, are calcium-dependent endopeptidases involved in angiogenesis and are fundamental for ocular extracellular matrix and photoreceptor outer segment homeostasis[122]. Recently, it has been seen that basolateral secretion of MMP is related to AMD, and increased levels of MMP-2 and MMP-9 were also observed in the Bruch's membrane of AMD and DR patient eyes[123]. Thus, inhibitors of both cited MMPs might also exert an advantageous role by blocking capillary cell apoptosis, growth of vessels and reduce inflammatory-mediated permeability[124].

In addition to angiogenic and antiangiogenic factors, numerous inflammatory chemokines and cytokines were elevated in retinal diseases, such as PD and PDR. Among them, the most investigated were MCP-1, IL-6 and IL-8. It was seen that MCP-1 and IL-8 secretion levels are directly correlated to blood glucose levels, suggesting a crucial role in altered blood retinal barrier (BRB) activities of DR affected patients[125]. MCP-1 carries out chemoattractant activity for monocytes and lymphocytes to promote endothelial proliferation and may limit the impairment of neurosensory retina[126]. IL-6 and IL-8, were overexpressed in cultured RPE cells stimulated with IL-1b or TNF- β , suggesting that polarized release of growth factors/cytokines is favored in retinal diseases[127].

Additionally, it is interesting to cite the recently discussed role of somatostatin as a neuromodulator of retinal homeostasis, as hypothesized by its downregulation in the RPE of diabetic eyes[128]. Finally, several substrates of the serine protease HTRA1 were found in the RPE secretome, proposing a link between it and complement modulation and amyloid deposition in AMD etiopathogenesis. In detail, a cleavage of fibromodulin (90%), CLU (50%) and vitronectin (54%) involved in regulation of the complement pathway was seen, along with a cleavage of 2-macroglobulin (55%) and ADAM9 (54%) related to amyloid deposition as well as some cell surface protein cleavages including talin-1 (21%), fascin (40%) and chloride intracellular channel protein 1 (51%)[129].

Müller glia cell secretome and retinal diseases

Regarding the RPE, Müller cells can modulate trophic secretion depending on the healthy or pathological status of the retina[130]. The Müller cell physiological secretome mainly contains molecules that are crucial to increase BRB tightness, like thrombospondin-1 and PEDF[131]. In pathological circumstances, factor synthesis and secretion both shift towards an inflammatory environment. Under hyperglycemic conditions, IL-1b release by Müller cells is increased, leading to vascular impairment and cell death *via* a paracrine mechanism. Thus, by inhibiting IL-1b or knocking down its receptor, it was possible to decrease inflammation and photoreceptor/retinal vessel disruption in murine models, exerting a possible therapeutic role for ocular dystrophies related to chemokine expression and/or diabetes[132]. Furthermore, the proinflammatory IL-6 and TNF- α can be secreted by Müller cells, determining a possible promotion of both vascular dysfunction and angiogenesis, even if IL-6 may exert protective effects toward photoreceptor cells[133].

Stimulation of porcine and human Müller cells with IL-4, IL-6, IL-10, VEGF, INF- γ , TGF- β 1, TGF- β 2, TGF- β 3 and TNF- α resulted in a primarily proinflammatory phenotype with release of cytokines and factors of the complement system[134]. Additionally, Müller cells expressed proteins linked to biosynthesis and maturation of phagosomes. These findings underline the relevance of Müller cell signaling in chronic retinal inflammation[135]. Additionally, under hyperglycemia and hypoxic conditions, Müller cells shift PEDF secretion to VEGF, contributing to ocular vascular diseases[136]. Therefore, inhibition or knockdown of Müller cell-derived VEGF could reduce ischemia-induced impairment of the BRB, prevent ischemia-induced retinal neovascularization and decrease vascular leakage[137].

Recent evidence highlighted that in the diabetic retina expression of VEGF could be regulated by increasing the activity of the receptor for retinoic acid alpha, which also stimulates the expression of glial cell line-derived neurotrophic factor, with a final significant decrease of vascular leakage[138]. Nevertheless, in recent years, the neuroprotective effects of VEGFR-2 in Müller glia have also been described, suggesting its significance for cell survival and consequential viability of neuronal cells in the diabetic retina[139]. Interestingly, the secretome of Müller cells also contains increased levels of MMP-2 and MMP-9 in patients with PDR and AMD, respectively[140]. It was proposed that the stabilization of HIF-1a could raise the level of VEGF, inducing MMP-2 expression in neighboring endothelial cells, with consequent retinal neovascularization[141]. As MMPs regulate crucial cellular pathways through angiogenesis and apoptosis, their targeting could represent an important therapeutic strategy for ocular diseases. Moreover, new evidence has proven that Müller glia release neurotrophic factors, such as CLU, osteopontin and basigin, that support RGC survival. The latter two significantly enhance RGC survival *in vitro*, suggesting that the survival-promoting activity of the Müller cell secretome is multifactorial[142].

Recently, it was shown that human iPSC-derived multinucleated giant cells (hiMGCs) could represent an alternative to primary MGCs in understanding glial cell involvement in retinal disorders, including DR. Under culture with palmitate, a major free fatty acid with elevated plasma levels in diabetic patients, hiMGCs and primary MGCs expressed low transcript levels of *AQP4*, *RLPB1*, *SLC1A3*, *KCNJ1* and *KCJN10*. Furthermore, the analysis of the palmitate-treated hiMGC secretome evidenced an upregulation of proangiogenic factors powerfully related to DR, including ANG2, endoglin, IL-1b, CXCL8, MMP-9, PDGF-AA and VEGF[143]. One of the most interesting pieces of evidence regarding the Müller cell secretome was linked to the production of different EVs from endfeet and microvilli of retinal Müller cells in adult mice. In particular, VAMP5 was identified as a Müller cell-specific snap receptor member that is part of EVs and responsive to ischemia, with relevant changes between the secretomes of Müller cells and neurons *in vitro*[144].

Other glial cell secretome and retinal diseases

Undifferentiated rat RGC line RGC-5 can secrete numerous protein markers of RGCs, even if they are unable to react to glutamate or N-methyl-D aspartate. Furthermore, it has recently been highlighted that human nonpigmented ciliary epithelial (HNPE) cells could release several neuroproteins located in the aqueous humor, many of which can influence the activity of neuronal cells. Recent works identified about 130 unique proteins from the HNPE cell-conditioned SF-medium, most of which are involved in cell differentiation. These results led to the hypothesis that a differentiation

system of HNPE cell-conditioned SF-medium with RGC-5 cells can promote a differentiated phenotype in RGC-5 cells, functionally close to primary cultures of rat RGCs[145].

The secretome of retinoblastoma, the solid malignancy of the developing retina, is immunosuppressive and induces a protumoral phenotype. This conclusion was the result of complex analyses that identified the cytokine extracellular matrix metalloproteinase inducer and macrophage migration inhibitory factor, both characterized by detected immunosuppressive activity and secreted at high levels in retinoblastoma primary cell cultures. In addition, macrophages derived from peripheral blood mononuclear cells increased the expression of M2-like polarization markers following exposure to retinoblastoma-conditioned medium or recombinant migration inhibitory factor[146].

CROSS-LINK BETWEEN RETINAL AND MESENCHYMAL SECRETOMES: DIFFERENT MSC-DERIVED SECRETOMES AS INNOVATIVE APPROACHES FOR RETINAL DISEASES

The MSC secretome is currently studied extensively for the treatment of several retinal diseases. Its therapeutic potential lies in its richness of immunomodulatory, antiangiogenic and neurotrophic factors, preventing retinal degeneration and improving retinal morphology and function. Additionally, exosomes secreted by MSCs showed anti-inflammatory and antiapoptotic effects (Figure 1 and Table 3). Based on MSC origins and their particular secreted factors, several promising preclinical and clinical studies were initiated to explore the potential advantages of MSC secretome for the treatment of retinal diseases.

Novel evidence on the role of MSC secreted factors in retinal disease etiopathogenetic pathways

Novel evidence showed that MSC conditioned media inhibits abnormal neovascularization and decreases vaso-obliteration (promoting revascularization) in retinopathies by restoring neuronal Sema3E levels, which reduce pathological concentrations of IL-17A (and associated proinflammatory factors, such as IL-1b) in myeloid cells[147]. Among MSC released factors, PDGF secretion may play a crucial role in MSC-mediated RGC neuroprotection. These results were obtained from the arrest of PDGF signaling by small molecule PDGF inhibitors, neutralizing antibody or downstream phosphatidylinositol 3 kinase, which blocked RGC neuroprotection conferred by MSC co-culture. Furthermore, intravitreal injection of PDGF led to relevant optic nerve neuroprotection *in vivo* after experimental induction of high intraocular pressure[148]. Application of conditioned media obtained from MSCs protected against A β 1-42 oligomer-induced retinal pathology in RGCs of both rat and ARPE-19 cells, due to proteins associated with SIRT1/pAKT/pGSK3 β / β -catenin, tight junction proteins and the apoptosis pathway[149]. Furthermore, in recent years, the administration of EVs in models of neurological disorders has highlighted a relevant improvement of neurological dysfunction. In particular, miRNAs from MSC-EVs, as one of the central mediators that control various genes and decrease neuropathological change, have been identified in various neurological pathologies[150].

Bone marrow MSC-derived secretome regulates retinal cell neuroprotection

The BMMSC secretome protects retinal morphology, regulates autophagy-, proapoptotic- and proneuroptotic-related gene and protein expression and promotes the activation of antioxidant machinery, exerting a neuroprotective ability during retinal degeneration[151]. A recent expression analysis of about 1000 proteins exhibited high levels of paracrine factors secreted by hBMMSCs that might be fundamental in the neuroprotective effect of the stem cell secretome over *in vitro* retinal degeneration. These results support the hypothesis that the paracrine effect of hBMMSCs may slow photoreceptor death and be a therapeutic possibility in retinal photoreceptor degenerative diseases[152]. Additionally, rat BMMSCs cultured with the secretome from neonatal rat retinal cells were able to differentiate into RGC-like cells, exhibiting protein expression patterns similar to those of isolated RGCs such as Map2, nestin and Thy1.1[153].

Adipose MSC-derived secretome regulates retinal cell regeneration

Recent evidence showed an important therapeutic effect of hADSCs and its secretome on an *in vivo* model of sodium iodate retinal neurodegeneration. The studies highlighted that the hADSC secretome effects were particularly striking, especially in terms of photoreceptor regeneration and retinal function, as underlined by increased expression of retinal regeneration markers such as Pax6, Chx10, S-Opsin (Opn1sw), Nrl, Crx and GFAP[154]. Oxidatively stressed ARPE-19 cells treated with adipose MSC conditioned media and/or combined with nicotinamide, vasoactive intestinal peptide or both factors showed an improved recovery from the damaged status. Additionally, the same treatment could determine better protection of the neuroretinal architecture, mainly rods and cones, and a lower degree of glial cell activation[155].

The preclinical efficacy of adipose-derived stem cell concentrated conditioned medium (ASC-CCM) was recently tested in repetitive ocular blast injury mice, highlighting a significant rescue from retinal injury and a significant restoration of visual function, also associated with a significant reduction of neuroinflammation markers, retinal GFAP and OS. Furthermore, *in vitro*, oxidatively stressed Müller cells pre-incubated with ASC-CCM exhibited normalized levels of GFAP, viability and catalase activity[156]. Intravitreal injection of ASC-CCM was safe and efficient against the visual impairments of mild TBI. Blast mice treated with ASC-CCM exhibited improved vision at 5 mo but minimal effects at 10 mo, associated with alterations of GFAP and proinflammatory gene expression in retina. Thus, the unchanged glial response and the risk of retinal injury with live cells suggested that ASC-CCM might have better safety and efficacy than live cells for visual dysfunction therapy[157].

Table 3 Key molecules of the mesenchymal secretome derived from different cell types in the treatment and regeneration of retinal cells

MSC	MSC secretome	Müller glia cell secretome	RPE cell secretome	Other retinal cell secretome
IFN- γ [8]	IFN- γ [8]	TSP-1[131]	VEGF[68]	TSP-1[131]
VEGF[10]	VEGF[10]	PEDF[70]	TGF- β [69]	PEDF[70]
PIGF[81]	PIGF[81]	IL-1b[11]	PEDF[70]	IL-1b[11]
SDF-1[10]	SDF-1[10]	IL-1R[143]	MMPs[71]	IL-1R[143]
ANG2[10]	ANG2[10]	TNF- α [11]	NGF[72]	IL-6[11]
G-CSF[10]	G-CSF[10]	IL-4[134]	FGF-1[73]	TNF- α [11]
SCF[10]	SCF[10]	IL-6[134]	FGF-2[73]	IL-4[134]
PDGF[10]	PDGF[10]	IL-10[134]	FGF-5[73]	IL-10[134]
EGF[10]	EGF[10]	VEGF[10]	IGF-1[74]	VEGF[10]
HGF[10]	HGF[10]	IFN- γ [8]	BDNF[75]	IFN- γ [8]
IGF-1[10]	IGF-1[10]	TGF- β 1[34]	PDGF[76]	TGF- β 1[34]
TNF- α [11]	TNF- α [11]	TGF- β 2[34]	CTGF[77]	TGF- β 2[34]
IL-1b[11]	IL-1b[11]	TGF- β 3[34]	LEDGF[78]	TGF- β 3[34]
IL-2[11]	IL-2[11]	MMPs[71]	IL-1b[79]	MMPs[71]
IL-3[11]	IL-3[11]	CLU[121]	IL-2[79]	CLU[121]
IL-6[11]	IL-6[11]	SPP1[142]	IL-3[79]	SPP1[142]
IL-8[11]	IL-8[11]	BSG[142]	IL-6[79]	BSG[142]
CCL5[11]	CCL5 ¹¹	AQP4[143]	IL-8[79]	AQP4[143]
CCL2[11]	CCL2[11]	RLBP1[143]	TIMP[80]	RLBP1[143]
	TGF- β [18]	SLC1A3[143]	PIGF[81]	SLC1A3[143]
	FGF[18]	KCNJ1[143]	ANGIOGENIN[82]	KCNJ1[143]
	NGF- β [18]	KCNJ10[143]	EPO[83]	KCNJ10[143]
	CXCL12[19]	ANG2[143]	SOMATOSTATIN[84]	ANG2[143]
	miR-210[44]	ENDOGLIN[143]	APOA1[85]	ENDOGLIN[143]
	miR-200b-3p[44]	CXCL8[39]	APOE[116]	CXCL8[143]
	miR146a[46]	PDGF[10]	TGF- β [69]	PDGF[76]
	miR27a/b[46]	VAMP5[144]	BMP1[116]	VAMP5[144]
	miR-122-5p[46]		HTRA1[129]	TGF- β [69]
	miR-206[46]			NGF[72]
	MALAT[46] (lncRNA)			FGF-1[73]
				FGF-2[73]
				FGF-5[73]
				IGF-1[74]
				BDNF[75]
				PDGF[76]
				CTGF[77]
				LEDGF[78]
				IL-2[79]
				IL-3[79]
				IL-8[79]

secretome on retinal degeneration, by modulating OS, autophagy and programmed cell death. This scenario could be of particular interest especially for MSC secretome-only factors, such as FGF, CXCL12, CCL511, NGF-, described miRNAs and the lncRNA MALAT, whose complementary action might play a functional compensation role towards retinal cell alterations.

However, many challenges still exist, including the specific characterization of secretome released factors to further target therapy to the pathology profile, better manipulation of the retinal secretome or from other cell sources for noteworthy therapeutic effect, improving methods for intraocular administration of secretome factors and developing personalized combinations of trophic factors involved in different pathological pathways (inflammation, reactive oxygen species, angiogenesis, proliferation) to evaluate the collective therapeutic potential. Nevertheless, the possible improvement of new efficient pharmaceutical formulations related to the secretome of MSCs and retinal cells, with the addition of exogenous factors or drugs without the necessity to deliver cells into the eye may represent a novel milestone towards a personalized approach to retinal disease.

FOOTNOTES

Author contributions: Donato L wrote the paper; Donato L, Scimone C, Alibrandi S, Mordà D, Scalinci SZ and Rinaldi C made relevant contributions to conception and design of the article; D'Angelo R and Sidoti A made critical revisions related to important intellectual content of the manuscript.

Conflict-of-interest statement: The authors declare no conflicts of interest for this article.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <https://creativecommons.org/licenses/by-nc/4.0/>

Country/Territory of origin: Italy

ORCID number: Luigi Donato [0000-0002-5292-426X](https://orcid.org/0000-0002-5292-426X); Concetta Scimone [0000-0002-4824-1693](https://orcid.org/0000-0002-4824-1693); Simona Alibrandi [0000-0003-3569-5394](https://orcid.org/0000-0003-3569-5394); Rosalia D'Angelo [0000-0001-6510-3121](https://orcid.org/0000-0001-6510-3121).

S-Editor: Zhang H

L-Editor: Filipodia

P-Editor: Zhang XD

REFERENCES

- Williams T, Salmanian G, Burns M, Maldonado V, Smith E, Porter RM, Song YH, Samsonraj RM. Versatility of mesenchymal stem cell-derived extracellular vesicles in tissue repair and regenerative applications. *Biochimie* 2023; **207**: 33-48 [PMID: [36427681](https://pubmed.ncbi.nlm.nih.gov/36427681/) DOI: [10.1016/j.biochi.2022.11.011](https://doi.org/10.1016/j.biochi.2022.11.011)]
- McLaughlin C, Datta P, Singh YP, Lo A, Horchler S, Elcheva IA, Ozbolat IT, Ravnic DJ, Koduru SV. Mesenchymal Stem Cell-Derived Extracellular Vesicles for Therapeutic Use and in Bioengineering Applications. *Cells* 2022; **11** [PMID: [36359762](https://pubmed.ncbi.nlm.nih.gov/36359762/) DOI: [10.3390/cells11213366](https://doi.org/10.3390/cells11213366)]
- Harrell CR, Volarevic V, Djonov V, Volarevic A. Therapeutic Potential of Exosomes Derived from Adipose Tissue-Sourced Mesenchymal Stem Cells in the Treatment of Neural and Retinal Diseases. *Int J Mol Sci* 2022; **23** [PMID: [35562878](https://pubmed.ncbi.nlm.nih.gov/35562878/) DOI: [10.3390/ijms23094487](https://doi.org/10.3390/ijms23094487)]
- Ganguly A, Swaminathan G, Garcia-Marques F, Regmi S, Yarani R, Primavera R, Chetty S, Bermudez A, Pitteri SJ, Thakor AS. Integrated transcriptome-proteome analyses of human stem cells reveal source-dependent differences in their regenerative signature. *Stem Cell Reports* 2023; **18**: 190-204 [PMID: [36493779](https://pubmed.ncbi.nlm.nih.gov/36493779/) DOI: [10.1016/j.stemcr.2022.11.006](https://doi.org/10.1016/j.stemcr.2022.11.006)]
- Zhang Y, Yang H, He F, Zhu X. Intra-articular injection choice for osteoarthritis: making sense of cell source-an updated systematic review and dual network meta-analysis. *Arthritis Res Ther* 2022; **24**: 260 [PMID: [36443838](https://pubmed.ncbi.nlm.nih.gov/36443838/) DOI: [10.1186/s13075-022-02953-0](https://doi.org/10.1186/s13075-022-02953-0)]
- Garcia GA, Oliveira RG, Dariolli R, Rudge MVC, Barbosa AMP, Floriano JF, Ribeiro-Paes JT. Isolation and characterization of farm pig adipose tissue-derived mesenchymal stromal/stem cells. *Braz J Med Biol Res* 2022; **55**: e12343 [PMID: [36477953](https://pubmed.ncbi.nlm.nih.gov/36477953/) DOI: [10.1590/1414-431X2022e12343](https://doi.org/10.1590/1414-431X2022e12343)]
- Cequier A, Vázquez FJ, Romero A, Vitoria A, Bernad E, García-Martínez M, Gascón I, Barrachina L, Rodellar C. The immunomodulation-immunogenicity balance of equine Mesenchymal Stem Cells (MSCs) is differentially affected by the immune cell response depending on inflammatory licensing and major histocompatibility complex (MHC) compatibility. *Front Vet Sci* 2022; **9**: 957153 [PMID: [36337202](https://pubmed.ncbi.nlm.nih.gov/36337202/) DOI: [10.3389/fvets.2022.957153](https://doi.org/10.3389/fvets.2022.957153)]
- Ding Y, Gong P, Jiang J, Feng C, Li Y, Su X, Bai X, Xu C, Liu C, Yang J, Fang J, Ji X, Chen Y, Li P, Guo L, Shao C, Shi Y. Mesenchymal stem/stromal cells primed by inflammatory cytokines alleviate psoriasis-like inflammation via the TSG-6-neutrophil axis. *Cell Death Dis* 2022; **13**: 996 [PMID: [36433947](https://pubmed.ncbi.nlm.nih.gov/36433947/) DOI: [10.1038/s41419-022-05445-w](https://doi.org/10.1038/s41419-022-05445-w)]
- Upadhyay TK, Trivedi R, Khan F, Pandey P, Sharangi AB, Goel H, Saeed M, Park MN, Kim B. Potential Therapeutic Role of Mesenchymal-Derived Stem Cells as an Alternative Therapy to Combat COVID-19 through Cytokines Storm. *Cells* 2022; **11** [PMID: [36078094](https://pubmed.ncbi.nlm.nih.gov/36078094/) DOI: [10.3390/cells11172686](https://doi.org/10.3390/cells11172686)]
- Jayaraman H, Ghone NV, Rajan RK, Dashora H. The Role of Cytokines in Interactions of Mesenchymal Stem Cells and Breast Cancer Cells. *Curr Stem Cell Res Ther* 2021; **16**: 443-453 [PMID: [33092514](https://pubmed.ncbi.nlm.nih.gov/33092514/) DOI: [10.2174/1574888X15666201022111942](https://doi.org/10.2174/1574888X15666201022111942)]

- 11 **Arabpour M**, Saghazadeh A, Rezaei N. Anti-inflammatory and M2 macrophage polarization-promoting effect of mesenchymal stem cell-derived exosomes. *Int Immunopharmacol* 2021; **97**: 107823 [PMID: 34102486 DOI: 10.1016/j.intimp.2021.107823]
- 12 **Boyle AJ**, McNiece IK, Hare JM. Mesenchymal stem cell therapy for cardiac repair. *Methods Mol Biol* 2010; **660**: 65-84 [PMID: 20680813 DOI: 10.1007/978-1-60761-705-1_5]
- 13 **Dama G**, Du J, Zhu X, Liu Y, Lin J. Bone marrow-derived mesenchymal stem cells: A promising therapeutic option for the treatment of diabetic foot ulcers. *Diabetes Res Clin Pract* 2023; **195**: 110201 [PMID: 36493913 DOI: 10.1016/j.diabres.2022.110201]
- 14 **Costa LA**, Eiro N, Fraile M, Gonzalez LO, Saá J, Garcia-Portabella P, Vega B, Schneider J, Vizoso FJ. Functional heterogeneity of mesenchymal stem cells from natural niches to culture conditions: implications for further clinical uses. *Cell Mol Life Sci* 2021; **78**: 447-467 [PMID: 32699947 DOI: 10.1007/s00018-020-03600-0]
- 15 **Zhang Z**, Alexanian AR. The neural plasticity of early-passage human bone marrow-derived mesenchymal stem cells and their modulation with chromatin-modifying agents. *J Tissue Eng Regen Med* 2014; **8**: 407-413 [PMID: 22674835 DOI: 10.1002/term.1535]
- 16 **L PK**, Kandoi S, Misra R, S V, K R, Verma RS. The mesenchymal stem cell secretome: A new paradigm towards cell-free therapeutic mode in regenerative medicine. *Cytokine Growth Factor Rev* 2019; **46**: 1-9 [PMID: 30954374 DOI: 10.1016/j.cytogfr.2019.04.002]
- 17 **Abdolmohammadi K**, Mahmoudi T, Alimohammadi M, Tahmasebi S, Zavvar M, Hashemi SM. Mesenchymal stem cell-based therapy as a new therapeutic approach for acute inflammation. *Life Sci* 2023; **312**: 121206 [PMID: 36403645 DOI: 10.1016/j.lfs.2022.121206]
- 18 **Nie WB**, Zhang D, Wang LS. Growth Factor Gene-Modified Mesenchymal Stem Cells in Tissue Regeneration. *Drug Des Devel Ther* 2020; **14**: 1241-1256 [PMID: 32273686 DOI: 10.2147/DDDT.S243944]
- 19 **Aboukheyr Es H**, Bigdeli B, Zhand S, Aref AR, Thiery JP, Warkiani ME. Mesenchymal stem cells induce PD-L1 expression through the secretion of CCL5 in breast cancer cells. *J Cell Physiol* 2021; **236**: 3918-3928 [PMID: 33145762 DOI: 10.1002/jcp.30135]
- 20 **Wruck W**, Graffmann N, Spitzhorn LS, Adjaye J. Human Induced Pluripotent Stem Cell-Derived Mesenchymal Stem Cells Acquire Rejuvenation and Reduced Heterogeneity. *Front Cell Dev Biol* 2021; **9**: 717772 [PMID: 34604216 DOI: 10.3389/fcell.2021.717772]
- 21 **Bucan V**, Vaslaitis D, Peck CT, Strauß S, Vogt PM, Radtke C. Effect of Exosomes from Rat Adipose-Derived Mesenchymal Stem Cells on Neurite Outgrowth and Sciatic Nerve Regeneration After Crush Injury. *Mol Neurobiol* 2019; **56**: 1812-1824 [PMID: 29931510 DOI: 10.1007/s12035-018-1172-z]
- 22 **Bogatcheva NV**, Coleman ME. Conditioned Medium of Mesenchymal Stromal Cells: A New Class of Therapeutics. *Biochemistry (Mosc)* 2019; **84**: 1375-1389 [PMID: 31760924 DOI: 10.1134/S0006297919110129]
- 23 **Yu B**, Zhang X, Li X. Exosomes derived from mesenchymal stem cells. *Int J Mol Sci* 2014; **15**: 4142-4157 [PMID: 24608926 DOI: 10.3390/ijms15034142]
- 24 **Wang ZG**, He ZY, Liang S, Yang Q, Cheng P, Chen AM. Comprehensive proteomic analysis of exosomes derived from human bone marrow, adipose tissue, and umbilical cord mesenchymal stem cells. *Stem Cell Res Ther* 2020; **11**: 511 [PMID: 33246507 DOI: 10.1186/s13287-020-02032-8]
- 25 **Xunian Z**, Kalluri R. Biology and therapeutic potential of mesenchymal stem cell-derived exosomes. *Cancer Sci* 2020; **111**: 3100-3110 [PMID: 32639675 DOI: 10.1111/cas.14563]
- 26 **Chen J**, Chen J, Cheng Y, Fu Y, Zhao H, Tang M, Lin N, Shi X, Lei Y, Wang S, Huang L, Wu W, Tan J. Mesenchymal stem cell-derived exosomes protect beta cells against hypoxia-induced apoptosis via miR-21 by alleviating ER stress and inhibiting p38 MAPK phosphorylation. *Stem Cell Res Ther* 2020; **11**: 97 [PMID: 32127037 DOI: 10.1186/s13287-020-01610-0]
- 27 **Gu X**, Li Y, Chen K, Wang X, Wang Z, Lian H, Lin Y, Rong X, Chu M, Lin J, Guo X. Exosomes derived from umbilical cord mesenchymal stem cells alleviate viral myocarditis through activating AMPK/mTOR-mediated autophagy flux pathway. *J Cell Mol Med* 2020; **24**: 7515-7530 [PMID: 32424968 DOI: 10.1111/jcmm.15378]
- 28 **Zhang L**, Jiao G, Ren S, Zhang X, Li C, Wu W, Wang H, Liu H, Zhou H, Chen Y. Exosomes from bone marrow mesenchymal stem cells enhance fracture healing through the promotion of osteogenesis and angiogenesis in a rat model of nonunion. *Stem Cell Res Ther* 2020; **11**: 38 [PMID: 31992369 DOI: 10.1186/s13287-020-1562-9]
- 29 **Sun Y**, Liu G, Zhang K, Cao Q, Liu T, Li J. Mesenchymal stem cells-derived exosomes for drug delivery. *Stem Cell Res Ther* 2021; **12**: 561 [PMID: 34717769 DOI: 10.1186/s13287-021-02629-7]
- 30 **González-González A**, García-Sánchez D, Dotta M, Rodríguez-Rey JC, Pérez-Campo FM. Mesenchymal stem cells secretome: The cornerstone of cell-free regenerative medicine. *World J Stem Cells* 2020; **12**: 1529-1552 [PMID: 33505599 DOI: 10.4252/wjsc.v12.i12.1529]
- 31 **Lin H**, Chen H, Zhao X, Chen Z, Zhang P, Tian Y, Wang Y, Ding T, Wang L, Shen Y. Advances in mesenchymal stem cell conditioned medium-mediated periodontal tissue regeneration. *J Transl Med* 2021; **19**: 456 [PMID: 34736500 DOI: 10.1186/s12967-021-03125-5]
- 32 **Gorgun C**, Africano C, Ciferri MC, Bertola N, Reverberi D, Quarto R, Ravera S, Tasso R. Preconditioned Mesenchymal Stromal Cell-Derived Extracellular Vesicles (EVs) Counteract Inflammation. *Cells* 2022; **11** [PMID: 36429124 DOI: 10.3390/cells11223695]
- 33 **Zhou T**, Rong M, Wang Z, Chu H, Chen C, Zhang J, Tian Z. Conditioned medium derived from 3D tooth germs: A novel cocktail for stem cell priming and early in vivo pulp regeneration. *Cell Prolif* 2021; **54**: e13129 [PMID: 34585454 DOI: 10.1111/cpr.13129]
- 34 **Uz M**, Büyükköz M, Sharma AD, Sakaguchi DS, Altinkaya SA, Mallapragada SK. Gelatin-based 3D conduits for transdifferentiation of mesenchymal stem cells into Schwann cell-like phenotypes. *Acta Biomater* 2017; **53**: 293-306 [PMID: 28213098 DOI: 10.1016/j.actbio.2017.02.018]
- 35 **Isildar B**, Ozkan S, Ercin M, Gezginici-Oktayoglu S, Oncul M, Koyuturk M. 2D and 3D cultured human umbilical cord-derived mesenchymal stem cell-conditioned medium has a dual effect in type 1 diabetes model in rats: immunomodulation and beta-cell regeneration. *Inflamm Regen* 2022; **42**: 55 [PMID: 36451229 DOI: 10.1186/s41232-022-00241-7]
- 36 **Drzeniek NM**, Mazzocchi A, Schlickeiser S, Forsythe SD, Moll G, Geißler S, Reinke P, Gossen M, Gorantla VS, Volk HD, Soker S. Bio-instructive hydrogel expands the paracrine potency of mesenchymal stem cells. *Biofabrication* 2021; **13** [PMID: 34111862 DOI: 10.1088/1758-5090/ac0a32]
- 37 **Romanazzo S**, Kopecky C, Jiang S, Doshi R, Mukund V, Srivastava P, Rnjak-Kovacina J, Kelly K, Kilian KA. Biomaterials directed activation of a cryostable therapeutic secretome in induced pluripotent stem cell derived mesenchymal stromal cells. *J Tissue Eng Regen Med* 2022; **16**: 1008-1018 [PMID: 36017672 DOI: 10.1002/term.3347]
- 38 **Liu C**, Xu Y, Lu Y, Du P, Li X, Wang C, Guo P, Diao L, Lu G. Mesenchymal stromal cells pretreated with proinflammatory cytokines enhance skin wound healing via IL-6-dependent M2 polarization. *Stem Cell Res Ther* 2022; **13**: 414 [PMID: 35964139 DOI: 10.1186/s13287-022-02934-9]
- 39 **Bruschi M**, Sahu N, Singla M, Grandi F, Agarwal P, Chu C, Bhutani N. A Quick and Efficient Method for the Generation of Immunomodulatory Mesenchymal Stromal Cell from Human Induced Pluripotent Stem Cell. *Tissue Eng Part A* 2022; **28**: 433-446 [PMID:

- 34693750 DOI: [10.1089/ten.TEA.2021.0172](https://doi.org/10.1089/ten.TEA.2021.0172)]
- 40 **Bunnell BA**. Adipose Tissue-Derived Mesenchymal Stem Cells. *Cells* 2021; **10**: 3433 [PMID: [34943941](https://pubmed.ncbi.nlm.nih.gov/34943941/) DOI: [10.3390/cells10123433](https://doi.org/10.3390/cells10123433)]
- 41 **Widjaja SL**, Salimo H, Yulianto I, Soetrisno. Proteomic analysis of hypoxia and non-hypoxia secretome mesenchymal stem-like cells from human breastmilk. *Saudi J Biol Sci* 2021; **28**: 4399-4407 [PMID: [34354424](https://pubmed.ncbi.nlm.nih.gov/34354424/) DOI: [10.1016/j.sjbs.2021.04.034](https://doi.org/10.1016/j.sjbs.2021.04.034)]
- 42 **He J**, Liu J, Huang Y, Tang X, Xiao H, Hu Z. Oxidative Stress, Inflammation, and Autophagy: Potential Targets of Mesenchymal Stem Cells-Based Therapies in Ischemic Stroke. *Front Neurosci* 2021; **15**: 641157 [PMID: [33716657](https://pubmed.ncbi.nlm.nih.gov/33716657/) DOI: [10.3389/fnins.2021.641157](https://doi.org/10.3389/fnins.2021.641157)]
- 43 **Eleuteri S**, Fierabracci A. Insights into the Secretome of Mesenchymal Stem Cells and Its Potential Applications. *Int J Mol Sci* 2019; **20** [PMID: [31533317](https://pubmed.ncbi.nlm.nih.gov/31533317/) DOI: [10.3390/ijms20184597](https://doi.org/10.3390/ijms20184597)]
- 44 **Baglio SR**, Rooijers K, Koppers-Lalic D, Verweij FJ, Pérez Lanzón M, Zini N, Naaijken B, Perut F, Niessen HW, Baldini N, Pegtel DM. Human bone marrow- and adipose-mesenchymal stem cells secrete exosomes enriched in distinctive miRNA and tRNA species. *Stem Cell Res Ther* 2015; **6**: 127 [PMID: [26129847](https://pubmed.ncbi.nlm.nih.gov/26129847/) DOI: [10.1186/s13287-015-0116-z](https://doi.org/10.1186/s13287-015-0116-z)]
- 45 **Dong L**, Pu Y, Zhang L, Qi Q, Xu L, Li W, Wei C, Wang X, Zhou S, Zhu J, Liu F, Chen X, Su C. Human umbilical cord mesenchymal stem cell-derived extracellular vesicles promote lung adenocarcinoma growth by transferring miR-410. *Cell Death Dis* 2018; **9**: 218 [PMID: [29440630](https://pubmed.ncbi.nlm.nih.gov/29440630/) DOI: [10.1038/s41419-018-0323-5](https://doi.org/10.1038/s41419-018-0323-5)]
- 46 **Basalova N**, Sagaradze G, Arbatskiy M, Evtushenko E, Kulebyakin K, Grigorieva O, Akopyan Z, Kalinina N, Efimenko A. Secretome of Mesenchymal Stromal Cells Prevents Myofibroblasts Differentiation by Transferring Fibrosis-Associated microRNAs within Extracellular Vesicles. *Cells* 2020; **9** [PMID: [32443855](https://pubmed.ncbi.nlm.nih.gov/32443855/) DOI: [10.3390/cells9051272](https://doi.org/10.3390/cells9051272)]
- 47 **Lam AT**, Lee AP, Jayaraman P, Tan KY, Raghothaman D, Lim HL, Cheng H, Zhou L, Tan AH, Reuveny S, Oh S. Multiomics analyses of cytokines, genes, miRNA, and regulatory networks in human mesenchymal stem cells expanded in stirred microcarrier-spinner cultures. *Stem Cell Res* 2021; **53**: 102272 [PMID: [33676128](https://pubmed.ncbi.nlm.nih.gov/33676128/) DOI: [10.1016/j.scr.2021.102272](https://doi.org/10.1016/j.scr.2021.102272)]
- 48 **Ulpiano C**, da Silva CL, Monteiro GA. Mesenchymal Stromal Cells (MSCs): A Promising Tool for Cell-Based Angiogenic Therapy. *Curr Gene Ther* 2021; **21**: 382-405 [PMID: [34533444](https://pubmed.ncbi.nlm.nih.gov/34533444/) DOI: [10.2174/1566523221666210917114353](https://doi.org/10.2174/1566523221666210917114353)]
- 49 **Chinnici CM**, Amico G, Monti M, Motta S, Casalone R, Petri SL, Spada M, Gridelli B, Conaldi PG. Isolation and characterization of multipotent cells from human fetal dermis. *Cell Transplant* 2014; **23**: 1169-1185 [PMID: [23768775](https://pubmed.ncbi.nlm.nih.gov/23768775/) DOI: [10.3727/096368913X668618](https://doi.org/10.3727/096368913X668618)]
- 50 **Miranda JP**, Camões SP, Gaspar MM, Rodrigues JS, Carvalheiro M, Bárçia RN, Cruz P, Cruz H, Simões S, Santos JM. The Secretome Derived From 3D-Cultured Umbilical Cord Tissue MSCs Counteracts Manifestations Typifying Rheumatoid Arthritis. *Front Immunol* 2019; **10**: 18 [PMID: [30804924](https://pubmed.ncbi.nlm.nih.gov/30804924/) DOI: [10.3389/fimmu.2019.00018](https://doi.org/10.3389/fimmu.2019.00018)]
- 51 **Alidoust Saharkhiz Lahiji M**, Safari F. Potential therapeutic effects of hAMSCs secretome on Panc1 pancreatic cancer cells through downregulation of SgK269, E-cadherin, vimentin, and snail expression. *Biologicals* 2022; **76**: 24-30 [PMID: [35216916](https://pubmed.ncbi.nlm.nih.gov/35216916/) DOI: [10.1016/j.biologicals.2022.02.001](https://doi.org/10.1016/j.biologicals.2022.02.001)]
- 52 **Liu A**, Zhang X, He H, Zhou L, Naito Y, Sugita S, Lee JW. Therapeutic potential of mesenchymal stem/stromal cell-derived secretome and vesicles for lung injury and disease. *Expert Opin Biol Ther* 2020; **20**: 125-140 [PMID: [31701782](https://pubmed.ncbi.nlm.nih.gov/31701782/) DOI: [10.1080/14712598.2020.1689954](https://doi.org/10.1080/14712598.2020.1689954)]
- 53 **Jiang D**, Chen FX, Zhou H, Lu YY, Tan H, Yu SJ, Yuan J, Liu H, Meng W, Jin ZB. Bioenergetic Crosstalk between Mesenchymal Stem Cells and various Ocular Cells through the intercellular trafficking of Mitochondria. *Theranostics* 2020; **10**: 7260-7272 [PMID: [32641991](https://pubmed.ncbi.nlm.nih.gov/32641991/) DOI: [10.7150/thno.46332](https://doi.org/10.7150/thno.46332)]
- 54 **Scuteri A**, Miloso M, Foudah D, Orciani M, Cavaletti G, Tredici G. Mesenchymal stem cells neuronal differentiation ability: a real perspective for nervous system repair? *Curr Stem Cell Res Ther* 2011; **6**: 82-92 [PMID: [21190538](https://pubmed.ncbi.nlm.nih.gov/21190538/) DOI: [10.2174/157488811795495486](https://doi.org/10.2174/157488811795495486)]
- 55 **Andrzejewska A**, Dabrowska S, Lukomska B, Janowski M. Mesenchymal Stem Cells for Neurological Disorders. *Adv Sci (Weinh)* 2021; **8**: 2002944 [PMID: [33854883](https://pubmed.ncbi.nlm.nih.gov/33854883/) DOI: [10.1002/advs.202002944](https://doi.org/10.1002/advs.202002944)]
- 56 **Das M**, Mayilsamy K, Mohapatra SS, Mohapatra S. Mesenchymal stem cell therapy for the treatment of traumatic brain injury: progress and prospects. *Rev Neurosci* 2019; **30**: 839-855 [PMID: [31203262](https://pubmed.ncbi.nlm.nih.gov/31203262/) DOI: [10.1515/revneuro-2019-0002](https://doi.org/10.1515/revneuro-2019-0002)]
- 57 **Hasan A**, Deeb G, Rahal R, Atwi K, Mondello S, Marei HE, Gali A, Sleiman E. Mesenchymal Stem Cells in the Treatment of Traumatic Brain Injury. *Front Neurol* 2017; **8**: 28 [PMID: [28265255](https://pubmed.ncbi.nlm.nih.gov/28265255/) DOI: [10.3389/fneur.2017.00028](https://doi.org/10.3389/fneur.2017.00028)]
- 58 **Cozene B**, Sadanandan N, Farooq J, Kingsbury C, Park YJ, Wang ZJ, Moscatello A, Saft M, Cho J, Gonzales-Portillo B, Borlongan CV. Mesenchymal Stem Cell-Induced Anti-Neuroinflammation Against Traumatic Brain Injury. *Cell Transplant* 2021; **30**: 9636897211035715 [PMID: [34559583](https://pubmed.ncbi.nlm.nih.gov/34559583/) DOI: [10.1177/09636897211035715](https://doi.org/10.1177/09636897211035715)]
- 59 **Cofano F**, Boido M, Monticelli M, Zenga F, Ducati A, Vercelli A, Garbossa D. Mesenchymal Stem Cells for Spinal Cord Injury: Current Options, Limitations, and Future of Cell Therapy. *Int J Mol Sci* 2019; **20** [PMID: [31159345](https://pubmed.ncbi.nlm.nih.gov/31159345/) DOI: [10.3390/ijms20112698](https://doi.org/10.3390/ijms20112698)]
- 60 **Pang QM**, Chen SY, Xu QJ, Fu SP, Yang YC, Zou WH, Zhang M, Liu J, Wan WH, Peng JC, Zhang T. Neuroinflammation and Scarring After Spinal Cord Injury: Therapeutic Roles of MSCs on Inflammation and Glial Scar. *Front Immunol* 2021; **12**: 751021 [PMID: [34925326](https://pubmed.ncbi.nlm.nih.gov/34925326/) DOI: [10.3389/fimmu.2021.751021](https://doi.org/10.3389/fimmu.2021.751021)]
- 61 **Gao L**, Peng Y, Xu W, He P, Li T, Lu X, Chen G. Progress in Stem Cell Therapy for Spinal Cord Injury. *Stem Cells Int* 2020; **2020**: 2853650 [PMID: [33204276](https://pubmed.ncbi.nlm.nih.gov/33204276/) DOI: [10.1155/2020/2853650](https://doi.org/10.1155/2020/2853650)]
- 62 **Chung JW**, Chang WH, Bang OY, Moon GJ, Kim SJ, Kim SK, Lee JS, Sohn SI, Kim YH; STARTING-2 Collaborators. Efficacy and Safety of Intravenous Mesenchymal Stem Cells for Ischemic Stroke. *Neurology* 2021; **96**: e1012-e1023 [PMID: [33472925](https://pubmed.ncbi.nlm.nih.gov/33472925/) DOI: [10.1212/WNL.00000000000011440](https://doi.org/10.1212/WNL.00000000000011440)]
- 63 **Dabrowska S**, Andrzejewska A, Lukomska B, Janowski M. Neuroinflammation as a target for treatment of stroke using mesenchymal stem cells and extracellular vesicles. *J Neuroinflammation* 2019; **16**: 178 [PMID: [31514749](https://pubmed.ncbi.nlm.nih.gov/31514749/) DOI: [10.1186/s12974-019-1571-8](https://doi.org/10.1186/s12974-019-1571-8)]
- 64 **Tatullo M**, Marrelli B, Zullo MJ, Codispoti B, Paduano F, Benincasa C, Fortunato F, Scacco S, Zavan B, Cocco T. Exosomes from Human Periapical Cyst-MSCs: Theranostic Application in Parkinson's Disease. *Int J Med Sci* 2020; **17**: 657-663 [PMID: [32210716](https://pubmed.ncbi.nlm.nih.gov/32210716/) DOI: [10.7150/ijms.41515](https://doi.org/10.7150/ijms.41515)]
- 65 **Peng H**, Li Y, Ji W, Zhao R, Lu Z, Shen J, Wu Y, Wang J, Hao Q, Wang W, Yang J, Zhang X. Intranasal Administration of Self-Oriented Nanocarriers Based on Therapeutic Exosomes for Synergistic Treatment of Parkinson's Disease. *ACS Nano* 2022; **16**: 869-884 [PMID: [34985280](https://pubmed.ncbi.nlm.nih.gov/34985280/) DOI: [10.1021/acsnano.1c08473](https://doi.org/10.1021/acsnano.1c08473)]
- 66 **Hoon M**, Okawa H, Della Santina L, Wong RO. Functional architecture of the retina: development and disease. *Prog Retin Eye Res* 2014; **42**: 44-84 [PMID: [24984227](https://pubmed.ncbi.nlm.nih.gov/24984227/) DOI: [10.1016/j.preteyeres.2014.06.003](https://doi.org/10.1016/j.preteyeres.2014.06.003)]
- 67 **Diomedede F**, Marconi GD, Fonticoli L, Pizzicanella J, Merciaro I, Bramanti P, Mazzon E, Trubiani O. Functional Relationship between Osteogenesis and Angiogenesis in Tissue Regeneration. *Int J Mol Sci* 2020; **21** [PMID: [32375269](https://pubmed.ncbi.nlm.nih.gov/32375269/) DOI: [10.3390/ijms21093242](https://doi.org/10.3390/ijms21093242)]

- 68 **Scimone C**, Donato L, Marino S, Alafaci C, D'Angelo R, Sidoti A. Vis-à-vis: a focus on genetic features of cerebral cavernous malformations and brain arteriovenous malformations pathogenesis. *Neurol Sci* 2019; **40**: 243-251 [PMID: 30523548 DOI: 10.1007/s10072-018-3674-x]
- 69 **Scimone C**, Donato L, Alibrandi S, Esposito T, Alafaci C, D'Angelo R, Sidoti A. Transcriptome analysis provides new molecular signatures in sporadic Cerebral Cavernous Malformation endothelial cells. *Biochim Biophys Acta Mol Basis Dis* 2020; **1866**: 165956 [PMID: 32877751 DOI: 10.1016/j.bbadis.2020.165956]
- 70 **Eslani M**, Putra I, Shen X, Hamouie J, Afsharkhamesh N, Besharat S, Rosenblatt MI, Dana R, Hematti P, Djalilian AR. Corneal Mesenchymal Stromal Cells Are Directly Antiangiogenic via PEDF and sFLT-1. *Invest Ophthalmol Vis Sci* 2017; **58**: 5507-5517 [PMID: 29075761 DOI: 10.1167/iovs.17-22680]
- 71 **Neidlinger-Wilke C**, Ekkerlein A, Goncalves RM, Ferreira JR, Ignatius A, Wilke HJ, Teixeira GQ. Mesenchymal stem cell secretome decreases the inflammatory response in annulus fibrosus organ cultures. *Eur Cell Mater* 2021; **42**: 1-19 [PMID: 34235715 DOI: 10.22203/eCM.v042a01]
- 72 **Karimi-Haghighi S**, Chavoshinezhad S, Safari A, Razeghian-Jahromi I, Jamhiri I, Khodabandeh Z, Khajeh S, Zare S, Borhani-Haghighi A, Dianatpour M, Pandamooz S, Salehi MS. Preconditioning with secretome of neural crest-derived stem cells enhanced neurotrophic expression in mesenchymal stem cells. *Neurosci Lett* 2022; **773**: 136511 [PMID: 35143889 DOI: 10.1016/j.neulet.2022.136511]
- 73 **Konala VBR**, Bhonde R, Pal R. Secretome studies of mesenchymal stromal cells (MSCs) isolated from three tissue sources reveal subtle differences in potency. *In Vitro Cell Dev Biol Anim* 2020; **56**: 689-700 [PMID: 33006709 DOI: 10.1007/s11626-020-00501-1]
- 74 **Sumapraja K**, Hestiantoro A, Liem IK, Boediono A, Z Jacob T. Effect of conditioned medium of umbilical cord-derived mesenchymal stem cells as a culture medium for human granulosa cells: An experimental study. *Int J Reprod Biomed* 2021; **19**: 1037-1044 [PMID: 35098005 DOI: 10.18502/ijrm.v19i12.10054]
- 75 **Llewellyn SH**, Faroni A, Iliut M, Bartlam C, Vijayaraghavan A, Reid AJ. Graphene Oxide Substrate Promotes Neurotrophic Factor Secretion and Survival of Human Schwann-Like Adipose Mesenchymal Stromal Cells. *Adv Biol (Weinh)* 2021; **5**: e2000271 [PMID: 33852181 DOI: 10.1002/adbi.202000271]
- 76 **Prakoewa CRS**, Rindiastuti Y, Wirohadidjojo YW, Komaratih E, Nurwasis, Dinaryati A, Lestari NMI, Rantam FA. Resveratrol promotes secretion of wound healing related growth factors of mesenchymal stem cells originated from adult and fetal tissues. *Artif Cells Nanomed Biotechnol* 2020; **48**: 1160-1167 [PMID: 32902361 DOI: 10.1080/21691401.2020.1817057]
- 77 **Alfaro MP**, Deskins DL, Wallus M, DasGupta J, Davidson JM, Nanney LB, A Guney M, Gannon M, Young PP. A physiological role for connective tissue growth factor in early wound healing. *Lab Invest* 2013; **93**: 81-95 [PMID: 23212098 DOI: 10.1038/labinvest.2012.162]
- 78 **Kubo E**, Shibata T, Singh DP, Sasaki H. Roles of TGF β and FGF Signals in the Lens: Tropomyosin Regulation for Posterior Capsule Opacity. *Int J Mol Sci* 2018; **19** [PMID: 30304871 DOI: 10.3390/ijms19103093]
- 79 **Chouw A**, Sartika CR, Miranda T, Faried A. Interleukins Profiling in Umbilical Cord Mesenchymal Stem Cell-Derived Secretome. *Stem Cells Cloning* 2022; **15**: 1-9 [PMID: 35444427 DOI: 10.2147/SCCAA.S356763]
- 80 **Madonna R**, Angelucci S, Di Giuseppe F, Doria V, Giricz Z, Görbe A, Ferdinandy P, De Caterina R. Proteomic analysis of the secretome of adipose tissue-derived murine mesenchymal cells overexpressing telomerase and myocardin. *J Mol Cell Cardiol* 2019; **131**: 171-186 [PMID: 31055035 DOI: 10.1016/j.yjmcc.2019.04.019]
- 81 **Scutera S**, Mitola S, Sparti R, Salvi V, Grillo E, Piersigilli G, Bugatti M, Alotto D, Schioppa T, Sozzani S, Musso T. Bartonella henselae Persistence within Mesenchymal Stromal Cells Enhances Endothelial Cell Activation and Infectibility That Amplifies the Angiogenic Process. *Infect Immun* 2021; **89**: e0014121 [PMID: 34031126 DOI: 10.1128/IAI.00141-21]
- 82 **Vašiček J**, Baláži A, Tirpáková M, Svoradová A, Ondruška L, Parkányi V, Chrenek P. Secretome Analysis of Rabbit and Human Mesenchymal Stem and Endothelial Progenitor Cells: A Comparative Study. *Int J Mol Sci* 2021; **22** [PMID: 34830165 DOI: 10.3390/ijms222212283]
- 83 **Tsiftoglou AS**. Erythropoietin (EPO) as a Key Regulator of Erythropoiesis, Bone Remodeling and Endothelial Transdifferentiation of Multipotent Mesenchymal Stem Cells (MSCs): Implications in Regenerative Medicine. *Cells* 2021; **10** [PMID: 34440909 DOI: 10.3390/cells10082140]
- 84 **González P**, Santos TM, Calil A, Corradi Perini C, Percegon LS, Silva IC, Kuligovski C, Aguiar AM, Câmara NO, Aita CA. Expression of pancreatic endocrine markers by prolactin-treated rat bone marrow mesenchymal stem cells. *Transplant Proc* 2010; **42**: 566-569 [PMID: 20304194 DOI: 10.1016/j.transproceed.2010.01.031]
- 85 **Jin X**, Dai G, Xuan L, Zhang M, Jiang H, Sui Y. Effects of Sodium Chlorophyllin Copper on APO-1 Expression in Bone Marrow Mesenchymal Stem Cells of Rats with Aplastic Anaemia. *J Immunol Res* 2022; **2022**: 6792866 [PMID: 35434141 DOI: 10.1155/2022/6792866]
- 86 **Shen Y**. Stem cell therapies for retinal diseases: from bench to bedside. *J Mol Med (Berl)* 2020; **98**: 1347-1368 [PMID: 32794020 DOI: 10.1007/s00109-020-01960-5]
- 87 **Ponnalagu M**, Subramani M, Jayadev C, Shetty R, Das D. Retinal pigment epithelium-secretome: A diabetic retinopathy perspective. *Cytokine* 2017; **95**: 126-135 [PMID: 28282610 DOI: 10.1016/j.cyto.2017.02.013]
- 88 **Donato L**, Scimone C, Alibrandi S, Scalinci SZ, Rinaldi C, D'Angelo R, Sidoti A. Epitranscriptome Analysis of Oxidative Stressed Retinal Epithelial Cells Depicted a Possible RNA Editing Landscape of Retinal Degeneration. *Antioxidants (Basel)* 2022; **11** [PMID: 36290689 DOI: 10.3390/antiox11101967]
- 89 **Flores-Bellver M**, Mighty J, Aparicio-Domingo S, Li KV, Shi C, Zhou J, Cobb H, McGrath P, Michelis G, Lenhart P, Bilousova G, Heissel S, Rudy MJ, Coughlan C, Goodspeed AE, Becerra SP, Redenti S, Canto-Soler MV. Extracellular vesicles released by human retinal pigment epithelium mediate increased polarised secretion of drusen proteins in response to AMD stressors. *J Extracell Vesicles* 2021; **10**: e12165 [PMID: 34750957 DOI: 10.1002/jev2.12165]
- 90 **Chen L**, Perera ND, Karoukis AJ, Feathers KL, Ali RR, Thompson DA, Fahim AT. Oxidative stress differentially impacts apical and basolateral secretion of angiogenic factors from human iPSC-derived retinal pigment epithelium cells. *Sci Rep* 2022; **12**: 12694 [PMID: 35882889 DOI: 10.1038/s41598-022-16701-6]
- 91 **Klettner A**, Kampers M, Töbelmann D, Roeder J, Dittmar M. The Influence of Melatonin and Light on VEGF Secretion in Primary RPE Cells. *Biomolecules* 2021; **11** [PMID: 33467052 DOI: 10.3390/biom11010114]
- 92 **Farjood F**, Vargis E. Physical disruption of cell-cell contact induces VEGF expression in RPE cells. *Mol Vis* 2017; **23**: 431-446 [PMID: 28761317]
- 93 **Warden C**, Brantley MA Jr. Glycine-Conjugated Bile Acids Protect RPE Tight Junctions against Oxidative Stress and Inhibit Choroidal Endothelial Cell Angiogenesis In Vitro. *Biomolecules* 2021; **11** [PMID: 33922434 DOI: 10.3390/biom11050626]

- 94 **Chen Q**, Tang L, Zhang Y, Wan C, Yu X, Dong Y, Chen X, Wang X, Li N, Xin G, Zhang M, Chen Z, Niu H, Huang W. STING up-regulates VEGF expression in oxidative stress-induced senescence of retinal pigment epithelium via NF- κ B/HIF-1 α pathway. *Life Sci* 2022; **293**: 120089 [PMID: 35007563 DOI: 10.1016/j.lfs.2021.120089]
- 95 **Baba T**, McLeod DS, Edwards MM, Merges C, Sen T, Sinha D, Luty GA. VEGF 165 b in the developing vasculatures of the fetal human eye. *Dev Dyn* 2012; **241**: 595-607 [PMID: 22275161 DOI: 10.1002/dvdy.23743]
- 96 **Rebustini IT**, Crawford SE, Becerra SP. PEDF Deletion Induces Senescence and Defects in Phagocytosis in the RPE. *Int J Mol Sci* 2022; **23** [PMID: 35887093 DOI: 10.3390/ijms23147745]
- 97 **Tombran-Tink J**. PEDF in angiogenic eye diseases. *Curr Mol Med* 2010; **10**: 267-278 [PMID: 20236057 DOI: 10.2174/156652410791065336]
- 98 **Araújo RS**, Silva GA. PIGF silencing combined with PEDF overexpression: Modeling RPE secretion as potential therapy for retinal neovascularization. *Mol Biol Rep* 2020; **47**: 4413-4425 [PMID: 32385771 DOI: 10.1007/s11033-020-05496-2]
- 99 **Pagan-Mercado G**, Becerra SP. Signaling Mechanisms Involved in PEDF-Mediated Retinoprotection. *Adv Exp Med Biol* 2019; **1185**: 445-449 [PMID: 31884652 DOI: 10.1007/978-3-030-27378-1_73]
- 100 **Hollborn M**, Bringmann A, Faude F, Wiedemann P, Kohen L. Signaling pathways involved in PDGF-evoked cellular responses in human RPE cells. *Biochem Biophys Res Commun* 2006; **344**: 912-919 [PMID: 16631611 DOI: 10.1016/j.bbrc.2006.03.185]
- 101 **Jung J**, Jeong J, Hong HS. Substance P improves MSC-mediated RPE regeneration by modulating PDGF-BB. *Biochem Biophys Res Commun* 2019; **515**: 524-530 [PMID: 31176487 DOI: 10.1016/j.bbrc.2019.05.186]
- 102 **Zhang H**, Shang Q, An J, Wang C, Ma J. Crocetin inhibits PDGF-BB-induced proliferation and migration of retinal pigment epithelial cells. *Eur J Pharmacol* 2019; **842**: 329-337 [PMID: 30395849 DOI: 10.1016/j.ejphar.2018.11.001]
- 103 **Lennikov A**, Mukwaya A, Fan L, Saddala MS, De Falco S, Huang H. Synergistic interactions of PIGF and VEGF contribute to blood-retinal barrier breakdown through canonical NF κ B activation. *Exp Cell Res* 2020; **397**: 112347 [PMID: 33130176 DOI: 10.1016/j.yexcr.2020.112347]
- 104 **Mesquita J**, Castro-de-Sousa JP, Vaz-Pereira S, Neves A, Passarinha LA, Tomaz CT. Vascular endothelial growth factors and placenta growth factor in retinal vasculopathies: Current research and future perspectives. *Cytokine Growth Factor Rev* 2018; **39**: 102-115 [PMID: 29248329 DOI: 10.1016/j.cytogfr.2017.11.005]
- 105 **Zhou AY**, Bai YJ, Zhao M, Yu WZ, Huang LZ, Li XX. Placental growth factor expression is reversed by antivascular endothelial growth factor therapy under hypoxic conditions. *World J Pediatr* 2014; **10**: 262-270 [PMID: 25124978 DOI: 10.1007/s12519-014-0502-0]
- 106 **Moon S**, Lee S, Caesar JA, Pruchenko S, Leask A, Knowles JA, Sinon J, Chaqour B. A CTGF-YAP Regulatory Pathway Is Essential for Angiogenesis and Barrierogenesis in the Retina. *iScience* 2020; **23**: 101184 [PMID: 32502964 DOI: 10.1016/j.isci.2020.101184]
- 107 **Yang H**, Huang Y, Chen X, Liu J, Lu Y, Bu L, Xia L, Xiao W, Chen M, Nie Q, Liu Z. The role of CTGF in the diabetic rat retina and its relationship with VEGF and TGF- β (2), elucidated by treatment with CTGFsiRNA. *Acta Ophthalmol* 2010; **88**: 652-659 [PMID: 20039857 DOI: 10.1111/j.1755-3768.2009.01641.x]
- 108 **Daftarian N**, Baigy O, Suri F, Kanavi MR, Balaghali S, Afsar Askı S, Moghaddasi A, Nourinia R, Abtahi SH, Ahmadi H. Intravitreal connective tissue growth factor neutralizing antibody or bevacizumab alone or in combination for prevention of proliferative vitreoretinopathy in an experimental model. *Exp Eye Res* 2021; **208**: 108622 [PMID: 34022176 DOI: 10.1016/j.exer.2021.108622]
- 109 **Hochmann S**, Kaslin J, Hans S, Weber A, Machate A, Geffarth M, Funk RH, Brand M. Fgf signaling is required for photoreceptor maintenance in the adult zebrafish retina. *PLoS One* 2012; **7**: e30365 [PMID: 22291943 DOI: 10.1371/journal.pone.0030365]
- 110 **Huang HW**, Yang CM, Yang CH. Fibroblast Growth Factor Type 1 Ameliorates High-Glucose-Induced Oxidative Stress and Neuroinflammation in Retinal Pigment Epithelial Cells and a Streptozotocin-Induced Diabetic Rat Model. *Int J Mol Sci* 2021; **22** [PMID: 34281287 DOI: 10.3390/ijms22137233]
- 111 **Mukherjee S**, Guidry C. The insulin-like growth factor system modulates retinal pigment epithelial cell tractional force generation. *Invest Ophthalmol Vis Sci* 2007; **48**: 1892-1899 [PMID: 17389525 DOI: 10.1167/iovs.06-1095]
- 112 **Puddu A**, Sanguineti R, Maggi D. Caveolin-1 Down-Regulation Reduces VEGF-A Secretion Induced by IGF-1 in ARPE-19 Cells. *Life (Basel)* 2021; **12**: 44 [PMID: 35054437 DOI: 10.3390/life12010044]
- 113 **Ainscough SL**, Feigl B, Malda J, Harkin DG. Discovery and characterization of IGFBP-mediated endocytosis in the human retinal pigment epithelial cell line ARPE-19. *Exp Eye Res* 2009; **89**: 629-637 [PMID: 19540231 DOI: 10.1016/j.exer.2009.06.005]
- 114 **Lee SC**, Kim SH, Koh HJ, Kwon OW. TGF-betas synthesized by RPE cells have autocrine activity on mesenchymal transformation and cell proliferation. *Yonsei Med J* 2001; **42**: 271-277 [PMID: 11456391 DOI: 10.3349/ymj.2001.42.3.271]
- 115 **Kelly TJ**, Brümmer A, Hooshdaran N, Tariveranmohabab M, Zamudio JR. Temporal Control of the TGF- β Signaling Network by Mouse ESC MicroRNA Targets of Different Affinities. *Cell Rep* 2019; **29**: 2702-2717.e7 [PMID: 31775039 DOI: 10.1016/j.celrep.2019.10.109]
- 116 **Meyer JG**, Garcia TY, Schilling B, Gibson BW, Lamba DA. Proteome and Secretome Dynamics of Human Retinal Pigment Epithelium in Response to Reactive Oxygen Species. *Sci Rep* 2019; **9**: 15440 [PMID: 31659173 DOI: 10.1038/s41598-019-51777-7]
- 117 **Bretz CA**, Divoky V, Prchal J, Kunz E, Simmons AB, Wang H, Hartnett ME. Erythropoietin Signaling Increases Choroidal Macrophages and Cytokine Expression, and Exacerbates Choroidal Neovascularization. *Sci Rep* 2018; **8**: 2161 [PMID: 29391474 DOI: 10.1038/s41598-018-20520-z]
- 118 **Xie H**, Zhang C, Liu D, Yang Q, Tang L, Wang T, Tian H, Lu L, Xu JY, Gao F, Wang J, Jin C, Li W, Xu G, Xu GT, Zhang J. Erythropoietin protects the inner blood-retinal barrier by inhibiting microglia phagocytosis via Src/Akt/cofilin signalling in experimental diabetic retinopathy. *Diabetologia* 2021; **64**: 211-225 [PMID: 33104828 DOI: 10.1007/s00125-020-05299-x]
- 119 **Zhang C**, Xie H, Yang Q, Yang Y, Li W, Tian H, Lu L, Wang F, Xu JY, Gao F, Wang J, Jin C, Xu G, Xu GT, Zhang J. Erythropoietin protects outer blood-retinal barrier in experimental diabetic retinopathy by up-regulating ZO-1 and occludin. *Clin Exp Ophthalmol* 2019; **47**: 1182-1197 [PMID: 31483932 DOI: 10.1111/ceo.13619]
- 120 **Biswal MR**, Wang Z, Paulson RJ, Uddin RR, Tong Y, Zhu P, Li H, Lewin AS. Erythropoietin Gene Therapy Delays Retinal Degeneration Resulting from Oxidative Stress in the Retinal Pigment Epithelium. *Antioxidants (Basel)* 2021; **10** [PMID: 34070383 DOI: 10.3390/antiox10060842]
- 121 **Samson FP**, He W, Sripathi SR, Patrick AT, Madu J, Chung H, Frost MC, Jee D, Gutsaeva DR, Jahng WJ. Dual Switch Mechanism of Erythropoietin as an Antiapoptotic and Pro-Angiogenic Determinant in the Retina. *ACS Omega* 2020; **5**: 21113-21126 [PMID: 32875248 DOI: 10.1021/acsomega.0c02763]
- 122 **Hoffmann S**, He S, Ehren M, Ryan SJ, Wiedemann P, Hinton DR. MMP-2 and MMP-9 secretion by rpe is stimulated by angiogenic molecules found in choroidal neovascular membranes. *Retina* 2006; **26**: 454-461 [PMID: 16603966 DOI: 10.1097/01.iae.0000238549.74626.33]

- 123 **Alge-Priglinger CS**, Kreutzer T, Obholzer K, Wolf A, Mempel M, Kernt M, Kampik A, Priglinger SG. Oxidative stress-mediated induction of MMP-1 and MMP-3 in human RPE cells. *Invest Ophthalmol Vis Sci* 2009; **50**: 5495-5503 [PMID: 19516002 DOI: 10.1167/iovs.08-3193]
- 124 **Kim J**, Kim JH, Do JY, Lee JY, Yanai R, Lee IK, Suk K, Park DH. Key Role of Microglial Matrix Metalloproteinases in Choroidal Neovascularization. *Front Cell Neurosci* 2021; **15**: 638098 [PMID: 33716674 DOI: 10.3389/fncel.2021.638098]
- 125 **Korhonen E**, Piippo N, Hytti M, Hyttinen JMT, Kaamiranta K, Kauppinen A. SQSTM1/p62 regulates the production of IL-8 and MCP-1 in IL-1 β -stimulated human retinal pigment epithelial cells. *Cytokine* 2019; **116**: 70-77 [PMID: 30685605 DOI: 10.1016/j.cyto.2018.12.015]
- 126 **Taghavi Y**, Hassanshahi G, Kounis NG, Koniari I, Khorramdelazad H. Monocyte chemoattractant protein-1 (MCP-1/CCL2) in diabetic retinopathy: latest evidence and clinical considerations. *J Cell Commun Signal* 2019; **13**: 451-462 [PMID: 30607767 DOI: 10.1007/s12079-018-00500-8]
- 127 **Klettner A**, Brinkmann A, Winkelmann K, Käckenmeister T, Hildebrandt J, Roeder J. Effect of long-term inflammation on viability and function of RPE cells. *Exp Eye Res* 2020; **200**: 108214 [PMID: 32898511 DOI: 10.1016/j.exer.2020.108214]
- 128 **Weir K**, Kim DW, Blackshaw S. A potential role for somatostatin signaling in regulating retinal neurogenesis. *Sci Rep* 2021; **11**: 10962 [PMID: 34040115 DOI: 10.1038/s41598-021-90554-3]
- 129 **An E**, Sen S, Park SK, Gordish-Dressman H, Hathout Y. Identification of novel substrates for the serine protease HTRA1 in the human RPE secretome. *Invest Ophthalmol Vis Sci* 2010; **51**: 3379-3386 [PMID: 20207970 DOI: 10.1167/iovs.09-4853]
- 130 **Donato L**, Alibrandi S, Scimone C, Rinaldi C, Dascola A, Calamuneri A, D'Angelo R, Sidoti A. The impact of modifier genes on cone-rod dystrophy heterogeneity: An explorative familial pilot study and a hypothesis on neurotransmission impairment. *PLoS One* 2022; **17**: e0278857 [PMID: 36490268 DOI: 10.1371/journal.pone.0278857]
- 131 **Ishida T**, Yoshida T, Shinohara K, Cao K, Nakahama KI, Morita I, Ohno-Matsui K. Potential role of sirtuin 1 in Müller glial cells in mice choroidal neovascularization. *PLoS One* 2017; **12**: e0183775 [PMID: 28886036 DOI: 10.1371/journal.pone.0183775]
- 132 **Salimiaghdam N**, Singh L, Schneider K, Chwa M, Atilano SR, Nalbandian A, Limb GA, Kenney MC. Effects of fluoroquinolones and tetracyclines on mitochondria of human retinal MIO-M1 cells. *Exp Eye Res* 2022; **214**: 108857 [PMID: 34856207 DOI: 10.1016/j.exer.2021.108857]
- 133 **Weigelt CM**, Fuchs H, Schönberger T, Stierstorfer B, Strobel B, Lamla T, Ciossek T, Bakker RA, Redemann NH. AAV-Mediated Expression of Human VEGF, TNF- α , and IL-6 Induces Retinal Pathology in Mice. *Transl Vis Sci Technol* 2021; **10**: 15 [PMID: 34520511 DOI: 10.1167/tvst.10.11.15]
- 134 **El-Hodiri HM**, Campbell WA, Kelly LE, Hawthorn EC, Schwartz M, Jalligampala A, McCall MA, Meyer K, Fischer AJ. Nuclear Factor I in neurons, glia and during the formation of Müller glia-derived progenitor cells in avian, porcine and primate retinas. *J Comp Neurol* 2022; **530**: 1213-1230 [PMID: 34729776 DOI: 10.1002/cne.25270]
- 135 **Schmalen A**, Lorenz L, Grosche A, Pauly D, Deeg CA, Hauck SM. Proteomic Phenotyping of Stimulated Müller Cells Uncovers Profound Pro-Inflammatory Signaling and Antigen-Presenting Capacity. *Front Pharmacol* 2021; **12**: 771571 [PMID: 34776983 DOI: 10.3389/fphar.2021.771571]
- 136 **Zwanzig A**, Meng J, Müller H, Bürger S, Schmidt M, Pankonin M, Wiedemann P, Unterlauff JD, Eichler W. Neuroprotective effects of glial mediators in interactions between retinal neurons and Müller cells. *Exp Eye Res* 2021; **209**: 108689 [PMID: 34216615 DOI: 10.1016/j.exer.2021.108689]
- 137 **Bürger S**, Meng J, Zwanzig A, Beck M, Pankonin M, Wiedemann P, Eichler W, Unterlauff JD. Pigment Epithelium-Derived Factor (PEDF) Receptors Are Involved in Survival of Retinal Neurons. *Int J Mol Sci* 2020; **22** [PMID: 33396450 DOI: 10.3390/ijms22010369]
- 138 **Nishikiori N**, Osanai M, Chiba H, Kojima T, Mitamura Y, Ohguro H, Sawada N. Glial cell-derived cytokines attenuate the breakdown of vascular integrity in diabetic retinopathy. *Diabetes* 2007; **56**: 1333-1340 [PMID: 17470563 DOI: 10.2337/db06-1431]
- 139 **Fu S**, Dong S, Zhu M, Sherry DM, Wang C, You Z, Haigh JJ, Le YZ. Müller Glia Are a Major Cellular Source of Survival Signals for Retinal Neurons in Diabetes. *Diabetes* 2015; **64**: 3554-3563 [PMID: 26068541 DOI: 10.2337/db15-0180]
- 140 **Abu El-Asrar AM**, Nawaz MI, Ahmad A, Siddiquei MM, Allegaert E, Gikandi PW, De Hertogh G, Odenakker G. CD146/Soluble CD146 Pathway Is a Novel Biomarker of Angiogenesis and Inflammation in Proliferative Diabetic Retinopathy. *Invest Ophthalmol Vis Sci* 2021; **62**: 32 [PMID: 34293080 DOI: 10.1167/iovs.62.9.32]
- 141 **Miyata Y**. [Structure Activity Relationship Study of Polymethoxylated Flavones Targeted Retinal Müller Cells for Prevention of Retinal Diseases]. *Yakugaku Zasshi* 2021; **141**: 41-45 [PMID: 33390446 DOI: 10.1248/yakushi.20-00177-2]
- 142 **Ruzafa N**, Pereiro X, Lepper MF, Hauck SM, Vecino E. A Proteomics Approach to Identify Candidate Proteins Secreted by Müller Glia that Protect Ganglion Cells in the Retina. *Proteomics* 2018; **18**: e1700321 [PMID: 29645351 DOI: 10.1002/pmic.201700321]
- 143 **Couturier A**, Blot G, Vignaud L, Nanteau C, Slembrouck-Brec A, Fradot V, Acar N, Sahel JA, Tadayoni R, Thuret G, Sennlaub F, Roger JE, Goureau O, Guillonnet X, Reichman S. Reproducing diabetic retinopathy features using newly developed human induced-pluripotent stem cell-derived retinal Müller glial cells. *Glia* 2021; **69**: 1679-1693 [PMID: 33683746 DOI: 10.1002/glia.23983]
- 144 **Demais V**, Pohl A, Wunderlich KA, Pfaller AM, Kaplan L, Barthélémy A, Dittrich R, Puig B, Giebel B, Hauck SM, Pfrieger FW, Grosche A. Release of VAMP5-positive extracellular vesicles by retinal Müller glia in vivo. *J Extracell Vesicles* 2022; **11**: e12254 [PMID: 36043482 DOI: 10.1002/jev2.12254]
- 145 **Yang MH**, Krishnamoorthy RR, Jong SB, Chu PY, Yang YH, Chen WC, Chen SC, Dibas A, Chung TW, Tyan YC. Protein profiling of human nonpigmented ciliary epithelium cell secretome: the differentiation factors characterization for retinal ganglion cell line. *J Biomed Biotechnol* 2011; **2011**: 901329 [PMID: 21860587 DOI: 10.1155/2011/901329]
- 146 **Cuadrado-Vilanova M**, Liu J, Paco S, Aschero R, Burgueño V, Sirab N, Pascual-Pasto G, Correa G, Balaguer-Lluna L, Castillo-Ecija H, Perez-Jaume S, Muñoz-Aznar O, Roldan M, Suñol M, Schaiquevich P, Aerts I, Doz F, Cassoux N, Lubieniecki F, Benitez-Ribas D, Lavarino C, Mora J, Chantada GL, Catala-Mora J, Radvanyi F, Carcaboso AM. Identification of immunosuppressive factors in retinoblastoma cell secretomes and aqueous humor from patients. *J Pathol* 2022; **257**: 327-339 [PMID: 35254670 DOI: 10.1002/path.5893]
- 147 **Noueihed B**, Rivera JC, Dabouz R, Abram P, Omri S, Lahaie I, Chemtob S. Mesenchymal Stromal Cells Promote Retinal Vascular Repair by Modulating Sema3E and IL-17A in a Model of Ischemic Retinopathy. *Front Cell Dev Biol* 2021; **9**: 630645 [PMID: 33553187 DOI: 10.3389/fcell.2021.630645]
- 148 **Johnson TV**, DeKorver NW, Levasseur VA, Osborne A, Tassoni A, Lorber B, Heller JP, Villasmil R, Bull ND, Martin KR, Tomarev SI. Identification of retinal ganglion cell neuroprotection conferred by platelet-derived growth factor through analysis of the mesenchymal stem cell secretome. *Brain* 2014; **137**: 503-519 [PMID: 24176979 DOI: 10.1093/brain/awt292]
- 149 **Kuo SC**, Chio CC, Yeh CH, Ma JT, Liu WP, Lin MT, Lin KC, Chang CP. Mesenchymal stem cell-conditioned medium attenuates the retinal pathology in amyloid- β -induced rat model of Alzheimer's disease: Underlying mechanisms. *Aging Cell* 2021; **20**: e13340 [PMID: 33783931]

DOI: [10.1111/accel.13340](https://doi.org/10.1111/accel.13340)]

- 150 **Jafarinia M**, Farrokhi MR, Ganjalikhani Hakemi M, Cho WC. The role of miRNAs from mesenchymal stem/stromal cells-derived extracellular vesicles in neurological disorders. *Hum Cell* 2023; **36**: 62-75 [PMID: [36261702](https://pubmed.ncbi.nlm.nih.gov/36261702/) DOI: [10.1007/s13577-022-00813-2](https://doi.org/10.1007/s13577-022-00813-2)]
- 151 **Usategui-Martín R**, Puertas-Neyra K, Galindo-Cabello N, Hernández-Rodríguez LA, González-Pérez F, Rodríguez-Cabello JC, González-Sarmiento R, Pastor JC, Fernandez-Bueno I. Retinal Neuroprotective Effect of Mesenchymal Stem Cells Secretome Through Modulation of Oxidative Stress, Autophagy, and Programmed Cell Death. *Invest Ophthalmol Vis Sci* 2022; **63**: 27 [PMID: [35486068](https://pubmed.ncbi.nlm.nih.gov/35486068/) DOI: [10.1167/iovs.63.4.27](https://doi.org/10.1167/iovs.63.4.27)]
- 152 **Usategui-Martín R**, Puertas-Neyra K, García-Gutiérrez MT, Fuentes M, Pastor JC, Fernandez-Bueno I. Human Mesenchymal Stem Cell Secretome Exhibits a Neuroprotective Effect over In Vitro Retinal Photoreceptor Degeneration. *Mol Ther Methods Clin Dev* 2020; **17**: 1155-1166 [PMID: [32514411](https://pubmed.ncbi.nlm.nih.gov/32514411/) DOI: [10.1016/j.omtm.2020.05.003](https://doi.org/10.1016/j.omtm.2020.05.003)]
- 153 **Xu C**, Lu H, Li F, Su G. Protein Expression Profile on Differentiation of Bone Marrow Mesenchymal Stem Cells into Retinal Ganglion-Like Cells. *J Comput Biol* 2020; **27**: 1329-1336 [PMID: [31841640](https://pubmed.ncbi.nlm.nih.gov/31841640/) DOI: [10.1089/cmb.2019.0024](https://doi.org/10.1089/cmb.2019.0024)]
- 154 **Dezfuly AR**, Safaee A, Amirpour N, Kazemi M, Ramezani A, Jafarinia M, Dehghani A, Salehi H. Therapeutic effects of human adipose mesenchymal stem cells and their paracrine agents on sodium iodate induced retinal degeneration in rats. *Life Sci* 2022; **300**: 120570 [PMID: [35469914](https://pubmed.ncbi.nlm.nih.gov/35469914/) DOI: [10.1016/j.lfs.2022.120570](https://doi.org/10.1016/j.lfs.2022.120570)]
- 155 **Alonso-Alonso ML**, Srivastava GK, Usategui-Martín R, García-Gutierrez MT, Pastor JC, Fernandez-Bueno I. Mesenchymal Stem Cell Secretome Enhancement by Nicotinamide and Vasoactive Intestinal Peptide: A New Therapeutic Approach for Retinal Degenerative Diseases. *Stem Cells Int* 2020; **2020**: 9463548 [PMID: [32676122](https://pubmed.ncbi.nlm.nih.gov/32676122/) DOI: [10.1155/2020/9463548](https://doi.org/10.1155/2020/9463548)]
- 156 **Jha KA**, Rasiah PK, Gentry J, Del Mar NA, Kumar R, Adebisi A, Reiner A, Gangaraju R. Mesenchymal stem cell secretome protects against oxidative stress-induced ocular blast visual pathologies. *Exp Eye Res* 2022; **215**: 108930 [PMID: [35016886](https://pubmed.ncbi.nlm.nih.gov/35016886/) DOI: [10.1016/j.exer.2022.108930](https://doi.org/10.1016/j.exer.2022.108930)]
- 157 **Rasiah PK**, Jha KA, Gentry J, Del Mar NA, Townsend T, Torgbe KE, Reiner A, Gangaraju R. A Long-Term Safety and Efficacy Report on Intravitreal Delivery of Adipose Stem Cells and Secretome on Visual Deficits After Traumatic Brain Injury. *Transl Vis Sci Technol* 2022; **11**: 1 [PMID: [36180031](https://pubmed.ncbi.nlm.nih.gov/36180031/) DOI: [10.1167/tvst.11.10.1](https://doi.org/10.1167/tvst.11.10.1)]
- 158 **Eiro N**, Sendon-Lago J, Cid S, Saa J, de Pablo N, Vega B, Bermudez MA, Perez-Fernandez R, Vizoso FJ. Conditioned Medium from Human Uterine Cervical Stem Cells Regulates Oxidative Stress and Angiogenesis of Retinal Pigment Epithelial Cells. *Ophthalmic Res* 2022; **65**: 556-565 [PMID: [35584686](https://pubmed.ncbi.nlm.nih.gov/35584686/) DOI: [10.1159/000524484](https://doi.org/10.1159/000524484)]
- 159 **Sanie-Jahromi F**, Nowroozadeh MH, Khodabandeh Z, Soheili ZS, Khajehahmadi Z, Emadi Z, Talebnejad MR. Effects of the secretome of human Wharton's jelly mesenchymal stem cells on the proliferation and apoptosis gene expression of the retinal pigmented epithelium. *Exp Eye Res* 2021; **205**: 108528 [PMID: [33662356](https://pubmed.ncbi.nlm.nih.gov/33662356/) DOI: [10.1016/j.exer.2021.108528](https://doi.org/10.1016/j.exer.2021.108528)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

