

Correction

Correction: Pata et al. Sclerobanding (Combined Rubber Band Ligation with 3% Polidocanol Foam Sclerotherapy) for the Treatment of Second- and Third-Degree Hemorrhoidal Disease: Feasibility and Short-Term Outcomes. *J. Clin. Med.* 2022, 11, 218

Francesco Pata ^{1,2,*}, Luigi Maria Bracchitta ³, Giancarlo D'Ambrosio ^{4,†} and Salvatore Bracchitta ^{5,†}

¹ Department of Surgery, Nicola Giannettasio Hospital, 87064 Corigliano-Rossano, Italy

² La Sapienza University, 00185 Rome, Italy

³ Primary Care Department, ATS Città Metropolitana, 20100 Milan, Italy; luigimaria.bracchitta@gmail.com

⁴ Department of General Surgery, Surgical Specialties and Organ Transplantation, La Sapienza University, 00161 Rome, Italy; giancarlo.dambrosio@uniroma1.it

⁵ Coloproctology Centre, Clinica del Mediterraneo, 97100 Ragusa, Italy; dott.salvatorebracchitta@gmail.com

* Correspondence: francesco.pata@gmail.com

† These authors share senior authorship.

1. Table Legend

In the original publication [1], there was a mistake in the legend for Table 1. “therange” should be changed to “the range”, and the sentence “Median is expressed as a median” should be changed to “Follow-up is expressed as a median”. The correct legend appears below.

Table 1. Demographic and postoperative outcomes of the patients included in the study. Age and ligations per procedure are expressed as a mean with the range between parentheses. Follow-up is expressed as a median.

2. Error in Table

In the original publication [1], there was a mistake in Table 1 as published. “Man” should be changed to “Men”, “56 (57.1%)” should be changed to “56 (57.7%)”, 42 (42.9%) should be changed to “41 (42.3%)”, “20 (20.4%)” should be changed to “20 (20.6%)”, and “78 (79.6%)” should be changed to “77 (79.4%)”. The corrected Table 1 appears below.

Table 1. Demographic and postoperative outcomes of the patients included in the study. Age and ligations per procedure are expressed as a mean with the range between parentheses. Follow-up is expressed as a median.

VARIABLE	N (%)
Patients	97
Men	56 (57.7%)
Women	41 (42.3%)
Age	52 years (20–84)
Goligher classification	
2nd degree	20 (20.6%)
3rd degree	77 (79.4%)
Ligations per procedure	2.7 (1–3)
Follow-up (median)	13 months (1–26)
Intraoperative complications	0 (0%)
30-day complications	4 (4.1%)
2nd degree	0 (0%)
3rd degree	4 (5.2%)
Readmission	0%
Mortality	0%



Citation: Pata, F.; Bracchitta, L.M.; D'Ambrosio, G.; Bracchitta, S. Correction: Pata et al. Sclerobanding (Combined Rubber Band Ligation with 3% Polidocanol Foam Sclerotherapy) for the Treatment of Second- and Third-Degree Hemorrhoidal Disease: Feasibility and Short-Term Outcomes. *J. Clin. Med.* 2022, 11, 218. *J. Clin. Med.* 2023, 12, 5159. <https://doi.org/10.3390/jcm12155159>

Received: 3 July 2023

Accepted: 20 July 2023

Published: 7 August 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

The authors state that the scientific conclusions are unaffected. This correction was approved by the Academic Editor. The original publication has also been updated.

Reference

1. Pata, F.; Bracchitta, L.M.; D'Ambrosio, G.; Bracchitta, S. Sclerobanding (Combined Rubber Band Ligation with 3% Polidocanol Foam Sclerotherapy) for the Treatment of Second- and Third-Degree Hemorrhoidal Disease: Feasibility and Short-Term Outcomes. *J. Clin. Med.* **2022**, *11*, 218. [[CrossRef](#)]

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.