

Corrigendum: 'Genome-wide association study of gastric cancer- and duodenal ulcer-derived *Helicobacter pylori* strains reveals discriminatory genetic variations and novel oncoprotein candidates'

Vo Phuoc Tuan^{1,2,†}, Koji Yahara^{3,*†}, Ho Dang Quy Dung¹, Tran Thanh Binh¹, Pham Huu Tung¹, Tran Dinh Tri¹, Ngo Phuong Minh Thuan¹, Vu Van Khien⁴, Tran Thi Huyen Trang⁵, Bui Hoang Phuc^{2,6}, Evariste Tshibangu-Kabamba², Takashi Matsumoto², Junko Akada², Rumiko Suzuki², Tadayoshi Okimoto⁷, Masaaki Kodama⁷, Kazunari Murakami⁷, Hirokazu Yano^{8,9,10}, Masaki Fukuyo^{8,9,11}, Noriko Takahashi^{8,9,12}, Mototsugu Kato^{13,14}, Shin Nishiumi^{15,16}, Takashi Azuma¹⁵, Yoshitoshi Ogura^{17,18}, Tetsuya Hayashi¹⁷, Atsushi Toyoda¹⁹, Ichizo Kobayashi^{8,9,12,20,*} and Yoshio Yamaoka^{2,21,*}

Microbial Genomics 2021;7:11, doi: 10.1099/mgen.0.000680

In the published version of this article, two errors of locus tags in Table 2 have been corrected as follows: HP0915 has been changed to HP0776, and HP_0915 has been changed to HP0915.

The authors apologise for any inconvenience caused.

Received 15 June 2023; Published 06 July 2023

Author affiliations: ¹Department of Endoscopy, Cho Ray Hospital, Ho Chi Minh, Vietnam; ²Department of Environmental and Preventive Medicine, Oita University Faculty of Medicine, Oita, Japan; ³Antimicrobial Resistance Research Center, National Institute of Infectious Diseases, Tokyo, Japan; ⁴Department of GI Endoscopy, 108 Central Hospital, Hanoi, Vietnam; ⁵Department of Molecular Biology, 108 Military Central Hospital, Hanoi, Vietnam; ⁶Department of Microbiology, Cho Ray Hospital, Ho Chi Minh, Vietnam; ⁷Department of Gastroenterology, Oita University Faculty of Medicine, Yufu, Oita, Japan; ⁸Department of Computational Biology and Medical Sciences, Graduate School of Frontier Sciences, University of Tokyo, Tokyo, Japan; ⁹Institute of Medical Science, University of Tokyo, Tokyo, Japan; ¹⁰Graduate School of Life Sciences, Tohoku University, Sendai, Japan; ¹¹Department of Molecular Oncology, Chiba University, Chiba, Japan; ¹²Department of Infectious Diseases, Kyorin University School of Medicine, Mitaka City, Tokyo, Japan; ¹³Division of Endoscopy, Hokkaido University Hospital, Sapporo, Hokkaido, Japan; ¹⁴Department of Gastroenterology, National Hospital Organization Hakodate Hospital, Hakodate, Hokkaido, Japan; ¹⁵Department of Gastroenterology, Graduate School of Medicine, Kobe University, Chuou-ku, Kobe, Hyogo, Japan; ¹⁶Department of Omics Medicine, Hyogo College of Medicine, Hyogo, Japan; ¹⁷Department of Bacteriology, Faculty of Medical Sciences, Kyushu University, Fukuoka, Japan; ¹⁸Division of Microbiology, Department of Infectious Medicine, Kurume University School of Medicine, Kurume, Fukuoka, Japan; ¹⁹Advanced Genomics Center, National Institute of Genetics, Shizuoka, Japan; ²⁰Research Center for Micro-Nano Technology, Hosei University, Tokyo, Japan; ²¹Department of Medicine, Gastroenterology Section, Baylor College of Medicine, Houston, TX, USA.

*Correspondence: Koji Yahara, k-yahara@nih.go.jp; Ichizo Kobayashi, zat14430@nifty.com; ikobaya@k.u-tokyo.ac.jp; Yoshio Yamaoka, yyamaoka@oita-u.ac.jp

†These authors contributed equally to this work

001062 © 2023 The Authors



This is an open-access article distributed under the terms of the Creative Commons Attribution License. The Microbiology Society waived the open access fees for this article.