Chronic Prostatitis Due to Yersinia pseudotuberculosis

Chronic bacterial prostatitis is commonly caused by *Escherichia coli* and is less frequently caused by *Klebsiella pneumoniae*, *Enterobacter*, *Proteus mirabilis*, and enterococci (6). Reported rare causes of prostatitis include *Candida* species, *Blastomyces dermatitidis*, *Histoplasma capsulatum*, *Mycobacterium tuberculosis*, and nontuberculous mycobacteria (8). We report the first case, to our knowledge, of chronic prostatitis due to *Yersinia pseudotuberculosis*.

A 55-year-old man had been suffering from recurrent urinary tract infections over the last 3 years. No prophylactic antibiotic treatment was administered. The pathogen was frequently *E. coli*, and in two cases it was *K. pneumoniae*. Findings from an intravenous pyelogram and abdominal ultrasound were unremarkable. In the last months preceding his referral to our center, he had three successive urinary tract infections due to *Y. pseudotuberculosis*.

Cultures of urethral urine, midstream urine, and prostatic secretions expressed by massage yielded heavy growth of *Y. pseudotuberculosis* from the last site and slight growth from the first two sites. Microscopic examination of the expressed prostatic secretions showed approximately 30 leukocytes per high-power field. Stool cultures did not grow *Yersinia* species. Prophylactic treatment with trimethoprim-sulfamethoxazole gave satisfactory results: the patient was free of symptoms, with negative urinary cultures.

The most common manifestation of *Y. pseudotuberculosis* infection in humans is mesenteric lymphadenitis accompanied by abdominal pain and fever (9). A septicemic form, occurring predominantly in patients with diabetes, hepatic cirrhosis, malignancy, and iron overload, has occasionally been described (5). Reported rare "atypical" manifestations of *Y. pseudotuberculosis* infection include interstitial nephritis (2), suppurative lymphadenitis (7), erythema nodosum, and nonsuppurative arthritis (1). To our knowledge, only one case of urinary tract

infection due to *Y. pseudotuberculosis* has been reported (3). Our case is the first reported case of prostatitis due to *Y. pseudotuberculosis*.

Epidemiologically, we could not identify any environmental risk factors that might be responsible for the patient's illness. Though several stool cultures did not yield *Yersinia* species, the feco-oral route is probably the main mode of acquisition, as with other enteric infections (4).

REFERENCES

- Butler, T. 1994. Yersinia infections: centennial of the discovery of the plague bacillus. Clin. Infect. Dis. 19:655–663.
- Cheong, H., E. Choi, I. Ha, H. Lee, and Y. Choi. 1995. Acute renal failure associated with Yersinia pseudotuberculosis infection. Nephron 70:319–323.
- Crchova, V., and C. Grondin. 1973. Urinary infection due to Yersinia pseudotuberculosis. Vie Med. Can. Fr. 2:3–5.
- Fukushima, H., M. Gomyada, S. Ishikura, T. Nishio, S. Moriki, J. Endo, S. Kaneko, and M. Tsubokura. 1989. Cat-contaminated environmental substances lead to *Yersinia pseudotuberculosis* infection in children. J. Clin. Microbiol. 27:2706–2709.
- Ljunberg, P., M. Valtonen, V. P. Harjola, S. S. Kaukoranta-Tolvanen, and M. Vaara. 1995. Report of four cases of *Yersinia pseudotuberculosis* septicemia and a literature review. Eur. J. Clin. Microbiol. Infect. Dis. 14:804–810.
- Meares, E. M., Jr. 1991. Prostatitis. Med. Clin. North Am. 75:405–424.
- Pouchot, J., V. Bortolotti, O. Sterkers, Y. Boussougant, and P. Vinceneux. 1995. Cervical suppurative lymphadenitis due to *Yersinia pseudotuberculosis*. Clin. Infect. Dis. 21:1063–1064.
- Sobel, J., and D. Kaye. 1995. Urinary tract infections, p. 662–690. In G. Mandell, T. Bennet, and R. Dolin (ed.), Principles and practice of infectious diseases, 4th ed. Churchill Livingstone, New York, N.Y.
- Tertti, R., R. Vuento, P. Mikkola, K. Granfors, A. Makela, and A. Toivanen. 1989. Clinical manifestations of Yersinia pseudotuberculosis infection in children. Eur. J. Microbiol. Infect. Dis. 8:587–591.

Bisharat Naiel, M.D. Raz Raul, M.D. Infectious Diseases Unit Ha'Emek Medical Center Afula 18101, Israel