



ORIGINAL WORK

Blood Pressure Affects the Early CT Perfusion Imaging in Patients with aSAH Reflecting Early Disturbed Autoregulation



Björn B. Hofmann^{1*} , Daniel M. Donaldson¹, Igor Fischer¹, Cihat Karadag¹, Milad Neyazi¹, Guilherme S. Piedade¹, Yousef Abusabha¹, Sajjad Muhammad¹, Christian Rubbert², Daniel Hänggi¹ and Kerim Beseoglu¹ 

© 2023 The Author(s)

Abstract

Background: Early computed tomography perfusion (CTP) is frequently used to predict delayed cerebral ischemia following aneurysmal subarachnoid hemorrhage (aSAH). However, the influence of blood pressure on CTP is currently controversial (HIMALAIA trial), which differs from our clinical observations. Therefore, we aimed to investigate the influence of blood pressure on early CTP imaging in patients with aSAH.

Methods: We retrospectively analyzed the mean transit time (MTT) of early CTP imaging within 24 h after bleeding prior to aneurysm occlusion with respect to blood pressure shortly before or after the examination in 134 patients. We correlated the cerebral blood flow with the cerebral perfusion pressure in the case of patients with intracranial pressure measurement. We performed a subgroup analysis of good-grade (World Federation of Neurosurgical Societies [WFNS] I–III), poor-grade (WFNS IV–V), and solely WFNS grade V aSAH patients.

Results: Mean arterial pressure (MAP) significantly correlated inversely with the mean MTT in early CTP imaging ($R = -0.18$, 95% confidence interval [CI] -0.34 to -0.01 , $p = 0.042$). Lower mean blood pressure was significantly associated with a higher mean MTT. Subgroup analysis revealed an increasing inverse correlation when comparing WFNS I–III ($R = -0.08$, 95% CI -0.31 to 0.16 , $p = 0.53$) patients with WFNS IV–V ($R = -0.2$, 95% CI -0.42 to 0.05 , $p = 0.12$) patients, without reaching statistical significance. However, if only patients with WFNS V are considered, a significant and even stronger correlation between MAP and MTT ($R = -0.4$, 95% CI -0.65 to 0.07 , $p = 0.02$) is observed. In patients with intracranial pressure monitoring, a stronger dependency of cerebral blood flow on cerebral perfusion pressure is observed for poor-grade patients compared with good-grade patients.

Conclusions: The inverse correlation between MAP and MTT in early CTP imaging, increasing with the severity of aSAH, suggests an increasing disturbance of cerebral autoregulation with the severity of early brain injury. Our results emphasize the importance of maintaining physiological blood pressure values in the early phase of aSAH and preventing hypotension, especially in patients with poor-grade aSAH.

Keywords: Aneurysmal subarachnoid hemorrhage, Early perfusion, Computed tomography, CT, Mean transit time, Blood pressure, Outcome

Introduction

In the past years, research into aneurysmal subarachnoid hemorrhage (aSAH) has been focused on the phase of delayed brain ischemia (DCI), with the occurrence of

*Correspondence: bjoern.hofmann@med.uni-duesseldorf.de

¹ Department of Neurosurgery, Medical Faculty and University Hospital Düsseldorf, Heinrich-Heine-University Düsseldorf, Düsseldorf, Germany
Full list of author information is available at the end of the article

vasospasm as the most prominent pathophysiological correlate [1]. Recently, it has been shown that early brain injury (EBI) is of great importance for the overall course of the disease [2]. Current studies indicate that the degree of cerebral hypoperfusion during the first hours after aSAH determines the severity of EBI, the occurrence of DCI, and the overall neurological outcome [2–7]. Several different perfusion imaging techniques and parameters have already been examined in the early evaluation of patients after bleeding, with computed tomography perfusion (CTP) imaging emerging as the most feasible technique [8–11]. As a result, CTP imaging is increasingly being used as a standard test during the initial and follow-up assessments of patients [12–14]. Particularly, the mean transit time (MTT) is a sensitive surrogate parameter to detect DCI-related perfusion impairment and is often used as a radiological follow-up parameter [8–11, 15–17]. In addition to its use as a follow-up parameter, the MTT of CTP imaging in the early phase after bleeding can be used as a predictor of DCI and neurologic outcome [6, 18–20]. Although an association between systemic blood pressure and cerebral perfusion in the light of altered cerebral autoregulation in EBI appears to be self-evident, robust data on the influence of blood pressure on the CTP in aSAH are lacking. In the DCI phase, so far only a trend for the cerebral blood flow (CBF) could be shown (e.g., in the HIMALAIA (Hypertension Induction in the Management of Aneurysmal subArachnoid haemorrhage with secondary Ischaemia) trial) [21, 22], but especially for the MTT, this influence is unclear, which differs with our clinical observations.

We hypothesize that the mean arterial pressure (MAP), possibly depending on the severity of SAH, correlates especially with the MTT in early perfusion imaging and that higher MAP is associated with shorter MTT, indicating better cerebral perfusion.

Methods

All procedures were in accordance with the ethical standards of the institutional committee and with the 1964 Helsinki declaration and its later amendments. For this type of study, written formal consent was not required. The local ethics committee approved the study (study ID: 5760R). Data will be made available on reasonable request. We prepared the article according to the Strengthening the Reporting of Observational Studies in Epidemiology guidelines.

Study Design and Study Population

A retrospective cohort study was performed for all patients with aSAH admitted to our tertiary neurovascular center from 2010 to 2015. The following inclusion criteria were applied: (1) SAH documented by an initial

nonenhanced CT, (2) CTP imaging within 24 h after admission, and (3) documented blood pressure values within, at most, 60 min before or after the CT measurement. Patients were excluded from the study based on the following criteria: nonaneurysmal SAHs and unknown onset of aSAH (determined by self-reported time of the headache event or third-party report of sudden neurologic deterioration or unconsciousness). If a prolonged headache was present or the time of bleeding could not be determined with certainty, the patients were excluded from the analysis as well as patients with unevaluable early CTP imaging (e.g., due to severe motion artifacts), patients with pretreated aneurysms or recurrent SAH at the time of early CT imaging, patients with a history of previous cranial neurosurgical or endovascular intervention, and patients with recorded mean blood pressure <30 mmHg or >130 mmHg at the time of interest.

The remaining patients were initially dichotomized for subgroup analysis based on World Federation of Neurosurgical Societies (WFNS) grade (I–III vs. IV–V). Additionally, patients with WFNS grade V were evaluated separately to map only the most severe hemorrhages. For each patient, the average MTT was calculated from the early CT perfusion and analyzed for correlation with the recorded MAP value. According to the common model of cerebral autoregulation, further evaluation of CBF versus cerebral perfusion pressure (CPP) was performed for patients with external ventricular drainage (EVD) or intracranial pressure probe at the time of blood pressure recording.

SAH Management

If an in-house native CT scan showed evidence of an aSAH, supplemental CT angiography and perfusion imaging were performed immediately in the same session. Patients in a peripheral hospital with evidence of SAH in a native CT scan and suspected aneurysm were transferred to our neurovascular center as soon as possible. They received CTP and CT angiography promptly on arrival. Treatment of patients with SAH in the early phase after hemorrhage was based on an in-house guideline following the applicable SAH guidelines [23]. In all patients, systolic blood pressure was maintained below 140 mm Hg before aneurysm occlusion. All patients with a Glasgow Coma Scale (GCS) \leq 12 (patients with a poor WFNS grade) were intubated and ventilated before aneurysm repair, and the partial pressure of carbon dioxide was maintained between 30 and 35 mm Hg [11]. An EVD was implanted in intubated patients and patients with a poor WFNS grade [11]. Further standardized therapy was performed, as extensively described in earlier

publications [11, 24]. Neurologic outcome was assessed by modified Rankin Scale (mRS) score at discharge and after 6 months.

CT Perfusion Analysis

Computed tomography perfusion scans within 24 h after hemorrhage and before aneurysm repair were considered as early CTP imaging. CTP was performed as previously described: CTP data were acquired with a multislice CT scanner (SOMATOM Volume Zoom, Definition Flash or AS; Siemens Erlangen, Germany, 80 kV, 120 mAs, two adjacent slices, 10-mm slice thickness, one image over 35 s) [4, 6, 11, 25]. Three seconds after starting the CT, a contrast agent (30 ml, 400 mg iodine/ml) immediately followed by a saline chaser (30 ml), were injected with a flow rate of 5 ml/s. A peripheral intravenous line of ≤ 18 gauge in a cubital vein or a high-flow central venous catheter was required for contrast agent administration. Slices were positioned at the level of the central parts of lateral ventricles, parallel to a plane through the orbital floor and the external auditory meatus, thereby sampling the anterior, middle, and posterior cerebral artery territories, as well as the anterior and posterior border zones [26]. Calculation of the parameter images (MTT, CBF, among cerebral blood volume and time-to-maximum (Tmax)) was performed by using STROKETOOL-CT (Digital Image Solutions, Frechen Germany), which processes data using singular value decomposition of the arterial input function matrix. We used the software Angiotux CT 2D (ECCET 2006, Beck A. Aurich V.) for standardized parameter value extraction from the different cortical brain regions. A 10-mm wide band is automatically delineated along the cortex as a region of interest (ROI), omitting outer cerebrospinal fluid spaces, the rostral falx cerebri, and the superior sagittal sinus. The automated definition of the ROI was reviewed, and potential deviations were corrected as needed. A running average spanning 10° of the ROI was computed in 2° steps for each perfusion parameter, yielding 180 measurements per parameter per CTP scan.

Blood Pressure Measurement

Vital parameters, such as blood pressure, are automatically documented and digitally stored with time stamps accurate to the second whenever a patient is connected to a stationary monitoring unit in either our emergency trauma room or neurosurgical intensive care unit. The blood pressure measurement is either a noninvasive measurement using an adequately sized blood pressure cuff on one of the upper arms or an invasive measurement from an arterial line (radial or brachial artery, occasionally femoral artery) with the transducer positioned at

heart level. The emergency trauma room of our tertiary care facility is equipped with a CT scanner capable of perfusion imaging, which allows for brief time intervals between blood pressure measurements and CTP imaging in primarily admitted patients.

If the patient was primarily admitted to a peripheral hospital, the patient was directly transferred to our neurosurgical intensive care unit. The aforementioned system documents no values during patient transport to the CT. Blood pressure values recorded directly before or after transport were recorded and checked individually for plausibility. Of these two values, the single blood pressure value used for analysis was the one that was closer in time to the CTP imaging. If the pretransport and post-transport blood pressure values had the same time interval to CTP imaging, the pretransport values were preferred, as these were considered less susceptible to interference. Systolic, diastolic, MAP (MAP = diastolic + $1/3$ [systolic-diastolic]), and CPP (if available) values with the shortest time interval to the CTP imaging were extracted from the database for evaluation.

Statistical Analysis

Patients' characteristics and clinical data were analyzed using RStudio, Version 1.3.1093 and Python Version 3.9.7. Our primary aim was to analyze the correlation of cerebral perfusion and blood pressure early after SAH. To achieve this, we calculated the average MTT over the whole cortex from the early CTP for each patient and correlated it with the recorded MAP by using the Pearson correlation coefficient, as the data are interval scaled. Normal distribution and the assumption of linearity were established by plotting the data in a histogram and scatter plot, respectively. Correlation between mean MTT and WFNS, Fisher score, and mRS was tested using the Kendall rank correlation coefficient because ordinal items in each scale were equal to or less than 6, according to Khamis et al. [27]. In the analysis of the 6-month mRS data, patients lost in the follow-up were excluded. Student's *t*-test was used to compare the mean of two groups. The significance level was set to 0.05. Dependency between CPP and CBF was modeled using linear regression.

Results

Patients

From the initial 702 patients with SAH during the study period, a total of 568 were excluded due to various factors, including those with missing pretreatment early CTP imaging, unknown onset of SAH, known history of cranial neurosurgical intervention or, in most cases, lack

Table 1 Patients' characteristics distinguished by subgroup

Parameter	Overall (N = 134)	WFNS I–III (n = 70)	WFNS IV–V (n = 64)	WFNS V (n = 34)
Age, mean (\pm SD) (yr)	55 (\pm 11)	53 (\pm 12)	58 (\pm 10)	56 (\pm 10)
Sex, n (%)				
Female	91 (68)	47 (67)	44 (69)	23 (68)
Male	43 (32)	23 (33)	20 (31)	11 (32)
Fisher grade, n (%)				
1	14 (10)	13 (19)	1 (2)	1 (3)
2	4 (3)	4 (6)	0 (0)	0 (0)
3	75 (56)	44 (63)	31 (48)	17 (50)
4	41 (31)	9 (13)	32 (50)	16 (47)
mRS discharge, n (%)				
0–2	54 (40)	45 (64)	9 (14)	4 (12)
3–5	56 (42)	21 (30)	35 (55)	17 (50)
6	24 (18)	4 (6)	20 (31)	13 (38)
mRS 6 months, n (%)				
0–2	74 (55)	50 (71)	24 (38)	11 (32)
3–5	27 (20)	11 (16)	16 (25)	7 (21)
6	24 (18)	4 (6)	20 (31)	14 (41)
Missing	9 (7)	5 (7)	4 (6)	2 (6)
Average MTT, mean (\pm SD) (ds)	40 (\pm 12)	38 (\pm 12)	42 (\pm 12)	40 (\pm 12)
MAP, mean (\pm SD) (mm Hg)	87 (\pm 21)	91 (\pm 19)	84 (\pm 18)	81 (\pm 18)

ds decisecond, MAP mean arterial pressure, MTT mean transit time, mRS modified Rankin scale, SD standard deviation, WFNS World Federation of Neurosurgical Societies

of timely blood pressure recording in relation to the time of CTP imaging. Thus, after applying the inclusion and exclusion criteria, 134 patients were included in the analysis. Patients' characteristics distinguished by subgroup are depicted in Table 1.

Blood Pressure Data

The MAP was 87 mm Hg (\pm 21; systolic [s]: 124 \pm 24; diastolic [d]): 65 \pm 19) for all patients, 91 mm Hg (\pm 19; s: 129 \pm 25; d: 66 \pm 17) for patients with WFNS I–III, 84 mm Hg (\pm 22; s: 119 \pm 21; d: 64 \pm 20) for patients with WFNS IV–V and 81 mm Hg (\pm 18; s: 116 \pm 21; d: 62 \pm 18) in the subgroup including only patients with WFNS V. In the subgroup comparison, there was a significant difference for systolic (95% confidence interval [CI] 1.38 to 17.1, $p=0.02$), but not for diastolic (95% CI -4.3 to 8.5, $p=0.5$) and mean blood pressure (95% CI -0.54 to 13.4, $p=0.07$) when comparing patients with WFNS I–III versus patients with WFNS IV–V (Fig. 1). The comparison of WFNS I–III versus WFNS V showed a significant difference in systolic (95% CI 3.8 to 22.5, $p=0.006$) and mean (95% CI 1.6 to 17, $p=0.02$), but not in diastolic blood pressure (95% CI -3.6 to 11.1, $p=0.3$; Fig. 1a).

The median time interval from the time of blood pressure measurement to CTP imaging was

9 min (mean = 5.8 \pm 24 min) for all patients, 9 min (mean = 5.8 \pm 22 min) for patients with WFNS I–III, 9 min (mean = 5.8 \pm 25 min) for patients with WFNS IV–V, and 5.5 min (mean = 3.3 \pm 23 min) for patients with WFNS V.


MTT of Early CTP Imaging

The mean MTT was 4.0 s (\pm 1.2) for all patients, 3.8 s (\pm 1.2) for patients with WFNS I–III, 4.2 s (\pm 1.2) for patients with WFNS IV–V, and 4 s (\pm 1.2) for patients with WFNS V. The difference between patients with WFNS grade I–III vs. WFNS grade IV–V was significant (CI, -8.3 to -0.2 ; $p=0.04$), whereas the difference between WFNS grade I–III patients versus WFNS V patients was not significant (CI, -7.2 to 2.8; $p=0.4$; Fig. 1b).

The MTT of early CTP in the whole study population significantly correlated with the initial WFNS ([T] = 0.17, $p < 0.01$), the mRS at discharge ([T] = 0.19 $p < 0.01$), and the mRS after 6 months ([T] = 0.17 $p < 0.01$), but not with the Fisher grade ([T] = 0.06 $p = 0.37$).

Correlation Between MTT of Early CTP Imaging and MAP

The mean MTT in early CTP correlated significantly with the MAP in the entire study population ($R = -0.18$, 95% CI -0.34 to 0.01, $p = 0.042$; Fig. 2a). Consequently,



a lower MAP was associated with higher MTT. When the dichotomized WFNS subgroups were analyzed, a nonsignificant correlation for patients with WFNS I–III ($R = -0.08$, 95% CI -0.31 to 0.16 , $p = 0.53$) was observed (Fig. 2b, left panel). Likewise, a nonsignificant but stronger trend for patients with WFNS IV–V ($R = -0.2$, 95% CI -0.42 to 0.05 , $p = 0.12$) was observed (Fig. 2b, middle panel). However, if only patients with WFNS V are considered, there is a significant and strong correlation of MAP with MTT ($R = -0.4$, 95% CI -0.65 to 0.07 , $p = 0.02$; Fig. 2b, right panel).

Correlation Between CBF of the Early CTP Imaging and CPP

When selecting patients with intracranial pressure measurement at the time of the documented blood pressure measurement, 16 good-grade patients and 40 poor-grade patients remained. The mean CBF in early CTP did not correlate significantly with the CPP in the subgroup of patients with good-grade SAH ($R = -0.47$, 95% CI -1.5 to 0.6 , $p = 0.35$; Fig. 3, blue). Poor-grade patients showed a stronger, yet also nonsignificant, correlation of CBF with CPP ($R = 1.25$, 95% CI -0.3 to 2.8 , $p = 0.12$; Fig. 3, red).




Fig. 2 MTT correlates with blood pressure; the correlation increases with the severity of aSAH. **A** MTT of early CTP correlated significant with the MAP in the entire study population ($R = -0.18$, 95% CI -0.34 to -0.01 , $p = 0.042$). **B** The dichotomized subgroups showed a nonsignificant correlation ($R = -0.08$, 95% CI -0.31 to 0.16 , $p = 0.53$; left panel) for the good WFNS grade patients and for the poor WFNS grade patients ($R = -0.2$, 95% CI -0.42 to 0.05 , $p = 0.12$; middle panel). However, when only patients with WFNS V are considered, there is a significant and strong correlation of MAP with MTT ($R = -0.4$, 95% CI -0.65 to -0.07 , $p = 0.02$) (right panel). aSAH, aneurysmal subarachnoid hemorrhage, CI, confidence interval, CTP, computed tomography perfusion, ds, decisecond, MAP, mean arterial pressure, MTT, mean transit time, WFNS, World Federation of Neurosurgical Societies

Discussion

This retrospective study in patients with aSAH has three main findings:

1. MTT of early CTP inversely correlates with the MAP when analyzing the whole study population. A high MTT is associated with low mean blood pressure.




Fig. 3 CBF tends to correlate with CPP in poor-grade patients. The CBF of early CTP did not correlate significantly with the CPP in the subgroup of patients with good WFNS grade aSAH ($n = 16$; $R = -0.47$, 95% CI -1.5 to 0.6 , $p = 0.35$; Fig. 3, blue). Poor WFNS grade patients ($n = 40$) showed a stronger, yet also nonsignificant, correlation of CBF with CPP ($R = 1.25$, 95% CI -0.3 to 2.8 , $p = 0.12$; Fig. 3, red). Outliers were excluded for better graphical presentation. aSAH, aneurysmal subarachnoid hemorrhage, CBF, cerebral blood flow, CI, confidence interval, CPP, cerebral perfusion pressure, CTP, computed tomography perfusion, WFNS, World Federation of Neurosurgical Societies (Color figure online)

2. The more severe the aSAH, as measured by the WFNS grade, the stronger the inverse correlation of the MAP with the MTT.
3. There is a stronger correlation between CBF and CPP in patients with poor-grade aSAH than in patients with good-grade aSAH.

The MTT of CTP imaging is an important routine follow-up parameter in the monitoring of patients with aSAH. For example, in our neurovascular center, according to an in-house treatment protocol, if the MTT threshold of 41 deciseconds is exceeded, which, in our opinion, is a sign of a critical cerebral perfusion situation based on a large number of previous observations regarding MTT in patients with aSAH [25, 26, 28], a multimodal therapy is usually initiated. In addition, the MTT of CTP imaging is increasingly used in the phase of EBI as an early surrogate parameter of patient outcome [6, 15–18, 20, 29, 30]. Although an association between systemic blood pressure and cerebral perfusion in the light of altered cerebral autoregulation in EBI appears to be self-evident, robust data on this topic are lacking [21, 31]. Until this study, there has been no other study directly designed to investigate the association of blood pressure with CTP imaging in patients with aSAH. Based on our clinical observation, we hypothesized that the MAP correlates with the MTT of early CTP imaging, such that a lower MAP is associated with a longer MTT.

The results of the present study support our hypothesis, clearly showing an inverse correlation of MAP with MTT of early CTP in the EBI phase. This is already evident in the initial analysis of the entire study population (Fig. 2a). Although, in our view, the increasing inverse correlation with severity of aSAH is an even more striking observation (Fig. 2b). The inverse correlation increases in the study from patients with good-grade aSAH (WFNS I–III), with $R = -0.08$, to patients with poor-grade aSAH (WFNS IV–V), with $R = -0.2$ (Fig. 2b). However, this correlation does not reach the significance level. Hereby, we attribute the lack of significance of this observation to the reduced sample size caused by dichotomization. We assume that the inverse correlation would also become significant with a larger study population for these subgroups. In this context, it is even more impressive that a significant and even more substantial inverse correlation of MAP with MTT with $R = -0.4$ is found when only the patients with WFNS V are analyzed, resulting in an even smaller group size (Fig. 2b, right panel). Because, to our knowledge, the relationship between blood pressure and CTP imaging in the phase of EBI was not examined yet, our results can only be compared with observations related to the phase of DCI. The results of this study therefore extend the results of the prospective HIMALAIA (Hypertension Induction in the Management of Aneurysmal subArachnoid haemorrhage with secondary Ischaemia) trial, which could only show a nonsignificant effect of blood pressure on CBF of perfusion imaging in the DCI phase [21]. However, it should be noted that these results are only a side evaluation of a prospective trial aimed to compare the outcome of patients treated with and without induced hypertension after SAH [21, 22]. Furthermore, mainly the CBF of the CTP scans was the focus of this side evaluation [21]. Specific statements on the MTT, which, in our opinion, is of greater benefit in the monitoring of patients with aSAH, are not available [21]. It should also be noted that the study based its conclusion on data from only 13 patients with induced hypertension versus 12 patients without induced hypertension, as it was terminated prematurely because of difficult recruitment [21, 22].

Furthermore, the results of this study are well aligned with new insights into the pathophysiological processes after aSAH. Recent studies show that aSAH is followed by disturbed autoregulation affecting cerebral perfusion [32–36]. This has already been demonstrated several times in animal models and humans, in which the impairment of autoregulation has also been observed very early post bleeding in the phase of EBI [33–36]. Theoretically, impaired autoregulation is defined as the inability to maintain constant cerebral perfusion when changes in blood pressure occur [37, 38]. Consequently, a more

powerful direct influence of blood pressure on cerebral perfusion is assumed. However, proof of the association of blood pressure with brain perfusion in CTP imaging has not been demonstrated in patients with aSAH until today. Our study adds relevant data to this discussion.

In our view, the results of this study underline the assumed effects of disturbed autoregulation on blood pressure. Blood pressure differences, which generally do not significantly affect cerebral perfusion, have an uninhibited, more direct influence on cerebral perfusion. Therefore, a lower blood pressure correlates with a higher MTT, i.e., a surrogate for poor cerebral perfusion. Furthermore, the results of the subgroup analysis also fit this theory. Our study shows that the relationship between MAP and MTT in early CTP imaging increases with the severity of aSAH, as measured by the WFNS grade, as already discussed above (Fig. 2). The WFNS as a neurological classification scale hereby most likely reflects the severity of the EBI and consequently the severity of disturbance of cerebral autoregulation. Therefore, in patients with poor-grade aSAH, changes in MAP lead more directly to changes in MTT. Particularly in the case of the most severely affected patients with aSAH, in whom intracranial circulatory arrest can occur as described before [39, 40], there may be parallels to the cerebral autoregulatory dysfunction observed after cardiac arrest [41, 42]. In contrast, patients with good-grade aSAH are less affected and display a less extensive EBI, reflected in a weaker correlation between the MAP and the MTT.

To support this theory of impaired autoregulation, we examined the common model of cerebral autoregulation and investigated the correlation of CBF with CPP in our patient cohort [38, 43, 44]. Even if the significance level was not reached in this analysis due to the even smaller sample size, it shows a stronger correlation for the poor-grade patients than the good-grade patients. This reflects the more severe disturbance of cerebral autoregulation in the poor-grade patients and supports our assumption.

In summary, the data from this study, for the first time, not only show the association of blood pressure with brain perfusion as measured by MTT in the early phase after SAH but also show the increase of dependence regarding the severity of hemorrhage measured by WFNS grade. Because numerous studies have already demonstrated that prolonged MTT in the EBI phase is associated with a poor outcome [6, 29], it can be deduced that initial blood pressure management is of high importance, especially for patients with poor-grade aSAH. This should by no means lead to induced hypertension in untreated aneurysms in the phase of EBI, but attention should be paid to maintaining physiological blood pressure values.

Some study limitations are worth noting:

1. Data were retrospectively collected, and only patients with early CTP were included. It is unclear whether there was a selection bias in the choice of patients who received CTP imaging prior to the aneurysm repair. Nevertheless, the study population is relatively large for a single-center study. It is overall well-balanced, for example, regarding patients with good-grade and poor-grade SAH according to their WFNS score. We believe it is unlikely that our inclusion/exclusion criteria introduced a relevant bias.
2. Because of technical constraints, the blood pressure values are measured with a certain time offset to the early CTP imaging. Therefore, no digitally documented value is available from the exact imaging time. Especially for patients primarily admitted to an external hospital and then directly transferred to our intensive care unit, blood pressure values during transport and imaging are not recorded digitally. This results in a blind spot at the time of CTP imaging. This is one major limitation of this study. Yet, we aim for stable blood pressure without blood pressure peaks or dips during treatment. We therefore consider the blood pressure values recorded shortly before or after imaging to be representative. For subsequent prospective studies, however, a purely arterial blood pressure measurement with a recording of the blood pressure at the imaging time should be performed. The proven possibility of minimal variations in the CPP values due to changes in the patient's position (30° torso elevation vs. supine in the CT) should also be taken into account, although in our study, this positional change most likely affects all patients to the same extent [45]. Furthermore, influences on the results of this study by external factors such as stimulation of patients on transport to the CT scanner affecting MAP and intracranial pressure should also be considered.
3. Patients with an inserted EVD are, in most cases, sedated and often also ventilated. The influence of sedation on the result of this study cannot be ruled out. Yet, an influence of the intracranial pressure on the results of the study is considered unlikely. Patients without EVD are conscious and a normal intracranial pressure can therefore be assumed, and in patients with EVD the intracranial pressure is normalized to a physiological level. Further subgroup analysis of dichotomized subgroups of patients with versus without EVD at the imaging time is attached in the supplementary material.
4. Data from CTP imaging are difficult to compare one-to-one between different setups due to different CTP

protocols and postprocessing. In addition, diseases affecting the microvasculature such as atherosclerosis, hypertension, or diabetes may influence the MTT. Additionally, metabolic state, use of sedatives, and concomitant medication may potentially impact the MTT, as well. Therefore, the effect of these factors on the results of this study remains unclear. Furthermore, it is unclear whether there are differences in the MTT when blood pressure values are present spontaneously or are influenced by medication. Similarly, it is not clear whether mean MTT alone or a combination of different parameters of CTP best represents the relevant pathophysiological changes in patients with aSAH, which should be taken into account when interpreting the data.

Conclusions

In the present cohort of 134 patients with aSAH, MTT of early CTP inversely correlates with the MAP, indicating early impairment of cerebral autoregulation in the phase of EBI. A low MAP is associated with a prolonged MTT. The subgroup analysis shows that the inverse correlation becomes stronger with the severity of aSAH as measured by the WFNS, whereas, in particular, WFNS grade V patients demonstrated a significant, strong, inverse correlation of MAP and MTT.

These results emphasize the importance of maintaining physiological blood pressure values in the EBI phase and preventing hypotension, especially in patients with poor-grade aSAH. Further prospective studies should seek to confirm this influence of MAP on MTT in EBI and should verify whether the early blood pressure values correlate with the outcome of the patients post aSAH.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1007/s12028-023-01683-8>.

Abbreviations

aSAH: Aneurysmal subarachnoid hemorrhage; CBF: Cerebral blood flow; CPP: Cerebral perfusion pressure; CTA: Computed tomography angiography; CTP: Computed tomography perfusion, DCI: delayed cerebral ischemia; EBI: Early brain injury; EVD: External ventricular drainage; ICP: Intracranial pressure; MAP: Mean Arterial Pressure; MRI: Magnetic Resonance Imaging; mRS: Modified Rankin Scale; MTT: Mean Transit Time; RA: Resuscitation Area; SVD: Singular value decomposition.

Author details

¹ Department of Neurosurgery, Medical Faculty and University Hospital Düsseldorf, Heinrich-Heine-University Düsseldorf, Düsseldorf, Germany.

² Department of Diagnostic and Interventional Radiology, Medical Faculty and University Hospital Düsseldorf, Heinrich-Heine-University Düsseldorf, Düsseldorf, Germany.

Author Contributions

BBH and KB conceived of the presented idea and designed the study. BBH, KB, and CR collected and managed the data. DMD, IF, and BBH performed the statistical analysis. All authors discussed the results. BBH drafted the initial article and designed the figures. DMD, IF, CK, MN, GSP, YA, SM, CR, DH, and KB

revised the article and figures. BBH, DH, and KB supervised the project. The final manuscript was approved by all authors.

Source of Support

Open Access funding enabled and organized by Projekt DEAL. No funding was received for this research.

Conflict of interest

The authors declare that they have no conflicts of interest.

Ethical Approval/Informed Consent

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional committee and with the 1964 Helsinki declaration and its later amendments. For this type of study, formal consent is not required. The study was approved by the local ethics committee (study ID: 5760R).

Open Access

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 7 October 2022 Accepted: 26 January 2023

Published: 17 February 2023

References

1. Macdonald RL. Delayed neurological deterioration after subarachnoid haemorrhage. *Nat Rev Neurol*. 2014;10:44–58.
2. van Lieshout JH, Dibue-Adjei M, Cornelius JF, et al. An introduction to the pathophysiology of aneurysmal subarachnoid hemorrhage. *Neurosurg Rev*. 2018;41:917–30.
3. Neulen A, Pantel T, Kosterhon M, et al. Neutrophils mediate early cerebral cortical hypoperfusion in a murine model of subarachnoid haemorrhage. *Sci Rep*. 2019;9:8460.
4. Kamp MA, Heiroth HJ, Beseoglu K, Turowski B, Steiger HJ, Hanggi D. Early CT perfusion measurement after aneurysmal subarachnoid hemorrhage: a screening method to predict outcome? *Acta Neurochir Suppl*. 2012;114:329–32.
5. Schubert GA, Seiz M, Hegewald AA, Manville J, Thomé C. Acute hypoperfusion immediately after subarachnoid hemorrhage: a xenon contrast-enhanced CT study. *J Neurotrauma*. 2009;26:2225–31.
6. Etminan N, Beseoglu K, Heiroth HJ, Turowski B, Steiger HJ, Hanggi D. Early perfusion computerized tomography imaging as a radiographic surrogate for delayed cerebral ischemia and functional outcome after subarachnoid hemorrhage. *Stroke*. 2013;44:1260–6.
7. Tsuang F-Y, Chen J-Y, Lee C-W, et al. Risk profile of patients with poor-grade aneurysmal subarachnoid hemorrhage using early perfusion computed tomography. *World Neurosurg*. 2012;78:455–61.
8. Dankbaar JW, de Rooij NK, Rijdsdijk M, et al. Diagnostic threshold values of cerebral perfusion measured with computed tomography for delayed cerebral ischemia after aneurysmal subarachnoid hemorrhage. *Stroke*. 2010;41:1927–32.
9. Mir DIA, Gupta A, Dunning A, et al. CT perfusion for detection of delayed cerebral ischemia in aneurysmal subarachnoid hemorrhage: a systematic review and meta-analysis. *Am J Neuroradiol*. 2014;35:866.

10. Sanelli PC, Ugorec I, Johnson CE, et al. Using quantitative CT perfusion for evaluation of delayed cerebral ischemia following aneurysmal subarachnoid hemorrhage. *Am J Neuroradiol.* 2011;32:2047.
11. Hofmann BB, Fischer I, Engel A, et al. MTT heterogeneity in perfusion CT imaging as a predictor of outcome after aneurysmal SAH. *AJNR Am J Neuroradiol.* 2021;43:1387–95. <https://doi.org/10.3174/ajnr.A7169>
12. Rabinstein AA, Lanzino G, Wijdicks EFM. Multidisciplinary management and emerging therapeutic strategies in aneurysmal subarachnoid haemorrhage. *Lancet Neurol.* 2010;9:504–19.
13. Wintermark M, Ko NU, Smith WS, Liu S, Higashida RT, Dillon WP. Vasospasm after subarachnoid hemorrhage: utility of perfusion CT and CT angiography on diagnosis and management. *Am J Neuroradiol.* 2006;27:26.
14. Sviri GE, Britz GW, Lewis DH, Newell DW, Zaaroor M, Cohen W. Dynamic perfusion computed tomography in the diagnosis of cerebral vasospasm. *Neurosurgery.* 2006;59:319–25.
15. Shi D, Jin D, Cai W, et al. Serial low-dose quantitative CT perfusion for the evaluation of delayed cerebral ischaemia following aneurysmal subarachnoid haemorrhage. *Clin Radiol.* 2020;75:131–9.
16. Bacigaluppi S, Zona G, Secci F, et al. Diagnosis of cerebral vasospasm and risk of delayed cerebral ischemia related to aneurysmal subarachnoid haemorrhage: an overview of available tools. *Neurosurg Rev.* 2015;38:603–18.
17. Kanazawa R, Kato M, Ishikawa K, Eguchi T, Teramoto A. Convenience of the computed tomography perfusion method for cerebral vasospasm detection after subarachnoid hemorrhage. *Surg Neurol.* 2007;67:604–11.
18. Rubbert C, Patil KR, Beseoglu K, et al. Prediction of outcome after aneurysmal subarachnoid haemorrhage using data from patient admission. *Eur Radiol.* 2018;28:4949–58.
19. Rodriguez-Régent C, Hafsa M, Turc G, et al. Early quantitative CT perfusion parameters variation for prediction of delayed cerebral ischemia following aneurysmal subarachnoid hemorrhage. *Eur Radiol.* 2016;26:2956–63.
20. Beseoglu K, Etminan N, Hanggi D. The value of perfusion computed tomography (PCT) imaging after aneurysmal subarachnoid hemorrhage: a review of the current data. *Acta Neurochir Suppl.* 2015;120:35–8.
21. Gathier CS, Dankbaar JW, van der Jagt M, et al. Effects of induced hypertension on cerebral perfusion in delayed cerebral ischemia after aneurysmal subarachnoid hemorrhage: a randomized clinical trial. *Stroke.* 2015;46:3277–81.
22. Gathier CS, van den Bergh WM, van der Jagt M, et al. Induced hypertension for delayed cerebral ischemia after aneurysmal subarachnoid hemorrhage: a randomized clinical trial. *Stroke.* 2018;49:76–83.
23. Connolly ES Jr, Rabinstein AA, Carhuapoma JR, et al. Guidelines for the management of aneurysmal subarachnoid hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* 2012;43:1711–37.
24. Hofmann BB, Rubbert C, Turowski B, Hänggi D, Muhammad S. Treatment of unique bilateral distal fusiform superior cerebellar artery aneurysms with mini-flow diverter device implantation: case report. *J Neurol Surg Part A Cent Eur Neurosurg.* 2021. <https://doi.org/10.1055/s-0041-1739212>
25. Turowski B, Haenggi D, Wittsack J, Beck A, Moedder U [cerebral perfusion computerized tomography in vasospasm after subarachnoid hemorrhage: diagnostic value of MTT]. *RoFo: Fortschr Gebiete Röntgenstrahlen Nukl.* 2007;179:847–54.
26. Turowski B, Haenggi D, Wittsack HJ, Beck A, Aurich V [computerized analysis of brain perfusion parameter images]. *RoFo: Fortschr Gebiete Röntgenstrahlen Nukl.* 2007;179:525–9.
27. Khamis H. Measures of association: how to choose? *J Diagn Med Sonogr.* 2008;24:155–62.
28. Hänggi D, Turowski B, Beseoglu K, Yong M, Steiger HJ. Intra-arterial nimodipine for severe cerebral vasospasm after aneurysmal subarachnoid hemorrhage: influence on clinical course and cerebral perfusion. *Am J Neuroradiol.* 2008;29:1053.
29. Beseoglu K, Etminan N, Hänggi D. The value of perfusion computed tomography (PCT) imaging after aneurysmal subarachnoid hemorrhage: a review of the current data. In: Fandino J, Marbacher S, Fathi A-R, Muroi C, Keller E, editors. *Neurovascular events after subarachnoid hemorrhage: towards experimental and clinical standardisation.* Cham: Springer International Publishing; 2015. p. 35–8.
30. Murphy A, Manoel AL, Burgers K, et al. Early CT perfusion changes and blood-brain barrier permeability after aneurysmal subarachnoid hemorrhage. *Neuroradiology.* 2015;57:767–73.
31. Murphy A, de Oliveira Manoel AL, Macdonald RL, et al. Changes in cerebral perfusion with induced hypertension in aneurysmal subarachnoid hemorrhage: a pilot and feasibility study. *Neurocrit Care.* 2017;27:3–10.
32. Francoeur CL, Mayer SA. Management of delayed cerebral ischemia after subarachnoid hemorrhage. *Crit Care.* 2016;20:277.
33. Budohoski KP, Czosnyka M, Kirkpatrick PJ, Smielewski P, Steiner LA, Pickard JD. Clinical relevance of cerebral autoregulation following subarachnoid haemorrhage. *Nat Rev Neurol.* 2013;9:152–63.
34. Jaeger M, Schuhmann MU, Soehle M, Nagel C, Meixensberger J. Continuous monitoring of cerebrovascular autoregulation after subarachnoid hemorrhage by brain tissue oxygen pressure reactivity and its relation to delayed cerebral infarction. *Stroke.* 2007;38:981–6.
35. Rasmussen G, Hauerberg J, Waldemar G, Gjerris F, Juhler M. Cerebral blood flow autoregulation in experimental subarachnoid haemorrhage in rat. *Acta Neurochir.* 1992;119:128–33.
36. Hattinger E, Blasel S, Dettmann E, et al. Perfusion-weighted MRI to evaluate cerebral autoregulation in aneurysmal subarachnoid haemorrhage. *Neuroradiology.* 2008;50:929.
37. Aaslid R, Lindegaard KF, Sorteberg W, Nornes H. Cerebral autoregulation dynamics in humans. *Stroke.* 1989;20:45–52.
38. Armstead WM. Cerebral blood flow autoregulation and dysautoregulation. *Anesthesiol Clin.* 2016;34:465–77.
39. Eng CC, Lam AM, Byrd S, Newell DW. The diagnosis and management of a perianesthetic cerebral aneurysmal rupture aided with transcranial Doppler ultrasonography. *Anesthesiology.* 1993;78:191–4.
40. Hassler W, Steinmetz H, Pirschel J. Transcranial Doppler study of intracranial circulatory arrest. *J Neurosurg.* 1989;71:195–201.
41. Sundgreen C, Larsen FS, Herzog TM, Knudsen GM, Boesgaard S, Aldershvile J. Autoregulation of cerebral blood flow in patients resuscitated from cardiac arrest. *Stroke.* 2001;32:128–32.
42. Pham P, Bindra J, Chuan A, Jaeger M, Aneman A. Are changes in cerebrovascular autoregulation following cardiac arrest associated with neurological outcome? Results of a pilot study. *Resuscitation.* 2015;96:192–8.
43. Dewey RC, Pieper HP, Hunt WE. Experimental cerebral hemodynamics. Vasomotor tone, critical closing pressure, and vascular bed resistance. *J Neurosurg.* 1974;41:597–606.
44. Rose JC, Mayer SA. Optimizing blood pressure in neurological emergencies. *Neurocrit Care.* 2004;1:287–99.
45. Rosner MJ, Coley IB. Cerebral perfusion pressure, intracranial pressure, and head elevation. *J Neurosurg.* 1986;65:636–41.