

Research Article

Extender Supplementation with Glutathione (GSH) and Taurine Improves *In Vitro* Sperm Quality and Antioxidant Status of New Zealand Rabbits during Chilled Storage for up to 72 hours

Mohamed F. F. Bayomy,^{1,2} Sobhy E. Hassab El-Nabi,¹ Tahany A. El Kassas,¹ Zeinab I. Attia,³ Ayman M. Saeed,⁴ Heba S. A. Taha,⁵ Mahmoud Alagawany ,⁶ Livio Galosi ,⁷ Lucia Biagini,⁷ and Seham El-Kassas⁸

¹Zoology Department, Faculty of Science, Menoufia University, Shibin Al Kawm, Egypt

²General Biology Department, Center of the Basic Sciences, Misr University for Science and Technology (MUST), 6th of October City, Egypt

³Zoology Department, Faculty of Science, Tanta University, Tanta, Egypt

⁴Biotechnology Department, Animal Production Research Institute, Giza, Egypt

⁵Department of Genetics, Faculty of Agriculture, Zagazig University, Zagazig, Egypt

⁶Poultry Department, Agriculture Faculty, Zagazig University, Zagazig 44519, Egypt

⁷School of Bioscience and Veterinary Medicine, University of Camerino, Matelica 62024, Italy

⁸Animal, Poultry, and Fish Breeding and Production, Department of Animal Wealth Development, Faculty of Veterinary Medicine, Kafrelsheikh University, Kafr el-Sheikh, Egypt

Correspondence should be addressed to Livio Galosi; livio.galosi@unicam.it

Received 21 March 2023; Revised 26 August 2023; Accepted 30 August 2023; Published 12 September 2023

Academic Editor: Juan G. Chediack

Copyright © 2023 Mohamed F. F. Bayomy et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This study assessed the influence of supplementing the rabbit semen extender with various concentrations of glutathione (GSH) and taurine at 24, 48, and 72 h postchilling at 5°C. Semen samples were collected from 20 New Zealand bucks, and ejaculates with standard color, motility (>85%), about 0.5 mL volume, and $\sim 400 \times 10^6/\text{mL}$ concentration were used and diluted with extenders supplemented with 0.5, 1, and 2 mM of GSH and 1, 5, and 10 mM of taurine and chilled at 5°C. Nonsupplemented samples were used as a control. Sperm's progressive motility, acrosome reaction, and extracellular oxidative stress biomarkers such as MDA contents and GPx, SOD, and CAT concentrations and intracellular transcriptomic levels of *SOD* and *CAT* genes were assessed. GSH and taurine supplementation improved the sperm's kinetics by reducing cooling-associated stress, which was ascertained by lowering MDA concentration and increasing SOD, CAT, and GPx concentrations ($P < 0.05$). Increasing the levels of antioxidant enzymes in the extender was due to the increasing mRNA copies of the *SOD* and *CAT* genes ($P < 0.05$). Furthermore, GSH and taurine maintained the fructose levels in the extender and lowered the GPT levels, which implies sperm membrane stability is maintained through GSH and taurine supplementation. GSH and taurine supplementation to the extender had protective influences on the *in vitro* rabbit semen quality during chilled storage for up to 72 h, which were remarkable with increasing supplementation dose and cooling time at 5°C.

1. Introduction

With the increase in human population and the decrease in resources, especially with the global climatic changes, ensuring food and nutritional security is a fundamental

requirement [1–5]. Thus, increasing animal productivity is necessary, and this goal can be achieved by increasing the number of animals, breeds, and strains produced. Artificial insemination (AI) has been proven to be an efficient tool to improve the animal's genetic potential, increase its yielding

capacity, and selectively increase genetic gain through increasing male selection intensity [6].

Rabbit is an indispensable animal species to meet the nutritional requirements of human beings and for biomedical research. Rabbits are generally used as valuable meat sources due to their high feed conversion rate and fur production, and they are also used as pets [7]. Besides, in biomedical research, this species is for antibody production and as a model for human diseases such as hypercholesterolemia and atherosclerosis [8]. Moreover, gene-modified rabbits were usefully developed and used for human disease research using genetic engineering applications such as gene editing tools, gene knockout, and conventional pronuclear microinjection techniques [9, 10]. Thus, it is vital to maintain and increase the production of rabbit strains for these many purposes. Preserving rabbit strains can be achieved by repeated breeding, but numerous animals must avoid inbreeding depression [11, 12]. Another way is the cryopreservation of ova and embryos in liquid nitrogen, which requires scarifying animals, professional surgical skills, and a high cost [10]. Therefore, cryopreservation of semen by repeated collection and preservation in liquid nitrogen is more effective [10], but most rabbit production relies on using fresh or shortly stored semen for routine AI [13]. However, the oxidative stress caused by the excessive production of reactive oxygen species (ROS) [14–20], and the consequent cellular, cytoplasm, and genome damage alter sperm quality [21]. Thus, the effectiveness of AI using cooling semen depends on the ability of the extenders to provide a suitable environment for spermatozoa metabolism.

AI programs in rabbits are currently performed with fresh diluted semen collected within 6–18 hours, and bucks are typically kept on the same farm [22]. Improving semen storage capacity would allow for a longer interval between semen collection and female insemination, thereby improving AI performance and allowing the existence of farms without males [23, 24]. However, storing rabbit sperm for more than 24 to 48 hours decreases fertility [25–27]. Therefore, one of the primary goals is to extend the interval of liquid semen storage beyond 48 hours or to freeze the semen. Although satisfactory in-vitro results on sperm quality and fertilizing ability have been obtained with frozen semen in the last decade [28–30], the limited survival of sperm after freezing is a major drawback for the widespread use of frozen semen in AI programs. Semen cryopreservation induces damage to all sperm compartments related to the intra- and extra-ice formation and causes a ROS excess generated by dysfunctional mitochondria, responsible for sperm membrane injuries [10, 31–34]. These damages are responsible for the loss of sperm motility, viability, acrosomal, and DNA integrity which decreases the fertilizing capacity of frozen-thawed sperm [35]. Moreover, the freezing technique is more expensive and requires more equipment and technical expertise than the chilling technique. AI with cryopreserved rabbit sperm finds applications in experimental or genetic resource banking. Under this view, chilling semen between 24 and 48 hours is still essential and is applicable, as this technique is simple and inexpensive,

requiring no special equipment. Accordingly, more attempts are needed to improve semen extenders and storage conditions to prolong the time stored semen can maintain its functional status.

Many approaches have been utilized to preserve and improve semen quality and sperm viability, including the supplementation of antioxidant substances, such as glutathione (GSH), taurine, lycopene, cysteine, and glutamine directly to the animals or in semen extenders to increase the antioxidant abilities and lower the impacts of oxidative stress [36, 37]. GSH and taurine are the most common antioxidants, which have markedly improved the antioxidant characteristics of semen extenders. GSH maintains semen quality by keeping better integrity of sperm's nuclear and plasma membranes. Besides, it improves the fertilizing ability of spermatozoa following the freezing-thawing process [38]. The improving effects of GSH are due to its property as an electron donor to reduce the H_2O_2 production plus its oxidation to glutathione disulfide (GSSG). Moreover, the thiol group content of the amino acid cysteine in GSH penetrates the spermatozoa's plasma membrane enhancing the intracellular biosynthesis of GSH, which protects sperm's DNA, proteins, and its membrane-lipids content via scavenging of ROS [39]. The sulfonic amino acid taurine is a nonenzymatic scavenger that also protects the spermatozoa by inhibiting lipid peroxidation and protecting sperm cells against the accumulation of ROS [37]. These antioxidant properties of GSH and taurine may be allied to modification in the levels of antioxidant enzymes such as sodium oxide dismutase (SOD) and catalase (CAT) which in turn may be linked with the changes in the transcription levels of their genes. Thus, this investigation examined the mRNA levels of SOD and CAT genes. Additionally, taurine enhances the motility of spermatozoa by protecting the integrity of the plasma, nuclear, and mitochondrial membranes and maintaining the structure of the cytoskeleton of the flagella of spermatozoa [40].

In the previous study of Ahmad et al, the exogenous antioxidants supplementation of GSH at 0, 1, 2, 4, and 8 mM to rabbit semen extender stored at 5°C for 24 h improved the semen attributes, especially the lower doses [41]. However, to the best of our knowledge, no reports on the GSH effect on semen attributes using lower doses and longer storage times. Also, It has been shown that supplementing taurine at 0, 1.5, 7, 12.5, and 50 mM in a semen extender and stored at 37°C for 4 h did not alter the viability, morphology, and acrosome integrity of rabbit spermatozoa [42]. Therefore, it was recommended to use taurine in rabbit extender at lower temperatures for possible protective effects. Therefore, the current study aimed to test the effect of supplementing different concentrations of GSH and taurine on the antioxidant properties of semen extender and semen quality after different chilling times.

2. Materials and Methods

2.1. Animal Management and Semen Collection. The current study was conducted following the ethical and regulation procedures of the Department of Animal Wealth

Development, Faculty of Veterinary Medicine, Kafrelsheikh University, Egypt (approval number: KFS-2021/05).

This study used 20 rabbit bucks ($n=20$) of the New Zealand white rabbit breed (about 2-3 kg body weight and 1-1.5-year-old). Bucks were individually labeled and caged with free access to feed and water. Animals were housed in well-ventilated rooms with a natural daylight supply. Bucks were trained for two weeks for semen collection using an artificial vagina and a mature female as a mount stimulator. Then, ten ejaculates from each buck were collected (one ejaculate/week), during the ten-week trial, for a total of 200 ejaculates. In each collection, fresh semen samples were microscopically examined and evaluated after removing the gel clot. The spermatozoa activity was microscopically evaluated at 100x (to evaluate mass motility) and 400x (for individual motility) using a phase-contrast microscope (Micros Austria, Micros St. Veit Hunnenbrunn-Gewerbezone, Veit/Glan, Austria) [43]. A Neubauer hemocytometer was used to measure sperm concentration. Only ejaculates with a standard color, more than 85% motility, about 0.5 mL volume, and $\sim 400 \times 10^6/\text{mL}$ concentration were used for the study. During each time of collection, ejaculates were pooled ($n=20$ with an average volume of ejaculates $\sim 8.75 \pm 1.165$ mL) to avoid individual variations and get sufficient semen volume for different treatments [41].

2.2. Extender Preparation and Semen Processing and Evaluation. The pooled semen samples were diluted in a 1 : 2 basic extender [41] containing 20 mM Tris, 57 mM citric acid anhydrous, 5 mM fructose, 22 mM glucose, and 4.6 mM streptomycin along with 10 mL egg yolk and completed to 100 mL with distilled water with pH 6.8-7, and osmotic pressure was kept at pH 6.9, 300 mOsm. The diluted pooled semen samples were divided equally into 7 fractions (1 fraction/treatment). The diluted semen samples were then supplemented with 0 (control with diluent only without additives), 0.5, 1, and 2 mM glutathione (GSH; cat. no. Y0000517, Merck KGaA, Darmstadt, Germany) and 1, 5, and 10 mM taurine (cat. no. T0625, Merck KGaA, Darmstadt, Germany). The pH was adjusted to 6.8-7 with a dilution factor of 1 : 2 (semen: prepared extenders), and the diluted semen was kept at 5°C. Then, changes in semen quality were evaluated after 24, 48, and 72 h of chilling. Sperm motility, viability (% of live spermatozoa), acrosome reaction, fructose and glutamic pyruvic transaminase (GPT) concentrations, and antioxidant enzymes activity (catalase (CAT), superoxide dismutase (SOD), GSH, and malondialdehyde (MDA)) as oxidative biomarkers were assessed. Relative transcriptomic levels of SOD and CAT genes were also evaluated. All the evaluations were repeated ten times throughout the trial (ten weeks).

2.3. Assessment of Sperm's Viability and Progressive Motility. Sperm viability was evaluated for control and treated samples according to the literature [41]. Sperms were stained with propidium iodide (PI, 200 $\mu\text{g}/\text{mL}$ semen), and viability was morphologically assessed based on sperm's membrane

integrity differences: sperms with damaged membrane were penetrated by PI stain showing partial or complete pinkish fluorescence, which indicates dead sperm. At the same time, the live sperm exhibited no fluorescence over the sperm head. Slides were examined at 1,000x magnification using a phase-contrast microscope (Leica Microsystems Inc., Parkway North, Deerfield, IL 60015 United States), and 200 sperms/slide were counted and evaluated.

Sperms' progressive motility was evaluated by visually measuring the percentage of spermatozoa showing any movement of the flagellum using a phase-contrast microscope (Micros Austria, Micros St. Veit Hunnenbrunn-Gewerbezone, Veit/Glan, Austria), that was equipped with a warm stage. The motility was measured after 24, 48, and 72 h of chilling for control and different treatments. Briefly, 5 μL of every semen sample was incubated at 37°C for 20 min, placed on a prewarmed glass slide, and covered by a glass cover slider. Slides were examined at 400x. Sperm motility was assessed in five microscopic fields for each semen sample (in the five examined fields within 1 min). Sperm motility was evaluated as the percentage of spermatozoa that moved their flagellum. The average of the five fields was used to calculate the final sperm motility value [41, 44].

2.4. Analysis of Acrosome Reaction. Acrosome reaction was also evaluated after 24, 48, and 72 h of chilling in differently treated semen samples using Giemsa stain according to Watson's methods with some modifications [45]. Briefly, 30 μL from each semen sample was smeared on glass slides, air-dried, fixed with 10% neutral formal saline for 15 min, washed in running water for 20 min, and allowed to dry. After drying, slides were immersed in Giemsa stain solution overnight. After incubation, the slides were rinsed with distilled water and air-dried. The acrosome status of 100 spermatozoa for each sample was examined at 1000x in each stained smear, using a phase-contrast light microscope (Micros Austria, Micros St. Veit Hunnenbrunn-Gewerbezone, Veit/Glan, Austria). Purple heads characterized spermatozoa with intact acrosomes, whereas the damaged-acrosome spermatozoa had a pale lavender head. The percentage of acrosome-intact spermatozoa was calculated.

2.5. Assessing the Oxidative Stress Biomarkers in Seminal Plasma. Semen samples at 24, 48, and 72 postchilling were centrifuged at 1,000 \times g for 20 min, and the supernatant, composed of semen extender and seminal fluid, was used to assess oxidative stress biomarkers, using MDA contents and assessing the concentration of glutathione peroxidase (GPx), SOD, and CAT. Pellets of spermatozoa were washed by suspending them in cold fresh phosphate buffer saline, centrifuged at 3,000 \times g for 10 min, and then frozen in liquid nitrogen for RNA extraction. Lipid peroxidation was evaluated by quantifying the MDA content (nmol/mL) using a commercial kit (MD 2529, Bio-Diagnostic, Egypt, with a sensitivity of 0.1 nmol/ml and detection range up to 50 nmol/ml) according to manufacturer instructions. MDA

was measured with 2-thiobarbituric acid, monitoring the absorbance change at 532 nm with the spectrophotometer (Shanghai Spectrophotometer Co. Ltd., China). Antioxidant concentrations of CAT, SOD, and GPx were determined with commercial kits: CAT (CA 2517, Bio-Diagnostic, Egypt, with a sensitivity of 0.22 U/ml and detection range up to 200 U/ml), SOD (SOD 2521, Bio-Diagnostic, Egypt, with a sensitivity of 2.0 mU/ml and detection range up to 500 U/ml), and GPx (1.11.1.9, Bio-Assay Systems, United States, with a sensitivity of 1 mU/ml and detection range up to 500 mU/ml), according to manufacturer's instruction.

2.6. qRT-PCR of Relative Transcriptomic Levels of SOD and CAT Genes. Total RNA was extracted using TRI Reagent (Bioline, UK) following the manufacturer's protocol. Diluted semen samples were centrifuged at $3,000 \times g$ for 20 min to eliminate semen extender and seminal fluid. After that, a pellet of spermatozoa was washed many times using ice-cold PBS. Lysis buffer was added to the sperm pellet and kept at room temperature for 10 min. Then, 1 ml TRI reagent was added, and extraction was completed. Afterward, complementary DNA (cDNA) was synthesized by reverse transcription reactions by iNtRON Biotechnology cDNA kits (Power cDNA Synthesis Kit, cat. no. 25011, iNtRON Bio, Seongnam, Gyeonggi 462-120, US) according to the manufacturer protocol. Specific primers for antioxidant-related genes such as CAT, SOD, and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) genes, as shown in Table 1, were used. The amplification reaction was performed using the SensiFast SYBR Lo-Rox kit (Bioline, UK) in the Stratagene MX3000P real-time PCR system. 20 μ l as a total reaction volume contained 10 μ l SYBER green master mix, 2 μ l cDNA, primers 0.8 μ l (50 nm), and 6.4 μ l RNase/DNase-free H₂O [47]. The cycling conditions were preheating at 62°C for 1 min, 95°C for 5 min, and 40 cycles at 95°C for 2 sec, and the annealing temperatures are listed in Table 1. At the end of the PCR reaction, a melting curve with 0.5°C increments in the temperature range of 65–95°C was performed to ensure the specificity of the primers. Also, amplification efficiency for each primer was estimated using the formula $E = 100 \times (10^{-1/\text{slope}} - 1)$. Samples for each treatment were repeated twice. The relative mRNA expression of the genes was calculated using the comparative threshold cycle method $2^{-\Delta\Delta CT}$ [48], and the results were reported as fold change differences relative to the control group.

2.7. Statistical Analysis. Before data analysis, normality testing was performed by testing the Gaussian normal distribution with the Shapiro–Wilk normality test. Results of the current study were statistically analyzed using two-way ANOVA to test the effect of GSH and taurine supplementation, chilling time, and their interactions on semen quality, followed by Tukey's multiple comparison test using GraphPad Prism (GraphPrism© Software, La Jolla, CA, USA) [49–57]. Results were expressed as means \pm SEM and considered significantly different at $P < 0.05$.

3. Results

3.1. Impact of GSH and Taurine Supplementation on Sperm Kinetics at Different Postchilling Times. Table 2 displays the postchilling kinetics of rabbit sperm supplemented with GSH and taurine at different concentrations (0.5–2 mM GSH and 1–10 mM taurine). The addition of GSH and taurine induced improving effects of sperm kinetic, especially with increasing chilling time. At 24 h postchilling, adding GSH and taurine at different rates did not alter sperm kinetics. However, at 48 and 72 h postchilling, there was a noticeable reduction in percentages of motile sperm, % of live sperms, and % of the acrosome reaction in nontreated semen samples. An interesting improvement of these parameters was reported for treated semen samples with GSH and taurine ($P < 0.05$).

3.2. GSH and Taurine Supplementation Improved the Antioxidant Properties of Semen Extender by Altering the MDA Content and Levels of Antioxidant Enzymes. Figure 1 illustrates the MDA content in semen samples following GSH and taurine supplementation after different postchilling times. GSH and taurine supplementation at different concentrations was significantly correlated with lowering MDA concentration in semen samples preserved at 5°C for 24, 48, and 72 h ($P < 0.05$). At 24 h postchilling, 1, 5, and 10 mM taurine and 0.5, 1, and 2 mM GSH-supplemented semen samples had the lowest MDA content ($P < 0.05$). After 48 h of chilling at 5°C, all samples had less MDA content than those after 24 h of chilling ($P < 0.05$). However, 1 and 10 mM taurine supplementation had the lowest level of MDA compared to the other treatments and control ($P < 0.05$). Increasing the chilling time to 72 h was significantly associated with reducing MDA content in all treatments compared to the control (except 5 and 10 mM taurine as well as 0.5 mM GSH). GSH and taurine supplementation lowers oxidative stress in chilled semen samples, up to 72 h, by reducing lipid peroxidation, confirmed by decreasing MDA contents.

Figure 2 displays the alterations in SOD activities during different chilling times and GSH and taurine supplementation. Both chilling time and GSH and taurine supplementation markedly modified SOD activities in semen with distinct interaction ($P < 0.05$). After 24 h chilling, higher concentrations of GSH and taurine (1 and 2 mM GSH, and 5 and 10 mM taurine) significantly increased SOD activities in semen samples, compared to the lower concentration (0.5-mM GSH and 1 mM taurine) and nonsupplemented semen ($P < 0.05$). Increasing the chilling time to 48 and 72 h was associated with increasing SOD in all GSH and taurine-supplemented samples, compared to nonsupplemented semen samples ($P < 0.05$). GSH and taurine supplementation in semen extender were associated with increasing SOD and increased chilling time.

GPx activity in semen samples was also measured at different chilling times, and GSH and taurine supplementation (Figure 3). Nonsupplemented semen samples exhibited nearly fixed GPx levels over increasing chilling

TABLE 1: Primer sequences and annealing temperature.

Gene	Primer	Annealing	Slope	Efficiency	Reference
SOD	F: CCCGGTCTTTGTA CTCTCGT R: AAGGATGAAGAGAGGCACGT	60°C + preheating (at 62°C for 1 min)	-3.34	99.25	[46] NM_001082627
CAT	F: GCCCTGGTCAGTCTTGTAGT R: CCATCGGCATATGAACGGAT	62°C	-3.42	96.06	[46] XM_002709045
GAPDH	F: ATCTCGCTCCTGGAAGATGG R: CAAAGTGGATGTTGTGCGCA	60°C + preheating (at 62°C for 1 min)	-3.39	97.24	[46] NM_001082253

SOD = superoxide dismutase; CAT = catalase; GAPDH = glyceraldehyde-3-phosphate dehydrogenase (reference gene).

TABLE 2: Effect of different concentrations of glutathione and taurine supplementation (0.5, 1, and 2 mM GSH and 1, 5, and 10 mM taurine) on rabbit sperm motility, vitality, and acrosome reaction rate during 24, 48, and 72 h storage at 5°C.

	Postchilling time (h)	Sperm motility (%)	Sperm vitality (%)	Acrosome reaction rate (%)
C	24	87.5 ± 0.50 ^{Aa}	93.00 ± 2.00 ^{Aa}	85.00 ± 0.50 ^{Aa}
	48	74 ± 1.00 ^{Bb}	84.00 ± 4.00 ^{Bb}	67.00 ± 2.00 ^{Ba}
	72	50 ± 5.00 ^{Bc}	77.50 ± 2.50 ^{Cc}	47.50 ± 2.50 ^{Bb}
G0.5	24	88 ± 1.00 ^{Aa}	93.50 ± 1.50 ^{Aa}	88.00 ± 1.00 ^{Aa}
	48	77 ± 2.00 ^{Aa}	86.50 ± 1.50 ^{Aa}	77.50 ± 2.50 ^{ABa}
	72	73 ± 2.00 ^{Ab}	82.5 ± 2.50 ^{Aa}	62.50 ± 2.50 ^{Bb}
G1	24	93 ± 2.00 ^{Aa}	94.00 ± 1.00 ^{Aa}	89.00 ± 2.00 ^{Aa}
	48	82.5 ± 2.50 ^{Aa}	91.50 ± 1.50 ^{Aa}	78.50 ± 6.50 ^{ABa}
	72	66 ± 6.00 ^{Aa}	90.5 ± 2.50 ^{Aa}	67.00 ± 2.00 ^{Aa}
G2	24	93 ± 1.00 ^{Aa}	94.5 ± 0.50 ^{Aa}	93.50 ± 0.50 ^{Aa}
	48	87 ± 2.00 ^{Ab}	92.00 ± 1.00 ^{Bb}	87.00 ± 1.00 ^{ABa}
	72	83 ± 3.00 ^{Ab}	91.00 ± 2.00 ^{Bb}	82.50 ± 2.50 ^{Bb}
T1	24	92.5 ± 2.50 ^{Aa}	93.00 ± 2.00 ^{Aa}	98.50 ± 0.50 ^{Aa}
	48	87.5 ± 0.50 ^{Aa}	91.50 ± 1.50 ^{Aa}	81.00 ± 6.00 ^{Aa}
	72	74.5 ± 1.50 ^{Ab}	89.50 ± 3.50 ^{Ab}	61.00 ± 6.00 ^{Bb}
T5	24	93.5 ± 0.50 ^{Aa}	92.5 ± 2.50 ^{Aa}	89.50 ± 0.50 ^{Aa}
	48	77.5 ± 2.50 ^{Ab}	92.50 ± 0.50 ^{Aa}	83.00 ± 1.00 ^{Aa}
	72	73.5 ± 1.50 ^{Ab}	91.5 ± 1.50 ^{Aa}	72.50 ± 2.50 ^{Aa}
T10	24	91.5 ± 0.50 ^{Aa}	95.00 ± 0.00 ^{Aa}	92.00 ± 1.00 ^{Aa}
	48	84 ± 1.00 ^{Aa}	94.00 ± 1.00 ^{Aa}	88.00 ± 2.00 ^{Aa}
	72	79.5 ± 0.50 ^{Aa}	91.50 ± 3.50 ^{Aa}	82.00 ± 2.50 ^{Aa}

Uppercase and lowercase letters indicate statistical significance at $P < 0.05$ between chilling times and extender supplements (GSH and taurine). C represents the control samples (no treatment), and G1, G2, and G0.5 represent 1, 2, and 0.5 mM of GSH, respectively. T1, T5, and T10 denote 1, 5, and 10 mM of taurine, respectively.




FIGURE 1: MDA contents in semen samples following different concentrations of GSH and taurine and chilling for up to 72 h. C represents the control samples (no treatment) stored at 5°C for 24, 45, and 72 h, and G1, G2, and G0.5 represent 1, 2, and 0.5 mM of GSH treatments stored at 5°C for 24, 45, and 72 h, respectively. T1, T5, and T10 denote 1, 5, and 10 mM of taurine supplementation after different postchilling times (24, 48, and 72 h) at 5°C. Different uppercase and lowercase letters indicate statistical significances at $P < 0.05$ between different chilling times and extender supplements (GSH and taurine), respectively.




FIGURE 2: SOD activity in response to different concentrations of GSH and taurine and chilling for up to 72 h. C represents the control samples (no treatment) stored at 5°C for 24, 45, and 72 h, and G1, G2, and G0.5 represent 1, 2, and 0.5 mM of GSH treatments stored at 5°C for 24, 45, and 72 h, respectively. T1, T5, and T10 denote 1, 5, and 10 mM of taurine supplementation after different postchilling times (24, 48, and 72 h) at 5°C. Different uppercase and lowercase letters indicate statistical significances at $P < 0.05$ between different chilling times and extender supplements (GSH and taurine), respectively.

time. However, supplementing the semen extender with either GSH or taurine at a different concentration significantly modified GPx levels which were prominent with increasing chilling time ($P < 0.05$). In this context, all supplementing taurine and GSH doses except 0.5 mM GSH prominently increased GPx levels at 24 h postchilling. The same effect was also reported at 48 h and 72 h postchilling for all taurine and GSH doses except 0.5 mM GSH, which only was able to induce higher GPx after 72 h of chilling ($P < 0.05$).

CAT activities in supplemented semen samples and at different chilling times were evaluated (Figure 4). After 24 h of chilling, 1 mM GSH and the same taurine dose induced higher levels of CAT than the other supplements and control ($P < 0.05$). Increasing chilling time to 48 h was associated with higher activities of CAT in the samples supplemented with 1 mM and 2 mM GSH along with the 1 mM taurine compared to the other treated samples ($P < 0.05$). Nevertheless, at 72 h postchilling, all treatments had nearly the same CAT activities. Additionally, the effect of GSH and taurine supplementation was significantly modified by chilling time. The highest activities of CAT were measured at 24 h and 48 h postchilling in the case of 1 mM GSH and taurine. At the same time, no differences were noticed for the other supplemented groups.

3.3. GSH and Taurine Supplementation to Semen-Extender Maintained Fructose and Glutamic Pyruvic Transaminase (GPT) Concentrations at Different Postchilling Times. Figure 5 displays changes in fructose and GPT concentrations at different chilling times, with and without GSH and taurine supplementation in the semen extender. Fructose concentration increased with increasing chilling time in the nonsupplemented semen extender ($P < 0.05$). Conversely, semen extenders supplemented with different doses of GSH and taurine showed slight changes in fructose levels at different chilling times. At 48 and 72 h postchilling, the

fructose levels in nonsupplemented semen extender (control) were significantly higher than those in supplemented ones, especially if compared with those treated with 5 and 10 mM taurine ($P < 0.05$).

GPT levels were also influenced by supplementations of semen extender with GSH and taurine and chilling time ($P < 0.05$). In comparison with the nonsupplemented group (control), all supplemented semen extenders, either by GSH or taurine (at different doses), had significantly fewer GPT levels ($P < 0.05$). However, no significant differences in GPT levels were found among supplemented semen extenders. Besides, the GPT level in nonsupplemented extenders significantly increased with increasing chilling times ($P < 0.05$). However, it remained unchanged in the case of supplemented extenders with either GSH or taurine.

3.4. Relative mRNA Levels of SOD and CAT Genes at Different Postchilling Times and following GSH and Taurine Supplementation. Figure 6 illustrates the relative mRNA copies of both SOD and CAT genes in spermatozoa after different chilling times and GSH or taurine supplementation. For the SOD gene, compared to nonsupplemented semen and after 24 h cooling, increasing chilling time to 48 and 72 h was associated with significant increases in mRNA copies. This effect was also reported, especially at 48 h postchilling, for semen samples supplemented with different doses of either GSH or taurine ($P < 0.05$). The highest expression levels of SOD were determined in the case of 2 mM GSH and 1 mM taurine, followed by 0.5 mM and 5 mM taurine. Increasing the cooling time to 72 h was associated with higher expression levels of SOD, especially in the case of 1 mM GSH and 10 mM taurine, compared with nonsupplemented or GSH- and taurine-supplemented samples ($P < 0.05$).

The CAT gene's relative expression level was modulated with GSH/taurine supplementation and cooling times. Increasing chilling time to 72 h was linked with a significant




FIGURE 3: GPx activity in response to different concentrations of GSH and taurine and chilling for up to 72 h. C represents the control samples (no treatment) stored at 5°C for 24, 45, and 72 h, and G1, G2, and G0.5 represent 1, 2, and 0.5 mM of GSH treatments stored at 5°C for 24, 45, and 72 h, respectively. T1, T5, and T10 denote 1, 5, and 10 mM of taurine supplementation after different postchilling times (24, 48, and 72 h) at 5°C. Different uppercase and lowercase letters indicate statistical significances at $P < 0.05$ between different chilling times and extender supplements (GSH and taurine), respectively.





FIGURE 4: CAT activity in response to different concentrations of GSH and taurine and chilling for up to 72 h. C represents the control samples (no treatment) stored at 5°C for 24, 45, and 72 h, and G1, G2, and G0.5 represent 1, 2, and 0.5 mM of GSH treatments stored at 5°C for 24, 45, and 72 h, respectively. T1, T5, and T10 denote 1, 5, and 10 mM of taurine supplementation after different postchilling times (24, 48, and 72 h) at 5°C. Different uppercase and lowercase letters indicate statistical significances at $P < 0.05$ between different chilling times and extender supplements (GSH and taurine), respectively.


reduction of CAT mRNA copies in samples either non-supplemented or supplemented with GSH or taurine ($P < 0.05$), except for the integration with 1 mM taurine. The highest expression levels were measured at 24 and 48 h postchilling for both treated (except 1-mM taurine) or untreated samples. The expression level fluctuated with different GSH supplementation doses; the highest level was found in the case of 1 mM GSH compared to 2 mM and 0.5 mM GSH, while a different response was reported for taurine supplementation. Increasing the taurine dose was associated with lowering CAT mRNA levels at 48 and 72 h postchilling. At 24 h postchilling, minor increases were found with increasing taurine dose.

4. Discussion

Cooling and freezing/thawing procedures of semen markedly alter the quality of preserved sperm. In this regard, long-term cooling deteriorates sperm quality and, consequently, its fertilizing ability because of the excessive release of ROS such as superoxide anion radicals (O_2^-), hydrogen peroxide (H_2O_2), and lipid hydroperoxides [22, 58, 59]. Moreover, semen dilution during chilled storage reduces the normal physiological antioxidants included in the semen. The maintenance of the fertilizing capacity of rabbit semen for longer than 48 to 72 hours remains an important target for rabbit production because the farming system is based



(a)



(b)

FIGURE 5: Fructose (a) and GPT (b) levels in semen samples in response to different concentrations of GSH and taurine and chill for up to 72 h. C represents the control samples (no treatment) stored at 5°C for 24, 45, and 72 h, and G1, G2, and G0.5 represent 1, 2, and 0.5 mM of GSH treatments stored at 5°C for 24, 45, and 72 h, respectively. T1, T5, and T10 denote 1, 5, and 10 mM of taurine supplementation after different postchilling times (24, 48, and 72 h) at 5°C. Different uppercase and lowercase letters indicate statistical significances at $P < 0.05$ between different chilling times and extender supplements (GSH and taurine), respectively.

exclusively on AI programs, which are currently performed with fresh diluted semen within 6–12 hours from the collection. To improve semen extenders' efficacy and functionality, in the current study, we targeted increasing rabbit semen quality by GSH and taurine supplementation and tested their effects over different chilling times (24, 48, and 72 h).

Supplementing the extender with antioxidant substances such as GSH and taurine improved the semen quality, as confirmed by the significant increase in the percentages of live spermatozoa, motile sperms, and acrosome reactions. However, at 24 h postchilling, GSH and taurine did not alter these parameters compared to the control. This effect is probably related to the natural antioxidant potency of sperm to scavenge ROS production [41]. The positive effects of both GSH and taurine are increased with the supplementation dose and cooling time (at 48 and 72 h postchilling). This response might be linked to the antioxidant properties of both GSH and taurine that protect the sperm membrane from cooling-associated oxidative stress and the overproduction of ROS [58]. Spermatozoa have antioxidant defenses against the production of ROS, but dilution in semen extender and cooling process lowers this antioxidant

capability [60]. Therefore, it is necessary to use exogenous antioxidant supplements such as GSH and taurine [60]. The results obtained in this study agreed with the literature on GSH and taurine supplementation, which reported improved sperm motility, viability, and acrosomal reaction in rabbits [22, 41, 61], stallions [62], and goat bucks [63]. Besides, the current findings agreed with the findings of the previous studies, which stated increases of postthaw motility, viability, and acrosomal integrity of ram and mithun (*Bos frontalis*) semen diluted in tris-based extender supplemented with 25 and 50 mM taurine, respectively [40, 64]. Additionally, supplementing 2 mM of reduced glutathione in the cryopreservation media of boar semen improved sperm membrane integrity and postthaw sperm motility [38]. Moreover, using a 4 mM GSH-supplemented extender protected sperm motility, motion kinetics, acrosome integrity, and viability for 24 h 5°C [41]. Our findings disagreed with other authors [42, 65], who reported that GSH and a wide range of taurine supplementation (1.5–50 mM) did not affect rabbit sperm viability, morphology, and acrosomal reaction during cooling. Nevertheless, the reported inferior semen quality in the case of nontreated sperms and small doses of GSH and taurine might reflect the damage to the




FIGURE 6: The relative expression level of SOD and CAT in sperms in response to different concentrations of GSH and taurine and chilling for up to 72 h. C represents the control samples (no treatment) stored at 5°C for 24, 45, and 72 h, and G1, G2, and G0.5 represent 1, 2, and 0.5 mM of GSH treatments stored at 5°C for 24, 45, and 72 h, respectively. T1, T5, and T10 denote 1, 5, and 10 mM of taurine supplementation after different postchilling times (24, 48, and 72 h) at 5°C. Different uppercase and lowercase letters indicate statistical significances at $P < 0.05$ between different chilling times and extender supplements (GSH and taurine), respectively.

sperm because of excessive ROS production that exceeds the normal antioxidant potency of sperms [41]. The variations in results between different studies might be due to variations in the reduced levels of normal sperm antioxidant power [66], pH, and osmolality [67]. Besides the supplementation dose, species differences exist in withstanding cryopreservation and cooling-associated stress and thawing-freezing problems [68].

The improving effects of both GSH and taurine may also be due to the prevention of lipid peroxidation, as confirmed by lowering the MDA concentrations in GSH and taurine-treated semen compared to nontreated semen. This effect ascertains the protective role of both taurine and GSH against lipid peroxidation and sperm membrane damage.

The high content of polyunsaturated fatty acids (PUFA) in the sperm plasma membrane increases its chance to get deteriorated and damaged, and the consequent release of PUFA, which binds to oxygen, increases ROS production [69]. This finding is possibly explained by increasing the intracellular biosynthesis of GSH, which protects the proteins, DNA, and membrane lipids from ROS-induced damage by the direct radical-scavenging ability of the GSH [70]. The protective effect of GSH and taurine supplementation was also established by lowering the extracellular level of GPT, which is a good indicator of sperm quality, measuring sperm membrane stability [71]. Increasing the percentage of abnormal and dead sperm is usually associated with high levels of GPT in extracellular

fluid because the damaged sperm membrane releases these enzymes [72]. Moreover, GSH and taurine were correlated with maintaining fructose levels (all times) relative to the number of live spermatozoa in the extender. Increasing fructose levels at increasing cooling time in non-supplemented extenders might be linked with a lowering percentage of live spermatozoa compared to the supplementing extenders.

Another interesting finding is that increasing SOD, GPx, and CAT concentrations are linked with increasing the supplementation dose and the cooling time. This finding might explain the improving effect of the exogenous supplementation of GSH and taurine on sperm quality. Cooling and cryopreservation processes impair the sperm's functions and lower its fertilizing capacity by either overproduction of ROS or reducing the antioxidant defense system [60]. Interestingly, the increases in SOD, CAT, and GPx concentrations might be coupled with increasing the antioxidant production by sperm and its release into seminal plasma [73]. This finding confirms that the exogenous GSH and taurine supplementation in semen extender protect the sperm against the excessive production of ROS during chilling by sustaining equilibrium between oxidation and levels of antioxidants [58]. This effect might be correlated with increasing the ability of sperm to synthesize SOD, CAT, and GPx during oxidative stress [39]. This result agrees with other studies that stated an improvement in GSH and SOD activities following GSH supplementation to semen extender as it is a cofactor of the enzymatic antioxidant systems during cooling [41]. The increased levels of antioxidants in the extender might be related to the enhanced transcription levels of its related genes, as confirmed by upregulating the mRNA copies of *CAT* and *SOD* genes. The upregulations of *CAT* and *SOD* following GSH and taurine supplementation to semen extender might be linked to the APS kinase (APK) and glutathione S-transferases (GST) family genes modulation by GSH [74] and to the concentration of taurine in the mitochondria of many cells [75].

Therefore, evaluating the transcriptomic levels of these pathways following taurine and GSH supplementation is recommended. Despite the reported improving effects of GSH and taurine on semen quality, this study has some limitations, including predicting the real male fertility. Also, this study did not highlight the problem of spermatozoa that occurs after entering the female tract. Therefore, other tests such as sperm penetration, sperm viability, and tests examining the sperm's attachment and fertilizing capacity (pregnancy test) are recommended in the future. Moreover, extending this study to cover as long as possible, up to 5-6 days, is recommended.

5. Conclusions

Exogenous supplementation of the extender with GSH and taurine improved the quality of rabbit semen cooled at 5°C by increasing the percentage of live spermatozoa, motile sperms, and acrosomal reaction, as confirmed by maintaining fructose level and decreasing GPT in extracellular fluid. Moreover, both GSH and taurine reduced cooling-

associated oxidative stress by lowering MDA concentration and increasing SOD, CAT, and GPx levels, along with upregulating the transcription levels of SOD and CAT genes. These improving effects increased with the supplementation dose and the cooling time to 72 h. Therefore, based on these results, the 2 mM GSH and 10 mM taurine can be effectively supplemented with a rabbit's semen extender to enhance the in-vitro sperm quality and antioxidant status of New Zealand rabbits' sperm during chilled storage for up to 72 hours.

Data Availability

All data are included in this article.

Ethical Approval

The animal study was reviewed and approved by the Institutional Ethics Committee of Kafrelsheikh University, Egypt (approval number: KFS-2021/05).

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Authors' Contributions

M.F.F.B., S.E.H.E.-N., T.A.E.K., Z.I.A., and A.M.S. contributed to conceptualization; M.F.F.B., S.E.H.E.-N., T.A.E.K., Z.I.A., A.M.S., and S.E.-K. developed methodology; H.S.A.T. and M.A. created software; H.S.A.T. and M.A. validated the study; M.F.F.B., S.E.H.E.-N., T.A.E.K., Z.I.A., A.M.S., and S.E.-K. performed formal analysis; M.F.F.B., S.E.H.E.-N., T.A.E.K., Z.I.A., A.M.S., and S.E.-K. investigated the study; H.S.A.T. and M.A. collected resources; S.E.-K., L.G., and L.B. curated the study; M.F.F.B., S.E.H.E.-N., T.A.E.K., Z.I.A., A.M.S., H.S.A.T., M.A., L.G., and S.E.-K. wrote the study and prepared the original draft; M.A., S.E.-K., L.G., and L.B. wrote the study and reviewed and edited the manuscript; M.A. performed project administration; M.A. and L.B. contributed to funding acquisition. All authors have read and agreed to the published version of the manuscript.

Supplementary Materials

This figure represents the acrosomal reaction and the sperm's livability. For live and dead sperm, it has been a while since this slide was made, and for that reason, there are no clear differences between the live and dead sperm. In this study, we prepared and examined the slides directly, without taking photos. (*Supplementary Materials*)

References

- [1] K. Wiebe, S. Robinson, and A. Cattaneo, "Chapter 4- climate change, agriculture and food security: impacts and the potential for adaptation and mitigation," in *Sustainable Food and Agriculture*, C. Campanhola and S. Pandey, Eds., Academic Press, Cambridge, MA, USA, pp. 55-74, 2019.
- [2] S. Sechi, A. Di Cerbo, S. Canello et al., "Effects in dogs with behavioural disorders of a commercial nutraceutical diet on

- stress and neuroendocrine parameters,” *The Veterinary Record*, vol. 180, no. 1, p. 18, 2017.
- [3] A. Di Cerbo, S. Canello, G. Guidetti, C. Laurino, and B. Palmieri, “Unusual antibiotic presence in gym trained subjects with food intolerance; a case report,” *Nutricion Hospitalaria*, vol. 30, no. 2, pp. 395–398, 2014.
- [4] A. Di Cerbo, G. Guidetti, S. Canello, and R. Cocco, “A possible correlation between diet, serum oxytetracycline concentration, and onset of reproductive disturbances in bitches: clinical observations and preliminary results,” *Turkish Journal of Veterinary and Animal Sciences*, vol. 43, no. 4, pp. 523–531, 2019.
- [5] A. Di Cerbo, F. Pezzuto, A. Scarano, G. Guidetti, and S. Canello, “The contradictory world of tetracyclines,” *Panminerva Medica*, vol. 62, no. 2, pp. 116–117, 2020.
- [6] M. A. Silvestre, J. L. Yániz, F. J. Peña, P. Santolaria, and M. Castelló-Ruiz, “Role of antioxidants in cooled liquid storage of mammal spermatozoa,” *Antioxidants*, vol. 10, no. 7, p. 1096, 2021.
- [7] N. Fathy Megahed, E. S. M. Abdel-Kafy, S. El-Kassas et al., “Association of insulin receptor substrate 1 (IRS-1) gene polymorphism with growth and litter-related traits in NMER rabbits,” *Animal Biotechnology*, pp. 1–8, 2023.
- [8] J. Fan, S. Kitajima, T. Watanabe et al., “Rabbit models for the study of human atherosclerosis: from pathophysiological mechanisms to translational medicine,” *Pharmacology & Therapeutics*, vol. 146, pp. 104–119, 2015.
- [9] F. Matsuhisa, S. Kitajima, K. Nishijima, T. Akiyoshi, M. Morimoto, and J. Fan, “Transgenic rabbit models: now and the future,” *Applied Sciences*, vol. 10, no. 21, p. 7416, 2020.
- [10] K. Nishijima, S. Kitajima, F. Matsuhisa, M. Niimi, C. C. Wang, and J. Fan, “Strategies for highly efficient rabbit sperm cryopreservation,” *Animals*, vol. 11, no. 5, p. 1220, 2021.
- [11] M. Ragab, J. P. Sanchez, and M. Baselga, “Effective population size and inbreeding depression on litter size in rabbits. A case study,” *Journal of Animal Breeding and Genetics*, vol. 132, no. 1, pp. 68–73, 2015.
- [12] J. Casellas, D. Vidal-Roqueta, E. Flores et al., “Epistasis for founder-specific inbreeding depression in rabbits,” *Journal of Heredity*, vol. 102, no. 2, pp. 157–164, 2011.
- [13] E. Kubovicova, A. V. Makarevich, A. Balazi, J. Vasicek, and P. Chrenek, “Factors affecting rabbit sperm cryopreservation: a mini-review,” *Zygote*, vol. 30, no. 1, pp. 1–8, 2022.
- [14] F. Amidi, A. Pazhohan, M. Shabani Nashtaei, M. Khodarahmian, and S. Nekoonam, “The role of antioxidants in sperm freezing: a review,” *Cell and Tissue Banking*, vol. 17, no. 4, pp. 745–756, 2016.
- [15] M. R. Farag, C. Zizzadoro, M. Alagawany et al., “In ovo protective effects of chicoric and rosmarinic acids against Thiocloprid-induced cytotoxicity, oxidative stress, and growth retardation on newly hatched chicks,” *Poultry Science*, vol. 102, no. 4, Article ID 102487, 2023.
- [16] M. A. Hassan, A. M. Abo-Elmaaty, A. W. Zaglool et al., “*Origanum vulgare* essential oil modulates the AFB1-induced oxidative damages, nephropathy, and altered inflammatory responses in growing rabbits,” *Toxins*, vol. 15, no. 1, p. 69, 2023.
- [17] S. A. Mawed, C. Marini, M. Alagawany et al., “Zinc oxide nanoparticles (ZnO-NPs) suppress fertility by activating autophagy, apoptosis, and oxidative stress in the developing oocytes of female zebrafish,” *Antioxidants*, vol. 11, no. 8, p. 1567, 2022.
- [18] M. R. Farag, A. A. Moselhy, A. El-Mleeh et al., “Quercetin alleviates the immunotoxic impact mediated by oxidative stress and inflammation induced by doxorubicin exposure in rats,” *Antioxidants*, vol. 10, no. 12, p. 1906, 2021.
- [19] M. R. Farag, S. R. Khalil, A. W. Zaglool et al., “Thiacloprid induced developmental neurotoxicity via ROS-oxidative injury and inflammation in chicken embryo: the possible attenuating role of chicoric and rosmarinic acids,” *Biology*, vol. 10, no. 11, p. 1100, 2021.
- [20] A. Di Cerbo, F. Pezzuto, G. Guidetti, S. Canello, and L. Corsi, “Tetracyclines: insights and updates of their use in human and animal pathology and their potential toxicity,” *The Open Biochemistry Journal*, vol. 13, pp. 1–12, 2019.
- [21] E. Isachenko, V. Isachenko, I. I. Katkov, and S. Dessole, “Vitrification of mammalian spermatozoa in the absence of cryoprotectants: from past practical difficulties to present success,” *Reproductive BioMedicine Online*, vol. 6, no. 2, pp. 191–200, 2003.
- [22] D. Johnke, S. P. de Graaf, and R. Bathgate, “The effect of sperm concentration and storage vessel on quercetin-supplemented rabbit semen during chilled storage,” *Reproduction in Domestic Animals*, vol. 50, no. 4, pp. 567–573, 2015.
- [23] P. Gogol, “Motility parameters and intracellular ATP content of rabbit spermatozoa stored for 3 Days at 15°C,” *Folia Biologica*, vol. 61, no. 1, pp. 87–91, 2013.
- [24] I. E. S. El-Seadawy, “Preservability of rabbit semen after chilled storage in tris based extender enriched with different concentrations of Propolis ethanolic extract (PEE),” *Asian Pacific Journal of Reproduction*, vol. 6, no. 2, pp. 68–76, 2017.
- [25] M. Aksoy, C. Wrenzycki, D. Herrmann, and H. Niemann, “Oxidative stress induced changes of glutathione concentration in in vitro-produced bovine blastocysts,” *Reproduction, Fertility and Development*, vol. 19, no. 1, p. 187, 2007.
- [26] F. Lopez-Gatius, G. Sances, M. Sancho et al., “Effect of solid storage at 15°C on the subsequent motility and fertility of rabbit semen,” *Theriogenology*, vol. 64, no. 2, pp. 252–260, 2005.
- [27] M. Rosato and N. Iaffaldano, “Effect of chilling temperature on the long-term survival of rabbit spermatozoa held either in a tris-based or a jellified extender,” *Reproduction in Domestic Animals*, vol. 46, no. 2, pp. 301–308, 2011.
- [28] M. Di Iorio, M. A. Colonna, M. Miranda et al., “Initial cooling time before freezing affects post-thaw quality and reproductive performance of rabbit semen,” *Animal Science Journal*, vol. 89, no. 9, pp. 1240–1244, 2018.
- [29] M. Di Iorio, G. Rusco, R. Iampietro et al., “Finding an effective freezing protocol for Turkey semen: benefits of ficoll as non-permeant cryoprotectant and 1:4 as dilution rate,” *Animals*, vol. 10, no. 3, p. 421, 2020.
- [30] N. Iaffaldano, M. Di Iorio, M. P. Rosato, and A. Manchisi, “Cryopreservation of rabbit semen using non-permeable cryoprotectants: effectiveness of different concentrations of low-density lipoproteins (LDL) from egg yolk versus egg yolk or sucrose,” *Animal Reproduction Science*, vol. 151, no. 3–4, pp. 220–228, 2014.
- [31] A. Rekha, B. F. Zohara, F. Y. Bari, and M. G. S. Alam, “Comparisons of commercial Triladyl and locally manufactured extenders for the chilling of semen and their effects on pregnancy rates after transcervical AI in Bangladeshi Indigenous (*Ovis aries*) sheep,” *Animal Reproduction*, vol. 13, no. 4, pp. 735–742, 2016.
- [32] C. Pacelli, A. Di Cerbo, L. Lecce et al., “Effect of chicken bone extracts on metabolic and mitochondrial functions of K562 cell line,” *Pharmaceuticals*, vol. 13, no. 6, p. 114, 2020.

- [33] G. Guidetti, A. Di Cerbo, A. Giovazzino et al., "In vitro effects of some botanicals with anti-inflammatory and antitoxic activity," *Journal of Immunology Research*, vol. 2016, Article ID 5457010, 11 pages, 2016.
- [34] A. Di Cerbo, V. Rubino, F. Morelli et al., "Mechanical phenotyping of K562 cells by the Micropipette Aspiration Technique allows identifying mechanical changes induced by drugs," *Scientific Reports*, vol. 8, no. 1, p. 1219, 2018.
- [35] A. Saha, M. Asaduzzaman, and F. Y. Bari, "Cryopreservation techniques for ram sperm," *Veterinary Medicine International*, vol. 2022, Article ID 7378379, 16 pages, 2022.
- [36] A. R. M. El-Khawagah, M. M. M. Kandiel, and H. Samir, "Effect of quercetin supplementation in extender on sperm kinematics, extracellular enzymes release, and oxidative stress of Egyptian Buffalo bulls frozen-thawed semen," *Frontiers in Veterinary Science*, vol. 7, p. 604460, 2020.
- [37] M. N. Bucak and N. Tekin, "Protective effect of taurine, glutathione and trehalose on the liquid storage of ram semen," *Small Ruminant Research*, vol. 73, no. 1-3, pp. 103–108, 2007.
- [38] E. Estrada, M. M. Rivera del Álamo, J. E. Rodríguez-Gil, and M. Yeste, "The addition of reduced glutathione to cryopreservation media induces changes in the structure of motile subpopulations of frozen-thawed boar sperm," *Cryobiology*, vol. 78, pp. 56–64, 2017.
- [39] Z. Zhu, Z. Ren, X. Fan et al., "Cysteine protects rabbit spermatozoa against reactive oxygen species-induced damages," *PLoS One*, vol. 12, no. 7, Article ID e0181110, 2017.
- [40] M. N. Bucak, A. Ateşşahin, Ö. Varışlı, A. Yüce, N. Tekin, and A. Akçay, "The influence of trehalose, taurine, cysteamine and hyaluronan on ram semen: microscopic and oxidative stress parameters after freeze-thawing process," *Theriogenology*, vol. 67, no. 5, pp. 1060–1067, 2007.
- [41] E. Ahmad, Z. Naseer, and M. Aksoy, "Glutathione supplementation in semen extender improves rabbit sperm attributes during refrigeration," *World Rabbit Science*, vol. 29, no. 2, pp. 81–86, 2021.
- [42] D. Paal, F. Strejček, E. Tvrđá et al., "Taurine does not improve the quality of short-term stored rabbit spermatozoa in vitro," *Reproduction in Domestic Animals*, vol. 52, no. 6, pp. 1046–1051, 2017.
- [43] F. Ciribe, R. Panzarella, M. C. Pisu, A. Di Cerbo, G. Guidetti, and S. Canello, "Hypospermia improvement in dogs fed on a nutraceutical diet," *The Scientific World Journal*, vol. 2018, Article ID 9520204, 4 pages, 2018.
- [44] S. A. Abdelnour, M. A. Hassan, A. K. Mohammed et al., "The effect of adding different levels of curcumin and its nanoparticles to extender on post-thaw quality of cryopreserved rabbit sperm," *Animals*, vol. 10, no. 9, p. 1508, 2020.
- [45] P. F. Watson, "Use of a Giemsa stain to detect changes in acrosomes of frozen ram spermatozoa," *The Veterinary Record*, vol. 97, no. 1, pp. 12–15, 1975.
- [46] H. M. Saleh, Y. S. El-Sayed, S. M. Naser, A. S. Eltahawy, A. Onoda, and M. Umezawa, "Efficacy of alpha-lipoic acid against cadmium toxicity on metal ion and oxidative imbalance, and expression of metallothionein and antioxidant genes in rabbit brain," *Environmental Science and Pollution Research*, vol. 24, no. 31, pp. 24593–24601, 2017.
- [47] A. Gallo, R. Landi, V. Rubino et al., "Oxytetracycline induces DNA damage and epigenetic changes: a possible risk for human and animal health?" *PeerJ*, vol. 5, p. e3236, 2017.
- [48] X. Rao, X. Huang, Z. Zhou, and X. Lin, "An improvement of the $\Delta\Delta$ CT method for quantitative real-time polymerase chain reaction data analysis," *Biostatistics, bioinformatics and biomathematics*, vol. 3, no. 3, pp. 71–85, 2013.
- [49] A. Di Cerbo, S. Canello, G. Guidetti et al., "Adverse food reactions in dogs due to antibiotic residues in pet food: a preliminary study," *Veterinaria Italiana*, vol. 54, no. 2, pp. 137–146, 2018.
- [50] G. Blaiotta, N. Murru, A. Di Cerbo, M. Succi, R. Coppola, and M. Aponte, "Commercially standardized process for probiotic 'Italice' cheese production," *LWT- Food Science and Technology*, vol. 79, pp. 601–608, 2017.
- [51] A. Scarano, G. Murmura, G. Vantaggiato et al., "Delayed expansion of atrophic mandible (deam): a case report," *Oral Implantology*, vol. 10, no. 2, pp. 190–196, 2017.
- [52] F. Mazzeranghi, C. Zanotti, A. Di Cerbo et al., "Clinical efficacy of nutraceutical diet for cats with clinical signs of cutaneous adverse food reaction (CAFR)," *Polish Journal of Veterinary Sciences*, vol. 20, no. 2, pp. 269–276, 2017.
- [53] D. L. Dreger, B. W. Davis, R. Cocco et al., "Commonalities in development of pure breeds and population isolates revealed in the genome of the Sardinian fonni's dog," *Genetics*, vol. 204, no. 2, pp. 737–755, 2016.
- [54] A. Di Cerbo, S. Centenaro, F. Beribè et al., "Clinical evaluation of an antiinflammatory and antioxidant diet effect in 30 dogs affected by chronic otitis externa: preliminary results," *Veterinary Research Communications*, vol. 40, no. 1, pp. 29–38, 2016.
- [55] S. Destefanis, D. Giretto, M. C. Muscolo et al., "Clinical evaluation of a nutraceutical diet as an adjuvant to pharmacological treatment in dogs affected by Keratoconjunctivitis sicca," *BMC Veterinary Research*, vol. 12, no. 1, p. 214, 2016.
- [56] A. Di Cerbo and B. Palmieri, "Review: the market of probiotics," *Pakistan journal of pharmaceutical sciences*, vol. 28, no. 6, pp. 2199–2206, 2015.
- [57] T. Iannitti, B. Palmieri, A. Aspiro, and A. Di Cerbo, "A preliminary study of painless and effective transdermal botulinum toxin A delivery by jet nebulization for treatment of primary hyperhidrosis," *Drug Design, Development and Therapy*, vol. 8, pp. 931–935, 2014.
- [58] A. K. Bansal and G. S. Bilaspuri, "Impacts of oxidative stress and antioxidants on semen functions," *Veterinary Medicine International*, vol. 2011, Article ID 686137, 7 pages, 2011.
- [59] B. Palmieri, A. Pennelli, and A. Di Cerbo, "Jurassic surgery and immunity enhancement by alkylglycerols of shark liver oil," *Lipids in Health and Disease*, vol. 13, no. 1, p. 178, 2014.
- [60] P. Petruska, M. Capcarova, and P. Sutovsky, "Antioxidant supplementation and purification of semen for improved artificial insemination in livestock species," *Turkish Journal of Veterinary and Animal Sciences*, vol. 38, pp. 643–652, 2014.
- [61] M. J. Maya-Soriano, E. Taberner, M. Sabés-Alsina, M. Piles, and M. Lopez-Bejar, "Absence of beneficial effects on rabbit sperm cell cryopreservation by several antioxidant agents," *Zygote*, vol. 23, no. 1, pp. 1–10, 2015.
- [62] M. Zhandi and V. Ghadimi, "Effect of glutathione-supplemented INRA82 extender on miniature caspian stallion sperm quality during storage at 5°C," *Journal of Equine Veterinary Science*, vol. 34, no. 5, pp. 606–610, 2014.
- [63] A. Sarangi, P. Singh, M. Virmani et al., "Effect of antioxidants supplementation on the quality of Beetal buck semen stored at 4°C," *Veterinary World*, vol. 10, no. 10, pp. 1184–1188, 2017.
- [64] S. S. Alam, N. A. Hafiz, and A. H. Abd El-Rahim, "Protective role of taurine against genotoxic damage in mice treated with methotrexate and tamoxfine," *Environmental Toxicology and Pharmacology*, vol. 31, no. 1, pp. 143–152, 2011.
- [65] P. M. Papa, F. O. Papa, L. A. Oliveira, P. N. Guasti, C. Castilho, and I. C. Giometti, "Different extenders in the

- cryopreservation of bovine epididymal spermatozoa,” *Animal Reproduction Science*, vol. 161, pp. 58–63, 2015.
- [66] C. Del Prete, F. Ciani, S. Tafuri et al., “Effect of superoxide dismutase, catalase, and glutathione peroxidase supplementation in the extender on chilled semen of fertile and hypofertile dogs,” *Journal of Veterinary Science*, vol. 19, no. 5, pp. 667–675, 2018.
- [67] R. A. Oliveira, R. L. Piersanti, C. Wolf, M. A. O. Viu, and M. L. Gambarini, “Glutathione for the freezing of cooled equine semen, using different protocols,” *Animal Reproduction, Belo Horizonte*, vol. 11, no. 2, pp. 104–109, 2014.
- [68] H. Rodriguez-Martinez and A. D. Barth, “<I>In vitro</I> evaluation of sperm quality related to <I>in vivo</I> function and fertility,” *Reproduction in Domestic Ruminants*, vol. 6, no. 1, pp. 39–54, 2007.
- [69] M. N. Banday, F. A. Lone, F. Rasool, M. Rashid, and A. Shikari, “Use of antioxidants reduce lipid peroxidation and improve quality of crossbred ram sperm during its cryopreservation,” *Cryobiology*, vol. 74, pp. 25–30, 2017.
- [70] O. Uysal and M. Bucak, “Effects of oxidized glutathione, bovine serum albumin, cysteine and lycopene on the quality of frozen-thawed ram semen,” *Acta Veterinaria Brno*, vol. 76, no. 3, pp. 383–390, 2007.
- [71] I. Dogan, U. Polat, and Z. Nur, “Correlations between seminal plasma enzyme activities and semen parameters in seminal fluid of Arabian horses,” *Iranian Journal of Veterinary Research*, vol. 10, pp. 119–124, 2009.
- [72] M. Gündoğan, “Some reproductive parameters and seminal plasma constituents in relation to season in Akkaraman and Awassi rams,” *Turkish Journal of Veterinary and Animal Sciences*, vol. 30, pp. 95–100, 2006.
- [73] S. R. Khalil, C. Zheng, S. M. Abou-Zeid et al., “Modulatory effect of thymol on the immune response and susceptibility to *Aeromonas hydrophila* infection in Nile tilapia fish exposed to zinc oxide nanoparticles,” *Aquatic Toxicology*, vol. 259, Article ID 106523, 2023.
- [74] B. Zhu, K. Wang, Z. Liang, Z. Zhu, and J. Yang, “Transcriptome analysis of glutathione response: RNA-Seq provides insights into balance between antioxidant response and glucosinolate metabolism,” *Antioxidants*, vol. 11, no. 7, p. 1322, 2022.
- [75] Y. Li, Q. Peng, J. Shang et al., “The role of taurine in male reproduction: physiology, pathology and toxicology,” *Frontiers in Endocrinology*, vol. 14, Article ID 1017886, 2023.