

## TO THE EDITOR:

## The race is on: bispecifics vs CAR T cells in B-cell lymphoma

Ashwath Gurumurthi,<sup>1</sup> Jason Westin,<sup>1</sup> and Marion Subklewe<sup>2,3</sup><sup>1</sup>Department of Lymphoma and Myeloma, The University of Texas MD Anderson Cancer Center, Houston, TX; and <sup>2</sup>Gene Center and Department of Biochemistry and<sup>3</sup>Department of Medicine II, University Hospital, Ludwig-Maximilians-Universität München, Munich, Germany

T-cell therapeutic approaches have come of age over the last 5 years and dramatically changed the treatment paradigm for relapsed/refractory (R/R) indolent and aggressive B-cell lymphomas. This consists of bispecific T-cell engaging antibodies (BiAbs) and autologous, ex vivo, expanded T cells with a chimeric antigen receptor (CAR T). Both of these therapies result in target cell lysis independent of T-cell receptor specificity and major histocompatibility complex presentation of a tumor antigen.<sup>1</sup> BiAbs combine the antigen binding regions from 2 different single chain antigen binding domains to form an antibody construct binding, with one arm to a surface antigen on B cells, for example CD20, and the second arm to an activating component of the T-cell receptor, CD3, causing cytotoxic T-cell activation and lysis.

Tisagenlecleucel (tisa-cel), axicabtagene ciloleucel (axi-cel), and lisocabtagene maraleucel (liso-cel) are the 3 CAR-T therapies currently approved in the United States and Europe to treat adult patients with R/R large B-cell lymphoma (LBCL).<sup>2-4</sup> In addition, axi-cel and tisa-cel are approved for R/R follicular lymphoma (FL). Brexucabtagene autoleucel (brexu-cel), with an identical CAR design to axi-cel but generated using CD4 and CD8 selected T cells, is approved for R/R mantle cell lymphoma and R/R B-cell precursor acute lymphoblastic leukemia (BCP-ALL) in adults. All 4 CAR-T products are directed against the same epitope on CD19, the B-cell lineage marker, and are second generation constructs with an intracellular costimulatory region. They differ in their costimulatory domain, CD28 for axi-cel and brexu-cel, responsible for early, rapid expansion with limited long-term persistence and 4-1BB for tisa-cel and liso-cel responsible for greater long-term CAR-T persistence. The degree to which the CAR-T construct affects clinical efficacy and toxicity outcomes remains unclear.<sup>5</sup>

Besides blinatumomab, a CD19-directed T-cell engager approved for minimal residual disease positive and R/R BCP-ALL, mosunetuzumab and teclistamab are the only bispecific antibody constructs approved for patients with R/R FL and R/R multiple myeloma, respectively. Subcutaneous epcoritamab based on the EPCORE NHL-1 trial<sup>6</sup> and glofitamab from the NP30179 study<sup>7</sup> in R/R LBCL are expected to be approved shortly with odronextamab currently progressing through phase 2 studies.<sup>8</sup>

Results from the pivotal phase 2 trials for the CAR-T products in LBCL along with real-world registry studies from the United States and European groups are detailed in Table 1.<sup>2,3,9-14</sup> ZUMA-1 has the longest follow-up over 5 years confirming durability of response with axi-cel achieving a median overall survival (OS) of 25.8 months with 5-year OS rate of 43%, whereas tisa-cel in the JULIET study had a median OS of 11.1 months.<sup>3,15</sup> However, cross-trial comparisons can be misleading because of differences in study designs between ZUMA-1 and JULIET trials.<sup>5</sup>

Both the European and US real-world evidence consistently demonstrated in nonrandomized retrospective comparisons that axi-cel is associated with higher efficacy defined by objective response rate (ORR) and complete response rate (CRR), corresponding to greater progression free survival and OS in comparison with tisa-cel. The improved efficacy with axi-cel is counterbalanced with higher rates of cytokine release syndrome (CRS), any grade immune effector cell-associated neurotoxicity syndrome (ICANS), high grade ICANS ( $\geq$  grade 3) and higher 1- and 2-year nonrelapse mortality.<sup>10,11,14</sup>

Submitted 23 January 2023; accepted 2 April 2023; prepublished online on *Blood Advances* First Edition 10 April 2023. <https://doi.org/10.1182/bloodadvances.2022009066>.

Data are available upon request to the corresponding author, Marion Subklewe ([marion.subklewe@med.uni-muenchen.de](mailto:marion.subklewe@med.uni-muenchen.de)).

© 2023 by The American Society of Hematology. Licensed under [Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International \(CC BY-NC-ND 4.0\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), permitting only noncommercial, nonderivative use with attribution. All other rights reserved.

**Table 1. Efficacy and toxicity outcomes from pivotal phase 2 CAR-T trials compared against real world experience and phase 2 BiAb trials in R/R LBCL**

	Phase 2 CAR-T trials				Real-world experience of CAR-T (axi-cel vs tisa-cel)				Phase 2 expansion BiAb Trials		
	ZUMA-1 (axi-cel) <sup>2</sup>	JULIET (tisa-cel) <sup>3</sup>	TRANSCEND NHL 001 (liso-cel) <sup>9</sup>	UK <sup>10</sup>	German GLA DRST registry <sup>11</sup>	Spanish <sup>12</sup>	French DESCAR-T <sup>13*</sup>	US CAR-T consortium <sup>14</sup>	Epcoritamab <sup>5</sup>	Glofitamab <sup>7</sup>	Odronextamab <sup>8</sup>
Median age (y)	58	56	63	57 vs 63.5	60 vs 61	59 vs 62	62 vs 64	59 vs 67	68	66	67
Median vein-to-vein time	17 days	NR	37 days	40 days vs 50 days	35 days vs 55 days	41 days vs 52 days	NA	28 days vs 45 days		Not applicable	
ORR/CRR	82%/58%	52%/40%	73%/53%	44%/40% vs 32%/32%	74%/42% vs 63%/32%	60%/42% vs 54%/19%	80%/60% vs 66%/42%	52%/44% vs 41%/35%	63%/39%	52%/39%	53%/37%
12 mo OS/12 mo PFS	59%/44%	49%/NR	58%/44%	57%/42% vs 44%/28%	55%/35% vs 58%/24%	51%/41% vs 47%/33%	64%/47% vs 49%/33%	62%/42% vs 59%/32%	NA (too short follow-up)	50%/37%	NA (too short follow-up)
Any grade CRS/≥grade 3	93%/13%	58%/22	42%/2%	93%/8% vs 74%/8%	81%/10% vs 65%/13%	88%/8% vs 73%/6%	86%/9% vs 76%/5%	85%/9% vs 39%/1%	50%/2.5%	63%/4%	53%/0%
Any grade ICANS/≥grade 3	64%/28%	21%/12%	30%/10%	44%/20% vs 15%/4%	44%/16% vs 22%/7%	42%/18% vs 16%/5%	49%/14% vs 22%/3%	56%/38% vs 11%/1%	6%/0.6%	8%/3%	4%/0%
NRM	3%	0%	NA	9% vs 3%	10% vs 4%	7% vs 3%	NA	12% vs 8%	0.6% (ICANS related)	0%	2%

LBCL, large B cell lymphoma; NA, not available; NR, not reported; NRM, nonrelapse mortality; PFS, progression free survival.  
\*The French DESCAR-T study used propensity score matching between axi-cel and tisa-cel, which might affect comparison of headline figures.

Registry studies have consistently demonstrated a higher median age for patients treated with tisa-cel, suggesting that clinicians may choose tisa-cel because of perceptions regarding toxicity.<sup>10,11,14</sup> Interestingly, the ZUMA-7 trial had a preplanned subgroup analysis which compared axi-cel to standard of care in patients aged >65 years that identified that axi-cel achieved CRR and event free survival outcomes similar if not greater than that in younger patients, with similar toxicity levels in the overall population.<sup>16</sup> Given the high efficacy and a favorable safety profile identified in the TRANSFORM trial, including median OS rate not reached with a median follow-up of 17.5 months and low CRS and ICANS rates in line with tisa-cel outcomes, liso-cel may be a preferred choice for some patients.<sup>17</sup> Currently, the main limitations with liso-cel include an absence of real-world evidence to confirm efficacy and safety and a more complicated manufacturing process with separate CD4 and CD8 products, which can increase the chances of out-of-specification products.

Each BiAb product developed for the LBCL setting comes with its own advantages and disadvantages with respect to administration, such as mosunetuzumab and glofitamab have fixed schedule regimens, whereas epcoritamab and odronextamab are continued until disease progression or toxicity is detected. Regular 3-week administrations until disease progression is difficult for maintaining the patients' quality of life in comparison with the one-and-done approach with CAR T. Mosunetuzumab and epcoritamab have subcutaneous formulations, which may be convenient to administer in the outpatient setting.<sup>18</sup>

The efficacy of the BiAbs in R/R LBCL are promising, with subcutaneous epcoritamab and IV glofitamab and odronextamab demonstrating response rates nearing those of tisa-cel in JULIET.<sup>3,6-8</sup> This is highlighted in Table 1 using the headline figures from the phase 2 bispecific trials in LBCL, albeit it includes some patients treated after CAR-T relapse. Longer-term data is necessary to confirm durability of response with real-world evidence needed to verify whether it proves to be a viable alternative to CAR T.

A recent phase 2 trial of fixed duration IV mosunetuzumab in R/R FL for 8 to 17 cycles based on the response demonstrated an ORR of 80% and CRR of 60%, which is comparable with an ORR of 94% and CRR of 70% with axi-cel in ZUMA-5 and an ORR of 87% and CRR of 73% with tisa-cel in the ELARA trial.<sup>19-21</sup> Notably, the median age of participants for all these studies at ~60 years is much younger than the real world R/R FL population. Given the follow-up is <18 months for these studies, it is not yet possible to draw conclusions comparing the durability of efficacy.

The BiAbs are potentially an alternative to CAR T in R/R disease, particularly in older patients, because they have relatively low rates of high grade (≥grade 3) CRS and ICANS, and off-the-shelf availability has convenience for use in community oncology setting. However, initial community use may remain impractical, given the incidence of grade 2 CRS events, which may necessitate hospitalization in 15% for epcoritamab, 17% for mosunetuzumab in FL, 12% for glofitamab, and 19% for odronextamab in LBCL cases.<sup>6,7,19,22</sup> Reassuringly, most BiAb-associated CRS events occurred in cycle 1 during step-up dosing soon after infusion, from within a few hours to the next day, with a short duration lasting from 1 to 3 days, rarely requiring intensive care unit admissions with usually no requirement for vasopressor support or high flow

oxygen.<sup>7,19,22,23</sup> One potential solution would be to treat patients in cell therapy centers for the first 1 or 2 cycles, followed by a transition to the community oncology to continue therapy.

A challenge is determining the order in which to treat R/R LBCL: CAR T, BiAbs, or autologous stem cell transplantation. ZUMA-7 and TRANSFORM, using axi-cel and liso-cel, respectively, found that CAR-T therapy is superior to autologous stem cell transplantation at the first relapse of LBCL in response rates and long-term survival.<sup>24,25</sup> The BELINDA trial found tisa-cel to be equivalent to autologous stem cell transplantation in this setting, but there were concerns about the methodology of all 3 trials that make comparison difficult.<sup>26,27</sup> There are limited data on the sequencing of BiAbs after CAR T for relapsed disease and vice-versa. In the BiAb trials, 11% of patients with R/R LBCL were treated with epcoritamab, 40% with odronextamab, 12% with mosunetuzumab, and 33% with glofitamab after CAR-T relapse. Reassuringly, these patients showed similar responses to patients with no prior CAR-T exposure.<sup>6,7,22,28</sup> The reverse may be true with BiAbs used as a line of therapy before CAR T, given that they target different tumor associated antigen, CD20 for BiAbs and CD19 for CAR-T cells. This approach has recently been described by the French LYSA group, and more evidence is expected in the future as BiAbs will be used as earlier lines of therapy.<sup>29</sup>

The case for BiAbs as an alternative to CAR-T therapy in R/R lymphoma is compelling, given less frequent toxicity and the capacity to manufacture to scale an off-the-shelf product, thus, providing rapid access. However, the survival data with BiAbs remain immature in comparison with those of CAR-T therapy, which has a wealth of evidence proving durability in the ~40% of patients expected to have a long-term cure.

Perhaps the question of greatest interest is how to determine the order in which to sequence these therapies. The durability of CAR T in comparison to immature survival data with BiAbs currently favors a CAR T first followed by BiAbs for progressive disease. Given the modest efficacy of standard frontline chemo-immunotherapy for LBCL and FL, the sequencing question is best answered by using prognostic tools and identifying patients at high risk of relapse and then offering them CAR T or BiAbs at an earlier line of therapy. An example of this approach was applied in the ZUMA-12 trial in which axi-cel was used at interim therapy response in the first line setting resulting in excellent outcomes.<sup>30</sup> Similarly, there is a great deal of interest with multiple studies combining BiAbs with conventional chemoimmunotherapy or novel therapies, such as polatuzumab, immunomodulators or Bruton's tyrosine kinase inhibitors, for earlier lines of therapy for patients who are treatment naive and in the R/R setting to improve response rates and survival. We are yet to identify the best combination approach incorporating either BiAbs or CAR T in frontline therapy for the minority of patients who are at high risk of early relapse of aggressive and indolent lymphoma. However, the promise of this future appears exciting.

**Acknowledgments:** This work was supported by a Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) research grant provided within the Sonderforschungsbereich SFB-TRR 388/1 2021 – 452881907, and DFG research grant 451580403 (M.S.). The work was further supported by the Bavarian Elite Graduate Training Network, the Wilhelm-Sander Stiftung (project 2018.087.1), the Else-Kröner-Fresenius Stiftung, the

Bavarian Center for Cancer Research (BZKF), and research funding from Miltenyi Biotec BV & Co KG (M.S.). In addition, this work was financially supported by ASCO Conquer Cancer and the Schweitzer Family Foundation. The trial was in part supported by the National Institutes of Health through MD Anderson's Cancer Center support grant CA016672.

**Contributions:** M.S., J.W., and A.G. wrote the manuscript. Concept and content were based on a pros-and-cons session by M.S. and J.W. at a meeting about lymphoma in September 2022.

**Conflict-of-interest disclosure:** M.S. receives industry research support from Amgen, Bristol Myers Squibb/Celgene, Gilead, Janssen, Miltenyi Biotec, MorphoSys, Novartis, Roche, Seattle Genetics, and Takeda; serves as a consultant/adviser to AvenCell, CDR-Life, Ichnos Sciences, Incyte Biosciences, Janssen, Molecular Partners, and Takeda; and serves on the speakers' bureau at Amgen, AstraZeneca, Bristol Myers Squibb/Celgene, Gilead, GlaxoSmithKline, Janssen, Novartis, Pfizer, Roche, and Takeda. J.W. receives industry research support from Kite/Gilead, Bristol Myers Squibb, Novartis, Genentech/Roche, AstraZeneca, Janssen, ADC Therapeutics, MorphoSys/Incyte, and Kymera, and serves as a consultant/adviser to Kite/Gilead, Bristol Myers Squibb, Novartis, Genentech/Roche, AstraZeneca, Janssen, ADC Therapeutics, MorphoSys/Incyte, AbbVie, Seagen, MonteRosa, Umoja, and Iksuda. A.G. serves as an adviser to Kite Pharma.

**ORCID profiles:** A.G., [0000-0003-2335-7769](https://orcid.org/0000-0003-2335-7769); J.W., [0000-0002-1824-2337](https://orcid.org/0000-0002-1824-2337); M.S., [0000-0003-3905-0251](https://orcid.org/0000-0003-3905-0251).

**Correspondence:** Marion Subklewe, Gene Center and Department of Biochemistry, Department of Medicine II, University Hospital, Ludwig-Maximilians-Universität München, 81377 Munich, Germany; email: [marion.subklewe@med.uni-muenchen.de](mailto:marion.subklewe@med.uni-muenchen.de).

## References

1. Subklewe M. BiTEs better than CAR T cells. *Blood Adv*. 2021;5(2):607-612.
2. Locke FL, Ghobadi A, Jacobson CA, et al. Long-term safety and activity of axicabtagene ciloleucel in refractory large B-cell lymphoma (ZUMA-1): a single-arm, multicentre, phase 1–2 trial. *Lancet Oncol*. 2019;20(1):31-42.
3. Schuster SJ, Tam CS, Borchmann P, et al. Long-term clinical outcomes of tisagenlecleucel in patients with relapsed or refractory aggressive B-cell lymphomas (JULIET): a multicentre, open-label, single-arm, phase 2 study. *Lancet Oncol*. 2021;22(10):1403-1415.
4. Abramson JS, Palomba ML, Gordon LI, et al. Lisocabtagene maraleucel for patients with relapsed or refractory large B-cell lymphomas (TRANSCEND NHL 001): a multicentre seamless design study. *Lancet*. 2020;396(10254):839-852.
5. Westin JR, Kersten MJ, Salles G, et al. Efficacy and safety of CD19-directed CAR-T cell therapies in patients with relapsed/refractory aggressive B-cell lymphomas: observations from the JULIET, ZUMA-1, and TRANSCEND trials. *Am J Hematol*. 2021;96(10):1295-1312.
6. Thieblemont C, Phillips T, Ghesquieres H, et al. ABCL-422 subcutaneous epcoritamab in patients with relapsed or refractory large B-cell lymphoma (EPCORE NHL-1): pivotal results from a phase 2 study. *Clinical Lymphoma Myeloma Leukemia*. 2022;22:S379.

7. Dickinson MJ, Carlo-Stella C, Morschhauser F, et al. Glofitamab for relapsed or refractory diffuse large B-cell lymphoma. *N Engl J Med*. 2022;387(24):2220-2231.
8. Kim WS, Kim TM, Cho SG, et al. Odronektamab in patients with relapsed/refractory (R/R) diffuse large B-cell lymphoma (DLBCL): results from a prespecified analysis of the pivotal phase II study ELM-2. *Blood*. 2022;140(suppl 1):1070-1071.
9. Abramson JS, Palomba ML, Gordon LI, et al. Two-year follow-up of transcend NHL 001, a multicenter phase 1 study of lisocabtagene maraleucel (liso-cel) in relapsed or refractory (R/R) large B-cell lymphomas (LBCL). *Blood*. 2021;138(suppl 1):2840-2840.
10. Kuhn A, Roddie C, Kirkwood AA, et al. A national service for delivering CD19 CAR-Tin large B-cell lymphoma - the UK real-world experience. *Br J Haematol*. 2022;198(3):492-502.
11. Bethge WA, Martus P, Schmitt M, et al. GLA/DRST real-world outcome analysis of CAR T-cell therapies for large B-cell lymphoma in Germany. *Blood*. 2022;140(4):349-358.
12. Kwon M, Iacoboni G, Reguera JL, et al. Axicabtagene ciloleucel compared to tisagenlecleucel for the treatment of aggressive B-cell lymphoma. *Haematologica*. 2023;108(1):110-121.
13. Bachy E, Le Gouill S, Di Blasi R, et al. A real-world comparison of tisagenlecleucel and axicabtagene ciloleucel CAR T cells in relapsed or refractory diffuse large B cell lymphoma. *Nat Med*. 2022;28(10):2145-2154.
14. Riedell PA, Hwang WT, Nastoupil LJ, et al. Patterns of use, outcomes, and resource utilization among recipients of commercial axicabtagene ciloleucel and tisagenlecleucel for relapsed/refractory aggressive B cell lymphomas. *Transplant Cell Ther*. 2022;28(10):669-676.
15. Jacobson CA, Locke F, Ghobadi A, et al. Long-term (5 year) overall survival in zuma-1, the pivotal study of axicabtagene ciloleucel (axi-cel) in patients with refractory large b-cell lymphoma (LBCL). *Bone Marrow Transplant*. 2022;57(suppl 1):131-132.
16. Westin J, Locke FL, Dickinson M, et al. Clinical and patient (pt)-reported outcomes (PROs) in a phase 3, randomized, open-label study evaluating axicabtagene ciloleucel (axi-cel) versus standard-of-care (SOC) therapy in elderly pts with relapsed/refractory (R/R) large B-cell lymphoma (LBCL; ZUMA-7). *J Clin Oncol*. 2022;40(suppl 16):7548-7548.
17. Abramson JS, Solomon SR, Arnason JE, et al. Lisocabtagene maraleucel (liso-cel) versus standard of care (SOC) with salvage chemotherapy followed by autologous stem cell transplantation (ASCT) as second-line (2L) treatment in patients with relapsed or refractory large B-cell lymphoma (LBCL): primary analysis of the randomized, phase 3 transform study. *Blood*. 2022;140(suppl 1):1581-1583.
18. Gonzalez Barca E. Role of bispecific antibodies in relapsed/refractory diffuse large B-cell lymphoma in the CART era. *Front Immunol*. 2022;13:909008.
19. Budde LE, Sehn LH, Matasar M, et al. Safety and efficacy of mosunetuzumab, a bispecific antibody, in patients with relapsed or refractory follicular lymphoma: a single-arm, multicentre, phase 2 study. *Lancet Oncol*. 2022;23(8):1055-1065.
20. Jacobson CA, Chavez JC, Sehgal AR, et al. Axicabtagene ciloleucel in relapsed or refractory indolent non-Hodgkin lymphoma (ZUMA-5): a single-arm, multicentre, phase 2 trial. *Lancet Oncol*. 2022;23(1):91-103.
21. Fowler NH, Dickinson M, Dreyling M, et al. Tisagenlecleucel in adult relapsed or refractory follicular lymphoma: the phase 2 ELARA trial. *Nat Med*. 2022;28(2):325-332.
22. Bannerji R, Arnason JE, Advani RH, et al. Odronektamab, a human CD20×CD3 bispecific antibody in patients with CD20-positive B-cell malignancies (ELM-1): results from the relapsed or refractory non-Hodgkin lymphoma cohort in a single-arm, multicentre, phase 1 trial. *Lancet Haematol*. 2022;9(5):e327-e339.
23. Hutchings M, Mous R, Clausen MR, et al. Dose escalation of subcutaneous epcoritamab in patients with relapsed or refractory B-cell non-Hodgkin lymphoma: an open-label, phase 1/2 study. *Lancet*. 2021;398(10306):1157-1169.
24. Kamdar M, Solomon SR, Arnason J, et al. Lisocabtagene maraleucel versus standard of care with salvage chemotherapy followed by autologous stem cell transplantation as second-line treatment in patients with relapsed or refractory large B-cell lymphoma (TRANSFORM): results from an interim analysis of an open-label, randomised, phase 3 trial. *Lancet*. 2022;399(10343):2294-2308.
25. Locke FL, Miklos DB, Jacobson CA, et al. Axicabtagene ciloleucel as second-line therapy for large B-cell lymphoma. *N Engl J Med*. 2022;386(7):640-654.
26. Bishop MR, Dickinson M, Purtill D, et al. Second-line tisagenlecleucel or standard care in aggressive B-cell lymphoma. *N Engl J Med*. 2022;386(7):629-639.
27. Westin J, Sehn LH. CAR T cells as a second-line therapy for large B-cell lymphoma: a paradigm shift? *Blood*. 2022;139(18):2737-2746.
28. Budde LE, Assouline S, Sehn LH, et al. Single-agent mosunetuzumab shows durable complete responses in patients with relapsed or refractory B-cell lymphomas: phase I dose-escalation study. *J Clin Oncol*. 2022;40(5):481-491.
29. Crochet G, Audrey C, Bachy E, et al. CAR T-cell therapy remain effective in patients with relapse/refractory B-cell non-Hodgkin lymphoma after bispecific antibodies exposure: results of a lisa study based on the descar-T registry. *Blood*. 2022;140(suppl 1):4639-4641.
30. Neelapu SS, Dickinson M, Munoz J, et al. Axicabtagene ciloleucel as first-line therapy in high-risk large B-cell lymphoma: the phase 2 ZUMA-12 trial. *Nat Med*. 2022;28(4):735-742.