

## Sequences of Homologous $\beta$ -Lactamases from Clinical Isolates of *Serratia marcescens* with Different Substrate Specificities

NAOKI MATSUMURA,<sup>1\*</sup> SHINZABURO MINAMI,<sup>2</sup> AND SUSUMU MITSUHASHI<sup>1</sup>

*Episome Institute, 2220 Kogure, Fujimi-mura, Seta-gun Gunma,<sup>1</sup> and Research Laboratories, Toyama Chemical Co., Ltd., 2-4-1 Shimo-okui, Toyama-city, Toyama, 930,<sup>2</sup> Japan*

Received 12 March 1997/Returned for modification 22 July 1997/Accepted 22 October 1997

**Genes for two group 1  $\beta$ -lactamases, SRT-1 and SST-1, were sequenced. These  $\beta$ -lactamases were produced by clinical isolates of *Serratia marcescens*, isolates GN16694 and GN19450, respectively. The resulting enzymes were 96% identical. SRT-1 hydrolyzed oxyimino cephalosporins, but SST-1 hardly hydrolyzed them. At residue 213 in the third motif, which is conserved among group 1  $\beta$ -lactamases, SRT-1 and SST-1 had Lys and Glu, respectively. By site-directed mutagenesis, the substitution of Glu by Lys at residue 213 in SST-1 resulted in an enzyme that hydrolyzed oxyimino cephalosporins.**

With the extended use of  $\beta$ -lactams in clinical practice, organisms resistant to these antibiotics have increasingly been isolated. Resistance can be mediated by the production of  $\beta$ -lactamases, such as in derepressed mutants (18, 19), the acquisition of plasmids with a  $\beta$ -lactamase gene from other strains (2, 5, 8, 17, 25), and the expansion of the substrate specificity of the organism's own enzyme (16, 22, 23).

To protect the  $\beta$ -lactam ring from hydrolysis by  $\beta$ -lactamases, cephalosporins with a 7-oxyimino moiety on the side chain of the cephem nucleus have been synthesized, and these have been shown to be stable. Recently, it was reported that the group 1  $\beta$ -lactamase hydrolyzed oxyimino cephalosporins in *Enterobacter cloacae* (16) and *Citrobacter freundii* (23). The  $\beta$ -lactamase from *E. cloacae* P99 hardly hydrolyzes oxyimino cephalosporins, but this enzyme with an insertion of three residues just behind Arg 210 did hydrolyze them. Likewise, the  $\beta$ -lactamase from *E. cloacae* GC1, which was isolated clinically and which had a duplication of Ala-Val-Arg at the same position, showed the same hydrolytic activity as the P99  $\beta$ -lactamase with the three-residue insertion. The group 1  $\beta$ -lactamase from *C. freundii* GN346 hardly hydrolyzed oxyimino cephalosporins, but the artificial substitution of Glu for Lys at position 219 allowed the enzyme to hydrolyze oxyimino cephalosporins.

In a previous study, we reported that *Serratia marcescens* GN16694 is resistant to oxyimino cephalosporins. The organism was isolated clinically and produced SRT-1, a group 1  $\beta$ -lactamase (1) that hydrolyzed these cephalosporins and that had a molecular weight of 42,000 and a pI of 8.6 (14).

In the report described in this paper, we determined the nucleotide sequences of genes encoding SRT-1 and SST-1 (SST-1 was produced by the  $\beta$ -lactam-susceptible strain *S. marcescens* GN19450 and hardly hydrolyzed oxyimino cephalosporins) and describe the relationship between the extended substrate specificity and the differences in the amino acid residues in their sequences.

*S. marcescens* GN16694, which has a high level of resistance to oxyimino cephalosporins, is a clinical isolate (14). *S. marcescens* GN19450 is a clinical isolate and is susceptible to various  $\beta$ -lactams (6). *Escherichia coli* JM83 (24) was used for

DNA techniques. Plasmids pHSG398 (21) and pHSG399 (21) were used as the cloning vector and the vector for DNA sequencing, respectively. The following antimicrobial agents were used: streptomycin sulfate, cephaloridine, cefotaxime, ceftazidime, cefmenoxime, and cefuroxime.

Susceptibility testing, the purification of  $\beta$ -lactamases from *S. marcescens* GN16694 and GN19450 and from *E. coli* JM83/pGFS11, JM83/pGFR5, and JM83/pGFSK, and studies of their kinetics were based on a previously described method (14).  $\beta$ -Lactamase activities were assayed spectrophotometrically. The kinetic parameters were determined with a Lineweaver-Burk plot.

Plasmid DNA was prepared by the rapid alkaline extraction method (11). Restriction endonucleases were obtained from Nippon Gene Co. Ltd., Toyama, Japan, and the DNA-Ligation Kit was obtained from Takara Shuzo Co. Ltd., Kyoto, Japan. DNA techniques were done according to the manufacturer's recommendation. For the cloning of the genes for SRT-1 and SST-1, *Bam*HI-digested fragments of DNAs from *S. marcescens* GN16694 and GN19450, respectively, were ligated into the *Bam*HI site of pHSG398. These recombinants were introduced into *E. coli* JM83, and transformants with the gene for SRT-1 or SST-1 were selected on Luria-Bertani agar plates containing chloramphenicol (30  $\mu$ g/ml) and ceftazidime (6.25  $\mu$ g/ml) or cephaloridine (12.5  $\mu$ g/ml), respectively. These resultant transformants maintained a plasmid with a 4.0-kbp fragment carrying the gene for SRT-1 or SST-1. These recombinant plasmids with the gene for SRT-1 and SST-1 were termed pGFR5 and pGFS11, respectively. These fragments were subcloned into pHSG399 to sequence both strands.

Double-stranded plasmid DNA templates for the sequences were constructed by using the Kilo-sequence Deletion Kit (Takara Shuzo). The nucleotide sequences were determined by the dideoxy chain termination method (20) by using the TAKARA *Taq* Cycle Sequencing kit for the Shimadzu DNA Sequencer, version 2, M13 forward and reverse fluorescence primers (Takara Shuzo), and a DSQ-1000 DNA Sequencer (Shimadzu, Kyoto, Japan). The nucleotide sequence data were organized and analyzed by using DNASIS (Hitachi Software Engineering Co. Ltd., Yokohama, Japan).

The oligonucleotide primer composed of a 21-mer (CTGG ACGCCAAATCTTACGGC) was used for site-directed mutagenesis. This primer corresponded to the DNA sequence encoding amino acid residues of 210 to 216 in SRT-1, and an

\* Corresponding author. Mailing address: Research Laboratories, Toyama Chemical Co., Ltd., 2-4-1 Shimo-okui, Toyama-city, Toyama, 930 Japan. Phone: 81-764-31-8268. Fax: 81-764-31-8208.




FIG. 1. Alignment of amino acids of SRT-1, SST-1, and the group 1  $\beta$ -lactamase, indicated SR50, from *S. marcescens* SR50 (15). Dashes indicate residues identical to those of SRT-1. The position of the N-terminal amino acid of the mature enzymes is designated position 1. The amino acid sequence from positions -22 to -1 is assumed to be the signal peptide.

attempt was made to anneal the primer to the corresponding region of the SST-1 gene. Site-directed mutagenesis was carried out by using a modification of the manufacturer's information for Mutan-Express Km (Takara Shuzo) by using pGFS11 with the gene for SST-1 as a template.

Within the 4.0-kbp DNA fragments, a 1,290-bp segment containing the gene for SRT-1 and a 1,288-bp segment containing the gene for SST-1 were sequenced. These sequenced regions each embraced an open reading frame, and it was deduced that they were composed of 379 amino acids (Fig. 1). The amino acid sequence of the group 1  $\beta$ -lactamase (cephalosporinase) from *S. marcescens* SR50 that hardly hydrolyzes oxyimino cephalosporins has been presented previously (1, 15). The amino-terminal region from positions -22 to -1 was assumed to be the signal peptide because a typical sequence (A-X-A) was found, and this sequence was recognized by the signal peptidase. Consequently, it was supposed that in SRT-1 and SST-1 the mature proteins are composed of 357 amino acids. The amino acid sequence of SST-1 showed an identity of 96% to that of SRT-1. There was a high degree of similarity of the amino acid sequence of SST-1 to those of the group 1  $\beta$ -lactamases from *S. marcescens* SR50 (92% identity) (15), *C. freundii* (42% identity) (9, 22), and *Pseudomonas aeruginosa* (49% identity) (10).

Relative rates of hydrolysis by SRT-1 and SST-1 prepared from *S. marcescens* GN16694 and GN19450, respectively, showed the enzymes have very different hydrolytic profiles. SRT-1 also hydrolyzed cephaloridine and oxyimino cephalosporins such as ceftazidime and cefotaxime, showing a higher degree of hydrolysis of these drugs than of cephaloridine. On

the other hand, SST-1 hardly hydrolyzed oxyimino cephalosporins (data not shown).

To study the relationship between the expansion of the substrate specificity of SRT-1 in comparison with that of SST-1 and the differences in the amino acid residues in their sequences, the amino acid sequences in the region of the mature enzymes were compared. There were differences at 11 amino acid residues (Fig. 1). At positions 120, 164, 178, 197, 348, and 354, the amino acid residues in SRT-1 were different from those in SST-1, whereas those in SRT-1 were the same as those in the group 1  $\beta$ -lactamase of *S. marcescens* SR50. At positions 121, 149, 243, and 246, these residues in SRT-1 differed from those in SST-1 and the enzyme of *S. marcescens* SR50. In a comparison of the amino acid sequences of  $\beta$ -lactamases with the same substrate specificity produced by the same genus of bacilli, it was found that there is not necessarily complete identity among them (3, 9, 16, 22). Although SST-1 is the same as the enzyme from *S. marcescens* SR50 with regard to its low level of hydrolysis of oxyimino cephalosporins, there is disagreement between their amino acid sequences (92% identity). These findings suggest that the differences between these residues in SRT-1 and those in SST-1 and the enzyme of *S. marcescens* SR50 has little effect on the expansion of the substrate specificity toward oxyimino cephalosporins. However, at position 213, the amino acid residue was Lys in SRT-1, whereas it was Glu in SST-1 and the enzyme of *S. marcescens* SR50. Since Lys is a basic amino acid and Glu is acidic, resulting in a potentially different character, we anticipated that the Lys at position 213 contributed to the expansion of the substrate specificity in SRT-1, and the substitution of Glu in SST-1 into Lys was carried out by site-directed mutagenesis.

By site-directed mutagenesis of the gene encoding SST-1, a variant termed SST-K was obtained. This variant hydrolyzed oxyimino cephalosporins. DNA sequencing certified that the sequence of the region encoding residues 210 to 216 in SST-K was the same as that of the gene for SRT-1, and the amino acid at residue 213 was Lys.

Table 1 presents the susceptibilities of *E. coli* JM83 producing SST-K, SRT-1, and SST-1. The oxyimino cephalosporins MICs for *E. coli* JM83/pGFSK (the strain producing SST-K) were higher than those for parent strain (JM83/pHSG398) and the same as those for JM83/pGFR5 (the strain producing SRT-1). The MICs of cefotaxime, cefuroxime, and cefmenoxime for *E. coli* JM83/pGFS11 (the strain producing SST-1) were also elevated. The cause of resistance would be high affinities of SST-1 for these antibiotics (Table 2).

Table 2 presents the values of the kinetic parameters for SST-K, SRT-1, and SST-1. SST-K acquired an extended substrate specificity (to oxyimino cephalosporins), whereas SST-1 had a substrate specificity typical of those of a cephalosporinase (12, 13) and the group 1  $\beta$ -lactamase (1). The affinities of SST-K for cefuroxime, cefotaxime, and cefmenoxime were greatly reduced in comparison with those of SST-1. These

TABLE 1. Susceptibility of *E. coli* harboring the recombinant plasmids to  $\beta$ -lactams

<i>E. coli</i> strain ( $\beta$ -lactamase)	MIC ( $\mu$ g/ml)			
	Cefotaxime	Cefuroxime	Cefmenoxime	Ceftazidime
JM83/pGFSK <sup>a</sup> (SST-K)	3.13	200	0.78	50
JM83/pGFR5 (SRT-1)	3.13	200	0.78	50
JM83/pGFS11 (SST-1)	3.13	100	0.39	0.39
JM83/pHSG398	<0.05	3.13	<0.05	0.2

<sup>a</sup> The recombinant carrying the gene for SST-K.

TABLE 2. Values of kinetic parameters for SST-K, SRT-1, and SST-1

Substrate	SST-K		SRT-1		SST-1		
	$K_m$ ( $\mu$ M)	Relative $V_{max}^a$	$K_m$ ( $\mu$ M)	Relative $V_{max}^a$	$K_m$ ( $\mu$ M)	$K_i$ ( $\mu$ M)	Relative $V_{max}^a$
Cephaloridine	670	100	660	100	180	— <sup>b</sup>	100
Cefuroxime	370	340	330	350	—	0.3	<1
Cefotaxime	390	400	410	410	—	2.8	<1
Cefmenoxime	180	120	170	130	—	1.7	<1
Ceftazidime	180	20	150	20	—	120	<1

<sup>a</sup> Values relative to that for cephaloridine, which was set at 100.

<sup>b</sup> —, not tested.

kinetic parameters for SST-K were in good agreement with those for SRT-1. These results indicate that the change of Glu into Lys at residue 213 is a means for the hydrolysis of oxyimino cephalosporins.

The work presented here suggests that SRT-1 is a clinically occurring mutant of a group 1  $\beta$ -lactamase without the ability to hydrolyze oxyimino cephalosporins, and the expansion of the substrate specificity in SRT-1 toward oxyimino cephalosporins is attributable to the change of Glu into Lys at position 213. Table 3 presents four motifs that are conserved in amino acid sequences of group 1  $\beta$ -lactamases from various gram-negative bacteria (4, 8). Position 213 in SRT-1 and SST-1 corresponds to the third amino acid residue in the third motif. A change of Glu to Lys at this position allowed the enzyme to extend the substrate specificity in the case of SST-1. It was reported that the change of Glu to Lys at position 219 extended the substrate specificity of the group 1  $\beta$ -lactamase from *C. freundii* GN346 (23). Position 219 would also correspond to that in the third motif. As indicated in Table 3, the amino acid sequences of the third motif were highly conserved among the group 1  $\beta$ -lactamases from *E. coli*, *Enterobacter cloacae*, *P. aeruginosa*, *Klebsiella pneumoniae*, and *C. freundii*. Moreover, except for the enzyme from *E. cloacae*, the third amino acid residue in the third motif in their enzymes is Glu. In the future, it is suggested that the Glu at this position in these  $\beta$ -lactamases could be changed into Lys by some clinical

TABLE 3. Alignment of the four motifs conserved in amino acid sequences of group 1  $\beta$ -lactamases<sup>a</sup>

Source ( $\beta$ -lactamase)	Amino acid sequence at the following motif:				Refer- ence(s)
	First	Second	Third	Fourth	
<i>E. coli</i>	SVSK	YAN	DAEA	KTG	7
<i>E. cloacae</i>	SISK	YAN	DAQA	KTG	3
<i>P. aeruginosa</i>	SVSK	YSN	DAEG	KTG	10
<i>K. pneumoniae</i> (FOX-1)	SVSK	YSN	AAEA	KTG	8
<i>C. freundii</i>	SVSK	YAN	DAE(K) <sup>b</sup> A	KTG	9, 22
<i>S. marcescens</i> SR50	58-SLSK	144-YSN	211-DAES	308-KTG	15
<i>S. marcescens</i> GN19450 (SST-1)	SLSK	YSN	DAES	KTG	This study
<i>S. marcescens</i> GN16694 (SRT-1)	SLSK	YSN	DAKS	KTG	This study

<sup>a</sup> The four motifs are in alignment from the upper position in their amino acid sequences.

<sup>b</sup> In  $\beta$ -lactamases with the extended substrate specificity for oxyimino cephalosporins, E was substituted for K in the third motif.

action; consequently, they would acquire the expanded substrate specificities.

**Nucleotide sequence accession number.** The nucleotide sequences of SRT-1 and SST-1 have been given nucleotide sequence accession nos. AB008454 and AB008455, respectively.

We are grateful for financial support from Toyama Chemical Co., Ltd.

We thank S. Iyobe and H. Yamada for technical support and valuable advice.

## REFERENCES

- Bush, K., G. A. Jacoby, and A. A. Medeiros. 1995. A functional classification scheme for  $\beta$ -lactamases and its correlation with molecular structure. *Antimicrob. Agents Chemother.* **39**:1211–1233.
- Fosberry, A. P., D. J. Payne, E. J. Lawlor, and J. E. Hodgson. 1994. Cloning and sequence analysis of bla<sub>BIL-1</sub>, a plasmid-mediated class C  $\beta$ -lactamase gene in *Escherichia coli* BS. *Antimicrob. Agents Chemother.* **38**:1182–1185.
- Galleni, M., F. Lindberg, S. Normark, S. Cole, N. Honore, B. Joris, and J.-M. Frere. 1988. Sequence and comparative analysis of three *Enterobacter cloacae ampC*  $\beta$ -lactamase genes and their products. *Biochem. J.* **250**:753–760.
- Ghuysen, J.-M. 1991. Serine  $\beta$ -lactamases and penicillin-binding proteins. *Annu. Rev. Microbiol.* **45**:37–67.
- Horii, T., Y. Arakawa, M. Ohta, S. Ichiyama, R. Wacharotayankun, and N. Kato. 1993. Plasmid-mediated AmpC-type  $\beta$ -lactamase isolated from *Klebsiella pneumoniae* confers resistance to broad-spectrum  $\beta$ -lactams, including moxalactam. *Antimicrob. Agents Chemother.* **37**:984–990.
- Iyobe, S., M. Tsunoda, and S. Mitsuhashi. 1994. Cloning and expression in Enterobacteriaceae of the extended-spectrum  $\beta$ -lactamase gene from a *Pseudomonas aeruginosa* plasmid. *FEMS Lett.* **121**:175–180.
- Jaurin, B., and T. Grundstrom. 1981. *ampC* cephalosporinase of *Escherichia coli* K-12 has a different evolutionary origin from that of  $\beta$ -lactamases of the penicillinase type. *Proc. Natl. Acad. Sci. USA* **78**:4897–4901.
- Leiza, M. G., J. C. Perez-Diaz, J. Ayala, J. M. Casellas, J. Martinez-Beltran, K. Bush, and F. Baquero. 1994. Gene sequence and biochemical characterization of FOX-1 from *Klebsiella pneumoniae*, a new AmpC-type plasmid-mediated  $\beta$ -lactamase with two molecular variants. *Antimicrob. Agents Chemother.* **38**:2150–2157.
- Lindberg, F., and S. Normark. 1986. Sequence of the *Citrobacter freundii* OS60 chromosomal *ampC*  $\beta$ -lactamase gene. *Eur. J. Biochem.* **156**:441–445.
- Lodge, J. M., S. D. Minchin, L. J. V. Piddock, and S. J. Busby. 1990. Cloning, sequencing and analysis of the structural gene and regulatory region of the *Pseudomonas aeruginosa* chromosomal *ampC*  $\beta$ -lactamase. *Biochem. J.* **272**:627–631.
- Maniatis, T., E. F. Fritsch, and J. Sambrook. 1982. Molecular cloning: a laboratory manual. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Mitsuhashi, S., and M. Inoue. 1981. Mechanism of resistance to  $\beta$ -lactam antibiotics, p. 41–56. *In* S. Mitsuhashi (ed.), *Beta-lactam antibiotics*. Japan Sciences Press, Tokyo, Japan.
- Mitsuhashi, S. 1985. Resistance to  $\beta$ -lactam antibiotics in bacteria, p. 3–9. *In* J. Ishigami (ed.), *Recent advances in chemotherapy*. University of Tokyo Press, Tokyo, Japan.
- Matsumura, N., and S. Mitsuhashi. 1995. A  $\beta$ -lactamase from *Serratia marcescens* hydrolyzing the 2-carboxypenam T-5575. *Antimicrob. Agents Chemother.* **39**:2132–2134.
- Nomura, K., and T. Yoshida. 1990. Nucleotide sequence of the *Serratia marcescens* SR50 chromosomal *ampC*  $\beta$ -lactamase gene. *FEMS Microbiol. Lett.* **70**:295–300.
- Nukaga, M., S. Haruta, K. Tanimoto, K. Kogure, K. Taniguchi, M. Tamaki, and T. Sawai. 1995. Molecular evolution of a class C  $\beta$ -lactamase extending its substrate specificity. *J. Biol. Chem.* **270**:5729–5735.
- Papanicolaou, G. A., A. A. Medeiros, and G. A. Jacoby. 1990. Novel plasmid-mediated  $\beta$ -lactamase (MIR-1) conferring resistance to oxyimino- and  $\alpha$ -methoxy  $\beta$ -lactams in clinical isolates of *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* **34**:2200–2209.
- Sanders, C. C. 1987. Chromosomal cephalosporinases responsible for multiple resistance to newer  $\beta$ -lactam antibiotics. *Annu. Rev. Microbiol.* **41**:573–593.
- Sanders, C. C. 1992.  $\beta$ -Lactamases of gram-negative bacteria: new challenges for new drugs. *Clin. Infect. Dis.* **14**:1089–1099.
- Sanger, T., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
- Takeshita, S., M. Sato, M. Toba, W. Masahashi, and T. Hashimoto-Gotoh. 1987. High-copy-number plasmid vectors for *lacZ*  $\alpha$ -complemen-

- tation and chloramphenicol- or kanamycin-resistance selection. *Gene* **61**: 63–74.
22. **Tsukamoto, K., K. Tchibana, N. Yamazaki, Y. Ishii, K. Ujiie, N. Nishida, and T. Sawai.** 1990. Role of lysine-67 in the active site of class C  $\beta$ -lactamase from *Citrobacter freundii* GN346. *Eur. J. Biochem.* **188**:15–22.
  23. **Tsukamoto, K., R. Ohono, and T. Sawai.** 1990. Extension of the substrate spectrum by an amino acid substitution at residue 219 in the *Citrobacter freundii* cephalosporinase. *J. Bacteriol.* **172**:4348–4351.
  24. **Vieira, J., and J. Messing.** 1982. The pUC plasmids, an M13mp7-derived system for insertion mutagenesis and sequencing with synthetic universal primers. *Gene* **19**:259–268.
  25. **Watanabe, M., S. Iyobe, M. Inoue, and S. Mitsuhashi.** 1991. Transferable imipenem resistance in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* **35**:147–151.