



Published in final edited form as:

Immunity. 2023 September 12; 56(9): 2054–2069.e10. doi:10.1016/j.immuni.2023.07.017.

A REGULATORY CIRCUIT CONTROLLED BY EXTRANUCLEAR AND NUCLEAR RETINOIC ACID RECEPTOR α DETERMINES T CELL ACTIVATION AND FUNCTION

Alexandre LARANGE^{1,2,10}, Ikuo TAKAZAWA^{1,2,10}, Kiyokazu KAKUGAWA^{3,10}, Nicolas THIAULT^{1,2,10}, SooMun NGOI⁴, Meagan E. OLIVE⁵, Hitoshi IWAYA^{1,2}, Laetitia SEGUIN^{1,2}, Ildefonso VICENTE-SUAREZ^{1,2}, Stephane BECART^{1,2}, Greet VERSTICHEL^{1,2}, Ann BALANCIO^{1,2}, Amnon ALTMAN^{1,2}, John T. CHANG⁴, Ichiro TANIUCHI⁶, Bjorn LILLEMEIER⁷, Mitchell KRONENBERG^{1,4,8}, Samuel A. MYERS^{1,5,9,*}, Hilde CHEROUTRE^{1,2,3,11,*}

¹Center for Autoimmunity and Inflammation, La Jolla Institute for Immunology, La Jolla, CA 92037, USA

²Center for Cancer Immunotherapy, La Jolla Institute for Immunology, La Jolla, CA 92037, USA

³Laboratory for Immune Crosstalk, RIKEN Center for Integrative Medical Sciences, 1-7-22 Suehiro, Tsurumi-ku, Yokohama 230-0045, Japan

⁴School of Biological Sciences, University of California San Diego, La Jolla CA 92093, USA

⁵Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, MA 02142, USA

⁶Laboratory for Transcriptional Regulation, RIKEN Center for Integrative Medical Sciences, 1-7-22 Suehiro, Tsurumi-ku, Yokohama 230-0045, Japan

⁷Immunobiology and Microbial Pathogenesis Laboratory, IMPL-L, Salk Institute for Biological Studies, La Jolla, CA 92037, USA

⁸Center for Infectious Disease and Vaccine Research, La Jolla Institute for Immunology, La Jolla, CA 92037, USA

*Correspondence: sam@lji.org (S.A.M.) and hilde@lji.org (H.C.).

AUTHOR CONTRIBUTIONS

Conceptualization, H.C., A.L., K.K., I. TAKAZAWA, N.T. and S.A.M.; Investigation, A.L., K.K., I. TAKAZAWA, N.T., M.E.O., S.A.M., H.I., L.S., B.L., I.V.S., S.B., S.N., and J.T.C.; Supervision, H.C., S.A.M., A.A., J.T.C., B.L. and I.T.; Project Administration, H.C.; Formal Analysis, H.C., S.A.M., J.T.C., and B.L.; Validation, K.K., I. TAKAZAWA, N.T., M.E.O., S.N., B.L. and S.A.M.; Methodology, K.K., A.L. and B.L.; Resources, A.B., A.A., B.L. and J.T.C.; Visualization, A.A., K.K., I. TAKAZAWA, N.T., G.V., B.L., S.N., J.T.C. and S.A.M.; Writing Original, H.C., A.A., K.K., I. TAKAZAWA and S.A.M.; Writing-Review and Editing, H.C., M.K., I.T., A.A., G.V. and N.T.; Funding Acquisition, A.A., I. TAKAZAWA and H.C.

DECLARATION OF INTERESTS

The authors declare no competing interests.

MATERIALS AVAILABILITY

All unique/stable reagents generated in this study are available from the lead contact with a completed material transfer agreement.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

⁹Laboratory for Immunochemical Circuits, La Jolla Institute for Immunology, La Jolla, CA 92037, USA

¹⁰These authors contributed equally.

¹¹Lead contact: hilde@lji.org


SUMMARY

Ligation of Retinoic acid receptor alpha (RAR α) by RA promotes varied transcriptional programs associated with immune activation and tolerance, but genetic deletion approaches suggest impact of RAR α in TCR signaling. Here we examined whether RAR α would exert roles beyond transcriptional regulation. Specific deletion of the nuclear isoform of RAR α revealed an RAR α isoform in the cytoplasm of T cells. Extranuclear RAR α was rapidly phosphorylated upon TCR stimulation and recruited to the TCR signalosome. RA interfered with extranuclear RAR α signaling, causing suboptimal TCR activation while enhancing FOXP3⁺ regulatory T cell conversion. TCR activation induced expression of CRABP2, which translocates RA to the nucleus. Deletion of *Crabp2* led to increased RA in the cytoplasm and interfered with signalosome-RAR α resulting in impaired anti-pathogen immunity and suppressed autoimmune disease. Our findings underscore the significance of subcellular RA/RAR α signaling in T cells and identify extranuclear RAR α as a component of the TCR signalosome and a determinant of immune responses.

eTOC blurb:

Nuclear retinoic acid receptor alpha (RAR α) regulates gene transcription upon binding of retinoic acid (RA). Larange, Takazawa, Kakugawa, Thiault et al., identify an RAR α isoform in the cytoplasm of T cells (cRAR α), and reveal that cRAR α binds the TCR-ZAP-70 complex thereby regulating T cell proliferation and effector differentiation. RA counteracts TCR-engaged RAR α resulting in suboptimal TCR activation but enhanced Treg cell differentiation.

Graphical Abstract



Keywords

Retinoic acid receptor alpha; Retinoic acid; extranuclear; T cell receptor; ZAP-70; signal transduction; proliferation; effector differentiation; FOXP3⁺Regulatory T cells; cellular retinoic acid binding protein; nuclear receptor; autoimmune disease; anti-pathogen immunity

INTRODUCTION

Retinoic acid receptor (RAR) signaling greatly impacts immune responses, ranging from protective immunity to immune tolerance, tissue homing and lymph node organogenesis(1). Defects in RAR functions weaken immunity against infections and cancers and lead to immune deregulation and inflammatory diseases(1–4).

RARs consist of three isotypes (RAR α , RAR β and RAR γ) encoded by separate genes, and each can give rise to several transcripts and alternative protein isoforms. Similar to other nuclear receptors, RARs exhibit a modular organization, comprising a conserved DNA-binding domain (DBD) and C-terminal ligand-binding domain (LBD), whereas the N-terminal region and F-domain vary greatly between different RAR isoforms(1).

At steady state, nuclear RAR α participates in transcription factor complexes that silence target genes. Upon ligation with the vitamin A metabolite, retinoic acid (RA), and in response to post-translational modulations, RAR α undergoes conformational changes of its LBD. These changes result in the release of corepressors and recruitment of coregulators

that alter the activity of the transcription complex(5). RA is present in blood and tissues and as a small molecule it can freely enter cells, but requires active transportation by cellular retinoic acid binding protein 2 (CRABP2) to translocate to the nucleus(6). CRABP2 not only transports RA, but it interacts with RAR α and specifically directs and facilitates RA ligation(6). This transactivates RAR α bound to retinoic acid responsive elements (RARE) in regulatory regions of RA target genes and promotes transcription(6, 7).

In T cells, RA ligation of RAR α activates transcription of signature genes, including *Ccr9* and *a4b7* that direct T cell migration to the intestine(8, 9). RA/RAR α signaling greatly affects proliferation and CD4⁺ T helper polarization(3, 10) especially influencing the reciprocal regulation of TGF- β -driven polarization by counteracting the generation of inflammatory T helper 17 (Th17) cells while promoting the differentiation of suppressive FOXP3-expressing regulatory T (Treg) cells(11–16). Although inhibition of pro-inflammatory cytokines produced by activated bystander T cells may indirectly contribute to enhanced FOXP3⁺Treg cell conversion(17), RA/RAR α signaling also promotes FOXP3 expression of purified naïve T cells activated in an antigen presenting cell (APC)-free system(18). Similar to interleukin (IL)-2-mediated control of Treg/Th17 cell polarization, the influence of RA/RAR α on FOXP3 expression coincides with activation of the transcription factor STAT5 and suppression of STAT3-driven IL-17 expression(19–22). However, the enhancing effect of RA on FOXP3 induction persists even in the absence of IL-2, STAT5 or STAT3(23). Moreover, while the interaction of RA with RAR α amplified TGF β -dependent phosphorylated SMAD3-binding to the enhancer I in the *Foxp3* locus(24), absence of SMAD3 expression does not prevent the promoting effect of RA/RAR α signaling on FOXP3⁺ Treg cell conversion(25). All together, these observations suggest that the main mode of action by RA/RAR α is not solely explained by the transcriptional control of *Foxp3* expression.

TCR signal strength also influences the induction of Treg cells. Absence of or low CD28 co-stimulation or transient reduction in the phosphatidylinositol-3-kinase (PI3K)/AKT pathway, favors Treg cell over T effector differentiation(25, 26). Notably, conditional deletion or overexpression of RAR α , impacts PI3K/AKT signaling in response to TCR stimulation and causes important downstream consequences for T cell activation, proliferation and differentiation(3, 27). These immediate effects downstream of the TCR suggest that RAR α signaling during T cell activation might involve non-genomic and extranuclear actions.

Here we examined whether RAR α would exert roles beyond transcriptional regulation. Specific deletion of the nuclear isoform of RAR α revealed the presence of RAR α in the cytoplasm. TCR stimulation activated extranuclear RAR α and promoted its recruitment to the immunological synapse (IS). Participation of extranuclear RAR α in the signalosome was critical for signal transduction at the plasma membrane and to promote downstream signaling cascades that subsequently also affected nuclear RAR α -controlled gene transcription. Effective extranuclear RAR α signaling was also required for TCR-induced CRABP2 expression, thereby constituting a circuit that induces RA responsiveness and RAR α -controlled gene transcription. In contrast to RA-mediated transactivation of nuclear RAR α , RA negatively impacted non-genomic signaling by RAR α in the cytoplasm,

leading to sub-optimal activation, reduced proliferation and impaired effector differentiation, but enhanced FOXP3⁺Treg cell conversion. These data identify extranuclear RAR α as a critical participant in the TCR signalosome and a determinant of T cell fate that ultimately controls the inflammatory or regulatory nature of the immune response.

RESULTS

RA enhances Treg cell conversion independently of nuclear RAR α

T cells with a global deletion of *Rara*, which eliminates all RAR α isoforms, do not enhance FOXP3 expression in the presence of exogenously added RA(3, 25). In contrast, T cells with a specific deletion of nuclear RAR α 1 (nRAR α -deficient) isoform responded with enhanced FOXP3 expression (Figure 1A and Figure S1A). These data suggest that transcriptional control by nuclear RAR α might not be the main mechanism that regulates RA-mediated enhanced FOXP3 expression. To address this possibility, we analyzed T cells that overexpressed a dominant negative (DN) isoform of RAR α (RAR403^{CD4-Cre}) that blocks the transcriptional function of RAR α (28). As expected, the presence of DNRAR α interfered with the transcription of *Ccr9*(Figure S1B). In contrast, it did not affect the RA-mediated increase in TGF- β -induced FOXP3 expression in naïve T cells activated *in vitro* in an APC-free system (Figure 1B). Notably, when the addition of RA was delayed by 24 hours, RA no longer enhanced FOXP3 expression (Figure 1C). Taken together the results suggest that RA targets an RAR α -function during early TCR stimulation independently of transactivation of nuclear RARs.

T cells express an extranuclear isoform of RAR α

To investigate physical participation of RAR α in early T cell activation, we used confocal imaging of ovalbumin peptide (OVA_p) (SIINFEKL)-responsive OT-I TCR transgenic CD8 $\alpha\beta$ T cells activated *in vitro* with OVA_p-loaded DCs and stained with RAR α -specific antibodies together with anti-CD3 ζ and anti-ZAP70 antibodies. Consistent with an extranuclear presence, RAR α was detected at the IS in close proximity to CD3e and ZAP-70 (Figure 2A). Colocalization of RAR α and CD3e alone, was not observed, suggesting an indirect link between RAR α and the TCR itself. FLAG-tagged constructs for nuclear *Rara1* and *Rara2* transduced in primary T cells that were activated with anti-CD3/CD28 and subsequently analyzed by western blotting, showed that RAR α 1 and - α 2 remained in the nucleus upon TCR stimulation (Figure 2B), and thus that there must be another *Rara* gene expressed in T cells, which encodes this cytoplasmic molecule. High-throughput sequencing and analysis of double positive (DP) thymocyte and CD4 and CD8 $\alpha\beta$ spleen T cell mRNA (RNA-seq) revealed that in addition to *Rara1* and -2 transcripts, there were other *Rara* sequences with different 5' splice junctions (Figure 2C). Three transcripts (J3, J4 and J5) aligned with predicted sequences in the RefSeq database (www.ncbi.nlm.nih.gov/refseq)(29). *Rara* transcripts that predominate in thymocytes were distinct from those in mature T cells (Figure S2A). All alternative transcript shared the same 3' end sequence with *Rara1* and -2 but each one had a unique 5' end sequence (Figure 2D), indicating a transcriptional start site (TSS) distinct from that of *Rara1* or -2 (Figure S2B). This was supported by data from the Functional ANnotation Of the Mammalian genome (FANTOM)5 database(30), which indicated a TSS downstream of the *Rara2* TSS and

upstream of the 5' end of the variant transcripts (Figure 2E). Complete sequence analysis specified a GC-rich 5' end with one or two variable non-translated exons (Figure S2C) and conserved exons 9 to 15 shared with *Rara1* and -2 (Figure 2E). All transcript variants encoded the same protein with a putative AUG translational start in exon 10 (Figures 2E and S2D). The alternative RAR α differed from nuclear RAR α by truncation with a predicted molecular weight of approximately 38kDa. There were no typical transcriptional features like a functional DNA binding domain (DBD) or nuclear translocation sequences (NLS). A complete ligand binding domain (LBD) and an intact F-domain however, were present suggesting that, similar to nuclear RAR α , this isoform potentially interacts with binding partners to form dynamic active (open) and inactive (closed) protein complexes. *Gfp*-tagged constructs of alternative *Rara*, referred to as *Rara3* (RAR α 3-eGFP) and *Rara1* (RAR α 1-eGFP) as control were transduced into mouse MCC-T CD4 hybridoma T cells(31) and analyzed by imaging for protein expression. As expected GFP-tagged RAR α 1 localized almost exclusively to the nucleus (Figure 2F). In contrast, a major fraction of GFP-tagged RAR α 3 was detected in the cytoplasm (Figure 2F), indicating that *Rara3* encodes an RAR α isoform that can reside outside the nucleus. Analysis of the cytoplasmic and nuclear fractions of human T cells confirmed RAR α in the cytoplasm of a similar size as mouse extranuclear RAR α (Figure 2G and 2H). Overall, the data indicate that T cells express a splice isoform of RAR α that is expressed in the cytoplasm.

Extranuclear RAR α controls TCR signal transduction

Primary T cells transduced with either *Rara1* or *Rara3* constructs were stimulated *in vitro* with anti-CD3+CD28 and proximal TCR signaling events were assessed. T cells, which overexpressed RAR α 3 but not RAR α 1, displayed markedly enhanced phosphorylation of ZAP-70 and phospholipase C gamma-1 (PLC γ 1) relative to the mock control (Figures 3A and 3B). Notably, *in vitro* stimulation of OT-I TCR transgenic T cells with OVAp-loaded DCs followed by immunoprecipitation with RAR α -specific antibody and western blotting for ZAP-70, indicated that RAR α antibody precipitated ZAP-70 from lysates of activated but not resting OT-I T cells (Figure 3C). These data showed that extranuclear RAR α influenced TCR/ZAP-70 activation and moreover that activated ZAP-70 directly or indirectly interacted with RAR α .

TCR downstream signal transduction depends on cytoskeletal remodeling and micro-domain scaffolds that recruit signaling molecules including kinases and adaptors(32–35). To further assess the influence of RAR α on the TCR-ZAP-70 activation complex, we added RAR α inhibitors to CD3 ζ -GFP transduced primary T cells stimulated *in vitro* with anti-CD3+CD28. In addition to the reduced *Ccr9* gene transcription (Figure S3A), RAR α inhibitor also reduced CD3 ζ -GFP clustering efficiency at the plasma membrane (Figure 3D) and notably disrupted the RAR α /ZAP-70 activation complex resulting in weakened TCR signal transduction (Figure 3E). Conversely, specific deletion of nuclear RAR α 1, which eliminated *Ccr9* transcription (Figure S1), had no effect on early TCR activation (Figure 3F). Moreover, transient blocking with specific RAR α -, but not RAR γ -inhibitor, reduced T cell proliferation (Figure 3G), and addition of IL-2 did not restore this defect (Figure 3H). IL-2-independent proliferation of activated T cells is mediated by TCR-induced activation of the NOTCH/cMYC pathway(34, 36). Accordingly, inhibition of RAR α signaling decreased

TCR-dependent cleavage of NOTCH (Figure 3I), to a similar degree as γ -secretase inhibitor (Figure S3B) and comprised upregulation of *c-Myc* mRNA and c-MYC protein expression (Figure 3J and Figure S3C). Overexpression of DNRAR α did not affect TCR-driven proliferation, which was only blocked when also the inhibitor was added (Figure 3K). Also, overexpression of DNRAR α affected neither TCR-mediated NOTCH activation (Figure 3L) nor upregulation of *cMyc* expression (Figure 3M).

All together, these results demonstrate that RAR α participates in TCR signal transduction, independently of its classical genomic function as a transcription factor.

Extranuclear and nuclear RAR α are activated by distinct mechanisms

RARs are phosphoproteins that alter their affinity for interacting binding partners through conformational changes induced by phosphorylation of specific sites(37). To assess phosphorylation of RAR α following TCR stimulation, we performed a large-scale quantitative phosphoproteomic analysis of Jurkat E6.1 T cells activated short term with anti-CD3 and anti-CD28 antibodies. Mass spectrometric analysis of whole lysates identified TCR-induced phosphorylation of a serine residue (S445) located within the F-domain of RAR α (Figure 4A). Phosphorylation of RAR α S445 (pS445) immediately increased roughly four-fold upon TCR stimulation but returned to basal levels after 10 minutes and continued to decline for some time after the initial stimulation (Figure 4B). Serine 445 is followed by a proline, indicating a proline directed serine kinase is likely responsible for the TCR-induced phosphorylation of the F domain (38). Thus, TCR stimulation leads to rapid phosphorylation of RAR α . Moreover, the fast kinetics of this activation process underscore the proximity and participation of RAR α in the TCR-ZAP-70 signalosome.

Unlike modulation of RAR α in response to TCR stimulation, transactivation of nuclear RARs is a direct consequence of RA ligation(1). T cell activation also leads to transcriptional activation of nuclear RAR α (3, 10). Accordingly, CD8 $\alpha\beta$ - and CD4 T cells from transgenic mice that expressed a luciferase transgene (Luc) transcriptionally controlled by a retinoic acid responsive element (RARE)(39), showed a gradual increase of RAR-controlled luciferase expression when activated with anti-CD3/CD28 in an APC-free system (Figure 4C). Also, spleen T cells from OT-I/RARE-Luc double transgenic mice induced increasing levels of luciferase after infection with OVA antigen-expressing *Listeria monocytogenes* (Lm-OVA) (Figure 4D). In contrast to the rapid activation of RAR α in the TCR/ZAP-70 complex, RA-dependent transactivation of nuclear RAR α was a late event measurable after 24 hours and peaked several days later (Figure 4C and 4D). RA is present at low physiological concentrations (<1nM) in blood and in most tissues and freely enters the cytoplasmic space of cells. Subcellular distribution and local concentration of RA are regulated by cellular retinoic acid binding proteins (CRABPs). CRABP1 targets RA for degradation via the CYP26 pathway(6, 40), whereas binding of RA to CRABP2 induces translocation of the CRABP2/RA complex to the nucleus and results in transactivation of nuclear RARs and RAR-controlled gene transcription(41, 42). Analysis of naïve or anti-CD3+CD28 stimulated T cells showed no CRABP1 expression (Figure 4E and 4F), but *Crabp1* mRNA was rapidly induced when triggered by IL-7 cytokine (Figure 4F). Expression of *Crabp2* mRNA was also insignificant at steady state but strongly induced

by TCR stimulation (Figure 4G–4I), whereas blocking of proximal TCR signaling at the plasma membrane with γ -secretase inhibitor (Figure 4J) or further downstream with ERK inhibitor (Figure 4K), interfered with CRABP2 induction. Both *Crabp2* mRNA and CRABP2 protein (Figure 4H and 4I), remained highly expressed during later stages of activation indicating that activation of the TCR promotes constant translocation of RA from the cytoplasm to the nucleus. TCR signals counteracted RA degradation by suppressing IL-7-induced CRABP1 expression (Figure 4F) whereas addition of IL-7 suppressed TCR-induced CRABP2 expression (Figure 4L). Altogether, the dose and subcellular location of RA is highly regulated in T cells. Furthermore, TCR activation induces CRABP2 expression, which removes RA from the cytoplasm and facilitates ligation of RA with nuclear RAR α .

RA counteracts extranuclear RAR α signaling in T cells

T cells with a genetic deletion for *Crabp2* failed to efficiently induce RA responsiveness, as indicated by reduced *Ccr9* expression when activated in the presence of a pharmacological dose (10 nM) of RA (Figure 5A and S4A). Also, RARE-Luc T cells on a *Crabp2*-deficient background showed inefficient *luciferase* gene transcription when stimulated with anti-CD3 and -CD28 in regular culture medium containing a physiological dose of RA (Figure 5B). TCR $\gamma\delta$ and TCR $\alpha\beta$ thymocytes and Foxp3⁺ natural Treg cell develop normally in CRABP2-deficient mice (Figure S4B), indicating that CRABP2 function influences mature T cells and enables RA-responsiveness associated with TCR $\alpha\beta$ stimulation. Accordingly, blocking of proximal TCR signaling diminished CRABP2/RA-dependent *luciferase* gene transcription (Figure 5C). Notably, RAR α inhibitors interfered with TCR-induced CRABP2 expression (Figure 5D), thereby linking the TCR-mediated activation of extranuclear RAR α with RA-responsiveness and the RA-dependent transactivation of nuclear RAR α .

To examine if RA also influences RAR α in the signalosome, we analyzed TCR signal transduction events in WT T cells activated *in vitro* with anti-CD3 and -CD28 together with a pharmacological dose (10nM) of RA. Contrary to its promoting effect on gene transcription, RA interfered with signaling events at the plasma membrane, including phosphorylation of phospholipase C gamma-1 (PLC γ 1) and AKT S473 (Figure 5E). Conversely, the absence or impaired transactivation of nuclear RAR α in *Rara1*-deletion mutant (Figure 5F) or dn*Rara* T cells (Figure 5G), respectively, had no measurable effect on proximal TCR/CD28 signaling. Addition of RA after 24 hours of stimulation, when CRABP2 was induced, no longer interfered with proximal signaling (Figure 5H), indicating that by translocating RA to the nucleus, CRABP2 eliminated the inhibitory effect of RA on early TCR signaling. Accordingly, in CRABP2-deficient T cells, RA added after 24 hours of stimulation continued to interfere with TCR signal transduction (Figure 5H).

Sub-optimal TCR activation promotes Treg cell conversion(25, 26). It is thus conceivable that the mechanism that drives the RA-enhanced FOXP3 Treg cell differentiation of naïve T cells might be due to the negative influence of RA on proximal TCR signaling. In agreement with this, RA-enhanced Treg cell conversion coincided with decreased activation-induced proliferation (Figure 5J). This was even more pronounced when RA accumulated in the cytoplasm of activated CRABP2-deficient T cells (Figure 5I). Notably, with inefficient

nuclear translocation and impaired transactivation of nuclear RAR α , RA still enhanced Treg cell conversion of CRABP2-deficient T cells (Figure 5J), even when added 24 hours after the initial stimulation (Figure 5K). Overall, these results indicated that the absence of CRABP2 expression or a large dose of RA interfere with the function of RAR α in the TCR signalosome resulting in suboptimal activation and reduced proliferation but enhanced Treg cell conversion *in vitro*.

Extranuclear RAR α signaling controls TCR-induced proliferation *in vivo*

To examine the significance of extranuclear RAR α signaling and the impact of RA interference during T cell responses *in vivo*, we compared the response of CRABP2-deficient and WT T cells in a pathogen infection. Naïve OT-I TCR transgenic T cells on a CRABP2-deficient background and WT OT-I T cells were co-transferred into C57BL/6 recipient mice that were subsequently infected with Lm-OVA (Figure 6A). When analyzed 7 days later, the frequency of CRABP2-deficient OT-I T cells was much reduced compared to WT OT-I cells (Figure 6B), indicating a cell-intrinsic defect consistent with impaired TCR-driven proliferation and or less survival. To test if the activation defect was due to the accumulation of RA in the cytoplasm in the absence of TCR-induced CRABP2 expression, we co-transferred WT and CRABP2-deficient naïve OT-I T cells into vitamin A-depleted (Vit A⁻) recipient animals, which had not received vitamin A since before birth. Vit A⁻ recipient mice were subsequently infected with Lm-OVA and analyzed 7 days later. As expected, without RA, homing of WT OT-I T cells to the intestinal epithelium was impaired (Figure 6C). Conversely, in the absence of RA, Lm-OVA responding CRABP2-deficient OT-I T cells in the spleen expanded vigorously, even to a greater extent than WT OT-I cells (Figure 6D). The frequency of recipient-derived OVA-reactive CD8 T cells did not differ between Vitamin A-sufficient or -depleted conditions (Figure S5), indicating that the reversed frequency ratio of donor WT and CRABP2-deficient OT-I T cells in VitA⁻ mice was not due to excessive cell death of the WT cells in the absence of RA. Re-supplementing RA restored intestinal homing of the OT-I T cells (Figure 6C) and prevented again the relative increase in number of Lm-OVA responding CRABP2-deficient OT-I T cells compared to WT OT-I cells (Figure 6E). The results underscored the significance of activation induced CRABP2 to prevent accumulation of RA in the cytoplasm and interference of RA with RAR α signaling in the TCR signalosome. The impaired proliferation and/or survival of the Lm-responding CRABP2-deficient T cells is in contrast to the vigorous expansion of Lm-responding *dnRara* T cells(43) and underscores the significance of participation of extranuclear RAR α in the TCR signalosome as a critical determinant of the magnitude of the T cell response *in vivo*.

Extranuclear RAR α signaling controls effector differentiation *in vivo*

To investigate the impact of cell intrinsic RAR α signaling during T cell activation and effector differentiation *in vivo*, we induced EAE in mice with a T cell-specific deletion of *Crabp2*, using distal *Lck-Cre* for specific deletion of *Crabp2* in mature T cells. Mice with a conditional deletion of *Crabp2* and WT mice were immunized subcutaneously with myelin oligodendrocyte glycoprotein (MOG₃₅₋₅₅) peptide in complete Freund's adjuvant and pertussis toxin on day 0 and day 2. Onset of disease was similar for *Crabp2* conditional deletion mutant mice and WT mice however disease did not progress in *Crabp2* conditional

deletion mutant animals (Figure 7A and Figure S6A). In contrast by day 25, some of the WT mice had developed severe symptoms, including paralyzed hind limbs (Figures 7B and 7C). Analysis of the mice at the peak of disease in the WT animals, showed more cell infiltration, including more T cells, in the spinal cord (Figure 7D and Figure S6B) and an overall higher histological disease score (Figure 7E) compared to *Crabp2* conditional deletion mutant mice. Moreover, the proportion of FOXP3⁺ROR γ t⁻ Treg cells was increased in several of the *Crabp2* conditional deletion mutant mice compared to the WT mice (Figure S6C), whereas in contrast, the absolute number (Figure 7F) and proportion of IFN γ -producing cells (Figure S6D) was reduced in the *Crabp2* conditional deletion mutant mice. Unlike Lm-responding dn*Rara* effector cells, however, which showed conversion of IFN γ -expressing cells to ROR γ t and IL-17 expressing cells(43), *Crabp2* conditional deletion mutant CD4 T cells in this EAE model did not convert more to inflammatory Th17 cells (Figure S6C). Instead comparison of the ratios of effector cells (IFN γ ⁺) over non-effector cells (IFN γ ⁻ and IL-17⁻) at day 25 of disease induction, indicated a significant overall decrease in inflammatory effector differentiation of the CRABP2 *Crabp2* conditional deletion mutant cells compared to WT cells (Figure 7G).

The global weakened effector response in *Crabp2* conditional deletion mutant mice highlights the essential role of extranuclear RAR α as a critical TCR-induced component of the signalosome required for T cell activation, expansion and optimal effector differentiation *in vivo*.

DISCUSSION

Here we identified an isoform of RAR that plays decisive roles in the mechanisms that control T cell activation *in vitro* and *in vivo*. Alternative RAR α is expressed in the cytoplasm and rapidly phosphorylated upon TCR stimulation. Extranuclear RAR α interacts with a ZAP-70-activation complex and participates in signal transduction during TCR-driven proliferation and effector differentiation. In contrast to RA-dependent transcriptional activation of nuclear RARs, RA negatively influences RAR α at the plasma membrane, which results in suboptimal activation and enhanced Treg cell differentiation. Transactivation of nuclear RAR α by RA, is important to control effector gene expression programs and depends on nuclear translocation of RA by TCR-induced CRABP2 expression. The non-genomic and genomic functions of RAR α have distinct kinetics, with actions of cytoplasmic RAR α occurring much faster. The decisive effects of extranuclear and nuclear RAR α signaling in T cells underscore the importance of their spatial and temporal bipartite regulation in controlling appropriate T cell responses to immune challenges.

Blocking of RAR α function with inhibitors known to interfere with structural dynamics of nuclear RARs also disrupted the RAR α /ZAP-70 complex, impaired phosphorylation of AKT and other signaling components downstream of the TCR and interfered with rapidly-induced actin-based events at the plasma membrane, including TCR/CD3-microdomain clustering(44, 45) and non-canonical ligand-independent cleavage of NOTCH(34, 36, 46, 47). These findings suggest that, similar to transcription-complex-building by nuclear RAR α , the mode of action of signalosome RAR α might also consist of a structural

interactome that coordinates functions of various cytoplasmic binding partners downstream of the TCR. These non-genomic roles of extranuclear RAR α are consistent with previously described findings showing that early T cell activation was influenced by the absence or overexpression of RAR α (3, 4, 27) and negatively impacted by RA(27).

Cytoplasmic RAR α in T cells is encoded by multiple transcripts with variable 5' non-translated ends. Each transcript encodes the same truncated RAR α isoform, which lacks typical transcriptional features, including a complete DBD and a functional NLS. Conversely, the conserved LBD, which interacts with binding partners and the regulatory F-domain are fully present. The F domain is unique and highly variable in length and sequence in different RAR proteins(5). Although its function is not fully understood, post-translational modifications of the F domain have been reported to counteract the transcriptional activity of nuclear RAR α and RA(5). Several putative phosphorylation sites for proline-directed kinases (Pdk) were previously identified in the F domain of RAR α (48). Phosphorylation of these sites is independent of RA ligation, and instead RA prevents the phosphorylation of the F region(48). Signals that lead to F domain phosphorylation had not been identified, but here we showed that phosphorylation of a predicted Pdk phosphorylation site (serine 445) in the F domain occurred within minutes of TCR stimulation. Phosphorylation of the F domain coincided with the interaction of RAR α with activated ZAP-70, suggesting that TCR-induced modulation of the F domain might facilitate the dynamic interaction of RAR α LBD with a ZAP-70-containing activation complex.

The current findings expand the repertoire of non-genomic actions by nuclear receptor superfamily members(49) to include participation of extranuclear RAR α in the TCR signalosome. Rapid, non-transcriptional functions for RAR α in actin remodeling were described in human platelets(50) and in neurons during the cleavage of amyloid preprotein by secretases(51), a process that parallels that of TCR-mediated NOTCH cleavage. In mouse embryonic fibroblasts and human mammary cancer cells, RAR α cooperates with G protein alpha Q (G α q) and directs Rho-GTPases and p38MAPK signaling(52), and in neuroblastoma cells extranuclear RAR α interacts with phosphatidylinositol 3-kinase subunit, p85 and recruits the catalytic subunit, p110 to the plasma membrane resulting in AKT and ERK1/2 phosphorylation(53). All these observations are consistent with participation of extranuclear RAR α in protein complexes that affect cytoskeleton remodeling and downstream signaling cascades in various cell types, similar to its role in the TCR activation process.

RA targets cytoplasmic RAR α resulting in diminished TCR signal strength and enhanced Treg cell induction, consistent with other studies linking reduced TCR signals with Treg cell differentiation(25, 54–56).

In contrast to the rapid TCR-mediated activation of extranuclear RAR α , transactivation of nuclear RAR α is much later and requires ligation by RA(1). Translocation and ligation are promoted by CRABP2(6, 41). We show here that CRABP2 is not expressed in resting T cells, but rapidly induced upon activation. TCR-induced CRABP2 not only established RA-responsiveness but it also prevented accumulation and interference of RA with extranuclear RAR α signaling. In the absence of CRABP2 induction, RA translocation and transactivation

of nuclear RAR α is greatly impaired. However, accumulation of RA in the cytoplasm of CRABP2-deficient T cells did not impact the RA-enhanced TGF β -induced Treg cell conversion, underscoring that transcriptional control by nuclear RARs is not the main mechanism for RA-mediated enhanced Foxp3 expression. Instead the data are consistent with a negative impact of a high dose of RA in the cytoplasm on the non-genomic function of extranuclear RAR α in the TCR signalosome, which led to suboptimal activation and impaired effector differentiation and survival but favored RA-enhanced Treg cell conversion. In WT T cells, TCR stimulation resulted in CRABP2 induction, which transports RA to the nucleus and away from the TCR signalosome. Addition of RA after CRABP2 induction no longer led to reduced signaling and no longer enhanced Treg cell conversion. In contrast, in pre-activated CRABP2-deficient T cells, RA accumulation in the cytoplasm counteracted extranuclear RAR α signaling and continued to promote Treg cell conversion, even when added after 24hrs of pre-activation. Interference with extranuclear RAR α signaling not only favored Treg cell conversion by weakening TCR signaling, but also hampered TCR-induced CRABP2 expression and consequently reduced transactivation of nuclear RAR α . Lack of TCR-induced CRABP2 expression significantly compromised the extranuclear RAR α action in TCR-induced activation and effector differentiation which caused a global suppression of EAE in *Crabp2* conditional deletion mutant mice.

The non-genomic role of RAR α increases the scope of its biological functions in T cells beyond its role as a regulator of gene transcription. A paradigm unfolds by which extranuclear RAR α signaling integrates early TCR-induced signaling events at the plasma membrane with subsequent nuclear events, coordinated by CRABP2. In transporting RA from the cytoplasm to the nucleus, TCR-induced CRABP2 functions as an intrinsic trait of the spatial and temporal regulation of RAR α signaling during T cell activation, enabling continuing participation of extranuclear RAR α in the TCR signalosome and promoting RA-dependent transactivation of nuclear RAR α . The findings reveal a fundamental process for extranuclear and nuclear RAR α signaling in control of both regulatory and effector arms of adaptive T cell immunity.

Limitation of the Study:

The cRAR α sequence is imbedded in the common sequence shared by all RAR α isoforms. Because deletion of a common exon is lethal, we attempted alternative strategies to specifically interfere with cRAR α expression. We targeted the putative promoter for *Rara3* located in the intron that separates the unique *Rara1* and -2 N-terminal sequences from the common C-portion. All mutant mice however still expressed cRAR α encoded by *Rara3* transcripts initiated from the *Rara2* promoter. Next, we altered the internal translational start codon of cRAR α to a neutral amino acid codon. The alternative codon remained active when only one nucleotide was replaced, whereas the exchange of two nucleotides led to expression of cRAR α starting from an alternative CUG start codon upstream of the internal methionine codon. Therefore, the results suggested that specific targeting of cRAR α was not feasible without also affecting expression or function of nRAR α and/or viability of the mutant mice.

STAR Method

RESOURCE AVAILABILITY

Lead contact—Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Hilde Cheroutre (hilde@lji.org).

Materials availability—Mice and embryonic stem cells used in this study are commercially available. Newly generated plasmids in this study are available from the lead contact with a completed material transfer agreement.

Data and code availability

- RNA-seq data have been deposited at GEO and are publicly available as of the date of publication. The accession number is listed in the key resources table.
- Original western blot images, raw flow cytometry data, (RT)-PCR, and microscopy data reported in this paper will be shared by the lead contact upon request.
- This paper does not report original code.
- Additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

In vivo Experimental Models

- All *in vivo* models used in this study were performed in mice.
- Age matched male and female mice were both analyzed separately and compared except where specifically indicated otherwise.
- Gender- and age-matched male and female animals were used between the ages of 2 months to 4 months.
- Mice were maintained and bred under specific pathogen-free conditions in the animal facility of the La Jolla Institute for Immunology. Animal care and experimentation were consistent with the guidelines of the US National Institutes of Health and were approved by the Institutional Animal Care and Use Committee of the La Jolla Institute for Immunology. All experiments were performed in accordance with the approved protocols.

In vivo T cell activation model.

- RARE-Luc transgenic mice were crossed to OVA-responsive OT-I TCR transgenic mice to generate OT-I/RARE-Luc double transgenic mice.
- In one case age-matched male and female mice were separately infected intravenous with ActA⁻ Lm-OVA and analyzed until 6 days post-infection for luciferase expression in harvested spleen cells.

- In another case, naïve CD45.2⁺ WT and CD45.1⁺ CRABP2-deficient OT-I cells isolated from CD45.2⁺ OT-I WT or from CD45.1⁺ CRABP2-deficient OT-I male or female age-matched donor mice were adoptively transferred to gender- and age-matched normal or vitamin A-free CD45.1⁺CD45.2⁺ C57BL/6 recipient mice that were infected intravenously with ActA⁻ Lm-OVA.

Donor and endogenous spleen and intestinal T cells isolated from the recipient mice were analyzed 7 days later. In some cases, vitamin A-free recipient mice were treated with DMSO or RA.

Experimental Autoimmune Encephalomyelitis (EAE) experiment

- Age-matched female C57BL/6 control mice or conditional *Crabp2* deletion mice between the ages of 2 months to 4 months, were analyzed per condition and per experiment. Littermates of 2 to 5 litters were randomly assigned to experimental groups.

In vitro Experimental Models

Primary cells and cell lines

- T cell hybridoma/cell lines
Jurkat (cl. E6–1, purchased from ATCC) and MCC-T CD4 T cell hybridoma cells were cultured in RPMI complemented with 10% FBS, penicillin, streptomycin and sodium pyruvate. Only low passage cells (<20) were used.
- Primary cells
Naïve sorted primary CD8αβ⁺ or CD4⁺ spleen T cells or total or sorted thymocytes or spleen dendritic cells isolated from male or female age-matched mice were harvested and used or analyzed in this study.

In vitro culture and activation of T cells

- For activation of naïve primary CD4⁺ or CD8αβ⁺ T cells isolated from the spleen of either female or male age-matched donor mice, 10⁵ cells were cultured in 96-well plates coated with anti-CD3ε (1 µg/ml, cl. 145–2C11, #16–0031–86, eBioscience) and soluble anti-CD28 (0.5 µg/ml, cl. 37.51, #16–0281–85, eBioscience).
- For Treg cell differentiation, naïve primary CD4⁺ spleen T cells isolated from age-matched male or female donor mice were activated with anti-mouse CD3ε and anti-mouse CD28 in the presence of human recombinant TGF-β 5 ng/ml and all *trans* retinoic acid (10 nM).
- For the short-term activation followed by the analysis of proximal signaling events, up to 10⁶ naïve primary CD4⁺ or CD8αβ⁺ T cells isolated from the spleen of either female or male age-matched donor mice were kept on ice for 10min in serum-free HBSS, then loaded with anti-CD3ε (10 µg/ml) and anti-CD28 (10 µg/ml) on ice for another 10–15min. Anti-hamster crosslinker (18 µg/ml) was added and the cells were immediately placed in a water-bath (37°C).

- For stimulation of pre-activated T cells, 10^5 naïve WT or *Crabp2* deficient spleen T cells isolated from male or female age-matched donor mice were cultured in 96-well plates coated with anti-CD3e (1 μ g/ml) and soluble anti-CD28 (0.5 μ g/ml) for 72h. Live activated T cells were sorted according to their FSC/SSC parameters. Equal numbers of pre-activated T cells were further activated *in vitro* according to the short-term activation protocol described hereinbefore.

METHOD DETAILS

Cell lines—Platinum-E (PLAT-E) (Cell Biolabs, San Diego, CA, USA) cells were cultured in DMEM supplemented with 10% FBS, penicillin, streptomycin, puromycin 1 μ g/ml and Balsticidin 10 μ g/ml. Jurkat (cl. E6–1, purchased from ATCC) and MCC-T CD4 T cell hybridoma cells(31) were cultured in RPMI complemented with 10% FBS, penicillin, streptomycin and sodium pyruvate. Only low passage cells (<20) were used.

Mice—C57BL/6 mice (WT) were bred in house. *Crabp2* deletion mutant mice(57) were a gift from V. Giguere (McGill University, Canada). *Crabp2* deletion mutant mice were backcrossed to the C57BL/6 genetic background for at least 10 generations. RARE-Luciferase (DR5) transgenic mice(39) and DNRAR α mice(28) were a gift from R. Noelle (Dartmouth-Hitchcock Medical College). The DNRAR α mice, that express a dn*Rara* transgene preceded by a floxed stop, were crossed with CD4cre(58) or E8Icre (also known as CD8cre) (59) transgenic mice in order to obtain littermates expressing or not the cre recombinase in mature CD4 and/or CD8 T cells. RARE-Luc transgenic mice were crossed to OVA-responsive OT-I TCR transgenic mice to generate OT-I/RARE-Luc double transgenic mice. S. Schoenberger provided the Act-mOVA mice(60). *Rara1* deletion mutant mice(61) were purchased from the Jackson laboratory. *Crabp2* -flox mice were generated with mutant ES cell clones with LoxP sites flanking *Crabp2* exons, obtained from EUCOMM. *Crabp2*-flox mice were crossed with distal Lck *Cre* transgenic mice to generate mature T cell specific *Crabp2* deletion mutant mice. Gender-matched and age-matched (from 2 to 4month-old) animals were used. Mice were maintained and bred under specific pathogen-free conditions in the animal facility of the La Jolla Institute for Immunology. Animal care and experimentation were consistent with the guidelines of the US National Institutes of Health and were approved by the Institutional Animal Care and Use Committee of the La Jolla Institute for Immunology.

T cell isolation, cell sorting and Cell Trace labeling—Splenic CD8 T cells or CD4 T cells were purified by magnetic negative selection with the MACS CD8 α^+ T cell isolation kit or naïve CD4 T cell isolation kit according to the manufacturer's protocol (#130–104-453, Miltenyi Biotec). Naïve T cells were sorted with a FACSAria (Becton Dickinson). In some cases, sorted cells were labeled with Cell Trace reagent (Invitrogen) according to the manufacturer's protocol. The division index, which is the average number of cell divisions that a cell in the original population has undergone, was calculated using the FlowJo software (Tristar). LIVE/DEAD™ Fixable Yellow Dead Cell Stain Kit, for 405 nm excitation, Thermo Fisher, L34968.

DC isolation—Spleens from naïve mice were perfused with 1 mg/ml collagenase D in the presence of DNase I. After perfusion the spleens were cut into small fragments and then digested with frequent mixing for 25min at room temperature. To disrupt DC-T cell complexes, EDTA (0.1 M) was added, and mixing continued for 5min. Cells were enriched using CD11c-beads (Miltenyi).

In vitro stimulation of T cells—For activation of T cells, 10^5 naïve CD4 or CD8 T cells were cultured in 96-well plates coated with anti-CD3 ϵ (1 μ g/ml, cl. 145–2C11, #16–0031-86, eBioscience) in the presence of soluble anti-CD28 (0.5 μ g/ml, cl. 37.51, #16–0281-85, eBioscience). For Treg cell differentiation, naïve CD4⁺ (APC Rat anti-mouse, #553051, BD) T cells were isolated with naïve CD4⁺ T Cell Isolation Kit, mouse, #130–104-453, Miltenyi Biotec) were activated with anti-mouse CD3 ϵ and anti-mouse CD28 in the presence of human recombinant TGF- β 5 ng/ml, (#240-B-002, R&D) and all *trans* retinoic acid (RA) (10 nM, #R2625, Sigma).

For the short-term activation followed by the analysis of the signaling events, up to 10^6 cells were rested on ice for 10min in serum-free HBSS, then loaded with anti-CD3 ϵ (10 μ g/ml) and anti-CD28 (10 μ g/ml) on ice for 10–15min. Anti-hamster crosslinker (18 μ g/ml, Jackson Immunoresearch) was added and the cells were immediately placed in a water-bath (37°C).

For stimulation of pre-activated T cells, 10^5 naïve WT or *Crabp2* deletion T cells were cultured in 96-well plates coated with anti-CD3 ϵ (1 μ g/ml) in the presence of soluble anti-CD28 (0.5 μ g/ml) for 72h. Live activated T cells were then sorted according to their FSC/SSC parameters. Equal numbers of T cells were then activated according to the short-term activation protocol described hereinbefore.

The RAR antagonist LE540 (iRARs, 5 μ M, Waco chemicals), the RAR α -selective antagonist Ro 41–5253 (iRAR α , 5 μ M, Enzo lifescience), the RAR γ -selective antagonist MM11253 (iRAR γ , 5 μ M, Tocris bioscience), MEK1/2 inhibitor U0126 (iERK, 10 μ M, InvivoGen), the PI3K inhibitor LY294002 (iPI3K, 20 μ M, InvivoGen), the γ -secretase inhibitor GSI (Tocris bioscience), RA (10 nM, Sigma) or rhIL-2 (40 U/ml, Peprotech), AKT-inhibitor (AKT Inhibitor VIII, Cayman Chemical Company, Inc, # 14870, at 0.3 μ M), were used in some cultures. When appropriate, DMSO (0.05 to 0.1%) was used as a control.

RNA-sequencing—After total cellular RNA extraction, 10^6 purified DP thymocytes or spleen T cells were collected to prepare total RNA using TRIzol reagent (Invitrogen). 200–300 ng total RNA was used for library construction with a SureSelect Strand Specific RNA-Seq Library Preparation kit (Agilent Technologies) according to the manufacturer's protocol. Sequencing was performed by the genomics facility at RIKEN IMS with an Illumina HiSeq 1500. Sequences retrieved in RNA-seq experiments were aligned on the UCSC mouse genome assembly (GRCm38/mm10) using the HISAT2 (hierarchical indexing for spliced alignment of transcripts 2). And the obtained BAM files were then sorted by Samtools and visualized by Integrative Genome viewer (IGV). Raw fastq files for the RNA-seq libraries were deposited at the Gene Expression Omnibus (GEO) database under the accession number GEO: GSE112609.

RNA-seq data of human naïve CD8 (CD3⁺ CD8⁺ CD45RA⁺ CD127⁺ CCR7⁺) or CD4 (CD3⁺ CD4⁺ CD45RA⁺ CD127⁺ CCR7⁺) T cells sorted from peripheral blood mononuclear cells (PBMC) were obtained through the Database of Immune Cell Expression, Expression quantitative trait loci (eQTLs) and Epigenomics (DICE); <https://dice-database.org/>.

RT-PCR—Total RNA was obtained as described above. cDNA was synthesized using SuperScript IV (Thermo Fisher Scientific) and an oligo dT primer, then each *Rara* or control transcript was measured with combinations of the following primers.

F1: CAGTCAGTGGTTACAGCACACCGTC

F2: GGACTCCGCTTTGGAATGGCTC

F3: GATGTCCAGGCCCAAGTAGAAGCCAG

R1: GCTGCAATCTGCTGCTCATGCCTACAC

R2: GGACTGACCTGCTGCAATCTGCTG

R3: CTGGCTTCTACTTGGGCCTGGACATC

R4: CAGGCTGTGAAAGACTCCTGCGGCT

R5: CAGCATGTGTTATGCCAGGCTGTGAAAG

R6: CCAACAGCATGTGTTATGCCAGGCTG

R7: TTCACCTCACTTCCTTCTCGGGAG

R9: CCCCATAGTGGTAGCCGGATGATTTG

R15: TCAGTGGA AACCCAGCAGGAACAGGTG

Hprt F: GTCGTGATTAGCGATGATGAACC

Hprt R: ATGACATCTCGAGCAAGTCTTTCAG

Quantitative RT-PCR—RNA was extracted with TRIzol (Invitrogen) and cDNA was synthesized with the iScript cDNA Synthesis kit (Bio-Rad). A 480 Real-Time PCR System (Roche) was used for real-time PCR. Values were normalized to the amount of *Gapdh*, *β-actin* or *Hprt* in each sample and multiplied by 10,000. The primers for quantitative PCR were as follows:

Crabp2 F: CCCAGAGCTTTTAGCATTTC

Crabp2 R: GAAGATCTAAAGAGAAAGCCACCT

c-Myc F: TTGAAGGCTGGATTTCTTTGGGC

c-Myc R: TCGTCGCAGATGAAATAGGGCTGT

Gapdh F: CAGATGCCTGCTTCACCA

Gapdh R: ATGGCCTTCCGTGTTTCCT

β-actin F: CCAGAAGGACTGTTATGTGGGA

β-actin R: GACTCCGTGTTCAATGGGATAC

Hprt F: GTCGTGATTAGCGATGATGAACC

Hprt R: ATGACATCTCGAGCAAGTCTTTTCAG

Immunofluorescence staining and flow cytometry—Cells were pre-incubated with anti-CD16-CD32 (2.4G2 mAb prepared in-house) to block binding of antibodies to the Fc receptor, and then stained in cold PBS containing 0.5% (vol/vol) FBS and 0.05% (wt/vol) sodium azide with the relevant labeled antibodies and tetramers. The following antibodies were used: anti-CD4 (cl. RMA4–5), anti-TCR $\gamma\delta$ (cl. GL3), anti CD45.1 (cl. A20), anti-CD44 (cl. IM7), anti-CD8 α (cl. 53–6.7) all from BD; anti-IFN γ (cl. XMG1.2), anti-IL-17A (cl. TC11–18H10.1), anti-CD8 α (cl. 53–6.7), anti-CD4 (cl. GK1.5) all from Biolegend; anti-FOXP3 (cl. FJK-16s), anti-CCR9 (cl. eBioCW-1.2), anti-CD45.2 (cl. 104), anti-ROR γ t (cl. AFKJS-9), anti-CD45 (cl. 30-F11), anti-CD25 (cl. PC61.5), anti-CD4 (cl. RMA4–5), anti-c-MYC (cl. 9E10) all from Thermo Fisher Scientific. A Fix/Perm kit was used according to the manufacturer's directions (eBioscience) for intracellular staining of c-MYC or FOXP3, after surface staining. For detection of cell death, cells were stained with the Live/Dead fixable dead cell stain kit according to the manufacturer's protocol (Invitrogen) and cells were analyzed immediately after staining. All stained cells were processed on an LSR-II (Becton Dickinson) and the data analyzed using FlowJo software (TreeStar). Dead cells were excluded from the analysis.

Western-blot analysis—Cells were lysed in cold triton lysis buffer (137 mM NaCl, 20 mM Tris base at pH 7.4, 10% glycerol and 1% Triton X-100) supplemented with a protease and phosphatase inhibitor mixture (Roche) for 20min on ice, and centrifuged at 15,000 rpm for 15min at 4°C. In some experiments, nuclear/cytoplasmic fractionation was performed according to the manufacturer's protocol (Pierce). Equal amounts of denatured proteins were loaded onto an SDS-PAGE gel (Biorad) and transferred onto a PVDF membrane (Invitrogen). The membranes were then incubated with antibodies to ZAP-70 (cl. D1C10E), GAPDH (cl. 14C10), Lamin B1 (cl. D9V6H), NOTCH1 (cl. D1E11), RAR α (#2554), RAR γ 1 (cl. D3A4) (all from Cell Signaling Technology), CRABP2 (#11250, One World Lab), RAR α (R12–3424, Assay Biotechnology Company), HSP90 (cl. AC16, Santa Cruz), FLAG (cl. M2, Sigma Aldrich), AKT (#9272) or with antibodies to the phosphorylated forms of AKT (Ser473) (#9271), PLC γ 1 (#2821), ZAP-70 (cl. 65E4) or ERK1/2 (cl. D13.14.4E) (all from Cell Signaling Technology). Immuno-reactive bands were detected by chemiluminescence (ECL solution, Santa Cruz).

Immunoprecipitation—OT-I CD8 T cells were activated with H-2K^b-SIINFEKL tetramers in order to avoid non-specific binding of anti-CD3 stimulating Abs to the beads used for immunoprecipitation. Cells were lysed in cold triton lysis buffer (137 mM NaCl, 20

mM Tris base at pH 7.4, 10% glycerol and 1% Triton X-100) supplemented with protease and phosphatase inhibitor mixture (Roche) for 20min on ice and centrifuged at 15,000 rpm for 15min at 4°C. Cell lysates were pre-cleared with A/G beads (Pierce) prior to immunoprecipitation. Equal amounts (at least three hundred micrograms) of protein were immunoprecipitated with 2 µg of antibodies to RAR α (#sc-551, Santa Cruz) or with a rabbit IgG overnight at 4°C, followed by capture with 25 µl of protein A/G beads (Pierce). Denatured proteins were then loaded onto an SDS-PAGE gel (Biorad) and transferred onto a PVDF membrane (Invitrogen). Immunoblotting was performed with antibodies to ZAP-70 (cl. D1C10E) (Cell Signaling Technology). Immuno-reactive bands were detected by chemiluminescence (ECL solution, Santa Cruz).

Retroviral plasmids—Sequences of interest were inserted into an MSCV-IRES-GFP or into an MSCV-IRES-Puromycin (for GFP-tagged proteins) plasmid (Addgene, #20672 and #68469).

The sequences used for overexpression code for these protein sequences:

RAR α 1-FLAG:

MASNSSCPTPGGGHLNGYPVPPYAFFFPMLGGLSPPGALTSLQHQLPVSGYSTPSP
 ATIETQSSSSEEIVSPSPPLPRIYKPCFVCQDKSSGYHYGVSACEGCKGFFRRSIQK
 NMVYTCHRDKNCIINKVTRNRCQYCRLQKCFDVGMSKESVRNDRNKKKKEAPKPE
 CSESYTLTPEVGELIEKVRKAHQETFPALCQLGKYTTNNSSEQRVSLDIDLWDFSEL
 STKCIKTVEFAKQLPGFTTLTIADQITLLKAACLDILILRICTRYTPEQDTMTFSDGLT
 LNRTQMHNAGFGPLTDLVFAFANQLLPLEMDDAETGLLSAICLICGDRQDLEQPKV
 DMLQEPLLEALKVYVRKRRPSRPHMFPKMLMKITDLRSISAKGAERVITLKMEIPGS
 MPPLIQEMLENSEGLDITLSGQSGGGTRDGGGLAPPPGSCSPSLSPSSHRSSPATQSPD
YKDDDDK

RAR α 2-FLAG:

MYESVEVGGLTPAPNPFLVDFYNQNRACLLQEKGLPAPGPYSTPLRTPLWNGSNHS
 IETQSSSSEEIVSPSPPLPRIYKPCFVCQDKSSGYHYGVSACEGCKGFFRRSIQKN
 MVYTCHRDKNCIINKVTRNRCQYCRLQKCFDVGMSKESVRNDRNKKKKEAPKPEC
 SESYTLTPEVGELIEKVRKAHQETFPALCQLGKYTTNNSSEQRVSLDIDLWDFSELS
 TKCIKTVEFAKQLPGFTTLTIADQITLLKAACLDILILRICTRYTPEQDTMTFSDGLTL
 NRTQMHNAGFGPLTDLVFAFANQLLPLEMDDAETGLLSAICLICGDRQDLEQPKVD
 MLQEPLLEALKVYVRKRRPSRPHMFPKMLMKITDLRSISAKGAERVITLKMEIPGSM
 PPLIQEMLENSEGLDITLSGQSGGGTRDGGGLAPPPGSCSPSLSPSSHRSSPATQSPDYK
DDDDK

RAR α 3:

MVYTCHRDKNCIINKVTRNRCQYCRLQKCFDVGMSKESVRNDRNKKKKEAPKPEC
 SESYTLTPEVGELIEKVRKAHQETFPALCQLGKYTTNNSSEQRVSLDIDLWDFSELS
 TKCIKTVEFAKQLPGFTTLTIADQITLLKAACLDILILRICTRYTPEQDTMTFSDGLTL
 NRTQMHNAGFGPLTDLVFAFANQLLPLEMDDAETGLLSAICLICGDRQDLEQPKVD

MLQEPLLEALKVYVRKRRPSRPHMFPKMLMKITDLRSISAKGAERVITLKMEIPGSM
PPLIQEMLENSEGLDLSGQSGGGTRDGGGLAPPPGSCSPSLSPSSHRSSPATQSP

RAR α 1-eGFP:

MASNSSSCPTPGGGHLNGYPVPPYAFFFPMLGGLSPPGALTSLQHQLPVSGYSTPSP
ATIQSSSSEEIVSPSPPLPRIYKPCFVCQDKSSGYHYGVSACEGCKGFFRRSIQK
NMVYTCHRDKNCIINKVTRNRCQYCRLLKCFDVGMSKESVRNDRNKKKKEAPKPE
CSESYTLTPEVGELIEKVRKAHQETFPALCQLGKYTTNNSSEQRVSLDIDLWDKFSEL
STKCIKTVEFAKQLPGFTTLTIADQITLLKAACLDILILRICTRYTPEQDTMTFSDGLT
LNRTQMHNAGFGPLTDLVFAFANQLLPLEMDDAETGLLSAICLICGDRQDLEQPKV
DMLQEPLLEALKVYVRKRRPSRPHMFPKMLMKITDLRSISAKGAERVITLKMEIPGS
MPPLIQEMLENSEGLDLSGQSGGGTRDGGGLAPPPGSCSPSLSPSSHRSSPATQSPG
ASLEMVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGK
LPVPWPTLVTTLYGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKT
RAEVKFEQDGLVNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMADKQKNGIKVN
FKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSALS KDPNEKRDMVL
LEFVTAAGITLGMDELYK

RAR α 3-eGFP:

MVYTCHRDKNCIINKVTRNRCQYCRLLKCFDVGMSKESVRNDRNKKKKEAPKPEC
SESYTLTPEVGELIEKVRKAHQETFPALCQLGKYTTNNSSEQRVSLDIDLWDKFSELS
TKCIKTVEFAKQLPGFTTLTIADQITLLKAACLDILILRICTRYTPEQDTMTFSDGLTL
NRTQMHNAGFGPLTDLVFAFANQLLPLEMDDAETGLLSAICLICGDRQDLEQPKVD
MLQEPLLEALKVYVRKRRPSRPHMFPKMLMKITDLRSISAKGAERVITLKMEIPGSM
PPLIQEMLENSEGLDLSGQSGGGTRDGGGLAPPPGSCSPSLSPSSHRSSPATQSPGAS
LEMVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGKLP
VPWPTLVTTLYGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIFFKDDGNYKTRA
EVKFEQDGLVNRIELKGIDFKEDGNILGHKLEYNYNSHNVYIMADKQKNGIKVNFKI
RHNIEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSALS KDPNEKRDMVLL
VTAAGITLGMDELYK

Retroviral transduction—Retroviral plasmids were transfected into the Platinum-E Retroviral Packaging cell line (Cell Biolabs) using the JetPrime transfection reagent (Polyplus Transfection) according to the manufacturer's instructions. Retrovirus-containing supernatants were collected 48hrs or 72hrs after transfection and filtered through a 0.45 μ m filter. For infection, cells were spin-infected with retroviral supernatants containing 5 μ g/ml polybrene (Sigma Aldrich) for 2h at room temperature at 2000 rpm. Supernatants were gently removed 4–5hrs later and replaced with adapted culture media.

Microscopy analysis—Purified DCs were activated with 1 μ g/ml poly(I:C) (Invivogen) for 4hrs at 37°C and then pulsed with 0.25 μ g/ μ l OVA257–264 peptide (AnaSpec Inc.) for 1hr at 37°C. Activated DCs and naïve OT-I cells were co-cultured for 40min at a 1:1 ratio at 37°C. During the last 10min, cells were transferred onto poly-L-lysine-coated microscopy slides. DC-T conjugates were fixed with 4% paraformaldehyde (PFA), quenched (50

mM NH₄Cl), permeabilized (0.3% Triton-X-100) and blocked (0.01% saponin /0.25% fish gelatin /anti-CD16/32). Cells were then stained with biotinylated anti-CD3 ϵ (Biolegend), anti-RAR α (cl. Ralpha 10, Thermo Fisher) and anti-ZAP-70 (cl. M20, Santa Cruz) at 4°C overnight. Staining with secondary antibodies and streptavidin, listed in the key resources table, was carried out at room temperature for 1hr. Images were recorded using a FV1000 laser scanning confocal microscope (Olympus). Images of at least 100 synapses were recorded and the number of CD3 only, CD3-ZAP-70 and CD3-ZAP-70-RAR α synapses was quantified. CD3-RAR α synapses were not observed.

T cells transduced with GFP-tagged RAR α 1 or RAR α 3 were fixed (PBS1X PFA 2%). Staining with DAPI was then performed for 1hr at 4°C. Samples were imaged on an Olympus Fluoview FV10i confocal microscope with a 60x oil immersion objective and acquired using FV10i SW. Cytosolic/nuclear localization was based on DAPI staining and quantified using Imaris software. CD8 T cells transduced with CD3 ζ -GFP were activated with anti-CD3/CD28 for 30min. Clustering of CD3 ζ -GFP was quantified following microscopic (Olympus) imaging of entire cells.

Mass spectrometry-based phosphoproteomic analysis—Jurkat E6.1 cells (ATCC) were passaged in RPMI + 10% heat inactivated FBS and kept at cell densities between 2e5 and 1e6 cell per mL. Jurkat cells were activated using plate bound anti-CD3 and anti-CD28 antibodies at 3.33 μ g/mL in 96 well plates. At select time points cells were transferred to cold PBS to stop signaling, washed twice with cold PBS and flash frozen. The zero timepoint was performed in biological duplicate, the others in triplicate. Cell pellets were prepared and were then analyzed for quantitative phosphoproteomics as previously described in (62, 63). Iron immobilized metal affinity chromatography (IMAC) was used to enrich for phosphopeptides, and tandem mass tags (TMT) 11-plex reagents (Thermo Scientific) were used for quantitation. Data was processed by Spectrum Mill (Agilent/Broad Institute). Mass spectra were manually annotated.

Bacterial infection—Acta deficient *Listeria monocytogenes* expressing OVA (ActA⁻ Lm-OVA)(64) was prepared from cultures in brain-heart–infusion broth. Bacteria were washed and resuspended in 1X PBS before intravenous infection (2.5×10^5 CFU per mouse).

Diet Studies—At day 14.5 of gestation, pregnant females were fed with either vitamin A-deficient or -sufficient diet (Dyets Inc.) and maintained on the diet until weaning of the litters. Weanlings were maintained on either diet until the experiment. In some experiments, mice were administered DMSO or 250 μ g all trans retinoic acid intraperitoneally every other day for 7 days prior to immunization and until euthanasia of the mice.

Preparation of intraepithelial T cells—Small intestines were removed and separated from Peyer’s patches and cut longitudinally and into pieces of 0.5 cm in length. Pieces were shaken for 30min in magnesium-free, calcium-free Hank’s balanced-salt solution supplemented with 1 mM dithiothreitol (SIGMA Aldrich), 5mM EDTA (SIGMA-Aldrich) and 5% (vol/vol) FCS. Cells were collected from the washes and were passed over a discontinuous 40–70% (vol/vol) gradient of Percoll (Pharmacia Biotech) for 20min at

1100g. Intraepithelial T cells were then isolated from the Percoll gradient interface and washed free of Percoll.

Experimental Autoimmune Encephalomyelitis (EAE) experiment—MOG_{35–55} peptide (200 µg/mouse, Anaspec) was mixed with an equal volume of incomplete Freund's adjuvant (#263910, 200µl/mouse, BD) and M.Tuberculosis (#231141, 400µg/mouse, BD), then, subcutaneously injected to each mouse. Diluted pertussis toxin from *B. pertussis* (#180, 400 ηg/mouse, List Biological Laboratories Inc) was intraperitoneally injected at day 0 and day 2. Clinical score was evaluated on a daily basis from day 7(65). All experiments were performed in accordance with the approved protocol by the institutional animal care committee.

Histology—Spines were cut open and spinal cords were extracted and fixed with 4% zinc formalin and dehydrated with 70%, 80%, 90% and 100% isopropanol and Pro-Par Clearant (Anatech) by Tissue-Tek VIP (SAKURA), then, embedded in Paraffin and sectioned. Tissue sections were stained with Hematoxylin and Eosin, then, scanned by Axio Scan Z1 (ZEISS) and analyzed by ZEN software (ZEISS).

Other Software—Graphic abstract created with [BioRender.com](https://www.biorender.com).

QUANTIFICATIONS AND STATISTICAL ANALYSIS

Quantifications

- Quantitative phosphoproteomics analysis was performed as previously described in(62, 63). Iron immobilized metal affinity chromatography (IMAC) was used to enrich for phosphopeptides, and tandem mass tags (TMT) 11-plex reagents (Thermo Scientific) were used for quantitation. Data was processed by Spectrum Mill (Agilent/Broad Institute). Mass spectra were manually annotated.
- Signaling intensity was measured by quantifying the optical density of immuno-reactive bands of proteins analyzed by western blot, detected by chemiluminescence (ECL solution, Santa Cruz). ImageJ software from NIH image (Version 1.51 23 April 2018) was used to quantify the intensity of each band in the negative picture after performing a gray scale 8-bit transformation on the scans and calibrating the threshold.

Statistics

- Statistical comparisons were performed using a two-tailed unpaired Student t-test, Mann-Whitney test or, when appropriate, a two-tailed paired Student t-test, with the Prism software (Graphpad).
- When applicable, variances were calculated and were similar between compared groups.
- For *in vivo* experiments, animals were gender- and age-matched and divided in groups where n= >3. The specific number of animals used per group and per experiment is indicated in the figure and/or figure legend.

- No statistical method was used to predetermine the number of animal sample size.
- No sample or animal was excluded from analysis.
- No randomization method was used to determine how samples/animals were allocated to experimental groups and processed.
- For *in vitro* experiments, the minimal sample size chosen was n=3 independent repeats, each independent experiment was performed in triplicates (3 technical replicates), except for signaling studies where technical replicates are not feasible.

ADDITIONAL RESOURCES

- Sequences retrieved in RNA-seq experiments were aligned on the UCSC mouse genome assembly (GRCm38/mm10) using the HISAT2 (hierarchical indexing for spliced alignment of transcripts 2). And the obtained BAM files were then sorted by Samtools and visualized by Integrative Genome viewer (IGV). Raw fastq files for the RNA-seq libraries were deposited at the Gene Expression Omnibus (GEO) database under the accession number GEO: GSE112609.
- Analysis of high-throughput sequencing identified other *Rara* sequences with different 5' splice junctions. The transcripts were then aligned with predicted sequences in the RefSeq database (www.ncbi.nlm.nih.gov/refseq).
- All alternative transcript shared the same 3' end sequence with *Rara1* and -2 but each one had a unique 5' end sequence. To identify a putative *Rara3* promoter region and transcriptional start site (TSS) data from the Functional ANnotation Of the Mammalian genome (FANTOM)5 database was used, which indicated a TSS downstream of the *Rara2* TSS and upstream of the 5' end of the variant transcripts.
- RNA-seq data of human naïve CD8 $\alpha\beta$ ⁺ (CD3⁺ CD8⁺ CD45RA⁺ CD127⁺ CCR7⁺) or CD4⁺ (CD3⁺ CD4⁺ CD45RA⁺ CD127⁺ CCR7⁺) peripheral blood T cells were from the Database of Immune Cell Expression, Expression quantitative trait loci (eQTLs) and Epigenomics (DICE); <https://dice-database.org/>.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We thank Y. Wang-Zhu and Gabriela Aguila for help with breeding and typing of the mice; R. Noelle for providing the RAR α DN mice; C. Kim, D. Hinz, L. Nosworthy, K. Van Gunst and A. Jose for cell sorting. Z. Mikulski for help with microscopy. Sara McArdle and Kenneth Kim for help with the histology analysis and scoring. This work was supported by the US National Institutes of Health (U01 AI125957 and R01 AI106298 (HC)). Support from the Philippe Foundation (AL), CCFA Fellowship 697275 (IT) and MK by the P01 DK46763.

This is manuscript 1634 from the La Jolla Institute for Immunology.

REFERENCES

1. Larange A, Cheroutre H. Retinoic Acid and Retinoic Acid Receptors as Pleiotropic Modulators of the Immune System. *Annu Rev Immunol.* 2016;34:369–94. [PubMed: 27168242]
2. di Masi A, Leboffe L, De Marinis E, Pagano F, Cicconi L, Rochette-Egly C, et al. Retinoic acid receptors: from molecular mechanisms to cancer therapy. *Mol Aspects Med.* 2015;41:1–115. [PubMed: 25543955]
3. Hall JA, Cannons JL, Grainger JR, Dos Santos LM, Hand TW, Naik S, et al. Essential role for retinoic acid in the promotion of CD4(+) T cell effector responses via retinoic acid receptor alpha. *Immunity.* 2011;34(3):435–47. [PubMed: 21419664]
4. Hall JA, Grainger JR, Spencer SP, Belkaid Y. The role of retinoic acid in tolerance and immunity. *Immunity.* 2011;35(1):13–22. [PubMed: 2177796]
5. Farboud B, Privalsky ML. Retinoic acid receptor-alpha is stabilized in a repressive state by its C-terminal, isotype-specific F domain. *Mol Endocrinol.* 2004;18(12):2839–53. [PubMed: 15331758]
6. Dong D, Ruuska SE, Levinthal DJ, Noy N. Distinct roles for cellular retinoic acid-binding proteins I and II in regulating signaling by retinoic acid. *J Biol Chem.* 1999;274(34):23695–8. [PubMed: 10446126]
7. Schug TT, Berry DC, Shaw NS, Travis SN, Noy N. Opposing effects of retinoic acid on cell growth result from alternate activation of two different nuclear receptors. *Cell.* 2007;129(4):723–33. [PubMed: 17512406]
8. Iwata M, Hirakiyama A, Eshima Y, Kagechika H, Kato C, Song SY. Retinoic acid imprints gut-homing specificity on T cells. *Immunity.* 2004;21(4):527–38. [PubMed: 15485630]
9. Ohoka Y, Yokota A, Takeuchi H, Maeda N, Iwata M. Retinoic acid-induced CCR9 expression requires transient TCR stimulation and cooperativity between NFATc2 and the retinoic acid receptor/retinoid X receptor complex. *J Immunol.* 2011;186(2):733–44. [PubMed: 21148038]
10. Pino-Lagos K, Guo Y, Brown C, Alexander MP, Elgueta R, Bennett KA, et al. A retinoic acid-dependent checkpoint in the development of CD4+ T cell-mediated immunity. *J Exp Med.* 2011;208(9):1767–75. [PubMed: 21859847]
11. Benson MJ, Pino-Lagos K, Rosemblatt M, Noelle RJ. All-trans retinoic acid mediates enhanced T reg cell growth, differentiation, and gut homing in the face of high levels of co-stimulation. *J Exp Med.* 2007;204(8):1765–74. [PubMed: 17620363]
12. Coombes JL, Siddiqui KR, Arancibia-Carcamo CV, Hall J, Sun CM, Belkaid Y, et al. A functionally specialized population of mucosal CD103+ DCs induces Foxp3+ regulatory T cells via a TGF-beta and retinoic acid-dependent mechanism. *J Exp Med.* 2007;204(8):1757–64. [PubMed: 17620361]
13. Mucida D, Park Y, Kim G, Turovskaya O, Scott I, Kronenberg M, et al. Reciprocal TH17 and regulatory T cell differentiation mediated by retinoic acid. *Science.* 2007;317(5835):256–60. [PubMed: 17569825]
14. Mucida D, Cheroutre H. TGFbeta and retinoic acid intersect in immune-regulation. *Cell Adh Migr.* 2007;1(3):142–4. [PubMed: 19262136]
15. Schambach F, Schupp M, Lazar MA, Reiner SL. Activation of retinoic acid receptor-alpha favours regulatory T cell induction at the expense of IL-17-secreting T helper cell differentiation. *Eur J Immunol.* 2007;37(9):2396–9. [PubMed: 17694576]
16. Sun CM, Hall JA, Blank RB, Bouladoux N, Oukka M, Mora JR, et al. Small intestine lamina propria dendritic cells promote de novo generation of Foxp3 T reg cells via retinoic acid. *J Exp Med.* 2007;204(8):1775–85. [PubMed: 17620362]
17. Hill JA, Hall JA, Sun CM, Cai Q, Ghyselinck N, Chambon P, et al. Retinoic acid enhances Foxp3 induction indirectly by relieving inhibition from CD4+CD44hi Cells. *Immunity.* 2008;29(5):758–70. [PubMed: 19006694]
18. Mucida D, Pino-Lagos K, Kim G, Nowak E, Benson MJ, Kronenberg M, et al. Retinoic acid can directly promote TGF-beta-mediated Foxp3(+) Treg cell conversion of naive T cells. *Immunity.* 2009;30(4):471–2; author reply 2–3. [PubMed: 19371709]

19. Laurence A, Tato CM, Davidson TS, Kanno Y, Chen Z, Yao Z, et al. Interleukin-2 signaling via STAT5 constrains T helper 17 cell generation. *Immunity*. 2007;26(3):371–81. [PubMed: 17363300]
20. Mathur AN, Chang HC, Zisoulis DG, Stritesky GL, Yu Q, O'Malley JT, et al. Stat3 and Stat4 direct development of IL-17-secreting Th cells. *J Immunol*. 2007;178(8):4901–7. [PubMed: 17404271]
21. Yang XO, Panopoulos AD, Nurieva R, Chang SH, Wang D, Watowich SS, et al. STAT3 regulates cytokine-mediated generation of inflammatory helper T cells. *J Biol Chem*. 2007;282(13):9358–63. [PubMed: 17277312]
22. Yao Z, Kanno Y, Kerenyi M, Stephens G, Durant L, Watford WT, et al. Nonredundant roles for Stat5a/b in directly regulating Foxp3. *Blood*. 2007;109(10):4368–75. [PubMed: 17227828]
23. Elias KM, Laurence A, Davidson TS, Stephens G, Kanno Y, Shevach EM, et al. Retinoic acid inhibits Th17 polarization and enhances FoxP3 expression through a Stat-3/Stat-5 independent signaling pathway. *Blood*. 2008;111(3):1013–20. [PubMed: 17951529]
24. Xu L, Kitani A, Strober W. Molecular mechanisms regulating TGF-beta-induced Foxp3 expression. *Mucosal Immunol*. 2010;3(3):230–8. [PubMed: 20404810]
25. Nolting J, Daniel C, Reuter S, Stuelten C, Li P, Sucov H, et al. Retinoic acid can enhance conversion of naive into regulatory T cells independently of secreted cytokines. *J Exp Med*. 2009;206(10):2131–9. [PubMed: 19737861]
26. Etemire E, Krull M, Hasenberg M, Reichardt P, Gunzer M. Transiently reduced PI3K/Akt activity drives the development of regulatory function in antigen-stimulated Naive T-cells. *PLoS One*. 2013;8(7):e68378. [PubMed: 23874604]
27. Friesen LR, Gu B, Kim CH. A ligand-independent fast function of RARalpha promotes exit from metabolic quiescence upon T cell activation and controls T cell differentiation. *Mucosal Immunol*. 2021;14(1):100–12. [PubMed: 32518366]
28. Rajaii F, Bitzer ZT, Xu Q, Sockanathan S. Expression of the dominant negative retinoid receptor, RAR403, alters telencephalic progenitor proliferation, survival, and cell fate specification. *Dev Biol*. 2008;316(2):371–82. [PubMed: 18329011]
29. Leroy P, Krust A, Zelent A, Mendelsohn C, Garnier JM, Kastner P, et al. Multiple isoforms of the mouse retinoic acid receptor alpha are generated by alternative splicing and differential induction by retinoic acid. *EMBO J*. 1991;10(1):59–69. [PubMed: 1846598]
30. Lizio M, Harshbarger J, Shimoji H, Severin J, Kasukawa T, Sahin S, et al. Gateways to the FANTOM5 promoter level mammalian expression atlas. *Genome Biol*. 2015;16:22. [PubMed: 25723102]
31. So T, Soroosh P, Eun SY, Altman A, Croft M. Antigen-independent signalosome of CARMA1, PKCtheta, and TNF receptor-associated factor 2 (TRAF2) determines NF-kappaB signaling in T cells. *Proc Natl Acad Sci U S A*. 2011;108(7):2903–8. [PubMed: 21282629]
32. Britton GJ, Ambler R, Clark DJ, Hill EV, Tunbridge HM, McNally KE, et al. PKCtheta links proximal T cell and Notch signaling through localized regulation of the actin cytoskeleton. *Elife*. 2017;6.
33. Dustin ML, Choudhuri K. Signaling and Polarized Communication Across the T Cell Immunological Synapse. *Annu Rev Cell Dev Biol*. 2016;32:303–25. [PubMed: 27501450]
34. Guy CS, Vignali KM, Temirov J, Bettini ML, Overacre AE, Smeltzer M, et al. Distinct TCR signaling pathways drive proliferation and cytokine production in T cells. *Nat Immunol*. 2013;14(3):262–70. [PubMed: 23377202]
35. Smith-Garvin JE, Koretzky GA, Jordan MS. T cell activation. *Annu Rev Immunol*. 2009;27:591–619. [PubMed: 19132916]
36. Palaga T, Miele L, Golde TE, Osborne BA. TCR-mediated Notch signaling regulates proliferation and IFN-gamma production in peripheral T cells. *J Immunol*. 2003;171(6):3019–24. [PubMed: 12960327]
37. Chebaro Y, Amal I, Rochel N, Rochette-Egly C, Stote RH, Dejaegere A. Phosphorylation of the retinoic acid receptor alpha induces a mechanical allosteric regulation and changes in internal dynamics. *PLoS Comput Biol*. 2013;9(4):e1003012. [PubMed: 23637584]

38. Lu KP, Liou YC, Zhou XZ. Pinning down proline-directed phosphorylation signaling. *Trends Cell Biol.* 2002;12(4):164–72. [PubMed: 11978535]
39. Svensson RU, Shey MR, Ballas ZK, Dorkin JR, Goldberg M, Akinc A, et al. Assessing siRNA pharmacodynamics in a luciferase-expressing mouse. *Mol Ther.* 2008;16(12):1995–2001. [PubMed: 18781145]
40. Won JY, Nam EC, Yoo SJ, Kwon HJ, Um SJ, Han HS, et al. The effect of cellular retinoic acid binding protein-I expression on the CYP26-mediated catabolism of all-trans retinoic acid and cell proliferation in head and neck squamous cell carcinoma. *Metabolism.* 2004;53(8):1007–12. [PubMed: 15281009]
41. Budhu AS, Noy N. Direct channeling of retinoic acid between cellular retinoic acid-binding protein II and retinoic acid receptor sensitizes mammary carcinoma cells to retinoic acid-induced growth arrest. *Mol Cell Biol.* 2002;22(8):2632–41. [PubMed: 11909957]
42. Sessler RJ, Noy N. A ligand-activated nuclear localization signal in cellular retinoic acid binding protein-II. *Mol Cell.* 2005;18(3):343–53. [PubMed: 15866176]
43. Brown CC, Esterhazy D, Sarde A, London M, Pullabhatla V, Osma-Garcia I, et al. Retinoic acid is essential for Th1 cell lineage stability and prevents transition to a Th17 cell program. *Immunity.* 2015;42(3):499–511. [PubMed: 25769610]
44. Billadeau DD, Nolz JC, Gomez TS. Regulation of T-cell activation by the cytoskeleton. *Nat Rev Immunol.* 2007;7(2):131–43. [PubMed: 17259969]
45. Yokosuka T, Sakata-Sogawa K, Kobayashi W, Hiroshima M, Hashimoto-Tane A, Tokunaga M, et al. Newly generated T cell receptor microclusters initiate and sustain T cell activation by recruitment of Zap70 and SLP-76. *Nat Immunol.* 2005;6(12):1253–62. [PubMed: 16273097]
46. Dongre A, Surampudi L, Lawlor RG, Fauq AH, Miele L, Golde TE, et al. Non-Canonical Notch Signaling Drives Activation and Differentiation of Peripheral CD4(+) T Cells. *Front Immunol.* 2014;5:54. [PubMed: 24611064]
47. Steinbuck MP, Arakcheeva K, Winandy S. Novel TCR-Mediated Mechanisms of Notch Activation and Signaling. *J Immunol.* 2018;200(3):997–1007. [PubMed: 29288204]
48. Rochette-Egly C, Adam S, Rossignol M, Egly JM, Chambon P. Stimulation of RAR alpha activation function AF-1 through binding to the general transcription factor TFIID and phosphorylation by CDK7. *Cell.* 1997;90(1):97–107. [PubMed: 9230306]
49. Unsworth AJ, Flora GD, Gibbins JM. Non-genomic effects of nuclear receptors: insights from the anucleate platelet. *Cardiovasc Res.* 2018;114(5):645–55. [PubMed: 29452349]
50. Rondina MT, Freitag M, Pluthero FG, Kahr WH, Rowley JW, Kraiss LW, et al. Non-genomic activities of retinoic acid receptor alpha control actin cytoskeletal events in human platelets. *J Thromb Haemost.* 2016;14(5):1082–94. [PubMed: 26848712]
51. Jarvis CI, Goncalves MB, Clarke E, Dogruel M, Kalindjian SB, Thomas SA, et al. Retinoic acid receptor-alpha signalling antagonizes both intracellular and extracellular amyloid-beta production and prevents neuronal cell death caused by amyloid-beta. *Eur J Neurosci.* 2010;32(8):1246–55. [PubMed: 20950278]
52. Piskunov A, Rochette-Egly C. A retinoic acid receptor RARalpha pool present in membrane lipid rafts forms complexes with G protein alphaQ to activate p38MAPK. *Oncogene.* 2012;31(28):3333–45. [PubMed: 22056876]
53. Masia S, Alvarez S, de Lera AR, Baretino D. Rapid, nongenomic actions of retinoic acid on phosphatidylinositol-3-kinase signaling pathway mediated by the retinoic acid receptor. *Mol Endocrinol.* 2007;21(10):2391–402. [PubMed: 17595318]
54. Haxhinasto S, Mathis D, Benoist C. The AKT-mTOR axis regulates de novo differentiation of CD4+Foxp3+ cells. *J Exp Med.* 2008;205(3):565–74. [PubMed: 18283119]
55. Merkenschlager M, von Boehmer H. PI3 kinase signalling blocks Foxp3 expression by sequestering Foxo factors. *J Exp Med.* 2010;207(7):1347–50. [PubMed: 20603315]
56. Sauer S, Bruno L, Hertweck A, Finlay D, Leleu M, Spivakov M, et al. T cell receptor signaling controls Foxp3 expression via PI3K, Akt, and mTOR. *Proc Natl Acad Sci U S A.* 2008;105(22):7797–802. [PubMed: 18509048]
57. Fawcett D, Pasceri P, Fraser R, Colbert M, Rossant J, Giguere V. Postaxial polydactyly in forelimbs of CRABP-II mutant mice. *Development.* 1995;121(3):671–9. [PubMed: 7720575]

58. Lee PP, Fitzpatrick DR, Beard C, Jessup HK, Lehar S, Makar KW, et al. A critical role for Dnmt1 and DNA methylation in T cell development, function, and survival. *Immunity*. 2001;15(5):763–74. [PubMed: 11728338]
59. Maekawa Y, Minato Y, Ishifune C, Kurihara T, Kitamura A, Kojima H, et al. Notch2 integrates signaling by the transcription factors RBP-J and CREB1 to promote T cell cytotoxicity. *Nat Immunol*. 2008;9(10):1140–7. [PubMed: 18724371]
60. Ehst BD, Ingulli E, Jenkins MK. Development of a novel transgenic mouse for the study of interactions between CD4 and CD8 T cells during graft rejection. *Am J Transplant*. 2003;3(11):1355–62. [PubMed: 14525595]
61. Li E, Sucov HM, Lee KF, Evans RM, Jaenisch R. Normal development and growth of mice carrying a targeted disruption of the alpha 1 retinoic acid receptor gene. *Proc Natl Acad Sci U S A*. 1993;90(4):1590–4. [PubMed: 7679509]
62. Katrancha SM, Shaw JE, Zhao AY, Myers SA, Cocco AR, Jeng AT, et al. Trio Haploinsufficiency Causes Neurodevelopmental Disease-Associated Deficits. *Cell Rep*. 2019;26(10):2805–17 e9. [PubMed: 30840899]
63. Mertins P, Tang LC, Krug K, Clark DJ, Gritsenko MA, Chen L, et al. Reproducible workflow for multiplexed deep-scale proteome and phosphoproteome analysis of tumor tissues by liquid chromatography-mass spectrometry. *Nat Protoc*. 2018;13(7):1632–61. [PubMed: 29988108]
64. Pamer EG. Immune responses to *Listeria monocytogenes*. *Nat Rev Immunol*. 2004;4(10):812–23. [PubMed: 15459672]
65. Bittner S, Afzali AM, Wiendl H, Meuth SG. Myelin oligodendrocyte glycoprotein (MOG35–55) induced experimental autoimmune encephalomyelitis (EAE) in C57BL/6 mice. *J Vis Exp*. 2014(86).
66. Schmiedel BJ, Singh D, Madrigal A, Valdovino-Gonzalez AG, White BM, Zapardiel-Gonzalo J, et al. Impact of Genetic Polymorphisms on Human Immune Cell Gene Expression. *Cell*. 2018;175(6):1701–15 e16. [PubMed: 30449622]
67. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol*. 2019;37(8):907–15. [PubMed: 31375807]

HIGHLIGHTS

- A cytoplasmic isoform of RAR α (cRAR α) is activated by the TCR and binds TCR-ZAP-70
- RA counteracts cRAR α for suboptimal T cell activation and enhanced Treg conversion
- Effective cRAR α -TCR signaling induces CRABP2 for RA translocation to the nucleus
- Defects in CRABP2 expression reduce anti-pathogen immunity and autoimmune disease




FIGURE 1: RA enhances Treg cell conversion independently of nuclear RAR α . (Related data shown in Figure S1)

(A-B) FOXP3 intracellular staining of (A) nRAR α -deficient or (B) DNRAR α (RAR403^{CD4-Cre}) spleen CD4 T cells stimulated for 96hrs with α -CD3/CD28 with or without TGF- β or RA (10 nM). (C) Naïve WT spleen CD4 T cells stimulated with α -CD3/CD28 with or without TGF- β . After 24hrs RA (10 nM) was added or not and cells were cultured for another 72hrs and stained for intracellular FOXP3. Shown representative histograms of 3 independent experiments (left) and statistical analyses of FOXP3⁺ T cell frequency and mean fluorescence intensity (MFI) (right). Statistical significance calculated with two-tailed Student t-test. Shown mean \pm SEM.




FIGURE 2: T cells express an extranuclear isoform of RARα. (Related data shown in Figure S2) (A) CD3, ZAP-70 and RARα staining of OT-I T cells co-cultured with OVA-loaded DCs for 40min. DAPI marks DNA. Representative cells are shown. Frequencies of synapse types from 1 representative experiment out of 2. (B) Naive spleen T cells transduced with FLAG-tagged RARα1 or RARα2 activated with α-CD3/CD28 for 5 or 30min. Localization of FLAG-tagged RARα1 and RARα2 assessed by western blotting. HSP90 as control. Data are representative of 3 experiments. (C) Mouse DP thymocytes and spleen T cells RNA-seq data aligned on UCSC mm10. Rara-mapped short-reads around exon 9 (gray background) shown. Numbers indicate short-reads of exons. J3, J4 and J5 are new splice junctions. (D) Junction usage with exon 9 in common. Forward (F) and reverse (R) primers indicated. RT-PCR of DN and DP thymocytes and CD4 and CD8 T cells with primers as shown. F1+R9 and F2+R9 primers detect *Rara1* and -2. F3 primer is of 5' end of exon E6 of XM_006532597.2. (E) Transcription start site(s) (TSS) of new *Rara* transcripts using FANTOM5 database. Forward and reverse transcription direction marked by red and blue peaks. New exon sequences shown. New RARα protein sequences with internal ATG

translational starts. Domains in RAR α 1, - α 2 and - α 3 are shown. **(F)** CD4 T hybridoma cells transduced with RAR α 1-GFP or RAR α 3-GFP analyzed by confocal microscopy and localization of GFP assessed. Images show 1 focal plane of representative cells. Graph represents mean \pm SEM from 150 or more cells from 1 representative experiment out of 2 (each dot represents 15–20 cells). **(G-H)** RAR α expression in cytoplasm (C) and nucleus (N) of human (Jurkat) **(G)** and primary mouse T cells **(H)**. Expression of RAR γ , Lamin B1 and HSP90 as control for N or C fractions, respectively. Data are representative from 2 (h) and 3 (m) independent experiments.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




FIGURE 3: Extranuclear RAR α controls TCR signal transduction. (Related data shown in Figure S3)

(A-B) Primary mouse spleen T cells transduced with (A) *Rara3*- or (B) *Rara1* cDNA, were activated with α -CD3/CD28 for 5min. Phosphorylation of ZAP-70 and PLC γ assessed by western blotting. HSP90 used as control. Representative of 2 experiments. (C) OT-I CD8 T cells stimulated with H2K^b-OVA tetramers. Cell-lysates immunoprecipitated with α -RAR α . Precipitates were assessed for ZAP-70 by western blotting. HSP90 in whole cell lysates as control. Shown 1 representative out of 3 experiments. (D) Spleen T cells transduced with CD3 ζ -GFP activated with α -CD3/CD28 for 30min. CD3 ζ -GFP clustering was quantified. Representative cells shown. Graph is mean \pm SEM from > 150 cells from 3 experiments. A large cluster was empirically defined. (E) OT-I T cells stimulated with H2K^b-OVA tetramers with or without RAR α inhibitor (Ro41-5253). Cell-lysates were immunoprecipitated with α -RAR α and blotted for ZAP-70 or RAR α . (F) WT or

RAR α 1-deficient spleen T cells activated with α -CD3/CD28 for 5min. Phosphorylation of PLC γ and ERK was assessed. HSP90 as control. Shown representative western blot out of 3. **(G-H)** Cell trace-labeled spleen T cells treated with DMSO, a pan-RAR antagonist LE540 (iRARs), an RAR α -specific antagonist Ro 41-5253 (iRAR α) or an RAR γ antagonist MM11253 (iRAR γ), activated with α -CD3/CD28 without **(G)** or with rIL-2 **(H)**. Proliferation assessed after 72hrs. Shown representative data. Graph means \pm SEM of division index calculated from 3 experiments expressed as percentage of control. **(I)** Primary spleen T cells treated with DMSO or RAR antagonist were activated with α -CD3/CD28 for 4hrs. Cleaved NOTCH1-intracellular domain assessed by western blotting. Shown 1 representative out of 3. **(J)** Primary spleen T cells with DMSO or inhibitor activated with α -CD3/CD28 for 4hrs and analyzed for *c-Myc* mRNA (upper panel) and c-MYC protein (lower panel). Graph represents means \pm SEM from 3 experiments. **(K)** Cell trace-labeled *dnRara* cre- or cre+ T cells with DMSO or iRAR α and activated with α -CD3/CD28. Proliferation assessed after 72hrs. Shown representative experiment. Graph represents means \pm SEM of division index from 3 experiments expressed as a percentage of control. **(L)** *dnRara* cre- or cre+ spleen T cells activated with α -CD3/CD28 for 4hrs. Cleaved NOTCH1 was assessed. Shown 1 representative western blot out of 3. **(M)** *dnRara* cre- or cre+ spleen T cells activated with α -CD3/CD28 for 4hrs. *c-Myc* mRNA assessed. Graph shows means \pm SEM from 2 experiments. Where appropriate band intensities were quantified by Relative Optical Density (ROD) relative to control.




FIGURE 4: Extranuclear and nuclear RAR α are activated by distinct mechanisms
(A) Higher-energy C-trap dissociation (HCD) MS2 mass spectrum identifying TCR-induced phosphorylation of RAR α at S445. Sequence of a 22 amino acid F domain peptide with phosphorylated serine is in lower case. Difference between measured and theoretical m/z expressed as parts per million (ppm). Product ions leading to the peptide sequence are shown, with *b*- (N-terminal) and *y*-series (C-terminal) ions indicated with left and right facing signs. **(B)** Phosphorylation dynamics of RAR α pS445 in the first 30min of T cell stimulation, relative to unstimulated, determined by tandem mass tag (TMT)-based quantification. Error bars show standard deviation for double or triple biological replicates. **(C)** RARE-Luc CD8 or CD4 T cells activated with α -CD3/CD28 for 24hrs, 48hrs and 72hrs. Luciferase activity was measured. The means \pm SEM of 2 experiments are shown. **(D)** Three OT-I TCR transgenic RARE-Luc mice infected i.v. with ActA⁻ Lm-OVA/day analyzed until 6 days post-infection for luciferase expression in spleen cells. Median fluorescence intensity of luciferase expression per mouse is shown. Data are from 1

experiment out of 2. **(E)** RNA-Seq data from human naïve CD8 or CD4 (CD3⁺ CD45RA⁺ CD127⁺ CCR7⁺) peripheral blood T cells stimulated with α -CD3/CD28 for 4hrs analyzed for *CRABP1* expression. *CRABP1* mRNA in transcripts per million (TPM) for each donor (n=88). P value calculated by Mann-Whitney U test. Datasets are from Database of Immune Cell Expression, expression quantitative trait loci (eQTLs) and epigenomics (DICE); <https://dice-database.org/>. **(F)** Mouse spleen CD8 T cells activated with α -CD3/CD28 or IL-7 or both. *Crabp1* mRNA was quantified. Data are means \pm SEM of 4 experiments. **(G)** Expression of *CRABP2* mRNA in activated human T cells assessed as in **(E)**. **(H-I)** CD8 or CD4 mouse spleen T cells activated with α -CD3/CD28 and *Crabp2* mRNA **(H)** and protein **(I)** were evaluated. Relative expression (mRNA) or western blot (Protein) of 1 of 3 experiments. **(J)** Expression of CRABP2 in CD4 mouse spleen T cells activated for 24hrs with α -CD3/CD28 alone or together with various doses of γ -secretase inhibitor (GSI). Quantification of the optical band density relative to control HSP90. Representative western blot of two independent experiments. **(K)** Spleen T cells activated with α -CD3/CD28 for 24hrs in DMSO or with MEK/ERK inhibitor U0126 (iERK). *Crabp2* mRNA was evaluated. Means \pm SEM of *Crabp2* mRNA are shown of 3 experiments. **(L)** Mouse spleen T cells activated with α -CD3/CD28 or IL-7 or both. mRNA of *Crabp2* was quantified. Data are means \pm SEM of 4 experiments.




FIGURE 5: RA counteracts extranuclear RAR α signaling in T cells. (Related data shown in Figure S4)

(A) WT or CRABP2-deficient spleen T cells activated with α -CD3/CD28 and RA for 72hrs and assessed for CCR9 expression. Graphs show means \pm SEM of CCR9⁺ T cells from 3 experiments. (B) WT- or CRABP2-deficient-RARE-Luc spleen T cells activated with α -CD3/CD28 for 48hrs and assessed for luciferase activity. Shown are means \pm SEM of luciferase activity per 50 μ g of protein for 3 experiments. (C) RARE-Luc CD8 (left) or CD4 (right) T cells activated with α -CD3/CD28 for 24hrs in DMSO, or a MEK/ERK inhibitor U0126 (iERK) or a PI3K inhibitor LY294002 (iPI3K). Luciferase activity was measured. Data are shown as means \pm SEM of 3 experiments. (D) WT or CRABP2-deficient spleen T cells activated with α -CD3/CD28 for 24hrs in DMSO and for WT T cells also with pan-RAR antagonist (iRARs) or RAR α antagonist (iRAR α). CRABP2 expression is evaluated. Representative western blot of 2 independent experiments. (E) Naïve CD4 T cells in DMSO, Retinoic Acid (RA) or an AKT inhibitor (AKT Inhibitor VIII) and activated with α -CD3/CD28 for indicated times. Phosphorylation of PLC γ and AKT was assessed. (F-G) Naïve CD4 T cells from nRAR α -deficient (F) or DNRAR α mice (G), activated with α -CD3/CD28 for indicated times were assessed for phosphorylation of PLC γ and AKT. (H) WT or CRABP2-deficient spleen T cells activated with α -CD3/CD28 for 72hrs were sorted and re-activated for 1min in DMSO or DMSO + (10nM) RA and PLC γ phosphorylation evaluated. Total PLC γ was measured as control. Data in (E-H) are representative from 3

experiments. **(I)** Proliferation of cell trace-labeled WT or CRABP2-deficient CD4 T cells activated with α -CD3/CD28 alone or with TGF- β or TGF- β + RA for 72hrs was evaluated. Histograms show a representative experiment and graphs show means \pm SEM of the division index from 3 experiments, expressed as a percentage of control. **(J)** Naïve WT or CRABP2-deficient CD4 T cells activated with α -CD3/CD28 alone or with TGF- β or TGF- β + RA for 72hrs were assessed for FOXP3. Frequency of FOXP3⁺ cells is shown. A representative histogram and means \pm SEM of 3 experiments are shown. **(K)** Naïve WT or CRABP2-deficient CD4 T cells activated with α -CD3/CD28 alone or with TGF- β or TGF- β + RA for 72hrs after 24hrs pre-activation without TGF- β + RA. FOXP3 was assessed and frequency of FOXP3⁺ cells assessed. A representative histogram and means \pm SEM of 3 experiments are shown. Student t-test was used for the statistical analysis. Where appropriate band intensity was quantified by ROD relative to control.




Figure 6: Extranuclear RAR α signaling controls TCR-induced proliferation *in vivo*. (Related data shown in Figure S5)

(A) Experimental strategy. (B) Tracking of adoptively transferred naïve CD45.2⁺ WT and CD45.1⁺ CRABP2-deficient OT-I cells in spleen of CD45.1⁺CD45.2⁺ recipient mice 7 days after intravenous (i.v.) infection with ActA⁻ Lm-OVA. Percentage of WT (CD45.2⁺) and CRABP2-deficient (CD45.1⁺) OT-I cells among donor cells is shown. (C) Tracking of adoptively transferred naïve CD45.2⁺ WT OT-I cells in spleen and small intestine epithelium of CD45.1⁺CD45.2⁺ vitamin A-free recipient mice treated with DMSO or RA, 7 days after infected i.v. with ActA⁻ Lm-OVA. Percentage is shown of WT OT-I T cells among total CD8 T cells. (D) Frequency of donor WT and CRABP2-deficient OT-I cells in the spleen of vitamin A-free CD45.1⁺CD45.2⁺ recipient mice infected i.v. with ActA⁻ Lm-OVA and assessed 7 days later. Percentage of WT and CRABP2-deficient OT-I cells among donor cells is shown. (E) Frequency of donor WT and CRABP2-deficient OT-I cells in spleen of DMSO or RA treated vitamin A-free CD45.1⁺CD45.2⁺ recipient mice infected i.v. with ActA⁻ Lm-OVA and assessed 7 days later. Percentage of WT and CRABP2-deficient OT-I cells among donor cells is shown. Data are representative of 2 experiments.




FIGURE 7: Extranuclear RAR α signaling controls effector differentiation *in vivo*. (Related data shown in Figure S6)

(A-C) EAE was induced and the clinical score was evaluated daily. (A) Cumulative clinical score, (B) maximum and minimum clinical scores and (C) images of WT and *Crabp2* conditional deletion mutant mice at day 22. Red arrows point to the hind limb position. (D-G) Histopathology analyses at day25 after EAE induction. (D) H&E staining of spinal cords from WT or *Crabp2* conditional deletion mutant mice, (E) Histological score comprising inflammation severity and axon dilation, (F) IFN γ expression of FOXP3⁻CD45⁺TGF- β ⁺CD4 T cells from the spinal cord, (G) Ratio of IFN γ ⁺ cells over IFN γ ⁻ and IL-17⁻ cells. P value was calculated by Student t-test. Five female mice of 8 to 12 weeks old were analyzed per condition in 2 EAE experiments. Shown mean \pm SD.

KEY RESSOURCE TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
T cell stimulation anti-CD3e	Thermo Fisher Scientific	16-0031-86
T cell stimulation anti-CD28	Thermo Fisher Scientific	16-0281-86
WB: anti-ZAP-70 (cl. D1C10E)	Cell Signaling Tech.	3165T
WB: anti-GAPDH (cl. 14C10)	Cell Signaling Tech.	85925SF
WB: anti-Lamin B1 (cl. D9V6H)	Cell Signaling Tech.	13435S
WB: anti-NOTCH1 (cl. D1E11)	Cell Signaling Tech.	3608T
WB: anti-RAR α	Cell Signaling Tech.	2554
WB: anti-RAR γ 1 (cl. D3A4)	Cell Signaling Tech.	8965T
WB: anti-CRABP2	One World Lab	11250
WB: anti-RAR α	Assay Biotechnology Company	R12-3424
WB: anti-HSP90 (cl. AC16)	Santa Cruz	sc-101494
WB: anti-Flag (cl. M2)	Millipore Sigma	A2220
WB: anti-AKT	Cell Signaling Tech.	9272
WB: anti-Phospho-AKT	Cell Signaling Tech.	9271T
WB: anti-Phospho- PLC γ 1	Cell Signaling Tech.	2821
WB: anti-Phospho-ZAP-70 (cl. 65E4)	Cell Signaling Tech.	2717
WB: anti-Phospho-ERK1/2 (cl. D13.14.4E)	Cell Signaling Tech.	4370
Activation: H-2K ^b -SIINFEKL tetramer	In house	N/A
IP: anti-RAR α	Santa Cruz	sc-551
Flow /Conf. Im.: anti-CD16-CD32 (cl. 2.4G2)	In house	N/A
Flow: anti-CD4-APC (cl. RM4-5)	BD	553051
Flow: anti-TCRgd-FITC (cl. GL3)	BD	553177
Flow: anti-CD45.1-FITC (cl. A20)	BD	553775
Flow: anti-CD44-V450 (cl. IM7)	BD	560452
Flow: anti-CD8a-PercpCy5.5 (cl. 53-6.7)	BD	551162
Flow: anti- CD8a-PeCy7 (cl. 53-6.7)	BD	552877
Flow: anti-IFN γ -BV711 (cl. XMG1.2)	Biologend	505835
Flow: anti-Il-17-PeCy7 (cl. TC11-18H10.1)	Biologend	506922
Flow: anti-CD8a-AF700 (cl. 53-6.7)	Biologend	100730
Flow: anti-CD4-AF700 (cl. GK1.5)	Biologend	100430
Flow: anti-FOXP3-PercpCy5.5 (cl. FJK-16s)	Thermo Fisher Scientific	45-5773-82
Flow: anti-CCR9-FITC (cl. eBioCW-1.2)	Thermo Fisher Scientific	11-1991-85
Flow: anti-TCRb-APCeF780 (cl. H57-597)	Thermo Fisher Scientific	47-5961-82
Flow: anti-CD45.2-APC (cl.104)	Thermo Fisher Scientific	17-0454-82
Flow: anti-RORgt-Pe (cl. AFKJS-9)	Thermo Fisher Scientific	12-6988-80
Flow: anti-CD45-eF450 (cl. 30-F11)	Thermo Fisher Scientific	48-0451-82

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Flow: anti-FOXP3-AF488 (cl. FJK-16s)	Thermo Fisher Scientific	53-5773-82
Flow: anti-FOXP3-Pe (cl. FJK-16s)	Thermo Fisher Scientific	12-5773-82
Flow: anti-CD25-APC (cl. PC61.5)	Thermo Fisher Scientific	17-0251-82
Flow: anti-CD4-eF450 (cl. RM4-5)	Thermo Fisher Scientific	48-0042-82
Flow: anti-c-MYC-FITC (cl. 9E10)	Thermo Fisher Scientific	13-2511
Conf. Im.: anti- CD3e-biotin (cl. 2C11)	Biologend	100304
Conf. Im.: Goat anti-mouse IgG-DyLight™ 488	Biologend	405310
Conf. Im.: Rabbit anti-Goat IgG-Alexa Fluor 647	Life Technology	A21446
Conf. Im.: anti-ZAP70 (cl. M20)	Santa Cruz	sc-1526
Conf. Im.: anti-RARα (c. Ralpha10)	Thermo Fisher Scientific	MA1810A
Bacterial and virus strains		
Listeria monocytogenes expressing OVA (ActA ⁻ Lm-OVA)	(64)	Gift
Biological samples		
N/A		
Chemicals, peptides, and recombinant proteins		
LIVE/DEAD™ Fixable Yellow Dead Cell Stain	Thermo Fisher Scientific	L34959
EDTA	Thermo Fisher Scientific	15575020
Collagenase D	Millipore Sigma	11088858001
DNase I	Millipore Sigma	10104159001
Human recombinant TGF-β	R&D System	240-B-002
Trans retinoic acid	Millipore Sigma	R2625
RAR antagonist LE540	FUJIFILM Wako Pure Chemical Corporation	Distributor 123-04521 Barcode No 4987481365261
RARα-selective antagonist	Enzo lifescience	Ro 41-5253
RARγ-selective antagonist MM 11253	Tocris bioscience	3822
MEK1/2 inhibitor U0126	Invivogen	tlrl-u0126
PI3K inhibitor LY294002	Invivogen	tlrl-ly29
γ-secretase inhibitor GSI	Tocris bioscience	2870
Human recombinant IL-2	Peptotech	200-02
AKT Inhibitor VIII	Cayman Chemical Company, Inc	14870
DMSO	Millipore Sigma	D2438-5X
TRIzol reagent	Thermo Fisher Scientific	15596026
TGX FastCast Acrylamide Solutions for SDS-PAGE gel	Biorad	1610171
PVDF membrane	Santa Cruz	sc-3723
NaCl	Millipore Sigma	S9888
Tris base	Santa Cruz	sc-362305
Glycerol	Millipore Sigma	G5516
Triton X-100	Millipore Sigma	X100
Protease and phosphatase inhibitor mixture	Millipore Sigma	11697498001
ECL solution	Santa Cruz	sc-2048

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Polybrene	Millipore Sigma	TR-1003-G
Poly I:C (HMW)	InvivoGen	tlrl-pic
OVA257-264 peptide	AnaSpec Inc.	AS-62572
Poly-L-lysine	R&D System	3438-100-01
NH ₄ Cl	Millipore Sigma	A4514
Saponin	Millipore Sigma	47036
Fish gelatin	Millipore Sigma	G7041
ProLong™ Gold Antifade Mountant with DNA Stain DAPI	Thermo Fisher Scientific	P36935
Dithiothreitol	Millipore Sigma	D2438
Percoll	Cytiva	17089109
Freund's adjuvant	BD	263910
M Tuberculosis H37Ra Dessicated	BD	231141
Pertussis toxin	List Biological Laboratories Inc	180
Zinc formalin	Anatech Lt	175
Pro-Par Clearant	Anatech Ltd	510
Conf. Im.: Streptavidine-Cyanine3	Biologend	405215
Critical commercial assays		
MACS CD8α ⁺ T cell isolation kit	Miltenyi Biotec	130-096-543
MACS CD4 ⁺ T cell isolation kit	Miltenyi Biotec	130-104-453
CD11c-Microbeads Ultrapure	Miltenyi Biotec	130-125-835
SureSelectXT RNA Direct Library Preparation kit	Agilent Technologies	G7564A
Pierce™ NE-PER® Nuclear and Cytoplasmic Extraction Reagent Kit	Thermo Fisher Scientific	78833
JetPrime transfection reagent	Polyplus transfection	101000027
Tandem mass tags (TMT) 11-plex reagents	Thermo Fisher Scientific	A34808
Cell Trace Violet	Thermo Fisher Scientific	C34557
Deposited data		
Raw and analyzed data	This paper	GEO: GSE112609
Database of Immune Cell Expression, Expression quantitative trait loci (eQTLs) and Epigenomics (DICE)	(66)	https://dice-database.org
UCSC mouse genome assembly (GRCm38/mm10)	UCSC Genome Browser Group	https://genome.ucsc.edu/cgi-bin/hgGateway
Experimental models: Cell lines		
Platinum-E	Cell Biolabs	RV-101
Jurkat, Clone E6-1	ATCC	TIB-152
MCC-T CD4 T cell hybridoma cells	(31)	Gift
Mutant ES cell (LoxP sites flanking Crabp2 exons)	EUCOMM	HEPD0679_8_F03
Experimental models: Organisms/strains		
Mouse: C57BL/6J	Jackson Lab.	000664
Mouse: Crabp2 ^{-/-} (10 generation backcrossed to B6 background)	Fawcett et al., 1995	Gift

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Mouse: RARE-Luciferase (DR5)	(39)	Gift
Mouse: RAR403 (DOMINANT NEGATIVE)	(28)	Gift
Mouse: CD4-Cre	(58)	Gift
Mouse: E8I-cre	(59)	Gift
Mouse: OT-I TCRM	Jackson Lab.	003831
Mouse: Act-mOVA	(60)	Gift
Mouse: RAR α 1 ^{-/-}	Jackson Lab.	023845
Mouse: Crabp2 ^{flox/flox} Crabp2 ^{tm1e(EUCOMM)Hmgu}	This paper	N/A
Mouse: Distal LCK-Cre	Jackson Lab.	Jackson Lab.
Oligonucleotides		
RT-RARa F1: CAGTCAGTGGTTACAGCACACCGTC	IDT DNA	N/A
RT-RARa F2: GGACTCCGCTTTGGAATGGCTC	IDT DNA	N/A
RT-RARa F3: GATGTCCAGGCCCAAGTAGAAGCCAG	IDT DNA	N/A
RT-RARa R1: GCTGCAATCTGCTGCTCATGCCTACAC	IDT DNA	N/A
RT-RARa R2: GGACTGACCTGCTGCAATCTGCTG	IDT DNA	N/A
RT-RARa R3: CTGGCTTCTACTTGGGCCTGGACATC	IDT DNA	N/A
RT-RARa R4: CAGGCTGTGAAAGACTCCTGCGGCT	IDT DNA	N/A
RT-RARa R5: CAGCATGTGTTATGCCAGGCTGTGAAAG	IDT DNA	N/A
RT-RARa R6: CCAACAGCATGTGTTATGCCAGGCTG	IDT DNA	N/A
RT-RARa R7: TTCACCTCACTTCTTCTCGGGAG	IDT DNA	N/A
RT-RARa R9: CCCCATAGTGGTAGCCGGATGATTTG	IDT DNA	N/A
RT-RARa R15: TCAGTGGAAACCCAGCAGGAACAGGTG	IDT DNA	N/A
RT-Hprt F: GTCGTGATTAGCGATGATGAACC	IDT DNA	N/A
RT-Hprt R: ATGACATCTCGAGCAAGTCTTTCAG	IDT DNA	N/A
qRT-Crabp2 F: CCCAGAGCTTTTAGCATTTCC	IDT DNA	N/A
qRT-Crabp2 R: GAAGATCTAAAGAGAAAGCCACCT	IDT DNA	N/A
qRT-cMYC F: TTGAAGGCTGGATTTCCCTTGGGC	IDT DNA	N/A
qRT-cMYC R: TCGTCGCAGATGAAATAGGGCTGT	IDT DNA	N/A
qRT-GADPH F: CAGATGCCTGCTTACCA	IDT DNA	N/A
qRT-GADPH R: ATGGCCTTCCGTGTTCT	IDT DNA	N/A
qRT-bACTIN F: CCAGAAGGACTGTTATGTGGGA	IDT DNA	N/A
qRT-bACTIN R: GACTCCGTGTTCAATGGGATAC	IDT DNA	N/A
qRT-HPRT F: GTCGTGATTAGCGATGATGAACC	IDT DNA	N/A
qRT-HPRT R: ATGACATCTCGAGCAAGTCTTTCAG	IDT DNA	N/A
Recombinant DNA		
Plasmid: MSCV-IRES-GFP	Addgene	20672
Plasmid: MSCV-IRES-Puromycin	Addgene	68469
Plasmid: CD3 ζ -GFP	Katz et al., 2016	Gift
Software and algorithms		

REAGENT or RESOURCE	SOURCE	IDENTIFIER
FlowJo software v10.8.0	Becton Dickinson & Company	https://www.flowjo.com/solutions/flowjo/downloads
HISAT2	(67)	https://github.com/DaehwanKimLab/hisat2
Tools (written in C using htslib) for manipulating next-generation sequencing data	Samtools	https://github.com/samtools/samtools
Integrative Genomics Viewer	Broad Institute	https://software.broadinstitute.org/software/igv/
FACSDIVA	Becton Dickinson & Company	https://www.bdbiosciences.com/en-us/products/software/instrument-software/bd-facsdiva-software
Spectrum Mill	Agilent/Broad Institute	https://proteomics.broadinstitute.org/millhome.htm
ZEN software	ZEISS	https://www.zeiss.com/microscopy/en/products/software/zeiss-zen.html
Prism 8.4.2	GraphPad	https://www.graphpad.com/features
Other		
Penicillin, streptomycin	Thermo Fisher Scientific	15140122
Puromycin	Invivogen	ant-pr-1
Balsticidin	Invivogen	ant-bl-05

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript