

Published in final edited form as:

Cell. 2023 September 28; 186(20): 4472–4473. doi:10.1016/j.cell.2023.08.036.

A widely distributed gene cluster compensates for uricase loss in hominids

Yuanyuan Liu,

J. Bryce Jarman,

Yen S. Low,

Hannah E. Augustijn,

Steven Huang,

Haoqing Chen,

Mary E. DeFeo,

Kazuma Sekiba,

Bi-Huei Hou,

Xiandong Meng,

Allison M. Weakley,

Ashley V. Cabrera,

Zhiwei Zhou,

Gilles van Wezel,

Marnix H. Medema,

Calyani Ganesan,

Alan C. Pao,


Saurabh Gombar,

Dylan Dodd*

Graphical abstract

 $^{^{\}star}$ Correspondence: ddodd2@stanford.edu.

Liu et al. Page 2



In the original Graphical Abstract of this paper, a chemical structure for uric acid was incorrectly labeled as xanthine.

The references for the above article were also incorrectly labeled in the original version. In the original version, reference 26 should have been reference 1, and references 1–22 should have been references 2–23. In addition, reference 37 should have been reference 24; references 23–25 should have been references 25–27; and references 27–36 should have been references 28–37.

Neither of these errors affect the results of the paper or the interpretation of the data, and both have now been corrected. We apologize for any confusion these errors may have caused.