

# Predicting the contribution of climate change on North Atlantic underwater sound propagation

Luca Possenti<sup>1</sup>, Gert-Jan Reichart<sup>1,2</sup>, Lennart de Nooijer<sup>1</sup>, Frans-Peter Lam<sup>3</sup>, Christ de Jong<sup>3</sup>, Mathieu Colin<sup>3</sup>, Bas Binnerts<sup>3</sup>, Amber Boot<sup>4</sup> and Anna von der Heydt<sup>4,5</sup>

<sup>1</sup> Ocean Systems (OCS), Royal Netherlands Institute for Sea Research (NIOZ), Texel, The Netherlands

<sup>2</sup> Department of Earth Sciences—Faculty of Geosciences, Utrecht University, Utrecht, The Netherlands

<sup>3</sup> Acoustics & Sonar, Netherlands Organization for Applied Scientific Research (TNO), The Hague, The Netherlands

<sup>4</sup> Department of Physics, Institute for Marine and Atmospheric research Utrecht (IMAU)—Faculty of Science, Utrecht University, Utrecht, The Netherlands

<sup>5</sup> Centre for Complex Systems Studies, Utrecht University, Utrecht, The Netherlands

## ABSTRACT

Since the industrial revolution, oceans have become substantially noisier. The noise increase is mainly caused by increased shipping, resource exploration, and infrastructure development affecting marine life at multiple levels, including behavior and physiology. Together with increasing anthropogenic noise, climate change is altering the thermal structure of the oceans, which in turn might affect noise propagation. During this century, we are witnessing an increase in seawater temperature and a decrease in ocean pH. Ocean acidification will decrease sound absorption at low frequencies (<10 kHz), enhancing long-range sound propagation. At the same time, temperature changes can modify the sound speed profile, leading to the creation or disappearance of sound ducts in which sound can propagate over large distances. The worldwide effect of climate change was explored for the winter and summer seasons using the (2018 to 2022) and (2094 to 2098, projected) atmospheric and seawater temperature, salinity, pH and wind speed as input. Using numerical modelling, we here explore the impact of climate change on underwater sound propagation. The future climate variables were taken from a Community Earth System Model v2 (CESM2) simulations forced under the concentration-driven SSP2-4.5 and SSP5-8.5 scenarios. The sound modeling results show, for future climate change scenarios, a global increase of sound speed at different depths (5, 125, 300, and 640 m) except for the North Atlantic Ocean and the Norwegian Sea, where in the upper 125 m sound speed will decrease by as much as  $40 \text{ m s}^{-1}$ . This decrease in sound speed results in a new sub-surface duct in the upper 200 m of the water column allowing ship noise to propagate over large distances (>500 km). In the case of the Northeast Atlantic Ocean, this sub-surface duct will only be present during winter, leading to similar total mean square pressure level ( $\text{SPL}_{\text{tot}}$ ) values in the summer for both (2018 to 2022) and (2094 to 2098). We observed a strong and similar correlation for the two climate change scenarios, with an increase of the top

Submitted 19 June 2023  
Accepted 8 September 2023  
Published 10 October 2023

Corresponding author  
Luca Possenti, luca.possenti@nioz.nl

Academic editor  
Guobin Fu

Additional Information and  
Declarations can be found on  
page 20

DOI [10.7717/peerj.16208](https://doi.org/10.7717/peerj.16208)

© Copyright  
2023 Possenti et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

200 m  $SPL_{tot}$  and a slowdown of Atlantic Meridional Overturning Circulation (AMOC) leading to an increase of  $SPL_{tot}$  at the end of the century by 7 dB.

**Subjects** Science Policy, Climate Change Biology, Aquatic and Marine Chemistry, Environmental Impacts

**Keywords** Climate change, North Atlantic Ocean, Underwater acoustics, Shipping noise, Human impact, Sound propagation, Atlantic Meridional Overturning Circulation

## INTRODUCTION

The natural soundscape is altered by anthropogenic activities such as shipping, transport, oil and gas exploitation, defense activities, tourism, fishing, offshore wind farming, and on- and near-shore construction (*Richardson et al., 2013; Duarte et al., 2021*). Among these the main anthropogenic noise source in the oceans is shipping, which dominates the soundscape in the low-frequency range (10 Hz to 1 kHz) (*Wenz, 1962*). Under 300 Hz this effect increased in the past 50–60 years because regions exposed to intense ship traffic have experienced an increase in ambient noise. In these regions ambient noise increased by 3 dB decade<sup>-1</sup> (*Andrew et al., 2002; Andrew, Howe & Mercer, 2011; Chapman & Price, 2011; Erbe et al., 2019; Miksis-Olds, Bradley & Maggie Niu, 2013; Miksis-Olds & Nichols, 2016*), resulting in an absolute sound increase by 15 to 20 dB (*Andrew et al., 2002; McDonald, Hildebrand & Wiggins, 2006; McKenna et al., 2012*). A major component of this increase is given by the rise in the number of ships, which is estimated to have doubled in the period between 1965 to 2000 (from approximately 44,000 to 88,000) (*Hildebrand, 2009*). Future estimates suggest that with the current rate of growth in ship traffic and economic trading, ambient noise is projected to continue to rise globally, especially in the Arctic and around Africa (*United Nations, 2021*). At the same time, humankind has introduced more than 330 Petagram of CO<sub>2</sub> into the atmosphere since the industrial revolution (starting around 1,760) (*Canadell et al., 2007*). A substantial part of the added CO<sub>2</sub> has been absorbed by the ocean (about 25%, (*Watson et al., 2020*)), which affected the oceanic carbon system. Global average surface ocean pH decreased from 8.21 to 8.1, corresponding to a 29% increase in H<sup>+</sup> activity (*Doney et al., 2009; Fabry et al., 2008*). Future projections suggest that in the next decades ocean CO<sub>2</sub> uptake will continue, decreasing the ocean pH in a process known as Ocean Acidification (OA). OA is adversely affecting the ocean environment by lowering sound absorption ( $\alpha$ ) at frequencies below 10 kHz which is controlled by pH-dependent borate ion chemistry (*Francois & Garrison, 1982*). At higher frequencies (>10 kHz) absorption is not affected by pH because this mechanism depends primarily on the chemical relaxation of magnesium sulfate and pure water viscous absorption. Largest relative pH-driven reduction in sound absorption will occur in the low frequency range, reaching values as high as a 40% reduction under 500 Hz (*Hester et al., 2008; Brewer & Hester, 2009*). Such a reduction in absorption will allow sound to travel further in situations when absorption is the dominant component in propagation loss (PL). The contribution of sound absorption at low frequencies (<500 Hz) is minimal, making its present and future contribution to the propagation loss negligible (*Udovydchenkov et al., 2010; Reeder & Chiu, 2010; Joseph & Chiu, 2010*). However, when sound is trapped in a

channel or duct in the ocean that may form as a consequence of the ocean's thermal structure and propagates over large distances, the latter effect may become important.

In addition to ongoing ocean acidification the ocean soundscape is primarily affected by other climate-related processes such as ocean warming, changes in wind speed and storm intensity and frequency, increase in sea-ice melting and decrease in salinity (Ainslie *et al.*, 2021; Andrew *et al.*, 2002; Duarte *et al.*, 2021; Munk, 2011; Young, Zieger & Babanin, 2011). For the period between 1971 to 2020 the total heat system had a heat gain of  $381 \pm 61$  ZJ with an associated total heating rate of  $0.48 \pm 0.1$  W m<sup>-2</sup> and about 89% of this heat is stored in the ocean (Von Schuckmann *et al.*, 2023). The temperature increase over the entire profile, together with sea-ice melting, is projected to alter the ocean's sound speed (*c*) profile. For a Representative Concentration Pathway RCP8.5, Affatati, Scaini & Salon (2022) quantified a general increase of sound speed up to 20 m s<sup>-1</sup> (1.5%) at the end of the current century.

The impact of climate change on marine ecosystems has been widely researched but implications are largely unknown. These changes in sound propagation due to climate change may have a substantial effect on marine mammals with specialized auditory systems (Wartzok *et al.*, 2004). Marine mammals use sound for various functions such as competition to show territorial hegemony, predation, mating and warning of others about presence of predators (Au & Hastings, 2008).

Potential impacts of climate change on the ocean's soundscape have received relatively little attention even though they may affect biology profoundly. The last assessment by the IPCC of climate change impacts (Skea, Shukla & Kikic, 2022) did not acknowledge climate change related impacts on the ocean soundscape, whereas the IPCC report on oceans and the cryosphere only acknowledged noise in the context of increased human operations in the Arctic Ocean (Poloczanska *et al.*, 2018) related to sea-ice melting.

Here we investigate the correlation between climate-related changes and the future sound propagation. We predict the expected changes by the end of the century in the sound field produced by a single vessel at 125 Hz using two different climate scenarios: Shared Socioeconomic Pathways SSP5-8.5 and SSP2-4.5 (Riahi *et al.*, 2017; O'Neill *et al.*, 2014). We selected a 125 Hz source frequency because (together with 63 Hz) it is a frequency band specified by the European MFSD (Marine Strategy Framework Directive) to assess the changes in ambient noise (Van der Graaf *et al.*, 2012). In a later study de Jong *et al.* (2021) recommended to use also higher frequencies to monitor ambient noise (e.g., 1 kHz). We contrast different areas globally to investigate spatial differences in the underwater sound propagation and compare different scenarios.

## MATERIALS AND METHODS

### Climate change data

To calculate the future change in PL, we retrieved 1 year of three-dimensional (3-D) monthly mean fields of salinity (*S*), temperature (*T*), pH and two-dimensional (2-D) monthly mean fields of near-surface (2 m) air temperature, Atlantic Meridional Overturning Circulation (AMOC) stream function and wind speed from 2022 to 2099 from the Community Earth System Model v2 (CESM2; Danabasoglu *et al.*, 2020) as

simulated in the Coupled Model Intercomparison Project 6 (CMIP6; [Eyring et al., 2016](#)) using the concentration driven SSP2-4.5 ([Danabasoglu, 2019](#)) and SSP5-8.5 scenarios ([Danabasoglu, 2019](#)) from the simulation r11i1p1f1. The ocean circulation model in CESM2, POP2 ([Smith et al., 2010](#)), uses a nominal horizontal resolution of  $1^\circ$  (100 km) on a displaced Greenland grid with 60 non-equidistant layers in the vertical. To fill the  $T$ ,  $S$  and pH depth gaps we interpolated the profiles using a shape-preserving piecewise cubic interpolation leading to the final profiles with 1 m resolution.

### Calculation of propagation loss

To determine changes in the future sound propagation we calculated the sound field produced by a single typical merchant vessel. We used as input the monopole vessel Source Level (SL) of 170.1 dB re  $1 \mu\text{Pa m}$  using the model proposed by [MacGillivray & de Jong \(2021\)](#) at a single frequency (125 Hz) for a bulker with reference length and speed (211 m and 13.9 kn).

To analyze the PL we selected six locations globally, including the North Atlantic Ocean. Locations used are two in the Atlantic Ocean ( $45^\circ \text{ N}$ ,  $40^\circ \text{ W}$  and  $48^\circ \text{ N}$ ,  $14^\circ \text{ W}$ ), Pacific Ocean ( $50^\circ \text{ N}$ ,  $167^\circ \text{ E}$ ), Southern Ocean ( $55^\circ \text{ S}$ ,  $140^\circ \text{ E}$ ), Arctic Ocean ( $75^\circ \text{ N}$ ,  $140^\circ \text{ W}$ ) and Norwegian Sea ( $72^\circ \text{ N}$ ,  $1^\circ \text{ W}$ ).

In each location, we placed the sound source at 6 m depth and we calculated the PL using the RAM parabolic equation model ([Collins, 1995](#)). We used the ocean bathymetry available from the 2022 General Bathymetry Chart of the Oceans ([https://www.gebco.net/data\\_and\\_products/gridded\\_bathymetry\\_data/](https://www.gebco.net/data_and_products/gridded_bathymetry_data/)). In the model, we evaluated the effect of climate change and ocean acidification using sound speed and potential density ( $\sigma_0$ ) mean profiles for boreal winter and summer (2018 to 2022) and (2094 to 2098).

We calculated sound speed and potential density using the equations of [Roquet et al. \(2015\)](#) that require temperature, salinity and pressure as inputs. Also, we calculated sound absorption using the formula of [van Moll, Ainslie & van Vossen \(2009\)](#) requiring an input of temperature, salinity, sound frequency ( $f$ ), depth ( $z$ ) and pH.

The final PL was calculated using a constant bathymetry from the sound source with a vertical resolution for PL of 1 m to a maximum distance of 500 km. Also, we assumed at every location the same sediment composition of very fine silt (grain size  $8 \mu\text{m}$ ), being the median grainsize for sediments in deep waters ([Ainslie, 2010](#)).

To calculate the sound pressure level of the ship ( $\text{SPL}_{\text{ship}}$ ) as a function of range and depth, the PL has been subtracted from the SL and then added to the ambient noise sound pressure level ( $\text{SPL}_{\text{wind}}$ ) calculated using a composite wind model ([Ainslie, 2010](#)).

To calculate  $\text{SPL}_{\text{wind}}$ , we assumed a homogenous wind surface source factor calculated using seasonal average wind speed. Subsequently we calculated  $\text{SPL}_{\text{wind}}$  as a function of depth propagating wind noise using a constant sound speed profile. The final result is presented as SPL over distance from the ship when  $\text{SPL}_{\text{ship}}$  (ship noise) exceeds  $\text{SPL}_{\text{wind}}$  presented as the total mean square pressure ( $\text{SPL}_{\text{tot}}$ ) of the sum of  $\text{SPL}_{\text{ship}}$  and  $\text{SPL}_{\text{wind}}$ . Also, when determining the effect of sound absorption, the difference of  $\text{SPL}_{\text{tot}}$  in the two different runs (with and without sound absorption) was calculated.

**Table 1** Methods summary.

Letter name	Method	Input
Source level (SL)	<i>MacGillivray &amp; de Jong (2021)</i>	Vessel length: 211 m Vessel speed 13.9 kn
Propagation loss (PL)	RAM	Bathymetry GEBCO Temperature profile CESM2 Salinity profile CESM2 pH profile CESM2
Ambient noise ( $SPL_{wind}$ )	<i>Ainslie (2010)</i>	Temperature profile CESM2 Air temperature CESM2 pH profile CESM2 Wind speed CESM2 Salinity profile CESM2

**Note:**

Methods used to calculate the final sound pressure level ( $SPL_{tot}$ ) at 125 Hz for SSP5-8.5 and SSP2-4.5 scenarios.

Subsequently, we present  $SPL_{tot}$  over 500 km for the entire water column for the six selected locations in the summer and winter seasons for SSP5-8.5 and SSP2-4.5 climate scenarios. In the results and discussion sections we discuss the differences between the six locations and the two seasons. To help the reader the methods applied in this section are synthesized in [Table 1](#).


Also, we relate the changes in  $SPL_{tot}$  in the North Atlantic Ocean (45° N–40° W) with changes in AMOC strength for SSP2-4.5 and SSP5-8.5 smoothed using a 5 years moving median of the yearly mean  $SPL_{tot}$  in the surface 200 m between 475 to 500 km distance from the source.

The final values of  $SPL_{tot}$  aims at identifying regions with the largest impact of predicted climate change on sound propagation. In order to calculate the absolute changes in noise levels all ambient noise sources should ideally be incorporated and a more complete wind model would be required. However, we expect the impact of using a more complex wind model to be minimal, given the high uncertainty of the wind within the climate data.

## RESULTS

### The trend in future sound speed


By 2098 sound speed will have increased (compared to 2022) globally under the SSP5-8.5 scenario at all depths below 300 m, with a predicted maximum increase in the North Atlantic Ocean, Labrador Sea and Norwegian Sea while it will decrease in the top 125 m ([Fig. 1](#)). In the less extreme SSP2-4.5 scenario the surface sound speed is also expected to increase, albeit less than under the SSP5-8.5 scenario with a maximum increase in the Arctic Ocean at 5 m below sea surface and the North Atlantic Ocean under 300 m ([Fig. 2](#)). In other regions such as the South Atlantic, Indian and Central Pacific Oceans the changes in sound speed will be negligible. For SSP2-4.5 scenario, the surface sound speed in the North Atlantic Ocean and Norwegian Sea is projected to decrease by  $>30 \text{ m s}^{-1}$ . In other locations such as the Pacific Ocean at around 0° N at 125 m and the Southern Ocean in the top 5 m sound speed is expected to remain more or less similar.



**Figure 1** Sound speed difference between (2018 to 2022) and (2094 to 2098) for SSP5-8.5. Maps of the difference in 5 years mean of sound speed ( $c$ ) in  $\text{m s}^{-1}$  between (2018 to 2022) and (2094 to 2098) at (A) 5, (B) 125, (C) 300 and (D) 640 m depth calculated for SSP5-8.5. The black dots indicate the sound source locations.  
[Full-size !\[\]\(1679558f37f6db0dd8360a2a7e913e90\_img.jpg\) DOI: 10.7717/peerj.16208/fig-1](https://doi.org/10.7717/peerj.16208/fig-1)

Sound speed profiles at the locations selected for the PL analysis for the winter and the summer seasons show a general sound speed increase in the top 2,000 m for SSP5-8.5 and SSP2-4.5 (Fig. 3). For SSP5-8.5 in the North Atlantic Ocean ( $47^\circ \text{N}$ ,  $14^\circ \text{W}$ ) the sound speed will be similar in the top 100 m, increasing up to  $40 \text{ m s}^{-1}$  in the deeper water column. A smaller increase is observed at another location in the North Atlantic Ocean ( $45^\circ \text{N}$ ,  $40^\circ \text{W}$ ) with  $10 \text{ m s}^{-1}$ . The SSP2-4.5 shows the same overall trends, but with different values: for example at  $47^\circ \text{N}$ ,  $14^\circ \text{W}$  the increase in sound speed is  $40 \text{ m s}^{-1}$  and at the surface at  $45^\circ \text{N}$ ,  $40^\circ \text{W}$  it decreases by  $5 \text{ m s}^{-1}$ .

Focusing on specific locations in the North Atlantic Ocean (Figs. 3A and 3B,  $45^\circ \text{N}$ ,  $40^\circ \text{W}$  and  $47^\circ \text{N}$ ,  $14^\circ \text{W}$ ), in 2022 there are two sound speed minima present: at 150 m and from 150 to 500 m. At the end of the century, the surface minimum will be more marked and the deep sound channel will deepen to 1,500 m. The deep sound channel is located at the slowest sound speed where sound waves can travel long distances. The difference in sound speed from the sub-surface channel and the layer below in 2022 is  $1 \text{ m s}^{-1}$ , which will increase to 24 and  $20 \text{ m s}^{-1}$  for SSP2-4.5 and SSP5.8.5, respectively. In 2022 the Norwegian Sea ( $72^\circ \text{N}$ ,  $1^\circ \text{W}$ ) shows a surface sound speed minimum that in the SSP2-4.5 and SSP5-8.5 scenario will disappear in favor of a sub-surface duct at 100 m (Fig. 3C). The sub-surface duct is defined by the sound speed minimum in the top 500 m and sound can travel large distances because continually bent, or refracted, towards the region of




**Figure 2** Sound speed difference between (2018 to 2022) and (2094 to 2098) for SSP2-4.5. Maps of the difference in 5 years mean of sound speed ( $c$ ) in  $\text{m s}^{-1}$  between (2018 to 2022) and (2094 to 2098) at (A) 5, (B) 125, (C) 300 and (D) 640 m depth calculated for SSP2-4.5. The black dots indicate the sound source locations.  
[Full-size !\[\]\(fd7fe780e8fd8eece60268c87d0c3e04\_img.jpg\) DOI: 10.7717/peerj.16208/fig-2](https://doi.org/10.7717/peerj.16208/fig-2)

lower sound speed. In both climate scenarios tested a deep sound channel will develop at 1,000 m before 2098.

In the Arctic Ocean ( $75^\circ \text{N}$ ,  $140^\circ \text{W}$ ) in 2022, a surface sound speed minimum is present that will have deepened by 2098 to 85 m (Fig. 3D). In the more severe climate change scenario (SSP5-8.5) this change is more pronounced compared to the moderate scenario SSP2-4.5 (Fig. 3). During winter the difference in sound speed from the surface to 85 m will be 2 and  $10 \text{ m s}^{-1}$  for SSP2-4.5 and SSP5-8.5, respectively. In the North Pacific Ocean ( $50^\circ \text{N}$ ,  $167^\circ \text{E}$ ), profiles in 2022 and 2098 are similar, with a sub-surface sound speed minimum located around 115 m depth (Fig. 3E). This minimum is expected to weaken over time, with a sound speed decrease in summer in the upper 115 m by  $18 \text{ m s}^{-1}$ , which will decelerate to a decrease by 2098 of  $12 \text{ m s}^{-1}$  for SSP5-8.5 and SSP2-4.5. In the Southern Ocean (Fig. 2F,  $60^\circ \text{S}$ ,  $25^\circ \text{E}$ ), the surface sound speed will also increase by  $>10 \text{ m s}^{-1}$  with an increase for SSP5-8.5 during summer ( $>15 \text{ m s}^{-1}$ ). However, the absolute change of sound speed has a minor effect on sound propagation because it is not affecting stratification.


### Trends in future sound absorption

The other analyzed variable potentially impacted by anthropogenic carbon addition and climate change is absorption. At the analyzed frequency of 125 Hz by the end of the century at 5, 125, 300 and 640 m depth, changes in sound absorption are small, with a



**Figure 3** Sound speed profiles for the selected locations. Sound speed ( $c$ ) in  $\text{m s}^{-1}$  profiles over depth for the winter (dashed lines) and summer season (continuous lines) where in blue is boreal summer (2018 to 2022), in red winter (2018 to 2022), in yellow summer (2094 to 2098) and in purple winter (2094 to 2098) for SSP5-8.5 and green and azure for summer and winter (2094 to 2098) for SSP2-4.5 for (A) Northwest Atlantic Ocean ( $45^\circ \text{ N } 40^\circ \text{ W}$ ), (B) Northeast Atlantic Ocean ( $47^\circ \text{ N } 14^\circ \text{ W}$ ), (C) Norwegian Sea ( $72^\circ \text{ N } 1^\circ \text{ W}$ ), (D) Arctic Ocean ( $75^\circ \text{ N } 140^\circ \text{ W}$ ), (E) North Pacific Ocean ( $50^\circ \text{ N } 167^\circ \text{ E}$ ) and (F) Southern Ocean ( $60^\circ \text{ S } 25^\circ \text{ E}$ ). [Full-size !\[\]\(5fd6ef84f97f42d7f8b34275f1b65312\_img.jpg\) DOI: 10.7717/peerj.16208/fig-3](https://doi.org/10.7717/peerj.16208/fig-3)





**Figure 4** Absorption difference between (2018 to 2022) and (2094 to 2098) for SSP5-8.5. Maps of the difference in 5 years mean of sound absorption ( $\alpha$ ) in  $\text{dB km}^{-1}$  in (2094 to 2098) and (2018 to 2022), calculated using *van Moll, Ainslie & van Vossen (2009)* algorithm at (A) 5, (B) 125, (C) 300 and (D) 640 m depth calculated for SSP5-8.5. The black dots indicate the sound source locations.


Full-size DOI: 10.7717/peerj.16208/fig-4

minor decrease in absorption at 5 m around  $80^\circ \text{N}$  of no more than  $0.0016 \text{ dB km}^{-1}$  that is just 0.8 dB at 500 km from the source. The decrease in absorption will be smaller closer to the equator, with a small decrease of  $0.0004 \text{ dB km}^{-1}$  (0.2 dB at 500 km) at  $0^\circ \text{N}$ . Another aspect is that these changes will disappear over depth with values close to  $0 \text{ dB km}^{-1}$ , for example from 640 m downward in the Pacific Ocean (Figs. 4 and 5). The decrease of absorption will be smaller in SSP2-4.5 compared to SSP5-8.5 where the largest decrease will be in the Arctic Ocean at 5 m with changes in sound absorption of around  $0.0008 \text{ dB km}^{-1}$  (0.4 dB at 500 km).

The absorption values at the selected locations (Fig. 6) show a consistent decrease in absorption at every location with a maximum decrease in the Norwegian Sea ( $72^\circ \text{N}$ ,  $1^\circ \text{W}$ ) of  $0.0019 \text{ dB km}^{-1}$  (0.95 at 500 km) for SSP5-8.5 and  $0.0013 \text{ dB km}^{-1}$  (0.65 dB at 500 km) for SSP2-4.5.

### Expected changes in sound propagation

The sound speed profile changes will lead to changes in received SPL at the different locations. The largest changes will be heard in the North Atlantic Ocean where in 2098 ship noise will travel in the sub-surface duct making the top 200 m noisier than today, that for scenario SSP5-8.5 will lead to an increase larger than 50 dB when propagating over 500 km. In the Northwest Atlantic Ocean ( $45^\circ \text{N}$ ,  $40^\circ \text{W}$ ) today, sound propagates mainly




**Figure 5** Absorption difference between (2018 to 2022) and (2094 to 2098) for SSP2-4.5. Maps of the difference in 5 years mean of sound absorption ( $\alpha$ ) in  $\text{dB km}^{-1}$  in (2094 to 2098) and (2018 to 2022), calculated using *van Moll, Ainslie & van Vossen (2009)* algorithm at (A) 5, (B) 125, (C) 300 and (D) 640 m depth calculated for SSP2-4.5. The black dots indicate the sound source locations.

Full-size DOI: [10.7717/peerj.16208/fig-5](https://doi.org/10.7717/peerj.16208/fig-5)


after reflection by the sea-bottom in a weak surface duct, particularly during winter (Fig. 7). During summer, the PL due to the interaction with the sea-bottom, does not allow sound to propagate over large distances and at the surface, it mostly is heard at the convergence zones. Here the sound rays propagating in the sub-surface duct interact with the rays reflected by the sea-bottom. In contrast we observe that by the year 2098 a new sub-surface duct will allow sound in the top 200 m to propagate over large distances (>500 km). This new duct is a robust observation, observed in different model simulations, albeit that it is similar for the SSP2-4.5 scenario in which some rays will still be reflected by the sea-bottom, forming convergence zones every 45 km. In particular, at the depth of the sound speed minimum (50 m below surface) the  $\text{SPL}_{\text{tot}}$  will be louder with increases larger than 50 dB (Fig. 8).

The  $\text{SPL}_{\text{tot}}$  for the Northeast Atlantic Ocean ( $47^\circ \text{ N}$ ,  $14^\circ \text{ W}$ ) shows a similar sound propagation (Fig. 9) to that in the Northwest Atlantic Ocean (Fig. 7). In 2098, the convergence zones will still be present, but in the winter sound will also propagate in a sub-surface duct (0 to 200 m) over >500 km for SSP5-8.5 and SSP2-4.5.

In 2098 the Norwegian Sea ( $72^\circ \text{ N}$ ,  $1^\circ \text{ W}$ , Fig. 10) will also have a new sub-surface sound duct. In the SSP2-4.5 and SSP5-8.5 scenarios, sound will propagate *via* the sub-surface duct, with some rays leaving the duct interacting with the sea-bottom. Today only part of the rays propagate *via* the surface duct and a large part of the rays reach the sea-bottom.




**Figure 6** Absorption profiles for the selected locations. Sound absorption ( $\alpha$ ) in  $\text{dB km}^{-1}$  profiles over depth for the winter (dashed lines) and summer season (continuous lines) where in blue is boreal summer (2018 to 2022), in red winter (2018 to 2022), in yellow summer (2094 to 2098) and in purple winter (2094 to 2098) for SSP5-8.5 and green and azure for summer and winter (2094 to 2098) for SSP2-4.5 for (A) Northwest Atlantic Ocean ( $45^\circ \text{ N } 40^\circ \text{ W}$ ), (B) Northeast Atlantic Ocean ( $47^\circ \text{ N } 14^\circ \text{ W}$ ), (C) Norwegian Sea ( $72^\circ \text{ N } 1^\circ \text{ W}$ ), (D) Arctic Ocean ( $75^\circ \text{ N } 140^\circ \text{ W}$ ), (E) North Pacific Ocean ( $50^\circ \text{ N } 167^\circ \text{ E}$ ), and (F) Southern Ocean ( $60^\circ \text{ S } 25^\circ \text{ E}$ ). [Full-size !\[\]\(ba1b80118482ccef74a5d718ca4d7242\_img.jpg\) DOI: 10.7717/peerj.16208/fig-6](https://doi.org/10.7717/peerj.16208/fig-6)




**Figure 7** Predicted  $SPL_{tot}$  for the Northwest Atlantic Ocean ( $45^{\circ}$  N  $40^{\circ}$  W) from a single bulker and wind. Predicted  $SPL_{tot}$  for the Northwest Atlantic Ocean ( $45^{\circ}$  N  $40^{\circ}$  W) from a single bulker and wind where (A) is winter (2018 to 2022), (B) winter (2094 to 2099) for SSP2-4.5, (C) winter (2094 to 2098) for SSP5-8.5 (D) summer (2018 to 2022), (E) summer (2094 to 2098) for SSP2-4.5 and (F) summer (2094 to 2098) for SSP5-8.5.

Full-size DOI: [10.7717/peerj.16208/fig-7](https://doi.org/10.7717/peerj.16208/fig-7)



**Figure 8** The sound pressure level from the bulker and wind. The figure shows for the Northwest Atlantic Ocean ( $45^{\circ}$  N  $40^{\circ}$  W) the SPL from the bulker ( $SPL_{ship}$ ) and from the wind ( $SPL_{wind}$ ) at a single depth (50 m) for (2018 to 2022) (red), (2094 to 2098) for SSP2-4.5 (orange) and SSP5-8.5 (yellow). The  $SPL_{ship}$  was calculated as the difference between the source level (SL) and the propagation loss (PL) derived using a parabolic equation model. The plot shows the  $SPL_{wind}$  in black for (2018 to 2022) (continuous line), (2094 to 2099) SSP2-4.5 (dotted line) and SSP5-8.5 (dash-dotted line).

Full-size DOI: [10.7717/peerj.16208/fig-8](https://doi.org/10.7717/peerj.16208/fig-8)



**Figure 9** Predicted  $SPL_{tot}$  for the Northeast Atlantic Ocean ( $47^{\circ}$  N  $14^{\circ}$  W) from a single bulker and wind. Predicted  $SPL_{tot}$  for the Northeast Atlantic Ocean ( $47^{\circ}$  N  $14^{\circ}$  W) from a single bulker and wind where (A) is winter (2018 to 2022), (B) winter (2094 to 2098) for SSP2-4.5, (C) winter (2094 to 2098) for SSP5-8.5 (D) summer (2018 to 2022), (E) summer (2094 to 2098) for SSP2-4.5 and (F) summer (2094 to 2098) for SSP5-8.5.

Full-size DOI: [10.7717/peerj.16208/fig-9](https://doi.org/10.7717/peerj.16208/fig-9)


The other locations studied, such as the North Pacific Ocean ( $50^{\circ}$  N,  $167^{\circ}$  E, Fig. 11), with the same  $SPL_{ship}$  will generally become quieter by 2098. In 2022 sound propagates reaching large distances ( $>500$  km), while in 2098 at the surface the sound will be confined to the convergence zones. In the Arctic Ocean ( $75^{\circ}$  N  $140^{\circ}$  W, Fig. 12) the  $SPL_{tot}$  in 2098 will be similar to 2022, with sound propagating through the entire water column. Despite the increase of sound speed at the surface in the Southern Ocean (Fig. 13)  $SPL_{tot}$  is expected to remain similar to today.

In particular, for SSP5-8.5 and SSP2-4.5 the contribution of absorption (*i.e.*, related to ocean acidification) to the future changes in  $SPL_{tot}$  is negligible ( $<1$  dB), with a maximum contribution of only 0.8 dB at 500 km in the Arctic Ocean.

## DISCUSSION

### Global drivers of the future sound propagation

Our results show that the predicted climate change for 2098 results in an overall increase of sound speed in the top 125 m of the water column, except for parts of the North Atlantic Ocean, Labrador and Norwegian Seas. These results are consistent with *Affatati, Scaini & Salon (2022)* using the CESM version 1 Large Ensemble project (LENS, (*Kay et al., 2015*)) forced with the RCP8.5 climate change scenario, who found an increase in sound speed from 2006–2016 to 2090–2100 up to  $24 \text{ m s}^{-1}$  (1.5 %) in the polar regions. Consistent with our study, the only regions where they identified a decrease in sound speed, with a




**Figure 10** Predicted  $SPL_{tot}$  for the Norwegian Sea ( $72^{\circ}$  N  $1^{\circ}$  W) from a single bulker and wind. Predicted  $SPL_{tot}$  for the Norwegian Sea ( $72^{\circ}$  N  $1^{\circ}$  W) from a single bulker and wind where (A) is winter (2018 to 2022), (B) winter (2094 to 2099) for SSP2-4.5, (C) winter (2094 to 2098) for SSP5-8.5 (D) summer (2018 to 2022), (E) summer (2094 to 2098) for SSP2-4.5 and (F) summer (2094 to 2098) for SSP5-8.5.

Full-size DOI: [10.7717/peerj.16208/fig-10](https://doi.org/10.7717/peerj.16208/fig-10)

maximum of  $10 \text{ m s}^{-1}$ , were parts of the Labrador Sea and North Atlantic Ocean. However, in our study for both the climate scenarios (SSP2-4.5 and SSP5-8.5) the decrease was much larger with a maximum of  $20 \text{ m s}^{-1}$  for SSP2-4.5 in the North Atlantic Ocean. This difference is probably partly caused by the difference in temperature and salinity projections between CESM1 and CESM2 and that in this we used 5 years when [Affatati, Scaini & Salon \(2022\)](#) used a 10 years mean.

At the frequency considered here ( $125 \text{ Hz}$ ), results show that the changes in PL are largely driven by stratification rather than sound absorption. Despite the large decrease in absorption, that in some cases is  $>60\%$ , the final contribution of absorption to  $SPL_{tot}$  is negligible  $1.5 \text{ dB}$  at  $500 \text{ km}$ . At higher frequencies, the contribution of absorption could be more important. For example, at  $500 \text{ Hz}$  absorption will decrease between  $0.007$  and  $0.019 \text{ dB km}^{-1}$ , with a maximum decrease in the Norwegian Sea between  $0.019$  and  $0.013 \text{ dB km}^{-1}$  and in the North Atlantic between  $0.023$  and  $0.082 \text{ dB km}^{-1}$  at  $3 \text{ kHz}$ . In particular, previous studies ([Duda, 2017](#); [Joseph & Chiu, 2010](#); [Reeder & Chiu, 2010](#); [Udovydenkov et al., 2010](#)) showed that in some scenarios, changes in absorption can significantly alter sound propagation. For example: when sound is trapped in a duct where it propagates without interactions with the sea-surface (*i.e.*, a deep sound channel). [Duda \(2017\)](#) found that in the Beaufort Sea Pacific Water duct pH will decrease by  $0.2$  (from  $8.1$  to  $7.9$ ) within the next  $30\text{--}50$  years and a source of  $900 \text{ Hz}$  located in the duct will thus have a SPL of  $7 \text{ dB}$  higher and sound will consequently travel  $38\%$  further. However, other




**Figure 11** Predicted sound pressure level for the North Pacific Ocean ( $50^{\circ}$  N  $167^{\circ}$  E) from a single bulker and wind. Predicted  $SPL_{tot}$  for the North Pacific Ocean ( $50^{\circ}$  N  $167^{\circ}$  E) from a single bulker and wind where (A) is winter (2018 to 2022), (B) winter (2094 to 2098) for SSP2-4.5, (C) winter (2094 to 2098) for SSP5-8.5 (D) summer (2018 to 2022), (E) summer (2094 to 2098) for SSP2-4.5 and (F) summer (2094 to 2098) for SSP5-8.5. Full-size [DOI: 10.7717/peerj.16208/fig-11](https://doi.org/10.7717/peerj.16208/fig-11)

studies with a surface sound source found similar results to our study, with an absolute change in SPL smaller than 2 dB (Duda, 2017; Joseph & Chiu, 2010; Reeder & Chiu, 2010; Udovychenkov et al., 2010).

The changes in  $SPL_{tot}$  are mainly visible in the top 200 m in the North Atlantic Ocean and Norwegian Sea, where the sub-surface duct (0 to 200 m) will become more marked. These changes are probably caused in both regions by changes in temperature and salinity profiles (Figs. S1 and S2). In particular, the decrease of surface temperature will increase the proportion of radiated power trapped in the ocean contributing to the increase of the future  $SPL_{tot}$ . The opposite effect will characterize regions where temperatures are projected to increase (e.g., Pacific and Southern Ocean), making the surface layer quieter. Ainslie (2011) quantified this contribution in a reduction of the noise level by 8% for a temperature increase of  $0.1^{\circ}$  C.

### Effect of the AMOC on the North Atlantic Ocean sound propagation

We observed the largest changes in  $SPL_{tot}$  in the North Atlantic Ocean and the Norwegian Sea. In these regions, stratification and consequently the sound speed profiles are controlled by the AMOC (Ivanovic et al., 2018; Haskins et al., 2020; Jackson et al., 2020). The AMOC started slowing down in the middle of the late 20<sup>th</sup> century and is still projected to continue slowing down in the next decades (Boers, 2021; Bryden, Longworth & Cunningham, 2005; Delworth & Dixon, 2000; Lynch-Stieglitz, 2017; Visbeck et al., 2001).




**Figure 12** Predicted sound pressure level for the Arctic Ocean (75° N 140° W) from a single bulker and wind. Predicted  $SPL_{tot}$  for the Arctic Ocean (75° N 140° W) from a single bulker and wind where (A) is winter (2018 to 2022), (B) winter (2094 to 2098) for SSP2-4.5, (C) winter (2094 to 2098) for SSP5-8.5 (D) summer (2018 to 2022), (E) summer (2094 to 2098) for SSP2-4.5 and (F) summer (2094 to 2098) for SSP5-8.5.

Full-size  DOI: [10.7717/peerj.16208/fig-12](https://doi.org/10.7717/peerj.16208/fig-12)


This slowdown has been observed by direct measurements at the Rapid Climate Change array at 26.5° N (*Smeed et al., 2018*), from 2014 by the OSNAP observing system at higher latitudes (*Susan Lozier et al., 2017*) and by temperature-based and geochemical proxy reconstructions (*Rahmstorf et al., 2015; Caesar et al., 2018; Thornalley et al., 2018*). For the next decades, the Sixth Assessment of the United Nations Intergovernmental Panel on Climate Change (IPCC, Geneva, Switzerland) projected that in the 21<sup>st</sup> century this slowing down will continue (*Skea, Shukla & Kilkis, 2022*). Previous studies showed that the AMOC changes are insensitive to the climate scenario (*Weijer et al., 2020*), which is also the case for our study (*Fig. 14*). The similar decrease of AMOC strength for the two climate scenarios lead to similar correlations with the North Atlantic Ocean surface sound duct  $SPL_{tot}$  (*Fig. 15*). In fact the two variables show a strong negative correlation for both climate change scenarios used, with an  $R^2$  of 0.8 for SSP5-8.5 and 0.87 for SSP2-4.5.  $SPL_{tot}$  in the duct will increase constantly over time with a maximum increase of  $SPL_{tot}$  of 7 dB at the end of this century. The strong observed correlation between AMOC and sound propagation shows that the AMOC will be the main driver affecting the future  $SPL_{tot}$  in the North Atlantic Ocean. Also, the consistency in the trend for both SSP2-4.5 and SSP5-8.5 shows that the changes in climate variables are proportional to the cumulative carbon emissions (*Herrington & Zickfeld, 2014; Notz & Stroeve, 2016; Steinacher & Joos, 2016*), implying that the AMOC in the first decades of any SSP scenario is mostly determined by historical CO<sub>2</sub> emissions. The mechanism behind the observed AMOC slowing is the






**Figure 13** Predicted sound pressure level for the Southern Ocean ( $60^{\circ}$  S  $25^{\circ}$  E) from a single bulker and wind. Predicted  $SPL_{tot}$  for the Southern Ocean ( $60^{\circ}$  S  $25^{\circ}$  E) from a single bulker and wind where (A) is winter (2018 to 2022), (B) winter (2094 to 2099) for SSP2-4.5, (C) winter (2094 to 2098) for SSP5-8.5 (D) summer (2018 to 2022), (E) summer (2094 to 2098) for SSP2-4.5 and (F) summer (2094 to 2098) for SSP5-8.5.

Full-size DOI: [10.7717/peerj.16208/fig-13](https://doi.org/10.7717/peerj.16208/fig-13)



**Figure 14** Changes in Atlantic Meridional Overturning Circulation (AMOC) strength over time for SSP5-8.5 and SSP2-4.5. Volumetric flow rate in Sverdrup of the Atlantic Meridional Overturning Circulation (AMOC) derived in the CESM2 for two different climate scenarios for SSP5-8.5 (in blue) and SSP2-4.5 (in red) at  $26.5^{\circ}$  N. The model data have been smoothed using a 5-year moving median.

Full-size DOI: [10.7717/peerj.16208/fig-14](https://doi.org/10.7717/peerj.16208/fig-14)



**Figure 15** Correlation between Atlantic Meridional Overturning Circulation (AMOC) strength and the sound pressure level in the top 200 m. Volumetric flow rate in Sverdrup of the Atlantic Meridional Overturning Circulation (AMOC) derived in the CESM2 using CMIP6 climate models for two different climate scenarios SSP5-8.5 (in blue) and SSP2-4.5 (in red) at 26.5° N smoothed using a 5-year moving median vs top 200 m median of the predicted  $SPL_{tot}$  between 475 and 500 km distance for the Northwest Atlantic Ocean (45° N 40° W) from a single bulker. The regression equations are: for SSP5-8.5:  $SPL_{tot}$  (dB re 1  $\mu$ Pa) =  $-1.2 \pm 0.1 AMOC + 85.7 \pm 0.8$  ( $R^2 = 0.8$ ) and for SSP2-4.5:  $SPL_{tot}$  (dB re 1  $\mu$ Pa) =  $-1.2 \pm 0.1 AMOC + 87.9 \pm 0.8$  ( $R^2 = 0.87$ ). [Full-size !\[\]\(1663bb69f307a960345edb0e712f8c02\_img.jpg\) DOI: 10.7717/peerj.16208/fig-15](https://doi.org/10.7717/peerj.16208/fig-15)

melting of ice and changes in the hydrological cycle (Liu et al., 2020) with the consequence that surface water in the North Atlantic Ocean (>40° N) will become less saline and colder, hindering the sinking of high-density (more saline) surface water. To confirm the correlation between AMOC and the changes in sound propagation, Figs. S3 and S4 shows that the regions of the North Atlantic Ocean where sound speed decrease the seawater temperature also decrease. This weakened AMOC will slow down even more the future Arctic due to sea-ice loss, with less heat reaching the Arctic Ocean (Liu et al., 2020; Boers, 2021) which modifies the North Atlantic Ocean and Norwegian Sea seawater temperature and salinity profiles (Figs. S1 and S2). In the North Atlantic Ocean and Norwegian Sea the surface salinity will decrease and temperature will increase with the exception of the Northeast Atlantic Ocean where surface winter temperature will decrease. Other consequences are changes in the intensity and frequency of winter storms over Europe (Woollings et al., 2012) and in sea level (Pardaens, Gregory & Lowe, 2011). Therefore, understanding the changes of AMOC is key for predicting the local impact of climate change with important consequences for the society, marine life and industry. Our results are consistent with previous studies, for a RCP8.5 a scenario similar to SSP5-8.5. Liu et al. (2020) predicted by 2100 a temperature decrease up to 1.8 K between 48 to 60° N in the entire water column with the largest decrease at the surface. In our study, this change in the

temperature profile lead to the formation of a new sub-surface sound duct at 150 m and a deepening of the sound channel.

Likely the projected changes in sound propagation will affect maritime users that rely on sound. For example, navies have been concerned with climate change effects (*National Research Council, 2010*). Particularly when relying heavily on acoustic sensors and systems, their performance in the North Atlantic Ocean and the Norwegian Sea will likely be severely affected. Also, we think these changes will impact fauna, although the exact extent remains to be investigated. Likely most harm will be done to marine mammals, compromising hearing ability and inducing physiological and behavioral changes putting animals under stress. Hence, further modelling and fieldwork studies are necessary to accurately quantify these changes and fully elucidate the mechanisms behind the changes in sound propagation.

## CONCLUSIONS AND FUTURE STUDIES

The strong correlation we observe between modeled sound propagation and the predicted changes in AMOC suggest that this could provide an additional toolbox to monitor AMOC changes. At the moment the AMOC is measured using transport mooring arrays equipped with dynamic height and current meters (*Smeed et al., 2018; Lozier et al., 2017; Meinen et al., 2013*) and indirect measurements such as satellite altimetry coupled with *in situ* measurements (e.g., Argo floats) (*McCarthy et al., 2020*). Unfortunately, there are no long observational records (or quantitative paleo proxies) of AMOC and that leads to a large uncertainty in AMOC projections with the consequence that it will take several decades to detect a forced trend in the AMOC due to the influence of internal variability (*Baehr et al., 2007; Roberts & Palmer, 2012; Roberts, Jackson & McNeall, 2014*). Due to this uncertainty, the Fifth report of the Intergovernmental Panel on Climate Change (IPCC, GE, Switzerland) concluded that a weakening of AMOC is likely for all scenarios, but the predicted weakening ranges between 34% and 45% (*Weijer et al., 2020*). The current methods along with new acoustic measurements could help to improve these predictions, with the acoustic measurements being suitable for real time monitoring. Such acoustic monitoring could be carried out using the existing (*Stanistreet et al., 2017; Davis et al., 2017; Durette-Morin et al., 2019; Soldevilla et al., 2014*) and new passive acoustic measurements and help to directly link the strength of AMOC with its acoustic impact on the marine ecosystem. We also suggest to use a series of surface artificial sound sources placed in different locations in the North Atlantic Ocean and measure the PL with a series of acoustic buoys. These acoustic buoys need to be equipped with several hydrophones located at different depths to capture the formation of new ducts. Combining models for climate and sound propagation we showed that climate change will significantly change the propagation of ship noise, especially the north Atlantic Ocean and the Norwegian Sea. These results are consistent between high (SSP5-8.5) and lower emission scenarios (SSP2-4.5). This implies that in the next century not only enhanced marine traffic will potentially make the future oceans noisier, but also a change in sound propagation. In the most affected regions changes will make the top 200 m noisier up to 7 dB, with possible adverse effects on marine life and maritime users that rely on sound. In other regions, the

propagation of ship noise will be similar to today with some regions that will be slightly quieter (e.g., Pacific and Southern Oceans).

The most likely mechanism behind the observed change in the propagation of ship noise in the North Atlantic is a slowing down of the AMOC, which will change the sound speed profile creating a stronger sub-surface duct at 150 m. This new duct will allow ship noise to propagate over large distances (>500 km). A strong correlation is observed between  $SPL_{tot}$  and AMOC ( $R^2 = 0.8$  for SSP5-8.5 and  $R^2 = 0.87$  for SSP2-4.5), which might open the way for future studies to quantify AMOC changes using sound propagation.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

This publication is part of the SOUND-2 project with project number P17-07 of the research program AQUA which is financed by the Dutch Research Council (NWO). Anna von der Heydt and Amber Boot were supported under the program of the Netherlands Earth System Science Centre (NESSC), financially supported by the Ministry of Education, Culture and Science (OCW, Grantnr. 024.002.001). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

SOUND-2 Project: P17-07.

Research Program AQUA.

Dutch Research Council (NWO).

Program of the Netherlands Earth System Science Centre (NESSC).

Ministry of Education, Culture and Science: 024.002.001.

### Competing Interests

The authors declare that they have no competing interests.

### Author Contributions

- Luca Possenti conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Gert-Jan Reichart conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Lennart de Nooijer conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Frans-Peter Lam conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Christ de Jong conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Mathieu Colin conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.

- Bas Binnerts conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Amber Boot conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.
- Anna von der Heydt conceived and designed the experiments, authored or reviewed drafts of the article, and approved the final draft.

### Data Availability

The following information was supplied regarding data availability:

The project data is available at WCRP CMIP6 using the single experiment run r11i1p1f1 for ssp5-8.5 and ssp2-4.5: <https://esgf-node.llnl.gov/search/cmip6/>.

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.16208#supplemental-information>.

## REFERENCES

- Affatati A, Scaini C, Salon S. 2022.** Ocean sound propagation in a changing climate: global sound speed changes and identification of acoustic hotspots. *Earth's Future* **10(3)**:e2021EF002099 DOI [10.1029/2021EF002099](https://doi.org/10.1029/2021EF002099).
- Ainslie MA. 2010.** *Principles of sonar performance modelling*. Vol. 707. Cham: Springer.
- Ainslie M. 2011.** Potential causes of increasing low frequency ocean noise levels. In: *Proceedings of Meetings on Acoustics 161ASA*, vol. 12. Melville: Acoustical Society of America, 70004.
- Ainslie MA, Andrew RK, Howe BM, Mercer JA. 2021.** Temperature-driven seasonal and longer term changes in spatially averaged deep ocean ambient sound at frequencies 63–125 Hz. *The Journal of the Acoustical Society of America* **149(4)**:2531–2545 DOI [10.1121/10.0003960](https://doi.org/10.1121/10.0003960).
- Andrew RK, Howe BM, Mercer JA. 2011.** Long-time trends in ship traffic noise for four sites off the North American west coast. *The Journal of the Acoustical Society of America* **129(2)**:642–651 DOI [10.1121/1.3518770](https://doi.org/10.1121/1.3518770).
- Andrew RK, Howe BM, Mercer JA, Dzieciuch MA. 2002.** Ocean ambient sound: comparing the 1960s with the 1990s for a receiver off the California Coast. *Acoustics Research Letters Online* **3(2)**:65–70 DOI [10.1121/1.1461915](https://doi.org/10.1121/1.1461915).
- Au WWL, Hastings MC. 2008.** *Principles of marine bioacoustics*. Vol. 510. Cham: Springer.
- Baehr J, Haak H, Alderson S, Cunningham SA, Jungclaus JH, Marotzke J. 2007.** Timely detection of changes in the meridional overturning circulation at 26 N in the Atlantic. *Journal of Climate* **20(23)**:5827–5841 DOI [10.1175/2007JCLI1686.1](https://doi.org/10.1175/2007JCLI1686.1).
- Boers N. 2021.** Observation-based early-warning signals for a collapse of the Atlantic meridional overturning circulation. *Nature Climate Change* **11(8)**:680–688 DOI [10.1038/s41558-021-01097-4](https://doi.org/10.1038/s41558-021-01097-4).
- Brewer PG, Hester K. 2009.** Ocean acidification and the increasing transparency of the ocean to low-frequency sound. *Oceanography* **22(4)**:86–93 DOI [10.5670/oceanog.2009.99](https://doi.org/10.5670/oceanog.2009.99).
- Bryden HL, Longworth HR, Cunningham SA. 2005.** Slowing of the Atlantic meridional overturning circulation at 25 N. *Nature* **438(7068)**:655–657 DOI [10.1038/nature04385](https://doi.org/10.1038/nature04385).

- Caesar L, Rahmstorf S, Robinson A, Feulner G, Saba V. 2018. Observed fingerprint of a weakening Atlantic Ocean overturning circulation. *Nature* 556(7700):191–196 DOI 10.1038/s41586-018-0006-5.
- Canadell J, Quere CLe, Raupach M, Field C. 2007. Contributions to accelerating atmospheric CO<sub>2</sub> growth from economic activity, carbon intensity, and efficiency of natural sinks. *Proceedings of the National Academy of Sciences of the United States of America* 104(47):18866–18870 DOI 10.1073/pnas.0702737104.
- Chapman NR, Price A. 2011. Low frequency deep ocean ambient noise trend in the Northeast Pacific Ocean. *The Journal of the Acoustical Society of America* 129(5):EL161–EL165 DOI 10.1121/1.3567084.
- Collins MD. 1995. *User's guide for RAM versions 1.0 and 1.0 P*. Vol. 20375. Washington, D.C.: Naval Research Lab, 14.
- Danabasoglu G. 2019. NCAR CESM2 model output prepared for CMIP6 ScenarioMIP Ssp245. Washington, D.C.: Earth System Grid Federation DOI 10.22033/ESGF/CMIP6.7748.
- Danabasoglu G, Lamarque J-F, Bacmeister J, Bailey DA, DuVivier AK, Edwards J, Emmons LK, Fasullo J, Gettelman A. 2020. The community earth system model version 2 (CESM2). *Journal of Advances in Modeling Earth Systems* 12(2):e2019MS001916 DOI 10.1029/2019MS001916.
- Davis GE, Baumgartner MF, Bonnell JM, Bell J, Berchok C, Thornton JB, Brault S, Buchanan G, Charif RA, Cholewiak D. 2017. Long-term passive acoustic recordings track the changing distribution of North Atlantic right whales (*Eubalaena Glacialis*) from 2004 to 2014. *Scientific Reports* 7(1):13460 DOI 10.1038/s41598-017-13359-3.
- de Jong CAF, Binnerts B, Robinson S, Wang L. 2021. Guidelines for modelling ocean ambient noise. Report of the EU INTERREG Joint Monitoring Programme for Ambient Noise North Sea (JOMOPANS).
- Delworth TL, Dixon KW. 2000. Implications of the recent trend in the Arctic/North Atlantic Oscillation for the North Atlantic thermohaline circulation. *Journal of Climate* 13(21):3721–3727 DOI 10.1175/1520-0442(2000)013<3721:JOTRTI>2.0.CO;2.
- Doney SC, Fabry VJ, Feely RA, Kleypas JA. 2009. Ocean acidification: the other CO<sub>2</sub> problem. *Annual Review of Marine Science* 1(1):169–192 DOI 10.1146/annurev.marine.010908.163834.
- Duarte CM, Chapuis L, Collin SP, Costa DP, Devassy RP, Eguiluz VM, Erbe C, Gordon TAC, Halpern BS, Harding HR. 2021. The soundscape of the Anthropocene Ocean. *Science* 371:6529 DOI 10.1126/science.aba4658.
- Duda TF. 2017. Acoustic signal and noise changes in the beaufort sea Pacific water duct under anticipated future acidification of Arctic Ocean waters. *The Journal of the Acoustical Society of America* 142(4):1926–1933 DOI 10.1121/1.5006184.
- Durette-Morin D, Davies KTA, Johnson HD, Brown MW, Moors-Murphy H, Martin B, Taggart CT. 2019. Passive acoustic monitoring predicts daily variation in north Atlantic right whale presence and relative abundance in Roseway Basin, Canada. *Marine Mammal Science* 35(4):1280–1303 DOI 10.1111/mms.12602.
- Erbe C, Marley SA, Schoeman RP, Smith JN, Trigg LE, Embling CB. 2019. The effects of ship noise on marine mammals—a review. *Frontiers in Marine Science* 6:606 DOI 10.3389/fmars.2019.00606.
- Eyring V, Bony S, Meehl GA, Senior CA, Stevens B, Stouffer RJ, Taylor KE. 2016. Overview of the coupled model intercomparison project phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development* 9(5):1937–1958 DOI 10.5194/gmd-9-1937-2016.

- Fabry VJ, Seibel BA, Feely RA, Orr JC. 2008.** Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES Journal of Marine Science* **65**(3):414–432  
[DOI 10.1093/icesjms/fsn048](https://doi.org/10.1093/icesjms/fsn048).
- Francois RE, Garrison GR. 1982.** Sound absorption based on ocean measurements. part II: boric acid contribution and equation for total absorption. *The Journal of the Acoustical Society of America* **72**(6):1879–1890 [DOI 10.1121/1.388673](https://doi.org/10.1121/1.388673).
- Haskins RK, Oliver KIC, Jackson LC, Wood RA, Drijfhout SS. 2020.** Temperature domination of AMOC weakening due to freshwater hosing in two GCMs. *Climate Dynamics* **54**(1):273–286  
[DOI 10.1007/s00382-019-04998-5](https://doi.org/10.1007/s00382-019-04998-5).
- Herrington T, Zickfeld K. 2014.** Path independence of climate and carbon cycle response over a broad range of cumulative carbon emissions. *Earth System Dynamics* **5**(2):409–422  
[DOI 10.5194/esd-5-409-2014](https://doi.org/10.5194/esd-5-409-2014).
- Hester KC, Peltzer ET, Kirkwood WJ, Brewer PG. 2008.** Unanticipated consequences of ocean acidification: a noisier ocean at lower PH. *Geophysical Research Letters* **35**(19):65  
[DOI 10.1029/2008GL034913](https://doi.org/10.1029/2008GL034913).
- Hildebrand JA. 2009.** Anthropogenic and natural sources of ambient noise in the ocean. *Marine Ecology Progress Series* **395**:5–20 [DOI 10.3354/meps08353](https://doi.org/10.3354/meps08353).
- Ivanovic RF, Gregoire LJ, Burke A, Wickert AD, Valdes PJ, Ng HC, Robinson LF, McManus JF, Mitrovica JX, Lee L. 2018.** Acceleration of northern ice sheet melt induces AMOC slowdown and northern cooling in simulations of the early last deglaciation. *Paleoceanography and Paleoclimatology* **33**(7):807–824 [DOI 10.1029/2017PA003308](https://doi.org/10.1029/2017PA003308).
- Jackson LC, Roberts MJ, Hewitt HT, Iovino D, Koenigk T, Meccia VL, Roberts CD, Ruprich-Robert Y, Wood RA. 2020.** Impact of ocean resolution and mean state on the rate of AMOC weakening. *Climate Dynamics* **55**(7):1711–1732 [DOI 10.1007/s00382-020-05345-9](https://doi.org/10.1007/s00382-020-05345-9).
- Joseph JE, Chiu C-S. 2010.** A computational assessment of the sensitivity of ambient noise level to ocean acidification. *The Journal of the Acoustical Society of America* **128**(3):EL144–EL149  
[DOI 10.1121/1.3425738](https://doi.org/10.1121/1.3425738).
- Kay JE, Clara Deser AP, Mai A, Hannay C, Strand G, Julie Michelle Arblaster SCB, Danabasoglu G, Edwards J. 2015.** The community earth system model (CESM) large ensemble project: a community resource for studying climate change in the presence of internal climate variability. *Bulletin of the American Meteorological Society* **96**(8):1333–1349  
[DOI 10.1175/BAMS-D-13-00255.1](https://doi.org/10.1175/BAMS-D-13-00255.1).
- Liu W, Fedorov AV, Xie S-P, Hu S. 2020.** Climate impacts of a weakened Atlantic meridional overturning circulation in a warming climate. *Science Advances* **6**(26):eaaz4876  
[DOI 10.1126/sciadv.aaz4876](https://doi.org/10.1126/sciadv.aaz4876).
- Lozier MS, Bacon S, Bower AS, Cunningham SA, De Jong MF, De Steur L, De young B, Fischer J, Gary SF, Greenan BJW. 2017.** Overturning in the subpolar North Atlantic program: a new international ocean observing system. *Bulletin of the American Meteorological Society* **98**(4):737–752 [DOI 10.1175/BAMS-D-16-0057.1](https://doi.org/10.1175/BAMS-D-16-0057.1).
- Lynch-Stieglitz J. 2017.** The Atlantic meridional overturning circulation and abrupt climate change. *Annual Review of Marine Science* **9**(585):83–104  
[DOI 10.1146/annurev-marine-010816-060415](https://doi.org/10.1146/annurev-marine-010816-060415).
- MacGillivray A, de Jong C. 2021.** A reference spectrum model for estimating source levels of marine shipping based on automated identification system data. *Journal of Marine Science and Engineering* **9**(4):369 [DOI 10.3390/jmse9040369](https://doi.org/10.3390/jmse9040369).

- McCarthy GD, Brown PJ, Flagg CN, Goni G, Houpert L, Hughes CW, Hummels R, Inall M, Jochumsen K, Larsen KMH. 2020. Sustainable observations of the AMOC: methodology and technology. *Reviews of Geophysics* 58(1):e2019RG000654 DOI 10.1029/2019RG000654.
- McDonald MA, Hildebrand JA, Wiggins SM. 2006. Increases in deep ocean ambient noise in the Northeast Pacific west of San Nicolas Island, California. *The Journal of the Acoustical Society of America* 120(2):711–718 DOI 10.1121/1.2216565.
- McKenna MF, Ross D, Wiggins SM, Hildebrand JA. 2012. Underwater radiated noise from modern commercial ships. *The Journal of the Acoustical Society of America* 131(1):92–103 DOI 10.1121/1.3664100.
- Meinen CS, Speich S, Perez RC, Dong S, Piola AR, Garzoli SL, Baringer MO, Gladyshev S, Campos EJD. 2013. Temporal variability of the meridional overturning circulation at 34.5 S: results from two pilot boundary arrays in the South Atlantic. *Journal of Geophysical Research: Oceans* 118(12):6461–6478 DOI 10.1002/2013JC009228.
- Miksis-Olds JL, Bradley DL, Maggie Niu X. 2013. Decadal trends in Indian Ocean ambient sound. *The Journal of the Acoustical Society of America* 134(5):3464–3475 DOI 10.1121/1.4821537.
- Miksis-Olds JL, Nichols SM. 2016. Is low frequency ocean sound increasing globally? *The Journal of the Acoustical Society of America* 139(1):501–511 DOI 10.1121/1.4938237.
- Munk W. 2011. The sound of climate change. *Tellus A: Dynamic Meteorology and Oceanography* 63(2):190–197 DOI 10.1111/j.1600-0870.2010.00495.x.
- National Research Council. 2010. National Security Implications of Climate Change for US Naval Forces: Letter Report. Available at <https://nap.nationalacademies.org/catalog/12897/national-security-implications-of-climate-change-for-us-naval-forces>.
- Notz D, Stroeve J. 2016. Observed Arctic sea-ice loss directly follows anthropogenic CO<sub>2</sub> emission. *Science* 354(6313):747–750 DOI 10.1126/science.aag2345.
- O'Neill BC, Kriegler E, Riahi K, Ebi KL, Hallegatte S, Carter TR, Mathur R, van Vuuren DP. 2014. A new scenario framework for climate change research: the concept of shared socioeconomic pathways. *Climatic Change* 122(3):387–400 DOI 10.1007/s10584-013-0905-2.
- Pardaens AK, Gregory JM, Lowe JA. 2011. A model study of factors influencing projected changes in regional sea level over the twenty-first century. *Climate Dynamics* 36(9):2015–2033 DOI 10.1007/s00382-009-0738-x.
- Poloczanska E, Mintenbeck K, Portner HO, Roberts D, Levin LA. 2018. The IPCC special report on the ocean and cryosphere in a changing climate. In: *2018 Ocean Sciences Meeting*. Washington, D.C.: AGU.
- Rahmstorf S, Box JE, Feulner G, Mann ME, Robinson A, Rutherford S, Schaffernicht EJ. 2015. Exceptional twentieth-century slowdown in Atlantic Ocean overturning circulation. *Nature Climate Change* 5(5):475–480 DOI 10.1038/nclimate2554.
- Reeder DB, Chiu C-S. 2010. Ocean acidification and its impact on ocean noise: phenomenology and analysis. *The Journal of the Acoustical Society of America* 128(3):EL137–EL143 DOI 10.1121/1.3431091.
- Riahi K, Van Vuuren DP, Kriegler E, Edmonds J, O'Neill BC, Fujimori S, Bauer N, Calvin K, Dellink R, Fricko O. 2017. The shared socioeconomic pathways and their energy, land use, and greenhouse gas emissions implications: an overview. *Global Environmental Change* 42:153–168 DOI 10.1016/j.gloenvcha.2016.05.009.
- Richardson WJ, Greene CR Jr, Malme CI, Thomson DH. 2013. *Marine mammals and noise*. Cambridge, MA: Academic press.



- Roberts CD, Jackson L, McNeill D. 2014.** Is the 2004–2012 reduction of the Atlantic meridional overturning circulation significant? *Geophysical Research Letters* **41(9)**:3204–3210 DOI [10.1002/2014GL059473](https://doi.org/10.1002/2014GL059473).
- Roberts CD, Palmer MD. 2012.** Detectability of changes to the Atlantic meridional overturning circulation in the hadley centre climate models. *Climate Dynamics* **39(9)**:2533–2546 DOI [10.1007/s00382-012-1306-3](https://doi.org/10.1007/s00382-012-1306-3).
- Roquet F, Madec G, McDougall TJ, Barker PM. 2015.** Accurate polynomial expressions for the density and specific volume of seawater using the TEOS-10 standard. *Ocean Modelling* **90(3–4)**:29–43 DOI [10.1016/j.ocemod.2015.04.002](https://doi.org/10.1016/j.ocemod.2015.04.002).
- Skea J, Shukla P, Kılıç Ş. 2022.** *Climate change 2022: mitigation of climate change*. Cambridge, MA, USA: Cambridge University Press.
- Smeed DA, Josey SA, Beaulieu C, Johns WE, Moat BI, Frajka-Williams E, Rayner D, Meinen CS, Baringer MO, Bryden HL. 2018.** The North Atlantic Ocean is in a state of reduced overturning. *Geophysical Research Letters* **45(3)**:1527–1533 DOI [10.1002/2017GL076350](https://doi.org/10.1002/2017GL076350).
- Smith R, Jones P, Briegleb B, Bryan F, Danabasoglu G, Dennis J, Dukowicz J, Eden C, Fox-Kemper B, Gent P. 2010.** The parallel ocean program (POP) reference manual ocean component of the community climate system model (CCSM) and community earth system model (CESM). Available at <https://www2.cesm.ucar.edu/models/cesm1.0/pop2/doc/sci/POPRefManual.pdf>.
- Soldevilla MS, Rice AN, Clark CW, Garrison LP. 2014.** Passive acoustic monitoring on the North Atlantic right whale calving grounds. *Endangered Species Research* **25(2)**:115–140 DOI [10.3354/esr00603](https://doi.org/10.3354/esr00603).
- Stanistreet JE, Nowacek DP, Baumann-Pickering S, Bell JT, Cholewiak DM, Hildebrand JA, Hodge LEW, Moors-Murphy HB, Van Parijs SM, Read AJ. 2017.** Using passive acoustic monitoring to document the distribution of beaked whale species in the western North Atlantic Ocean. *Canadian Journal of Fisheries and Aquatic Sciences* **74(12)**:2098–2109 DOI [10.1139/cjfas-2016-0503](https://doi.org/10.1139/cjfas-2016-0503).
- Steinacher M, Joos F. 2016.** Transient earth system responses to cumulative carbon dioxide emissions: linearities, uncertainties, and probabilities in an observation-constrained model ensemble. *Biogeosciences* **13(4)**:1071–1103 DOI [10.5194/bg-13-1071-2016](https://doi.org/10.5194/bg-13-1071-2016).
- Susan Lozier M, Bacon S, Bower AS, Cunningham SA, de Jong MF, de Steur L, Deyoung B, Fischer J, Gary SF, Greenan BJW. 2017.** Overturning in the subpolar North Atlantic program: a new international ocean observing system. *Bulletin of the American Meteorological Society* **98(4)**:737–752 DOI [10.1175/BAMS-D-16-0057.1](https://doi.org/10.1175/BAMS-D-16-0057.1).
- Thornalley DJR, Oppo DW, Ortega P, Robson JI, Brierley CM, Davis R, Hall IR, Moffa-Sanchez P, Rose NL, Spooner PT. 2018.** Anomalously weak labrador sea convection and Atlantic overturning during the past 150 years. *Nature* **556(7700)**:227–230 DOI [10.1038/s41586-018-0007-4](https://doi.org/10.1038/s41586-018-0007-4).
- Udovydchenkov IA, Duda TF, Doney SC, Lima ID. 2010.** Modeling deep ocean shipping noise in varying acidity conditions. *The Journal of the Acoustical Society of America* **128(3)**:EL130–EL136 DOI [10.1121/1.3402284](https://doi.org/10.1121/1.3402284).
- United Nations. 2021.** *The second world ocean assessment*. New York, USA: United Nations.
- Van der Graaf AJ, Ainslie MA, André M, Brensing K, Dalen J, Dekeling RPA, Robinson S, Tasker ML, Thomsen F, Werner S. 2012.** *European marine strategy framework directive-good environmental status (MSFD GEES): report of the technical subgroup on underwater noise and other forms of energy*. Brussels: TSG Noise & Milieu Ltd.

- van Moll CAM, Ainslie MA, van Vossen R. 2009. A simple and accurate formula for the absorption of sound in seawater. *IEEE Journal of Oceanic Engineering* 34(4):610–616 DOI 10.1109/JOE.2009.2027800.
- Visbeck MH, Hurrell JW, Polvani L, Cullen HM. 2001. The North Atlantic oscillation: past, present, and future. *Proceedings of the National Academy of Sciences of the United States of America* 98(23):12876–12877 DOI 10.1073/pnas.231391598.
- Von Schuckmann K, Minière A, Gues F, Cuesta-Valero FJ, Kirchengast G, Adusumilli S, Straneo F, Michaël A, Allan RP, Paul MB. 2023. Heat stored in the earth system 1960–2020: where does the energy go? *Earth System Science Data* 15(4):1675–1709 DOI 10.5194/essd-15-1675-2023.
- Wartzok D, Poppper AN, Gordon J, Merrill J. 2004. Factors affecting the responses of marine mammals to acoustic disturbance. *Marine Technology Society Journal* 37(4):6–15 DOI 10.4031/002533203787537041.
- Watson AJ, Schuster U, Shutler JD, Holding T, Ashton IGC, Landschützer P, Woolf DK, Goddijn-Murphy L. 2020. Revised estimates of ocean-atmosphere CO<sub>2</sub> flux are consistent with ocean carbon inventory. *Nature Communications* 11(1):4422 DOI 10.1038/s41467-020-18203-3.
- Weijer W, Cheng W, Garuba OA, Hu A, Nadiga BT. 2020. CMIP6 models predict significant 21st century decline of the Atlantic meridional overturning circulation. *Geophysical Research Letters* 47(12):e2019GL086075 DOI 10.1029/2019GL086075.
- Wenz GM. 1962. Acoustic ambient noise in the ocean: spectra and sources. *The Journal of the Acoustical Society of America* 34(12):1936–1956 DOI 10.1121/1.1909155.
- Woollings T, Gregory JM, Pinto JG, Reyers M, Brayshaw DJ. 2012. Response of the North Atlantic storm track to climate change shaped by ocean-atmosphere coupling. *Nature Geoscience* 5(5):313–317 DOI 10.1038/ngeo1438.
- Young IR, Zieger S, Babanin AV. 2011. Global trends in wind speed and wave height. *Science* 332(6028):451–455 DOI 10.1126/science.1197219.