



Diagnosis, Treatment, and Prevention of Tuberculosis Among People Experiencing Homelessness in the United States: Current Recommendations

Suzanne M. Marks, MPH, MA¹ ; Julie L. Self, PhD, MPH¹;
Thara Venkatappa, PhD, MSc¹ ; Marilyn B. Wolff, PhD, MPH¹;
Peri B. Hopkins, MPH¹; Ryan J. Augustine, MPH¹; Awal Khan, PhD¹;
Noah G. Schwartz, MD¹; Kristine M. Schmit, MD, MPH¹;
and Sapna Bamrah Morris, MD, MBA¹

Abstract

Objective: Tuberculosis (TB) is a public health problem, especially among people experiencing homelessness (PEH). The Advisory Council for the Elimination of Tuberculosis issued recommendations in 1992 for TB prevention and control among PEH. Our goal was to provide current guidelines and information in one place to inform medical and public health providers and TB programs on TB incidence, diagnosis, and treatment among PEH.

Methods: We reviewed and synthesized diagnostic and treatment recommendations for TB disease and latent TB infection (LTBI) as of 2022 and information after 1992 on the magnitude of homelessness in the United States, the incidence of TB among PEH, the role of public health departments in TB case management among PEH, and recently published evidence.

Results: In 2018, there were 1.45 million estimated PEH in the United States. During the past 2 decades, the incidence of TB was >10 times higher and the prevalence of LTBI was 7 to 20 times higher among PEH than among people not experiencing homelessness. TB outbreaks were common in overnight shelters. Permanent housing for PEH and the use of rapid TB diagnostic tests, along with isolation and treatment, reduced TB exposure among PEH. The use of direct observation enhanced treatment adherence among PEH, as did involvement of social workers to help secure shelter, food, safety, and treatment for comorbidities, especially HIV and substance use disorders. Testing and treatment for LTBI prevented progression to TB disease, and shorter LTBI regimens helped improve adherence. Federal agencies and the National Health Care for the Homeless Council have helpful resources.

Conclusion: Improvements in TB diagnosis, treatment, and prevention among PEH are possible by following existing recommendations and using client-centered approaches.

Keywords

tuberculosis, latent tuberculosis infection, diagnosis, treatment, prevention, homelessness

Tuberculosis (TB) is a public health problem in the United States, especially among people experiencing homelessness (PEH). An individual or family who lacks a fixed, regular, and adequate nighttime residence is generally defined as experiencing homelessness.¹ The rate of TB is higher among PEH than among people not experiencing homelessness.²⁻⁴ TB transmission can occur in congregate settings (eg, shelters, correctional facilities), especially in crowded shelters and when outside ventilation is diminished.⁵⁻⁷ PEH often have a greater risk than people not experiencing homelessness of developing TB or having comorbidities that increase TB risk, if infected with latent TB infection (LTBI).⁸

The Advisory Council for the Elimination of Tuberculosis (ACET) issued recommendations in 1992 for TB prevention

¹ Division of Tuberculosis Elimination, National Center for HIV, Viral Hepatitis, STD, and TB Prevention, Centers for Disease Control and Prevention, Atlanta, GA, USA

Corresponding Author:

Suzanne M. Marks, MPH, MA, Centers for Disease Control and Prevention, National Center for HIV, Viral Hepatitis, STD, and TB Prevention, Division of Tuberculosis Elimination, MS US12-4, 1600 Clifton Rd NE, Atlanta, GA 30329, USA.
Email: smarks@cdc.gov

and control among PEH.⁹ Since 1992, guidelines for TB diagnosis, treatment, and prevention among adults and children have been updated. For up-to-date recommendations on TB disease and LTBI diagnosis, treatment, and prevention among PEH, the audience, who might not be public health practitioners, would need to review multiple documents. Our objective was to review and synthesize the current TB guidelines that are relevant to PEH. Specifically, we assessed and reported the current magnitude of homelessness in the United States, the TB incidence among PEH, the applicability of current diagnostic and treatment guidelines for TB disease and LTBI to PEH, the role of public health departments in TB case management among PEH, and the recently published evidence for each of these topics. Our goal was to inform medical and public health providers, TB programs, and other service providers on the present TB incidence among PEH and on updated diagnosis and treatment protocols for TB among PEH.

Methods

To state the problem, we summarized the magnitude of homelessness by reviewing US Department of Housing and Urban Development (HUD) publications. To describe TB among PEH, we summarized statistics from reports on the US National TB Surveillance System, which includes data on homelessness within the past year.¹⁰ We identified and reviewed current recommendations as of 2022 relevant to PEH on diagnosis, treatment, and prevention of TB and LTBI for US populations. We then searched PubMed for publications published from January 1993 through April 2022 that provided evidence supporting the recommendations by using the following terms: “homeless or homelessness” AND “tuberculosis or latent tuberculosis infection” AND “treatment” AND “United States.” Among 260 results, we found 55 articles after excluding recommendation documents and duplicate articles from the same study or those included in systematic reviews. We selected systematic reviews that included ≥ 1 US study rather than single-site US studies. We retained some references prior to 1992 from the 1992 ACET recommendations.⁹

Results

Homelessness

For the most recent year (2018) for which estimates were not adversely affected by the COVID-19 pandemic, HUD reported an estimated 1.45 million PEH in the United States.¹¹ HUD also estimated that the number of PEH on a single night (“point-in-time”) in January 2020 was 580 466, of whom 61% were sheltered.¹²

The ACET 1992 recommendations advised health care and social service providers to ask their clients if they are experiencing homelessness or if they lack a fixed, regular, and adequate nighttime residence.⁹ To facilitate TB diagnosis

and treatment of PEH, public health departments are recommended to maintain, and regularly update, listings of single-room occupancy hotels and homeless shelters so that addresses of PEH can be checked against these listings. Shelters should maintain lists of people staying in their facilities and share clinical data on PEH.⁹ The National Health Care for the Homeless Program of the Bureau of Primary Health Care at the Health Resources and Services Administration (HRSA) funds health centers serving PEH and collects and shares health services data on PEH.¹³

TB Among PEH

ACET recommends assessment of the incidence of TB among PEH, which requires accurate PEH population estimates.⁹ In 2020, 290 TB cases among PEH (4.3% of TB cases in the United States) were reported to the US National TB Surveillance System.¹⁰ From 2007 through 2016 in the United States, the number of TB cases was 11 times greater among PEH (36 per 100 000 PEH) than among people not experiencing homelessness (3.1 TB cases per 100 000 US population).^{3,10} In 2021, estimates of TB incidence among PEH from 2011 through 2016 were also published for large HUD continuum of care program areas.³

Although data have not been systematically collected among all PEH, the estimated LTBI prevalence from individual studies ranged from 18% to 54% among PEH,¹⁴⁻¹⁶ which is 7 to 20 times higher than the estimated 2.7% to 5.0% LTBI prevalence in the general US population.¹⁷ This higher LTBI prevalence places PEH at greater risk for TB progression than the general population.

From 2009 through 2015, 8 of 21 TB outbreaks involved overnight facilities for PEH.¹⁸ Of 457 people included in these TB outbreaks, 204 (45%) had experienced homelessness in the year before diagnosis.¹⁸ Large genotypic clusters of TB cases gathered from data from 2009 through 2018 disproportionately included PEH, suggesting recent TB transmission.¹⁹

TB Diagnoses Among PEH

Rapid diagnosis of pulmonary TB, isolation, and treatment can reduce transmission in crowded congregate settings such as shelters, hospitals, and correctional facilities. The diagnostic process starts by determining whether TB disease symptoms are present, performing chest radiographs and, if results are abnormal, conducting additional tests.²⁰ Diagnostic tests for TB include analysis of sputum for acid-fast bacilli by microscopy and nucleic acid amplification testing (NAAT), which have turnaround times averaging 24 to 48 hours.²⁰⁻²³ Current American Thoracic Society/Infectious Diseases Society of America/Centers for Disease Control and Prevention (CDC) guidelines for diagnosis of TB disease among adults and children recommend performing diagnostic NAAT on the initial respiratory specimen from people with suspected pulmonary TB and on specimens collected from sites of suspected extrapulmonary TB.²⁰ One multisite

study found that incremental cost savings occurred with NAAT (vs no NAAT) among PEH to diagnose or rule out TB.²⁴ Chest radiography has also been used as a screening tool in shelters with a low incidence of TB.²⁵

Because shelter residents and staff (including volunteers) are at increased risk of TB infection and disease, the American Thoracic Society/Infectious Diseases Society of America/CDC and the US Preventive Services Task Force (USPSTF) guidelines recommend targeted testing and treatment for LTBI in this setting and population²⁶⁻²⁸ in accordance with updated TB diagnostic guidelines.²⁰ Among tests for TB infection, interferon-gamma release assays (IGRAs) performed on blood samples have been found to be more efficient than tuberculin skin tests (TSTs) in improving test completion and retention in LTBI treatment among PEH.²⁹⁻³¹ Testing frequency could range from one-time testing among PEH who are at low risk for TB exposure to annual testing among PEH who are at continued risk of exposure.²⁸ CDC and the USPSTF recommend that people with LTBI be evaluated and considered for treatment.^{28,32}

A study in a low-income US community concluded that mandating LTBI testing and treatment along with education and outreach decreased TB cases, transmission, deaths, and costs.³³ Parriott et al³⁴ conducted a systematic review of TB screening among PEH and found high proportions of PEH who had been successfully tested, with most PEH with LTBI successfully referred for treatment.

Case Management of PEH With TB

A timely public health department outreach visit to PEH with possible TB disease can facilitate access to housing and other health care resources (eg, treatment for mental health, substance use disorders, HIV) and development of a TB treatment plan, including scheduling the patient's first TB care visit.⁹ Adding details about a patient's physical description, activities, friends, and frequent gathering places in the medical record assists case managers in locating patients.

ACET recommends providing appropriate housing of PEH with infectious TB disease to allow directly observed initial TB therapy and to stop TB transmission in the community.⁹ Without housing, PEH often experience long hospitalizations for TB at significant cost to society.³⁵⁻⁴⁰ The Community Preventive Services Task Force and the US Interagency Council on Homelessness recommend Housing First programs to decrease homelessness, increase housing stability, and improve quality of life among PEH.^{41,42} Providing housing during treatment is also critical to ensuring TB treatment completion, prompting ACET to recommend the establishment of special shelters or other long-term care arrangements for PEH with TB disease.⁹

An analysis of TB cases from 1994 through 2003 found that, compared with TB patients not experiencing homelessness, PEH with TB had a higher prevalence of substance use (54% excess alcohol use, 29.5% noninjecting drug use, and 14% injecting drug use), with 34% of PEH tested having

coinfection with HIV.⁸ Shelter, food, safety, and comorbidities are likely to be top concerns for PEH; involvement of social workers (or other staff) on the treatment team to assist in solving these other problems is important for achieving successful TB treatment.⁹

TB and LTBI Treatment of PEH

TB disease treatment of PEH should follow CDC guidelines.⁴³ The use of shorter regimens (eg, 4-month drug-susceptible TB treatment) might facilitate treatment completion,⁴⁴ especially among people such as PEH who experience barriers to completion of longer 6- or 9-month regimens. Health care providers can assess interactions of TB medications with those for comorbidities to inform TB regimen choices.⁴⁵ Failure to take TB medications as prescribed can result in TB relapse, drug resistance, further TB transmission, and death. To facilitate adherence, the 1992 ACET recommendations stated that all TB treatments of PEH should be free of charge and transportation to clinics provided if needed.⁹ A patient-centered treatment approach that uses incentives and directly observed therapy, the standard of practice, allows careful monitoring for treatment adherence and drug side effects and results in high treatment completion rates.^{43,46-48} In-person directly observed therapy might reduce the likelihood of robbery or assault of PEH compared with self-administered treatment because TB medications might be mistaken by others to have street value.⁹ With adequate medical supervision, treatment can be given and observed by designated workers at the shelter or at another location.⁹ Directly observed therapy by mobile telephone or web-based video is feasible, has high uptake, has adherence rates similar to in-person directly observed therapy,^{49,50} including among PEH,⁵¹ and costs less than in-person directly observed therapy.^{52,53} If a patient with infectious TB disease (regardless of residential status) refuses treatment, temporary enforced isolation should be instituted in accordance with state and local public health laws and regulations.⁹ Medicaid reimbursement for some services might be available, depending on state regulations.

Shelter operators may accept hospitalized PEH with TB who have bacteriologic and clinical evidence of response to therapy (ie, 3 consecutive daily negative sputum smears and asymptomatic status).^{9,20}

Although homelessness has been associated with non-completion of LTBI treatment, the use of short rifamycin-based 3- or 4-month regimens compared with 6- or 9-month isoniazid-monotherapy regimens can facilitate successful completion of LTBI treatment, including among PEH,⁵⁴⁻⁵⁷ as does LTBI treatment from nurse-led community health workers.⁵⁸ Several studies have estimated that LTBI treatment is cost-effective compared with no LTBI treatment of PEH.^{59,60}

The following populations may be disproportionately represented among PEH and are recommended for TB prevention through TB infection testing and treatment: people with HIV,

people who inject drugs, people who had recent contact with patients with infectious TB disease, people with medical conditions that increase the risk of TB disease, and people with chest radiographs consistent with old, healed prior TB disease.^{9,20,61}

TB Prevention Among PEH

In addition to the treatment of LTBI and inadequately treated prior TB disease, identification of early or active cases, including the use of NAAT for diagnosis, and effective treatment of TB disease are the most important measures for preventing TB transmission.^{21,22,27,62-65} Staff workers and others who provide services at homeless shelters can assist in case finding by identifying people with productive, persistent cough (ie, lasting ≥ 3 weeks) and fever and ensuring that PEH with suspected TB disease are quickly evaluated by a health care provider. PEH with suspected TB disease should wear a surgical mask, be instructed to observe cough etiquette, and be transferred to respiratory isolation at a health care facility.²⁶

Public health departments should conduct thorough contact investigations of every infectious TB case.⁶⁶ Location-based investigations are the most successful among PEH, because naming of contacts by PEH often yields few names.⁶⁶ Contact investigations are usually based on TST or IGRA testing, followed by chest radiographs for those with TST reactions ≥ 5 mm or positive IGRA results. Because of the increased risk of TB in congregate settings and the high prevalence of immunosuppression among PEH, and because PEH are unlikely to return for the TST to be read, IGRAs are strongly recommended; however, TSTs are acceptable.²⁰ Screening PEH for TB using chest radiographs might be useful during outbreak investigations. The anticipated challenges for contact investigations involving PEH include difficulty locating patients and their contacts if they are mobile, have episodic incarceration, migrate between jurisdictions, have psychiatric illnesses (including substance use disorders) that hinder communication or participation, or have preexisting medical conditions (especially HIV).⁹

The number and density of people sharing the same breathing space in congregate settings are important factors influencing the likelihood of TB transmission.⁵ Supplemental upper room germicidal ultraviolet air disinfection may be useful to reduce ongoing transmission.⁶⁷⁻⁷² Shelter operators can consult with air quality experts to determine the adequacy of ventilation and to obtain recommendations for improvements.

Administrators of shelters for PEH are recommended to undertake the following actions (similar to those for outpatient health care settings) to prevent TB^{26,73,74}: (1) assign responsibility for TB infection control; (2) conduct periodic TB risk assessments; (3) develop and institute a written TB infection control plan to ensure prompt detection, airborne precautions, and treatment of people who have suspected or confirmed TB disease; (4) implement effective work practices for the management of staff and residents with suspected or confirmed TB disease; (5) ensure proper cleaning

and sterilization or disinfection of potentially contaminated surfaces; (6) train and educate shelter staff on TB, with a focus on prevention, transmission, and symptom identification; (7) maintain bed maps and track bed assignments, ideally in a searchable electronic format (eg, a spreadsheet) to facilitate contact investigations if a TB case is identified; (8) maintain as much space as possible between beds and position beds head to toe to reduce the possibility of transmission; (9) post signs and informational posters for client awareness and cough monitoring⁷⁵; (10) use a cough log to document people who are coughing, particularly at night, so that they can be referred for medical evaluation⁷⁴; and (11) coordinate efforts with the local or state public health department.

A workshop that included TB program staff, health care and social service providers for PEH, other health agency staff, and shelter administrators convened in 2015 and provided input to improve infection control in overnight shelters for PEH.⁷⁶ The group stated that addressing stigma (associated with poverty, experiences of discrimination or exclusion, and HIV) is also essential for optimizing TB prevention efforts among PEH.⁷⁶

HIV and TB

HIV is a major risk factor for the development of TB disease.^{77,78} An association among TB disease, HIV, and homelessness has been documented.^{27,43} The National Alliance to End Homelessness estimated that 3.4% of PEH were HIV positive in 2006,⁷⁹ compared with CDC's estimate of 0.4% of adults and adolescents in the general population⁸⁰ or an 8.5 times higher prevalence among PEH than among the general population. Among TB patients with HIV, PEH had higher odds of TB being attributed to recent transmission compared with TB patients not experiencing homelessness.⁸¹ Several studies have shown higher odds of TB-related mortality among people with HIV than among people without HIV.^{81,82}

The annual risk of TB disease due to reactivation of LTBI among people with untreated HIV has been estimated as 3% to 16% per year, which approximates the lifetime risk of TB for people with LTBI who are HIV negative.^{83,84} The advent of highly active antiretroviral therapy resulted in a reduced incidence of TB disease among people who are HIV positive.⁸⁵ Health care providers should implement routine HIV testing of PEH who are suspected to have TB, which can reduce mortality among immunosuppressed patients with TB and HIV if antiretroviral therapy is implemented.^{86,87}

People with HIV are recommended to be tested after diagnosis of HIV and annually if they have TB exposure risk.⁷⁸ Among people with HIV, rapid TB diagnosis using NAAT is essential, given the quick clinical progression to TB disease and risk of mortality among patients with advanced immunodeficiency.⁷⁸ With increasing immunodeficiency in people with HIV, chest radiographs are often atypical, or TB is extrapulmonary or disseminated, which makes TB diagnosis difficult and further increases the importance of NAAT.⁷⁸

Table 1. Summary of recommendations for tuberculosis (TB) diagnosis, treatment, and prevention among people experiencing homelessness (PEH), United States, 2022

Recommendation	Reference
1. TB programs should ascertain the housing status of people being evaluated for TB using as many modalities as possible (eg, interview, medical record review, homeless registry)	<ul style="list-style-type: none"> • CDC⁹ • CDC⁴¹
2. Public health departments should maintain and regularly update listings of single-room occupancy hotels and homeless shelters so that addresses of PEH with TB can be checked against these listings	<ul style="list-style-type: none"> • CDC⁹
3. Organizations that provide shelter and other types of emergency housing for PEH should develop institutional TB-control plans	<ul style="list-style-type: none"> • American Thoracic Society/CDC/Infectious Diseases Society of America²⁷
4. Homeless shelters should use administrative control interventions as a first line of defense to reduce TB exposure risk of shelter clients and staff	<ul style="list-style-type: none"> • Cole et al (Advisory Council for the Elimination of Tuberculosis and the National Tuberculosis Controllers Association)⁶ • Jensen et al (CDC)²⁶ • CDC⁷³ • Curry International Tuberculosis Center⁷⁴ • National Tuberculosis Controllers Association/CDC⁶⁶
5. TB programs should have written guidance or policies for investigating the index PEH with TB and sites of transmission	<ul style="list-style-type: none"> • CDC⁹
6. Targeted testing and treatment for TB infection is recommended for residents and staff of homeless shelters, following current TB diagnostic guidelines	<ul style="list-style-type: none"> • CDC⁹ • Lewinsohn et al (American Thoracic Society/Infectious Diseases Society of America/CDC)²⁰ • Jensen et al (CDC)²⁶ • US Preventive Services Task Force²⁸ • CDC, National Center for HIV, Viral Hepatitis, STD, and TB Prevention, Division of Tuberculosis Elimination⁶¹
7. Appropriate diagnostic tests (eg, sputum smears and cultures, NAAT of sputum and extrapulmonary specimens, chest radiographs, IGRAs, TSTs) should be used to evaluate people for TB; rapid diagnostic tests (eg, NAAT) are useful to quickly diagnose TB and prevent transmission	<ul style="list-style-type: none"> • CDC⁹ • Lewinsohn et al (American Thoracic Society/Infectious Diseases Society of America/CDC)²⁰ • CDC²¹ • CDC²² • CDC, National Center for HIV, Viral Hepatitis, STD, and TB Prevention, Division of Tuberculosis Elimination⁶¹ • National Institutes of Health⁷⁸ • CDC⁹
8. Shelters should maintain and share clinical data on PEH between shelters	<ul style="list-style-type: none"> • CDC⁹
9. Health care providers and organizations serving PEH should promptly notify the public health department of possible or confirmed TB cases among PEH or shelter staff	<ul style="list-style-type: none"> • CDC⁹ • Nahid et al (American Thoracic Society/CDC/Infectious Diseases Society of America)⁴³ • National Tuberculosis Controllers Association, CDC⁶⁶
10. PEH with newly diagnosed infectious TB disease should be appropriately housed to allow initial therapy to be directly observed and to preclude continuing transmission of TB in the community	<ul style="list-style-type: none"> • CDC⁹ • CDC⁴¹ • Nahid et al (American Thoracic Society/CDC/Infectious Diseases Society of America)⁴³ • Chaulk and Kazandjian (Public Health Tuberculosis Guidelines Panel)⁴⁶
11. TB programs should treat using a patient-centered approach to ensure that ancillary services, such as treatment for substance use disorders and evaluation and treatment of HIV, are provided to newly housed PEH with TB and to PEH in temporary housing facilities	<ul style="list-style-type: none"> • CDC⁹ • Nahid et al (American Thoracic Society/CDC/Infectious Diseases Society of America)⁴³ • Chaulk and Kazandjian (Public Health Tuberculosis Guidelines Panel)⁴⁶
12. Public health department staff should visit PEH with possible or confirmed TB, in the hospital or elsewhere, as soon as possible during diagnosis, to obtain patient consent on treatment plans	<ul style="list-style-type: none"> • CDC⁹ • National Tuberculosis Controllers Association, CDC⁶⁶
13. Before hospitalized PEH with TB are discharged, arrange for their first visit to the clinic or other place of intended outpatient care	<ul style="list-style-type: none"> • CDC⁹ • Nahid et al (American Thoracic Society/CDC/Infectious Diseases Society of America)⁴³

(Continued)

Table 1. (Continued)

Recommendation	Reference
14. Consider involving a social worker on the TB treatment team for PEH to facilitate access to food, shelter, and safety	<ul style="list-style-type: none"> • CDC⁹ • Chaulk and Kazandjian (Public Health Tuberculosis Guidelines Panel)⁴⁶ • National Tuberculosis Controllers Association, CDC⁶⁶
15. Use directly observed therapy and incentives to enhance treatment adherence	<ul style="list-style-type: none"> • CDC⁹ • Nahid et al (American Thoracic Society/CDC/Infectious Diseases Society of America)⁴³ • Chaulk and Kazandjian (Public Health Tuberculosis Guidelines Panel)⁴⁶ • CDC⁹
16. If TB clinics are not close to the location of PEH, transportation to the clinic should be provided or services should be brought to the PEH	<ul style="list-style-type: none"> • CDC⁹
17. If PEH with infectious TB refuse treatment, temporary, enforced isolation should be instituted in accordance with state and local public health laws and regulations	<ul style="list-style-type: none"> • CDC⁹ • National Tuberculosis Controllers Association, CDC⁶⁶
18. TB and LTBI treatment should follow CDC guidelines. Health care providers should prescribe short regimens when possible and avoid critical drug interactions	<ul style="list-style-type: none"> • CDC⁹ • Sterling et al (National Tuberculosis Controllers Association and CDC)³² • Nahid et al (American Thoracic Society/CDC/Infectious Diseases Society of America)⁴³ • Carr et al⁴⁴ • University of Liverpool⁴⁵
19. Public health departments and service providers for PEH should work together to conduct active case finding to identify TB cases and their contacts early, especially during outbreaks	<ul style="list-style-type: none"> • American Thoracic Society, CDC, Infectious Diseases Society of America²⁷ • National Tuberculosis Controllers Association, CDC⁶⁶
20. TB programs should conduct a thorough, location-based contact investigation for every patient, based upon testing with IGRA (or TST), followed by chest radiographs and NAAT for those with positive test results	<ul style="list-style-type: none"> • CDC⁹ • Lewinsohn et al (American Thoracic Society/Infectious Diseases Society of America/CDC)²⁰ • CDC²¹ • CDC²² • National Tuberculosis Controllers Association, CDC⁶⁶
21. All PEH initiating treatment for TB should be screened routinely for HIV	<ul style="list-style-type: none"> • CDC⁹ • Nahid et al (American Thoracic Society/CDC/Infectious Diseases Society of America)⁴³ • Branson et al (CDC)⁸⁷
22. The presence of HIV in a person with a positive IGRA or TST result is an indication for LTBI treatment following current guidelines; LTBI treatment should be started only after excluding pulmonary or extrapulmonary TB disease	<ul style="list-style-type: none"> • Lewinsohn et al (American Thoracic Society/Infectious Diseases Society of America/CDC)²⁰ • Sterling et al (National Tuberculosis Controllers Association and CDC)³²
23. PEH with HIV who have contact with infectious TB should receive a chest radiograph, be screened for signs and symptoms of TB, and be examined for evidence of extrapulmonary TB, regardless of IGRA or TST result; if abnormalities are noted, additional diagnostic studies for TB should be undertaken	<ul style="list-style-type: none"> • Lewinsohn et al (American Thoracic Society/Infectious Diseases Society of America/CDC)²⁰ • CDC²¹ • CDC²² • National Tuberculosis Controllers Association, CDC⁶⁶ • National Institutes of Health⁷⁸
24. TB programs can access CDC-funded Tuberculosis Centers of Excellence for Training, Education, and Medical Consultation for resources and training to prevent and manage TB among PEH	<ul style="list-style-type: none"> • CDC⁹ • CDC⁸⁸ • CDC⁸⁹
25. TB programs can consult with local and national programs focusing on PEH (eg, through the Bureau of Primary Health Care at the Health Resources and Services Administration, the National Health Care for the Homeless Council, and Ryan White programs for health care and supportive services for PEH)	<ul style="list-style-type: none"> • CDC⁹

Abbreviations: CDC, Centers for Disease Control and Prevention; IGRA, interferon-gamma release assay; LTBI, latent tuberculosis infection; NAAT, nucleic acid amplification test; TST, tuberculin skin test.

Table 2. Summary of recommendations for tuberculosis diagnosis, treatment, and prevention among people experiencing homelessness, United States, 2022, in chronological order of publication

Year	Name of recommendation	Reference
1992	Prevention and control of tuberculosis among homeless persons. Recommendations of the Advisory Council for the Elimination of Tuberculosis	CDC ⁹
1998	Directly observed therapy for treatment completion of pulmonary tuberculosis: consensus statement of the Public Health Tuberculosis Guidelines Panel	Chaulk and Kazandjian (Public Health Tuberculosis Guidelines Panel) ⁴⁶
2005	Guidelines for preventing the transmission of <i>Mycobacterium tuberculosis</i> in health-care settings	Jensen et al (CDC) ²⁶
2005	American Thoracic Society/Centers for Disease Control and Prevention/ Infectious Diseases Society of America: controlling tuberculosis in the United States	American Thoracic Society/CDC/Infectious Diseases Society of America ²⁷
2005	Guidelines for the investigation of contacts of persons with infectious tuberculosis	National Tuberculosis Controllers Association, CDC ⁶⁶
2006	Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings	Branson et al (CDC) ⁸⁷
2006	Tuberculosis (TB) risk assessment worksheet	CDC ⁷³
2009	Updated guidelines for the use of nucleic acid amplification tests in the diagnosis of tuberculosis	CDC ²²
2013	Availability of an assay for detecting <i>Mycobacterium tuberculosis</i> , including rifampin-resistant strains, and considerations for its use—United States, 2013	CDC ²¹
2013	Shelters and TB: what staff need to know	Curry International Tuberculosis Center ⁷⁴
2016	Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America clinical practice guidelines: treatment of drug-susceptible tuberculosis	Nahid et al (American Thoracic Society/ CDC/Infectious Diseases Society of America) ⁴³
2016	Screening for latent tuberculosis infection in adults: US Preventive Services Task Force recommendation statement	US Preventive Services Task Force ²⁸
2017	Official American Thoracic Society/Infectious Diseases Society of America/ Centers for Disease Control and Prevention clinical practice guidelines: diagnosis of tuberculosis in adults and children	Lewinsohn et al (American Thoracic Society/ Infectious Diseases Society of America/ CDC) ²⁰
2019	Community guide reference to Housing First recommendations	CDC ⁴¹
2020	Essential components of a public health tuberculosis prevention, control, and elimination program: recommendations of the Advisory Council for the Elimination of Tuberculosis and the National Tuberculosis Controllers Association	Cole et al ⁶
2020	Guidelines for the treatment of latent tuberculosis infection: recommendations from the National Tuberculosis Controllers Association and CDC, 2020	Sterling et al ³²
2020	Latent tuberculosis infection: a guide for primary health care providers	CDC, National Center for HIV, Viral Hepatitis, STD, and TB Prevention, Division of Tuberculosis Elimination ⁶¹
2022	Guidelines for the prevention and treatment of opportunistic infections in adults and adolescents with HIV	National Institutes of Health ⁷⁸
2022	Interim guidance: 4-month rifapentine–moxifloxacin regimen for the treatment of drug-susceptible pulmonary tuberculosis—United States, 2022	Carr et al (CDC) ⁴⁴

Abbreviation: CDC, Centers for Disease Control and Prevention.

Resources

Tables 1 and 2 list current US guidelines and recommendations^{6,9,20–22,26,28,32,41,43,44,46,61,65,77,86} relevant to TB diagnosis, treatment, and prevention among PEH. We list the recommendations in Table 1 and the recommendation documents chronologically in Table 2. CDC provides information and links to resources for TB programs, homeless shelters, and other service providers for PEH.^{88,89} CDC-funded

Tuberculosis Centers of Excellence for Training, Education, and Medical Consultation (TB COEs) are resources for public health departments and other entities to access expert TB medical consultation for providers caring for PEH.⁹⁰ The TB COEs have also developed educational materials and shelter staff training on TB.^{74,91,92}

The McKinney Homeless Assistance Act led to a national network of primary health care programs for PEH.⁹³ Community programs supported by HRSA's National Health Care for the

Homeless Program of the Bureau of Primary Health Care are appropriate partners for public health departments in preventing TB among PEH.⁹⁴ The National Health Care for the Homeless Council is a network of >10000 physicians, nurses, social workers, patients, and advocates providing support to >200 health centers and Health Care for the Homeless programs. The Council provides links to health care resources in each state for PEH.⁹⁵ Health care and supportive services are also available for people with HIV through HRSA's Ryan White Comprehensive AIDS Resources Emergency Act.⁹⁶

Discussion

The current US guidelines and recommendations relevant to TB diagnosis, treatment, and prevention call for health care providers serving PEH to (1) assess the magnitude of homelessness in their jurisdictions, (2) test PEH and shelter staff for TB and LTBI and treat those diagnosed using short treatment regimens, (3) identify TB among PEH rapidly through the use of TB diagnostics that can detect TB in 24 to 48 hours, (4) immediately report possible TB among PEH to the local public health department, (5) implement routine HIV testing of PEH who have LTBI or are being evaluated for TB disease, (6) provide temporary housing of PEH during TB treatment and link PEH to opportunities for permanent housing, and (7) conduct location-based contact investigations of infectious cases among PEH.

There is a paucity of recent studies on TB and PEH. More studies are needed to inform best practices in TB diagnosis, treatment, and prevention among PEH.

Recent improvements in health care access through the Affordable Care Act have provided greater health insurance coverage through Medicaid to people with low incomes in 39 US states.⁹⁷ These advancements enable greater access to primary health care for PEH, including LTBI testing and treatment.⁹⁸

Public Health Implications

PEH disproportionately have chronic and infectious diseases, including TB. Detection, treatment, and prevention of TB among PEH can benefit not only PEH but also society at large. Assessing homelessness among people suspected of having TB, providing housing to PEH with TB during treatment, using short 3- to 4-month regimens for LTBI, and providing directly observed therapy for TB treatment can increase successful treatment of TB among PEH. TB prevention among PEH is challenging but can be achieved.

Disclaimer

The findings and conclusions in this article are those of the authors and do not necessarily represent the official position of CDC.


Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

Suzanne M. Marks, MPH, MA  <https://orcid.org/0000-0003-3024-1940>

Thara Venkatappa, PhD, MSc  <https://orcid.org/0000-0002-1432-2719>

References

1. Homeless emergency assistance and rapid transition to housing: defining "homeless." *Fed Regist.* 2011;76(233):75994-76019. Codified at 24 CFR Parts 91, 582, and 583.
2. Bamrah S, Yelk Woodruff RS, Powell K, Ghosh S, Kammerer JS, Haddad MB. Tuberculosis among the homeless, United States, 1994-2010. *Int J Tuberc Lung Dis.* 2013;17(11):1414-1419. doi:10.5588/ijtld.13.0270
3. Self JL, McDaniel CJ, Bamrah Morris S, Silk BJ. Estimating and evaluating tuberculosis incidence rates among people experiencing homelessness, United States, 2007-2016. *Med Care.* 2021;59(suppl 2):S175-S181. doi:10.1097/MLR.0000000000001466
4. Beijer U, Wolf A, Fazel S. Prevalence of tuberculosis, hepatitis C virus, and HIV in homeless people: a systematic review and meta-analysis. *Lancet Infect Dis.* 2012;12(11):859-870. doi:10.1016/S1473-3099(12)70177-9
5. Nardell E, McInnis B, Thomas B, Weidhaas S. Exogenous reinfection with tuberculosis in a shelter for the homeless. *N Engl J Med.* 1986;315(25):1570-1575. doi:10.1056/NEJM198612183152502
6. Cole B, Nilsen DM, Will L, Etkind SC, Burgos M, Chorba T. Essential components of a public health tuberculosis prevention, control, and elimination program: recommendations of the Advisory Council for the Elimination of Tuberculosis and the National Tuberculosis Controllers Association. *MMWR Recomm Rep.* 2020;69(7):1-27. doi:10.15585/mmwr.rr6907a1
7. Powell KM, VanderEnde DS, Holland DP, et al. Outbreak of drug-resistant *Mycobacterium tuberculosis* among homeless people in Atlanta, Georgia, 2008-2015. *Public Health Rep.* 2017;132(2):231-240. doi:10.1177/0033354917694008
8. Haddad MB, Wilson TW, Ijaz K, Marks SM, Moore M. Tuberculosis and homelessness in the United States, 1994-2003. *JAMA.* 2005;293(22):2762-2766. doi:10.1001/jama.293.22.2762
9. Centers for Disease Control and Prevention. Prevention and control of tuberculosis among homeless persons. Recommendations of the Advisory Council for the Elimination of Tuberculosis. *MMWR Recomm Rep.* 1992;41(RR-5):13-23.
10. Centers for Disease Control and Prevention. Reported tuberculosis in the United States, 2020. 2021. Accessed December 14, 2021. <https://www.cdc.gov/tb/statistics/reports/2020/default.htm>
11. US Department of Housing and Urban Development, Office of Community Planning and Development. *Part 2: Estimates of Homelessness in the United States. The 2018 Annual Homeless Assessment Report (AHAR) to Congress.* September 2020. Accessed April 15, 2022. <https://www.huduser.gov/portal/sites/default/files/pdf/2018-AHAR-Part-2.pdf>

12. Henry M, de Sousa T, Roddey C, Gayen S, Bednar TJ; Abt Associates. *Part 1: Point-in-Time Estimates of Homelessness. The 2020 Annual Homeless Assessment Report (AHAR) to Congress*. US Department of Housing and Urban Development, Office of Community Planning and Development; January 2021. Accessed December 13, 2021. <https://www.huduser.gov/portal/sites/default/files/pdf/2020-AHAR-Part-1.pdf>
13. National Health Care for the Homeless Program, Bureau of Primary Health Care, Health Resources and Services Administration. Health Care for the Homeless data. Accessed April 15, 2022. <https://data.hrsa.gov/tools/data-reporting/special-populations>
14. Weinfurter P, Blumberg HM, Goldbaum G, et al. Predictors of discordant tuberculin skin test and QuantiFERON-TB Gold In-Tube results in various high-risk groups. *Int J Tuberc Lung Dis*. 2011;15(8):1056-1061. doi:10.5588/ijtld.10.0650
15. Kahwati LC, Feltner C, Halpern M, et al. *Screening for Latent Tuberculosis Infection in Adults: An Evidence Review for the US Preventive Services Task Force*. Evidence synthesis, no. 142. AHRQ pub no. 14-05212-EF-1. Agency for Health Care Research and Quality; 2016.
16. Yun LW, Reves RR, Reichler MR, et al. Outcomes of contact investigation among homeless persons with infectious tuberculosis. *Int J Tuberc Lung Dis*. 2003;7(12 suppl 3):S405-S411.
17. Centers for Disease Control and Prevention. Latent TB infection in the United States—published estimates. Accessed April 22, 2022. <https://www.cdc.gov/tb/statistics/ltbi.htm>
18. Mindra G, Wortham JM, Haddad MB, Powell KM. Tuberculosis outbreaks in the United States, 2009-2015. *Public Health Rep*. 2017;132(2):157-163. doi:10.1177/0033354916688270
19. Wortham JM, Li R, Althomsons SP, Kammerer S, Haddad MB, Powell KM. Tuberculosis genotype clusters and transmission in the U.S., 2009-2018. *Am J Prev Med*. 2021;61(2):201-208. doi:10.1016/j.amepre.2021.02.006
20. Lewinsohn DM, Leonard MK, LoBue PA, et al. Official American Thoracic Society/Infectious Diseases Society of America/Centers for Disease Control and Prevention Clinical Practice Guidelines: diagnosis of tuberculosis in adults and children. *Clin Infect Dis*. 2017;64(2):111-115. doi:10.1093/cid/ciw778
21. Centers for Disease Control and Prevention. Availability of an assay for detecting *Mycobacterium tuberculosis*, including rifampin-resistant strains, and considerations for its use—United States, 2013. *MMWR Morb Mortal Wkly Rep*. 2013;62(41):821-827.
22. Centers for Disease Control and Prevention. Updated guidelines for the use of nucleic acid amplification tests in the diagnosis of tuberculosis. *MMWR Morb Mortal Wkly Rep*. 2009;58(1):7-10.
23. Centers for Disease Control and Prevention. *Tuberculosis Laboratory Aggregate Report*. 6th ed. US Department of Health and Human Services; 2021.
24. Marks SM, Cronin W, Venkatappa T, et al. The health-system benefits and cost-effectiveness of using *Mycobacterium tuberculosis* direct nucleic acid amplification testing to diagnose tuberculosis disease in the United States. *Clin Infect Dis*. 2013;57(4):532-542. doi:10.1093/cid/cit336
25. Paquette K, Cheng MP, Kadatz MJ, Cook VJ, Chen W, Johnston JC. Chest radiography for active tuberculosis case finding in the homeless: a systematic review and meta-analysis. *Int J Tuberc Lung Dis*. 2014;18(10):1231-1236. doi:10.5588/ijtld.14.0105
26. Jensen PA, Lambert LA, Iademarco MF, Ridzon R. Guidelines for preventing the transmission of *Mycobacterium tuberculosis* in health-care settings, 2005. *MMWR Recomm Rep*. 2005;54(RR-17):1-141.
27. American Thoracic Society, Centers for Disease Control and Prevention, Infectious Diseases Society of America. American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America: controlling tuberculosis in the United States. *Am J Respir Crit Care Med*. 2005;172(9):1169-1227. doi:10.1164/rccm.2508001
28. US Preventive Services Task Force, Bibbins-Domingo K, Grossman DC, et al. Screening for latent tuberculosis infection in adults: US Preventive Services Task Force recommendation statement. *JAMA*. 2016;316(9):962-969. doi:10.1001/jama.2016.11046
29. Mazurek GH, Jereb J, Vernon A, et al. Updated guidelines for using interferon gamma release assays to detect *Mycobacterium tuberculosis* infection—United States, 2010. *MMWR Recomm Rep*. 2010;59(RR-5):1-25.
30. Collins JM, Onwubiko U, Holland DP. QuantiFERON-TB Gold versus tuberculin screening and care retention among persons experiencing homelessness: Georgia, 2015-2017. *Am J Public Health*. 2019;109(7):1028-1033. doi:10.2105/AJPH.2019.305069
31. Silva EN, Pereira A, de Araujo WN, Elias FTS. A systematic review of economic evaluations of interventions to tackle tuberculosis in homeless people. *Rev Panam Salud Publica*. 2018;42:e40. doi:10.26633/RPSP.2018.40
32. Sterling TR, Njie G, Zenner D, et al. Guidelines for the treatment of latent tuberculosis infection: recommendations from the National Tuberculosis Controllers Association and CDC, 2020. *MMWR Recomm Rep*. 2020;69(1):1-11. doi:10.15585/mmwr.rr6901a1
33. Rendleman NJ. Mandated tuberculosis screening in a community of homeless people. *Am J Prev Med*. 1999;17(2):108-113. doi:10.1016/s0749-3797(99)00052-5
34. Parriott A, Malekinejad M, Miller AP, Marks SM, Horvath H, Kahn JG. Care cascade for targeted tuberculosis testing and linkage to care in homeless populations in the United States: a meta-analysis. *BMC Public Health*. 2018;18(1):485. doi:10.1186/s12889-018-5393-x
35. Marks SM, Taylor Z, Burrows NR, Qayad MG, Miller B. Hospitalization of homeless persons with tuberculosis in the United States. *Am J Public Health*. 2000;90(3):435-438. doi:10.2105/ajph.90.3.435
36. LoBue PA, Cass R, Lobo D, Moser K, Catanzaro A. Development of housing programs to aid in the treatment of tuberculosis in homeless individuals: a pilot study. *Chest*. 1999;115(1):218-223. doi:10.1378/chest.115.1.218
37. Marks SM, Taylor Z, Miller BI. Tuberculosis prevention versus hospitalization: taxpayers save with prevention. *J Health Care Poor Underserved*. 2002;13(3):392-401. doi:10.1353/hpu.2010.0708
38. Baxter AJ, Tweed EJ, Katikireddi SV, Thomson H. Effects of Housing First approaches on health and well-being of adults who are homeless or at risk of homelessness: systematic review and meta-analysis of randomised controlled trials. *J Epidemiol*

- Community Health*. 2019;73(5):379-387. doi:10.1136/jech-2018-210981
39. Chhabra M, Spector E, Demuynck S, Wiest D, Buckley L, Shea JA. Assessing the relationship between housing and health among medically complex, chronically homeless individuals experiencing frequent hospital use in the United States. *Health Soc Care Community*. 2020;28(1):91-99. doi:10.1111/hsc.12843
 40. Riley ED, Vittinghoff E, Koss CA, et al. Housing First: unsuppressed viral load among women living with HIV in San Francisco. *AIDS Behav*. 2019;23(9):2326-2336. doi:10.1007/s10461-019-02601-w
 41. Centers for Disease Control and Prevention. TFFRS—social determinants of health: permanent supportive housing with Housing First (Housing First programs). Accessed November 8, 2022. <https://www.thecommunityguide.org/pages/tffrs-social-determinants-health-housing-first-programs.html>
 42. US Interagency Council on Homelessness. Housing First citations. Accessed March 17, 2021. <https://www.usich.gov/search/search-results/eyJyZXN1bHRfcGFnZSI6InNlYXJjaFwvc2VhcmNoLXJlc3VsdHMiLCJrZXI3b3JkeYl6IlwiSG91c2luZyBGaXJzdFwiIn0>
 43. Nahid P, Dorman SE, Alipanah N, et al. Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America clinical practice guidelines: treatment of drug-susceptible tuberculosis. *Clin Infect Dis*. 2016;63(7):e147-e195. doi:10.1093/cid/ciw376
 44. Carr W, Kurbatova E, Starks A, Goswami N, Allen L, Winston C. Interim guidance: 4-month rifapentine-moxifloxacin regimen for the treatment of drug-susceptible pulmonary tuberculosis—United States, 2022. *MMWR Morb Mortal Wkly Rep*. 2022;71(8):285-289. doi:10.15585/mmwr.mm7108a1
 45. University of Liverpool. Hep drug interactions: interaction checker. Accessed January 20, 2022. <https://www.hep-druginteractions.org>
 46. Chaulk CP, Kazandjian VA. Directly observed therapy for treatment completion of pulmonary tuberculosis: consensus statement of the Public Health Tuberculosis Guidelines Panel. *JAMA*. 1998;279(12):943-948. doi:10.1001/jama.279.12.943
 47. Davidson H, Schluger NW, Feldman PH, Valentine DP, Telzak EE, Laufer FN. The effects of increasing incentives on adherence to tuberculosis directly observed therapy. *Int J Tuberc Lung Dis*. 2000;4(9):860-865.
 48. Alipanah N, Jarlsberg L, Miller C, et al. Adherence interventions and outcomes of tuberculosis treatment: a systematic review and meta-analysis of trials and observational studies. *PLoS Med*. 2018;15(7):e1002595. doi:10.1371/journal.pmed.1002595
 49. Krueger K, Ruby D, Cooley P, et al. Videophone utilization as an alternative to directly observed therapy for tuberculosis. *Int J Tuberc Lung Dis*. 2010;14(6):779-781.
 50. Garfein RS, Doshi RP. Synchronous and asynchronous video observed therapy (VOT) for tuberculosis treatment adherence monitoring and support. *J Clin Tuberc Other Mycobact Dis*. 2019;17:100098. doi:10.1016/j.jctube.2019.100098
 51. Story A, Aldridge RW, Smith CM, et al. Smartphone-enabled video-observed versus directly observed treatment for tuberculosis: a multicentre, analyst-blinded randomised, controlled superiority trial. *Lancet*. 2019;393(10177):1216-1224. doi:10.1016/S0140-6736(18)32993-3
 52. Mirsaedi M, Farshidpour M, Banks-Tripp D, Hashmi S, Kujoth C, Schraufnagel D. Video directly observed therapy for treatment of tuberculosis is patient-oriented and cost-effective. *Eur Respir J*. 2015;46(3):871-874. doi:10.1183/09031936.00011015
 53. Beeler Asay GR, Lam CK, Stewart B, et al. Cost of tuberculosis therapy directly observed on video for health departments and patients in New York City; San Francisco, California; and Rhode Island (2017-2018). *Am J Public Health*. 2020;110(11):1696-1703. doi:10.2105/AJPH.2020.305877
 54. Moro RN, Sterling TR, Saukkonen J, et al. Factors associated with non-completion of follow-up: 33-month latent tuberculous infection treatment trial. *Int J Tuberc Lung Dis*. 2017;21(3):286-296. doi:10.5588/ijtld.16.0469
 55. Nwana N, Marks SM, Lan E, Chang AH, Holcombe M, Morris SB. Treatment of latent *Mycobacterium tuberculosis* infection with 12 once weekly directly-observed doses of isoniazid and rifampentine among persons experiencing homelessness. *PLoS One*. 2019;14(3):e0213524. doi:10.1371/journal.pone.0213524
 56. Onwubiko U, Wall K, Sales RM, Holland DP. Using directly observed therapy (DOT) for latent tuberculosis treatment—a hit or a miss? A propensity score analysis of treatment completion among 274 homeless adults in Fulton County, GA. *PLoS One*. 2019;14(6):e0218373. doi:10.1371/journal.pone.0218373
 57. Sandul AL, Nwana N, Holcombe JM, et al. High rate of treatment completion in program settings with 12-dose weekly isoniazid and rifampentine for latent *Mycobacterium tuberculosis* infection. *Clin Infect Dis*. 2017;65(7):1085-1093. doi:10.1093/cid/cix505
 58. Nyamathi A, Salem BE, Shin SS, et al. Effect of a nurse-led community health worker intervention on latent tuberculosis medication completion among homeless adults. *Nurs Res*. 2021;70(6):433-442. doi:10.1097/NNR.0000000000000545
 59. Linas BP, Wong AY, Freedberg KA, Horsburgh CR Jr. Priorities for screening and treatment of latent tuberculosis infection in the United States. *Am J Respir Crit Care Med*. 2011;184(5):590-601. doi:10.1164/rccm.201101-0181OC
 60. Jo Y, Shrestha S, Gomes I, et al. Model-based cost-effectiveness of state-level latent tuberculosis interventions in California, Florida, New York, and Texas. *Clin Infect Dis*. 2021;73(9):e3476-e3482. doi:10.1093/cid/ciaa857
 61. Centers for Disease Control and Prevention, National Center for HIV, Viral Hepatitis, STD, and TB Prevention, Division of Tuberculosis Elimination. *Latent Tuberculosis Infection: A Guide for Primary Health Care Providers*. Pub no. 22-0468. Centers for Disease Control and Prevention; 2020. Accessed March 21, 2022. <https://www.cdc.gov/tb/publications/ltbi/pdf/LTBIbooklet508.pdf>
 62. Golub JE, Mohan CI, Comstock GW, Chaisson RE. Active case finding of tuberculosis: historical perspective and future prospects. *Int J Tuberc Lung Dis*. 2005;9(11):1183-1203.
 63. Hamilton K, Tolfree R, Mytton J. A systematic review of active case-finding strategies for tuberculosis in homeless populations. *Int J Tuberc Lung Dis*. 2018;22(10):1135-1144. doi:10.5588/ijtld.17.0784
 64. Association of Public Health Laboratories. Laboratory considerations for use of cepheid Xpert MTB/RIF assay. 2013. Accessed November 8, 2022. https://www.aphl.org/AboutAPHL/publications/Documents/ID_2013Nov_Cepheid-Xpert-Fact-Sheet.pdf

65. Azevedo MJ, Conwill DE, Lawrence S, et al. Tuberculosis containment among the homeless in metropolitan Jackson, Mississippi. *J Miss State Med Assoc.* 2015;56(8):243-248.
66. National Tuberculosis Controllers Association, Centers for Disease Control and Prevention. Guidelines for the investigation of contacts of persons with infectious tuberculosis. Recommendations from the National Tuberculosis Controllers Association and CDC. *MMWR Recomm Rep.* 2005;54(RR-15):1-47.
67. Brickner PW, Vincent RL, Nardell EA, et al. Ultraviolet upper room air disinfection for tuberculosis control: an epidemiological trial. *J Health Saf Compliance Infect Control.* 2000;4(3):123-131.
68. Coffey CC, Hudnall JB, Martin SB. Improving the environmental controls at a homeless shelter to assist in reducing the probability of airborne transmission of *Mycobacterium tuberculosis*: a case study. *Indoor Built Environ.* 2009;18(2):168-182. doi:10.1177/1420326X09103008
69. Moffa M, Cronk R, Fejfar D, Dancausse S, Padilla LA, Bartram J. A systematic scoping review of environmental health conditions and hygiene behaviors in homeless shelters. *Int J Hyg Environ Health.* 2019;222(3):335-346. doi:10.1016/j.ijheh.2018.12.004
70. Nardell EA, Bucher SJ, Brickner PW, et al. Safety of upper-room ultraviolet germicidal air disinfection for room occupants: results from the Tuberculosis Ultraviolet Shelter Study. *Public Health Rep.* 2008;123(1):52-60. doi:10.1177/003335490812300108
71. Riley RL, Nardell EA. Clearing the air: the theory and application of ultraviolet air disinfection. *Am Rev Respir Dis.* 1989;139(5):1286-1294. doi:10.1164/ajrccm/139.5.1286
72. Reed NG. The history of ultraviolet germicidal irradiation for air disinfection. *Public Health Rep.* 2010;125(1):15-27. doi:10.1177/003335491012500105
73. Centers for Disease Control and Prevention, Division of Tuberculosis Elimination. Appendix B. Tuberculosis (TB) risk assessment worksheet. September 27, 2006. Accessed December 15, 2021. https://www.cdc.gov/tb/publications/guidelines/AppendixB_092706.pdf
74. Curry International Tuberculosis Center. Shelters and TB: what staff need to know. January 2013. Accessed December 15, 2021. https://www.currytbccenter.ucsf.edu/sites/default/files/shelters_and_tb_viewers_guide.pdf
75. Centers for Disease Control and Prevention. Cover your cough [poster]. Accessed November 16, 2022. https://www.cdc.gov/flu/pdf/protect/cdc_cough.pdf
76. Centers for Disease Control and Prevention, National Center for HIV, Viral Hepatitis, STD, and TB Prevention. Workshop on tuberculosis and homelessness: infection control measures in homeless shelters and other overnight facilities that provide shelter. Summary of the workshop held September 28-29, 2015. US Department of Health and Human Services, Centers for Disease Control and Prevention, Office of Infectious Diseases; 2018.
77. Corbett EL, Watt CJ, Walker N, et al. The growing burden of tuberculosis: global trends and interactions with the HIV epidemic. *Arch Intern Med.* 2003;163(9):1009-1021. doi:10.1001/archinte.163.9.1009
78. National Institutes of Health. Guidelines for the prevention and treatment of opportunistic infections in adults and adolescents with HIV. Updated February 17, 2022. Accessed May 13, 2022. <https://clinicalinfo.hiv.gov/en/guidelines/adult-and-adolescent-opportunistic-infection/mycobacterium-tuberculosis-infection-and>
79. National Coalition for the Homeless. HIV/AIDS and homelessness. July 2009. Accessed November 3, 2020. <http://www.nationalhomeless.org/factsheets/hiv.html>
80. Centers for Disease Control and Prevention. HIV prevalence estimates—United States, 2006. *MMWR Morb Mortal Wkly Rep.* 2008;57(39):1073-1076.
81. Schmit KM, Shah N, Kammerer S, Bamrah Morris S, Marks SM. Tuberculosis transmission or mortality among persons living with HIV, USA, 2011-2016. *J Racial Ethn Health Disparities.* 2020;7(5):865-873. doi:10.1007/s40615-020-00709-7
82. Beavers SF, Pascopella L, Davidow AL, et al. Tuberculosis mortality in the United States: epidemiology and prevention opportunities. *Ann Am Thorac Soc.* 2018;15(6):683-692. doi:10.1513/AnnalsATS.201705-405OC
83. Selwyn PA, Sckell BM, Alcabes P, Friedland GH, Klein RS, Schoenbaum EE. High risk of active tuberculosis in HIV-infected drug users with cutaneous anergy. *JAMA.* 1992;268(4):504-509. doi:10.1001/jama.1992.03490040080029
84. Shea KM, Kammerer JS, Winston CA, Navin TR, Horsburgh CR Jr. Estimated rate of reactivation of latent tuberculosis infection in the United States, overall and by population subgroup. *Am J Epidemiol.* 2014;179(2):216-225. doi:10.1093/aje/kwt246
85. Jones JL, Hanson DL, Dworkin MS, DeCock KM; Adult/Adolescent Spectrum of HIV Disease Group. HIV-associated tuberculosis in the era of highly active antiretroviral therapy. The Adult/Adolescent Spectrum of HIV Disease Group. *Int J Tuberc Lung Dis.* 2000;4(11):1026-1031.
86. Uthman OA, Okwundu C, Gbenga K, et al. Optimal timing of antiretroviral therapy initiation for HIV-infected adults with newly diagnosed pulmonary tuberculosis: a systematic review and meta-analysis. *Ann Intern Med.* 2015;163(1):32-39. doi:10.7326/M14-2979
87. Branson BM, Handsfield HH, Lampe MA, et al. Revised recommendations for HIV testing of adults, adolescents, and pregnant women in health-care settings. *MMWR Recomm Rep.* 2006;55(RR-14):1-17.
88. Centers for Disease Control and Prevention. TB and people experiencing homelessness. Reviewed October 19, 2021. Accessed May 13, 2022. <https://www.cdc.gov/tb/topic/populations/homelessness/default.htm>
89. Centers for Disease Control and Prevention. Find TB resources. Accessed May 20, 2022. <https://findtbrsources.cdc.gov/about>
90. Centers for Disease Control and Prevention. TB Centers of Excellence for training, education, and medical consultation. Accessed February 11, 2022. https://www.cdc.gov/tb/education/tb_coe/default.htm?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Ftb%2Feducation%2Ftrmc%2Fdefault.htm
91. Curry International Tuberculosis Center. Homelessness and TB toolkit. Accessed December 15, 2021. https://www.currytbccenter.ucsf.edu/sites/default/files/product_tools/homelessnessandttoolkit/index.html
92. Southeastern National Tuberculosis Center. Centers of Excellence (COE) TB training and education products. Accessed December 15, 2021. <https://sntc.medicine.ufl.edu/rtmccproducts.aspx>
93. The McKinney Homeless Assistance Act. Pub L No 100-77 (1982). 101 Stat 482. Accessed March 21, 2022. <https://www.congress.gov/100/statute/STATUTE-101/STATUTE-101-Pg482.pdf>

94. Health Resources and Services Administration. HRSA Health Center Program. Accessed February 11, 2022. <https://bphc.hrsa.gov>
95. National Health Care for the Homeless Council. HCH grantee directory. Accessed November 8, 2022. <https://nhhc.org/grantee-directory>
96. Health Resources and Services Administration. About the Ryan White HIV/AIDS Program. Accessed November 8, 2022. <https://ryanwhite.hrsa.gov/resources/provider-resources>
97. Kaiser Family Foundation. Status of state Medicaid expansion decisions: interactive map. Accessed February 10, 2022. <https://www.kff.medicicaid/issue-brief/status-of-state-medic-aid-expansion-decisions-interactive-map>
98. Gupta V, Sugg N, Butners M, Allen-White G, Molnar A. Tuberculosis among the homeless—preventing another outbreak through community action. *N Engl J Med.* 2015;372(16):1483-1485. doi:10.1056/NEJMp1501316