








RESEARCH ARTICLE

REVISED **Development of an online intervention for the Rehabilitation Exercise and psychological support After covid-19 Infection (REGAIN) trial [version 2; peer review: 2 approved]**

Stuart Ennis^{1,2}, Peter Heine^{1,2}, Harbinder Sandhu¹, Bart Sheehan³, Joyce Yeung ¹, David McWilliams ⁴, Christina Jones⁵, Charles Abraham⁶, Martin Underwood¹, Julie Bruce ¹, Kate Seers ⁷, Gordon McGregor ^{1,2,8}

¹Clinical Trials Unit, University of Warwick, Coventry, UK

²Cardiopulmonary Rehabilitation, University Hospitals Coventry & Warwickshire NHS Trust, Coventry, UK

³The Oxford Psychological Medicine Centre, Oxford University Hospitals NHS Foundation Trust, Oxford, UK

⁴Centre for Care Excellence, Coventry University, Coventry, UK

⁵ICUsteps Peer Support Charity, ICUsteps Peer Support Charity, London, UK

⁶School of Psychology, Deakin University, Melbourne, Australia

⁷Health Sciences, University of Warwick, Coventry, UK

⁸Centre for Sport, Exercise and Life Sciences, Coventry University, Coventry, UK

V2 **First published:** 23 Feb 2023, 3:10
<https://doi.org/10.3310/nihropenres.13371.1>

Latest published: 14 Jul 2023, 3:10
<https://doi.org/10.3310/nihropenres.13371.2>

Abstract

Background







Up to half of people hospitalised with COVID-19 report diverse and persistent symptoms affecting quality of life for months and sometimes years after discharge (long-COVID). We describe the development of an online group exercise and behavioural support intervention for people who continue to experience such physical and/or emotional health problems more than three months after hospital discharge.

Methods


Intervention development was informed by the Medical Research Council framework for complex interventions. Our multidisciplinary team of academics, clinicians, and people with long-COVID, had collective expertise in the development and testing of complex interventions. We integrated a bio-psycho-social model of care

Open Peer Review

Approval Status  

	1	2
version 2 (revision) 14 Jul 2023	 view	 view
		
version 1 23 Feb 2023	 view	 view

1. **Garry A. Tew** , York St John University, York, UK

2. **Stefan Tino Kulnik** , Ludwig Boltzmann Gesellschaft, Salzburg, Austria

Any reports and responses or comments on the article can be found at the end of the article.

drawing on rehabilitation literature for long-term health conditions and experiences from our pre-pilot study. Multiple stakeholder meetings were held to refine the intervention which was designed to be deliverable within the UK National Health Service. We adhere to TIDieR guidance for transparent and explicit reporting of telehealth interventions.

Results

The final REGAIN online exercise and behavioural support intervention consisted of an initial 1:1 consultation with a trained practitioner, followed by eight online group exercise, and six group support, sessions delivered over eight weeks. Participants could also access an online library of on-demand exercise and support videos.

Conclusions

The final REGAIN intervention, combining exercise and behavioural support, is fully manualised with clear pathways to delivery and implementation. It is currently being tested in a randomised controlled trial. The intervention, developed with extensive patient and stakeholder engagement, could be incorporated into existing NHS rehabilitation programmes, should it prove to be clinically and cost-effective for people with long-COVID.

Trial registration

International Standard Randomised Controlled Trial Number (ISRCTN) 11466448: Rehabilitation exercise and psychological support after COVID-19 infection: REGAIN.

Plain English summary

Long-COVID has many debilitating symptoms, such as breathlessness, muscle weakness and fatigue, which significantly affect peoples' physical and mental health and quality of life. Rehabilitation programmes can help people improve their quality of life in other medical conditions with similar symptoms. We developed a programme of physical and mental health rehabilitation, delivered online, specifically to support people with ongoing long-COVID symptoms more than three months after hospital discharge. The programme was developed by people with long-COVID along with clinicians and researchers. The programme described in this article is now being tested in a large research trial to see if it can help people with long-COVID.

Keywords

rehabilitation, long-COVID, physical activity, online support, intervention development, physical activity

Corresponding author: Gordon McGregor (gordon.mcgregor@warwick.ac.uk)

Author roles: **Ennis S:** Investigation, Methodology, Project Administration, Resources, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Heine P:** Data Curation, Investigation, Methodology, Project Administration, Resources, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Sandhu H:** Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Sheehan B:** Conceptualization, Funding Acquisition, Investigation, Methodology, Resources, Supervision, Validation, Writing – Review & Editing; **Yeung J:** Conceptualization, Funding Acquisition, Methodology, Resources, Supervision, Validation, Writing – Review & Editing; **McWilliams D:** Conceptualization, Funding Acquisition, Investigation, Methodology, Validation, Writing – Review & Editing; **Jones C:** Conceptualization, Data Curation, Funding Acquisition, Investigation, Methodology, Resources, Validation, Writing – Review & Editing; **Abraham C:** Conceptualization, Funding Acquisition, Writing – Review & Editing; **Underwood M:** Conceptualization, Data Curation, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Visualization, Writing – Review & Editing; **Bruce J:** Conceptualization, Data Curation, Funding Acquisition, Investigation, Methodology, Project Administration, Resources, Supervision, Validation, Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Seers K:** Conceptualization, Funding Acquisition, Investigation, Methodology, Resources, Supervision, Validation, Visualization, Writing – Review & Editing; **McGregor G:** Conceptualization, Funding Acquisition, Investigation, Methodology, Project Administration, Supervision, Validation, Writing – Original Draft Preparation, Writing – Review & Editing

Competing interests: JB, MU, GM, HS, BS, JY, DW are chief or co-investigators on multiple research grants funded by NIHR and other funding bodies. KS previously sat on the NIHR HS&RD funding board for 8 years. SE and GM are directors of Atrium Health Ltd, a provider of rehabilitation services for the National Health Service. HS is director of Health Psychology Services Ltd, providing psychological services for a range of health-related conditions. MU is chief investigator or co-investigator grants from Arthritis Research UK and is a co-investigator on grants funded by the Australian NHMRC and Norwegian MRC. He was an NIHR Senior Investigator until March 2021. He has received travel expenses for speaking at conferences from the professional organisations hosting the conferences. He is a director and shareholder of Clinvivo Ltd that provides electronic data collection for health services research. He is part of an academic partnership with Serco Ltd, funded by the European Social Fund, related to return to work initiatives. MU and JB receive some salary support from University Hospitals Coventry and Warwickshire NHS Trust. MU is a co-investigator on two current and one completed NIHR funded studies that are, or have had, additional support from Stryker Ltd. Until March 2020 he was an editor of the NIHR journal series, and a member of the NIHR Journal Editors Group, for which he received a fee. JB is a co-investigator on one NIHR funded study that receives support from Stryker Ltd.

Grant information: This research was funded by the National Institute for Health Research Health Technology Assessment Programme: COVID-19 Recovery & Learning (NIHR132046).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2023 Ennis S *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Ennis S, Heine P, Sandhu H *et al.* **Development of an online intervention for the Rehabilitation Exercise and psycholoGical support After covid-19 InfectioN (REGAIN) trial [version 2; peer review: 2 approved]** NIHR Open Research 2023, 3:10 <https://doi.org/10.3310/nihropenres.13371.2>

First published: 23 Feb 2023, 3:10 <https://doi.org/10.3310/nihropenres.13371.1>

REVISED Amendments from Version 1

- We have amended the text throughout to improve consistency and readability.
- We have amended Figure 2 for clarity and added Figure 3 (logic model) and appendix 1 (sample exercise session template).
- We have added co-author Professor Charles Abraham, School of Psychology, Deakin University, Melbourne, Australia. Conceptualization, funding acquisition, editing and review

Any further responses from the reviewers can be found at the end of the article

Plain English summary

Long-COVID has many debilitating symptoms, such as breathlessness, muscle weakness and fatigue, which significantly affect peoples' physical and mental health and quality of life. Rehabilitation programmes can help people improve their quality of life in other medical conditions with similar symptoms. We developed a programme of physical and mental health rehabilitation, delivered online, specifically to support people with ongoing long-COVID symptoms more than three months after hospital discharge. The programme was developed by people with long-COVID along with clinicians and researchers. The programme described in this article is now being tested in a large research trial to see if it can help people with long-COVID.

Introduction

'Long-COVID' is the term commonly used to describe symptoms which continue or develop after the initial acute infection with COVID-19 has resolved. It includes ongoing symptomatic COVID-19 (from 4 to 12 weeks) and post-COVID-19 syndrome (12 weeks or more), where symptoms cannot be explained by an alternative diagnosis¹. As of January 2023, there were approximately 2.1 million people (3.3% of the population) self-reporting long-COVID in the UK¹. Of those, 57% reported symptoms lasting at least one year, and 76% reported symptoms that adversely affected their daily activities¹. The most common symptoms were fatigue (71%), difficulty concentrating ('brain fog') (49%), breathlessness (47%), and muscle aches (46%)¹. Symptoms vary in severity within and between individuals, and commonly remit and relapse over time leading to considerable distress and a negative impact on mental wellbeing². People often feel dismissed by medical professionals, and abandoned after hospital discharge, and commonly seek validation and support elsewhere via peer behavioural support groups and social media^{3,4}. Long-COVID can have devastating psychological, physical and cognitive consequences that disrupt lives and livelihoods⁵.

To address long-COVID, an NIHR review⁶ suggested the need for a holistic, integrated treatment approach rather than symptom by symptom management. This was supported by UK COVID-19 guidelines which recommended pragmatic treatment for psychological, emotional, and physical health⁷. However, for people with long-COVID, direct face-to-face contact with medical professionals in general, and specifically specialist post-COVID-19 clinics, has been difficult to access,

and provision is highly variable. One method of countering these problems is to provide resources online.

The primary digital resource for people living with long-COVID in the UK is 'Your COVID Recovery'⁸ produced by the National Health Service (NHS). 'Your COVID Recovery' is a comprehensive NHS online resource providing self-management symptom advice and support. However, this resource lacks the social connection that patients often desire, is necessarily generic, and may not be versatile enough to meet the needs of everyone with long-COVID. Furthermore, awareness of the website is not widespread, and evaluation of patient experience of the intervention has not yet been published.

Currently, few studies have examined rehabilitation for people with long-COVID, particularly in an online setting, although there are reports that multidisciplinary rehabilitation can improve symptoms in long-COVID⁹ and enhance quality of life for people living with other long-term health conditions with similar symptom profiles, such as Chronic Obstructive Pulmonary Disease (COPD)¹⁰ and Severe Acute Respiratory Syndrome (SARS)¹¹. Therefore, a large-scale trial to test such an intervention suitable for online, remote delivery, is required.

This manuscript describes the design and development of an online exercise and behavioural support intervention for people with long-COVID. The REGAIN randomised controlled trial (RCT) was funded by the National Institute of Health Research COVID-19 Recovery and Learning Programme (NIHR132046) to evaluate the clinical and cost-effectiveness of an online exercise and psychological support intervention for people hospitalised with COVID-19 who still reported symptoms more than three months after hospital discharge¹².

Methods

Ethical approval

Ethical approval was received from the East of England-Cambridge South Research Ethics Committee (REC: 20/EE/0235).


Written informed consent for publication of the participants details and/or their images was obtained from the participants.

Overview of the development process

We followed the Medical Research Council framework¹³ for design of complex interventions (Figure 1). Our manuscript adheres to the recent Template for Intervention Development and Replication (TIDieR) for 'telehealth' complex interventions¹⁴. Work was conducted between August 2020 and January 2021.

Stage 1: Exploratory, background research (patient experience, evidence base, expert opinion, current practice)

A search of bibliographic databases (MEDLINE, PsycInfo and CINAHL) identified literature relating to behavioural or self-management needs for people living with long-COVID. Due to the relative paucity of long-COVID research, we also searched for literature on chronic fatigue syndrome (CFS), chronic obstructive pulmonary disease (COPD), and the 2002–2004 SARS epidemic. Collectively these conditions exhibit



PPI, Patient & Public Involvement; RCT, randomised control trial; SARS, severe acute respiratory syndrome; CFS - chronic fatigue syndrome; ME, Myalgic encephalomyelitis; ICU, intensive care unit; NHS, National Health Service.

Figure 1. Intervention development process.

several of the most common and debilitating symptoms of long-COVID (e.g. fatigue, lethargy, breathlessness, muscle weakness, brain fog).

Titles and abstracts of papers were assessed for relevance, prioritising recent systematic reviews of needs and interventions in long-COVID, CFS, COPD and SARS. Articles were examined to ensure that the needs identified by our stakeholder groups were addressed by research evidence where possible. Reports of interventions for conditions with similar symptoms helped to identify core components that may be effective for people living with long-COVID.

The literature review revealed limited evidence regarding the efficacy of rehabilitation interventions for COVID-19 survivors⁹.

One RCT (n=72) of respiratory rehabilitation reported improved respiratory function, quality of life (QoL) and anxiety in elderly people with COVID-19¹⁵, but there was little other data available. In other similar conditions, a meta-analysis of 65 RCTs involving 3822 participants reported that pulmonary rehabilitation programmes, combining exercise with various psychological support and education components, were beneficial in improving health-related quality of life (HR-QoL) and exercise capacity in people with COPD¹⁰. Similarly, a meta-analysis of eight RCTs (n=1518) concluded that people with CFS may generally benefit and feel less fatigued following exercise therapy, comparable to the effectiveness of cognitive behavioural therapy (CBT), with no evidence to suggest that exercise therapy may worsen outcomes¹⁶. Exercise training was also reported to be effective in improving cardiorespiratory

and musculoskeletal fitness in people recovering from SARS, although there was no reported effect on HR-QoL¹¹. Other studies highlighted the persistent mental and physical abnormalities in SARS survivors up to two years post-infection¹⁷. Exercise therapy alone, therefore, is likely insufficient to optimise recovery for people with long-COVID.

We concluded that to combat the multiple long-term physical and mental health consequences of COVID-19, a combined physical and psychological rehabilitation intervention was required. In order to make the programme as widely accessible as possible during COVID-19 'lock-downs' and reduced access to healthcare, the intervention was developed to be deliverable online.

Stage 2: Production of draft REGAIN intervention

To holistically appraise the needs of people with long-term COVID-19 symptoms following hospitalisation, a core intervention development team was formed. The team (n=22) had specialism and extensive experience in clinical academia, physiotherapy, psychiatry, health psychology, clinical exercise physiology, critical care medicine, complex intervention development and implementation, and online rehabilitation and behavioural support. Six people (male n=3, female n=3; age 33–62 years) with symptoms of long-COVID (3–9 months duration) were also involved throughout every stage of the development of the REGAIN intervention (patient and public involvement (PPI) group). Online meetings facilitated discussions informed by evidence from Stage 1 development. The team met frequently with the objective of identifying and formulating the key components of a complex intervention.

The research team identified key themes and priorities from the literature to form the outline for a series of online discussions held between August and October 2020 (Table 1). Consensus responses relating to core components for inclusion in the exercise and behavioural support intervention were collated into the following categories: online group exercise classes (frequency, intensity, time and type principles), concerns about post exertional malaise/chronic fatigue, exercise programme progression, behavioural support materials, and the feasibility of delivering the intervention online in a NHS setting to participants with diverse symptomology and varying degrees of information technology (IT) literacy. Findings from the stakeholder meetings are summarised in Table 1. A framework of a draft intervention was finalised for pre-pilot testing.

Post exertional malaise

During the development phase, the trial team were contacted by several organisations representing the Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS) community with concerns relating to the potential harms of unsupervised vigorous exercise for people with long-COVID, in particular the risk of post-exertional malaise¹⁸. We recognised the lived experience and expertise within these groups and sought to ensure that REGAIN practitioners were appropriately trained in risks relating to post-exertional malaise. Furthermore, early identification screening measures for post-exertional

malaise were incorporated into the intervention protocols, particularly during the initial one-to-one consultation and every subsequent exercise session. Practitioners were encouraged to monitor the progress of participants during and after exercise sessions in order to ensure that exercise intensity and volume were within individuals' capabilities and did not cause undue persistent fatigue or other problems.

Stage 3: Pre-pilot testing of the REGAIN intervention

To test acceptability and deliverability, the draft REGAIN intervention was feasibility tested in a pre-pilot study with eight people living with long-COVID. The online exercise and behavioural support sessions were delivered from a community NHS exercise rehabilitation facility (Atrium Health, Centre for Exercise and Health, Coventry). Feedback was gathered on the acceptability and practicality of the intervention from eight participants as well as the healthcare practitioners delivering the intervention. This included comments on the written and online trial materials (e.g. practitioner/participant manuals) and the online platforms used to deliver the intervention. This helped the trial team further refine and develop the intervention before proceeding to the main trial. The key findings from the pre-pilot stage are described in Table 2.

Stage 4: Refinement of the REGAIN intervention

On-demand exercise and educational videos. The intervention team, with the support of our patient partners, developed 14 'on-demand' exercise videos which used a similar format to the live online exercise sessions and were designed to account for all abilities. Lower intensity videos with postural stability and balance exercises were included for people with more pronounced levels of fatigue or disability. Some videos were themed (e.g., shopping, housework) to reintroduce activities of daily living and to add some additional context for those unaccustomed to exercise videos. We also included videos of other exercise modalities (e.g., Yoga, Pilates, and high intensity interval training (HIIT)), which had already been produced by the online physical activity provider (BeamFeelGood) used to deliver the intervention. Finally, five videos providing advice and education on lung function and care, including breathing techniques/exercises (Fit for Surgery team, University Hospital Birmingham) and six 'mindfulness' videos (Dr Gail Davies) were produced to give participants additional online resources. All videos were hosted on the online platform via a trial-specific protected user area. Participants were encouraged to access these exercise videos, unsupervised, between weekly group sessions and to progress to performing up to two additional exercise sessions per week as tolerated.

Practitioner-led online group exercise sessions

A series of 12 templates were designed for the online group exercise sessions. They ranged in modality from low intensity chair-based activity to higher intensity whole body exercise. These formed the basis of all online group exercise sessions and ensured a level of exercise programme standardisation throughout the trial. All the exercises within each template required little or no equipment and were easily modifiable to account for different ability levels within the group. Depending on the

Table 1. Key findings from intervention co-development meetings.

Component	Outcome/consensus
Online group exercise classes	<p>All agreed that one supervised exercise session per week of 20-60 minutes duration, dependent on fitness level, should be tested for acceptability during the pre-pilot phase. Additional pre-recorded exercise classes planned to provide “on- demand” video material for participants to follow unsupervised at home, up to 2-3 per week.</p> <p>Videos would include a variety of exercises to appeal to all, including: Yoga, Pilates, chair-based and breathing exercise videos. Concerns were expressed by patients that some exercises may not be tolerated by more debilitated patients. The need for robust safety/emergency procedures was also highlighted.</p> <p>People with long-COVID symptoms expressed a loss of confidence about being physically active and anxiety about relapse; also concern about over-exertion. An emphasis on fun over intensity was agreed for first few weeks to promote confidence and adherence.</p> <p>All agreed that light-moderate aerobic exercise (40–70% Heart Rate Reserve (HRR), rating of perceived exertion (RPE) - 11-14) was realistic for most participants after familiarisation. Gentle mobility, coordination, and balance exercises would be needed after long sedentary periods for some participants. Clinical exercise physiologists should test acceptability during the pre-pilot phase. All Long Covid patients agreed that IT literacy might be an obstacle to accessing sessions for older participants.</p>
Concern about post-exertional malaise (PEM)	<p>Members of trial team met with ME groups to discuss the dangers of post-exertional malaise (PEM) in the long-COVID population. The word exercise seemed to have negative connotations, assumed to only mean high intensity/exertion levels which could be detrimental to recovery. Use of term physical activity would be better received. Groups were reassured that (a) intensity would be low-moderate and at the comfort level of the individual and (b) that the risk of PEM would be minimised by closely monitoring participants before, during, and after each session, as well as assessing risk of PEM during the initial 1:1 consultation (c) a pre-determined ‘graded’ approach adopted in previous trials would not be applied to this population. Specific information was provided to participating clinicians during the training sessions highlighting the need to be aware of PEM and how to address it, and a section on PEM was incorporated into the Practitioner Manual provided to all clinicians taking part in REGAIN.</p>
Exercise progression	<p>It was agreed that exercise physiologists should produce 8-12 exercise session templates of different levels of intensity for standardised delivery across the trial.</p> <p>Classes would follow a circuit format with active recovery between exercises. A point was made that quite conceivably a group containing very low and very high ability participants was likely as practitioners would not always be able to arrange groups based on ability level. Therefore, a great degree of versatility was needed with exercise class design and delivery. People with long-COVID symptoms requested ample rest periods and water breaks. Following warm-up and mobility, duration of exercise should be progressed primarily as tolerated, for maximal therapeutic benefit. Researchers requested that exercise physiologists collect ratings of perceived exertion (RPE) to assess intervention fidelity.</p>
Behavioural support materials	<p>Mind-body disconnect was mentioned by people with long-COVID symptoms, so they felt it would be really useful to have an integrated intervention. They felt strongly that long Covid wasn't simply physical symptoms of fatigue and breathlessness, but also how this affected their thoughts and feelings.</p> <p>Many topics were deemed useful by patients, especially ‘challenging unhelpful thoughts’ and ‘anxiety management’. Pacing of activities was considered crucial.</p> <p>Talking to the trained practitioner before and after group exercise sessions in a supportive environment was thought to be important to ensure continued adherence and group cohesion. This would also promote peer support.</p> <p>Breathing was difficult for some and so being able to stop, take a breath and relax was important. Reinfection was also major source of anxiety/worry.</p> <p>The participant workbook was thought by patients to be a long document, but all content was relevant as this could be read in their own time. The written targets were welcomed (e.g. writing down goals), and making notes during the session was deemed useful. Patients preferred a paper copy to an online PDF. Patients suggested colour-coded sections of the workbook to make it less daunting, and tabs to make easy to find sections for each weekly session. Suggested workbook format included adding the aims of behavioural support, with short clear statements at the beginning of each section of the workbook.</p>
Feasibility of delivery within NHS setting	<p>Exercise physiologists and health psychologists highlighted potential obstacles to online delivery. These included: the need for a second practitioner, or ‘co-pilot’, for safety reasons, IT connectivity, participant retention, risk of busy classes, accurate intensity monitoring, and adequate behavioural intervention training for practitioners less experienced with these techniques.</p> <p>Actions were agreed to mitigate these issues: (1) deliver appropriate training for practitioners led by health psychologist and an experienced clinical research fellow; (2) provide comprehensive reference manuals for participants and practitioners to follow; and (3) ensure sufficient time was spent with each participant during the 1:1 consultation to familiarise them with the online platform and IT requirements.</p>

Table 2. Main findings from pre-pilot study testing REGAIN intervention delivery.

Component	Finding(s)	Implications for REGAIN Intervention
IT/online issues	<p>Participants not placing device in adequate position for practitioner to view during exercises. Some older participants had difficulty navigating Zoom/MS Teams, enabling microphone/camera, and accessing link to session.</p> <p>Some participants accessed the session through their Smartphone which limited functionality (couldn't see the rest of group).</p> <p>One person held smartphone throughout, so very jerky & too close; it was not possible for the practitioner to safely monitor exertion levels or technique.</p>	<p>During 1:1 consultation there is a need to go through set up - test the device and where the camera will be set up. etc. Log in to zoom and do test session. Practitioners and 'co-pilot' to log in early to group sessions as considerable time needed to let everyone into session.</p> <p>Amend workbook/telephone screening calls to indicate that a smartphone not good enough unless it can be 'cast' to a TV. Laptops & tablets preferred.</p>
Progression	<p>Participants found the progression more acceptable once they had gained confidence in the sessions and felt included in short 'debrief' post exercise with rest of group.</p>	<p>Class templates amended so that Practitioners could alter the duration of exercises and recovery periods to suit each group. Participants to be asked how they felt the day after each session, as well as post session, to guard against post exertional fatigue.</p>
Safety	<p>Participants would frequently talk or make noise throughout, inadvertently interrupting practitioners' instructions.</p> <p>Occasionally participants would disappear from view- going to toilet/answering door- practitioners were unsure if the patient was unwell.</p>	<p>All participants should be instructed to mute during exercise. Hand signals required to indicate whether easy, ok, too hard and another to say I am unwell/need to stop/etc.</p> <p>Set ground rules and instructions at start of each class requesting participants to signal before they leave the screen.</p>
Very debilitated participants	<p>One participant was chair bound in the same session as another with a relatively high level of fitness. Practitioners needed assistance from 'co-pilots' (via chat function) to adequately monitor all participants.</p>	<p>Need alternative chair-based option for all exercises in the session templates. Priority should be given to enjoyment while participants gain confidence. Frequent water breaks should be provided and the emphasis placed on the participant to take responsibility for their own exertion level</p>
One to one consultation	<p>Several participants were identified as having case level mental health disorder on anxiety/depression questionnaire. More discussion during the consultations revealed a higher level of emotional support needed than initially planned.</p> <p>Some participants spent longer overcoming IT issues than they did talking about their long-COVID.</p>	<p>Consultations should be at least an hour in order to introduce website and sign-up process, triage participant for exercise classes, and go through IT accessibility/optimal camera position etc. Regular meetings with psychologist planned throughout trial for ongoing advice/support for practitioners during trial.</p>
Behavioural education	<p>Participants required more support coping with anxiety/avoidance around social interaction and not just physical activity.</p> <p>Goal setting session was difficult as most participants experienced frequent setbacks and had unknown timescale of recovery.</p> <p>Some participants dominated the support sessions and others barely spoke. Practitioners were unsure how long to 'let it go' before interrupting.</p> <p>Practitioners were also unsure about PTSD issues that some participants may be facing, and giving advice outside of their competences.</p>	<p>Additional material included in the participant workbook relating to negative thought patterns.</p> <p>Practitioners should approach long COVID differently to traditional rehabilitation for long term conditions with no expectation of linear improvement each week</p> <p>Health psychologists to do additional training with practitioners regarding managing groups.</p>

group, different exercises were swapped in or out of the session template to cater for the specific needs. Generally, the templates included 7–8 exercises to be completed 2–3 times (i.e., 2–3 circuits). The warm-up and cool down lasted 5–10 minutes each.

Results

Overview of REGAIN intervention

The final REGAIN exercise and behavioural support intervention consisted of four main components delivered by clinical exercise physiologists or physiotherapists (REGAIN practitioners).

The final intervention structure and accompanying logic model are presented in Figure 2) and Figure 3 respectively.

1. Online one-to one video consultation and participant workbook
2. Eight-week exercise programme: practitioner-led live online group sessions once per week
3. Six practitioner-led live online group behavioural support sessions (during weeks 1–5 & week 8).
4. On-demand exercise sessions.

Online one-to-one consultation

The one-to one consultation with a REGAIN practitioner included several components and was designed to triage the participant for safe exercise, and familiarise them with the IT requirements:

- Clinical history including COVID-19 hospitalisation episode(s), resulting problems and other relevant medical history and co-morbidities

- Potential barriers to participation
- Demonstration of, and sign-up processes, for the online platform.
- Discussion regarding home (e.g. space, lighting) and device set-up for live group exercise and support sessions

Eight-week exercise programme

The online exercise rehabilitation programme delivered by REGAIN practitioners consisted of up to 30 minutes of light to moderate intensity exercise two to three times per week for eight weeks. Practitioners individualised group sessions to accommodate varying abilities, with progressive multi-modality exercise at a manageable intensity, monitored with breathlessness and perceived exertion scales. Participants were encouraged to attend one live online group exercise session each week, and additionally access the online, pre-recorded exercise videos once or twice each week. This equipment-free exercise programme aimed to improve cardiovascular fitness, strength, balance, co-ordination, and confidence. Sessions were undertaken




Figure 2. Final format and components of the ‘Rehabilitation Exercise and psycholoGical support After covid-19 InfectioN’ (REGAIN) intervention.




Figure 3. Logic model for the REGAIN intervention.

in discrete groups where participants remained in the same group over the eight-week programme. An example of an intermediate level exercise session is presented in Table 3.

Online group support sessions

Emerging evidence highlighted the importance of the biopsychosocial model^{19,20} in understanding the interrelationships among risk factors and multidimensional clinical and psychosocial COVID-19 outcomes. The support sessions were based on behaviour change theory (Michie’s COM-B model: mapping key processes and functions to enhance Capability, Opportunity, and Motivation)²¹, self-efficacy²², motivational interviewing²³, and group-based learning²⁴. These theoretical principles were used to inform the psychosocial content, structure and delivery of the sessions in combination with the British Psychological Guidance for management of long-COVID which was available at the time²⁵.

To enhance the psychological capability of participants (as per the COM-B model), support sessions, delivered by REGAIN practitioners, were designed to increase knowledge and understanding about COVID-19 and its impact on daily living. Feedback from patient partners highlighted the need for practitioners to address misinformation from various sources (e.g., friends/relatives, news, social media etc.) which often caused more anxiety and distress. Reduced processing and retention of information due to brain fog and poor

concentration also informed the design and content of the support sessions. Resources and activities could be paused at any time and revisited by participants via the provided study materials (PowerPoint presentations, written summaries of key topics, video clips and further reading). Motivation was incorporated into support sessions through exploration of self-identity and the meaning of COVID-19 to the participants and those around them (e.g., family). By discussing COVID-19 experiences in a group setting, participants could see that others were having the same issues and could share experiences and coping strategies. The aim was for participants to feel validated that their symptoms were not only real, but that they were not alone in experiencing them.

Opportunity was the third core principle incorporated into the support sessions: participants were permitted time to engage in sessions in a supportive environment. Group based support sessions were chosen because they can promote behaviour change through shared values and norms, influencing beliefs, and encouraging common motivations and behavioural patterning. We stressed the importance of engaging with all elements of the intervention. To help attenuate fatigue, group support sessions were no longer than 60 minutes. The six group support sessions below are further summarised in Table 4:

Session 1: Introduction, expectations, motivation and goal setting

Table 3. Example template for a live online REGAIN exercise session.

Session Plan – REGAIN session 5		Challenge:	Intermediate	Circuits:	2
Equipment:	chair (optional: weights of some kind e.g. water bottles, tins, dumbbells etc)				
Comments:	All exercises below are interspersed with active recovery (AR), e.g. marching, side-steps, heel digs etc				
	Exercise Name	Reps and/or Time	Teaching Points (Key safety points)	Progression	Alternatives (where appropriate)
1	Squat + one arm reach	45s on/30s AR	<input type="checkbox"/> Feet hip width apart <input type="checkbox"/> Back straight, head up, extend arm up as you stand up	<input type="checkbox"/> Increase ROM <input type="checkbox"/> Add weight	<input type="checkbox"/> Decrease ROM
2	Lateral raise super slow	45s on/30s AR	<input type="checkbox"/> Hold abdominals tight <input type="checkbox"/> Slow controlled movement <input type="checkbox"/> Arms up to shoulders	<input type="checkbox"/> Add weight <input type="checkbox"/> Pause at top	<input type="checkbox"/> Front raise <input type="checkbox"/> Seated <input type="checkbox"/> Alternate arms
3	Side lunge with super slow bicep curl	45s on/30s AR	<input type="checkbox"/> Toe down first to side <input type="checkbox"/> Slight bend in knee <input type="checkbox"/> When in lunge squeeze arms into curl	<input type="checkbox"/> Deeper lunge <input type="checkbox"/> Add Weight <input type="checkbox"/> Pause at top	<input type="checkbox"/> External support <input type="checkbox"/> Shallow lunge
4	High pull	45s on/30s AR	<input type="checkbox"/> Feet hip width apart <input type="checkbox"/> Slight bend in knees back straight <input type="checkbox"/> Stand back up bringing hands from knees to chest in 1 movt	<input type="checkbox"/> Add weight <input type="checkbox"/> increase ROM	<input type="checkbox"/> Decrease ROM <input type="checkbox"/> upright row w/out legs
5	Wide stance knee to opp. toe touch	45s on/30s AR	<input type="checkbox"/> Feet flat on floor, shoulder width apart <input type="checkbox"/> Back straight, head up <input type="checkbox"/> Knees soft <input type="checkbox"/> Rotate to reach down to opposing knee	<input type="checkbox"/> Increase ROM to ankle <input type="checkbox"/> wider stance	<input type="checkbox"/> Decrease ROM <input type="checkbox"/> seated
6	Standing overhead press	45s on/30s AR	<input type="checkbox"/> Feet hip width apart <input type="checkbox"/> Slight pelvic tilt to protect back, Abdominals tight <input type="checkbox"/> Extend both arms up to meet in arc overhead	<input type="checkbox"/> Add weight <input type="checkbox"/> Pause at top	<input type="checkbox"/> Alternate arms <input type="checkbox"/> Seated <input type="checkbox"/> Frontal raise
7	Seated leg lift over object	45s on/30s AR	<input type="checkbox"/> Sit towards edge of chair <input type="checkbox"/> Lean back into chair <input type="checkbox"/> Lift straight leg one at a time over water bottle	<input type="checkbox"/> both legs together	<input type="checkbox"/> bent leg <input type="checkbox"/> hold chair for balance

Session 2: Fear avoidance and pacing

Session 3: Recovery and sleep, sleep management strategies

Session 4: Management of emotions (perceived stigma, mood/unhelpful thoughts)

Session 5: Understanding stress and anxiety, and management strategies

Session 6: Managing setbacks and long-term behaviour change and future goals.

Behavioural support videos

Feedback from our patient partners and intervention practitioners suggested potential benefit of having a short video or conversation where topics could be explored and viewed during

the group sessions. We developed short introductory videos for each of the six group support topics. The videos featured discussions between the trial health psychologist and patient partners, filmed and edited by a professional production company. Practitioners played the relevant video at the start of each group support session with the intention that participants may recognise patterns of behaviour in themselves, thus stimulating support group discussion.

Intervention fidelity: practitioner training and quality control

To ensure standardisation of intervention delivery, all REGAIN practitioners completed a full day of training. This included practical and theoretical components, delivered by a clinical exercise research fellow and health psychologist. Guidance was provided on delivery of behavioural education sessions,

Table 4. Summary of REGAIN intervention components and theoretical underpinnings.

Intervention component	Aims	Theoretical Framework(s)	Behaviour Change Taxonomy
One to one consultation	To gather relevant information (clinical history) To assess motivations and potential barriers to engagement with the REGAIN Intervention To demonstrate and introduce the online platform To offer an opportunity to answer any questions related to the study or intervention and discuss home environment for engagement with the sessions	COM-B Self-efficacy Communication	Goal setting (behaviour) Restructuring the physical environment Instruction on how to perform a behaviour (online platform) Verbal persuasion about capability (if appropriate)
Support session 1 Introduction, expectations, motivation and goal setting	To introduce and familiarise participant with the programme Getting to know the group Explore personal and group expectations allowing time for planning, prioritising and setting achievable targets Explore motivation, understanding potential short term and long-term benefits of engagement in the programme	COM-B Social Learning Theory Bio-Psycho-Social	Goal setting (behaviour) Goal setting (outcome) Action planning
Group session 2 Fear avoidance and pacing	To introduce relationship between thoughts, feelings, and emotions related to avoidance of behaviour (for example fear of aggravating COVID-19 symptoms such as breathlessness/fatigue) Exploring reasons for fear avoidance and consequences on health outcomes Exploring strategies to overcome fear avoidance and introduce pacing of activities	COM-B Self-Efficacy Bio-Psycho-Social	Information about antecedents Problem solving
Group session 3 Recovery and sleep, sleep management strategies	To explore how sleep patterns may have changed since having COVID Understanding sleep Introduction of sleep management strategies	COM-B Bio-Psycho-Social	Information about health consequences Problem solving Re-attribution
Group session 4 Management of emotions (perceived stigma, mood/unhelpful thoughts)	To introduce and explore the impact of COVID on mood, thoughts and behaviour including unhelpful thought patterns Introduce skills for reframing thoughts, encouraging engagement with all components of the programme	COM-B Cognitive Behavioural Principles	Social support (emotional) Information about emotional consequences Monitoring of emotional Consequences Framing/reframing
Group session 5 Understanding stress and anxiety and management strategies	To understand and explore impact of stress on symptom management (acute vs chronic stress) Understand own stress response to COVID and hospitalisation Introduce stress management strategies	COM-B Bio-Psycho-Social	Information about social and environmental consequences Reduce negative emotions
Group Session 6 Managing setbacks and long-term behaviour change and future goals.	To consolidate learning from previous sessions and reinforce long term behaviour change To introduce management of setbacks and strategies of problem solving for long term change	COM-B Group Based Learning Self-management	Problem solving Review behaviour goal(s) Social support (emotional) Comparative imagining of future outcomes Self-talk Focus on past success

Intervention component	Aims	Theoretical Framework(s)	Behaviour Change Taxonomy
Practitioner handbook	To provide step-by-step guidance on delivering the REGAIN intervention To provide facilitation skills and motivational interviewing prompts and questions for group sessions	COM-B Motivational Interviewing	Feedback on behaviour (through quality assessment and observations of sessions) Credible source
Participant workbook	Provide information on programme structure, safety and contact information for further support if needed Provide information and content for each session, with clear aims, and tasks to complete to consolidate learning To allow a log of personal progress and reflection and point of reference during the programme and beyond	COM-B Self-Efficacy Bio-Psycho-Social	Self-monitoring of outcome(s) of behaviour Information about health consequences Behavioural practice/rehearsal
Live group exercise sessions	To increase ability to engage in physical activity To increase strength, flexibility, balance and aerobic capacity. To increase confidence in ability to move and engage in physical activity To encourage physical activity outside of live sessions. To strengthen group dynamics	COM-B Self-Efficacy Group Based Learning Social Learning Theory	Feedback on outcome(s) of behaviour Instruction on how to perform a behaviour Demonstration of the behaviour Graded tasks Body changes
On demand exercise Videos	To increase ability to engage in physical activity To increase strength, flexibility, balance and aerobic capacity. To increase confidence in ability to move and engage in physical activity To encourage physical activity outside of live sessions.	COM-B Self-Efficacy	Demonstration of the Behaviour Instruction on how to perform a behaviour Behavioural practice/rehearsal Body changes
Other on demand videos; breathing technique, mindfulness	To provide advice and education on relevant topics. To provide associated skills/behaviours to promote engagement in physical activity To increase confidence in ability to move and engage in physical activity	COM-B Self-Efficacy	Demonstration of the Behaviour Instruction on how to perform a behaviour Behavioural practice/Rehearsal Distraction

motivational interviewing techniques, and group facilitation skills. All practitioners completed an assessment after training, to gauge readiness to deliver the intervention and/or the need for further training or support. Practitioners were provided with comprehensive written instruction manuals, and met monthly with a senior health psychologist and other members of the intervention delivery team for ongoing support, advice and troubleshooting. All intervention components were within the scope of normal practice for the clinicians involved. The REGAIN intervention was designed for delivery by NHS clinical exercise physiologists or physiotherapists. All practitioners were subject to regular quality control reviews conducted by an independent health psychologist, clinical exercise

physiologist, and qualitative researcher using pre-determined checklist criteria. This ensured that exercise and behavioural support sessions were delivered in a standardised manner. All online sessions were recorded for independent review as part of a separate process evaluation.

Safety

Supervised sessions were led by practitioners experienced in the assessment, prescription, and delivery of exercise for multi-morbid clinical populations. Pre-exercise session online poll questions were completed by participants to capture any adverse events experienced since the previous supervised session. Practitioners also ended each session by inquiring about any

symptoms or adverse events the participants may have experienced. Participants with any issues were asked to remain online after the session or were contacted by telephone or email to explore further. If a participant failed to attend an intervention appointment, the practitioner attempted to contact them via telephone or email to ascertain their welfare.

Exercise carries a very small theoretical risk of complications. All participants were assessed during the one-to-one consultation for any underlying health conditions or severe complications related to COVID-19. Participants were excluded from the study at the eligibility stage where exercise was clearly contraindicated, as assessed by a clinical member of the research team. A further assessment was undertaken by the REGAIN practitioner, through discussion with the patient about their current health, at the time of the initial online intervention assessment. During all group online exercise and support sessions, a second practitioner ('co-pilot') was immediately available online to assist with any emergencies, or if a participant became unresponsive or left the session without prior notification.

Regulatory compliance

To ensure compliance with appropriate confidentiality and data protection regulation, the proposed online intervention delivery platform and associated data storage facilities were rigorously reviewed and approved by organisational Information Governance committees, further to submission of a comprehensive Data Protection Impact Assessment.

Discussion

Regular physical activity can benefit people recovering from illness by increasing cardiorespiratory fitness and muscle strength, and reducing breathlessness. However, overly vigorous or prolonged activity without adaptation for ability or adequate rest periods can be detrimental to recovery and motivation, particularly for people with post-viral fatigue. It is important to mediate the psychosocial obstacles to activity and include intervention components designed to reduce anxiety and improve confidence, thus supporting people to regain physical strength, mobility, and independence. Provision of psychological support has become more challenging due to online delivery and differing levels of IT literacy. Hence, during development and pre-pilot testing, it was consistently demonstrated that our intervention would need to be as accessible and well supported as possible to avoid poor uptake and maintain adherence.

The introduction of an hour-long one-to-one consultation prior to the group sessions, a 'co-pilot' facilitator during the exercise sessions, and a comprehensive printed participant workbook aimed to address participants' concerns about their condition and any additional fear of engaging with digital technology. During the pre-pilot phase, we discovered that not only did the support sessions enhance engagement with the exercise sessions, but equally, the exercise sessions enhanced engagement with the support sessions. that the opposite was

also true. We found substantial unmet need for this patient group in being able to talk about their traumatic experiences in hospital with COVID-19, and their need to validate their ongoing problems since discharge. Our patient stakeholders expressed their desire to return to activities of normal living, (e.g., returning to work, walking to collect their children from school), or resume basic domestic chores at home. This required not only improved aerobic endurance and muscle strength, but also the knowledge, confidence and skills to cope with the inevitable peaks and troughs of recovery. Exercise practitioners also expressed their desire to improve their own group facilitation skills to best support the complex emotional needs of patients traumatised by COVID-19 and hospitalisation. The final fully manualised complex intervention for the REGAIN trial was co-produced with patients, lay people and healthcare professionals, and has been rolled out for testing in the REGAIN randomised controlled trial.

Conclusion

The REGAIN trial will be the largest study to-date to test whether online group exercise rehabilitation and psychological support delivered by NHS professionals is clinically and cost-effective for people with long-COVID symptoms more than three months after hospitalisation, when compared to best practice usual care. The REGAIN intervention, developed with extensive patient and stakeholder engagement, could be incorporated into existing NHS rehabilitation programmes, should it prove to be clinically and cost-effective for people with long-COVID.

Data availability

Underlying data

Dryad: REGAIN trial intervention development process evaluation data, <https://doi.org/10.5061/dryad.n8pk0p30r26>.

This project contains the following underlying data:

- REGAIN_PPI_feedback_on_behavioural_component_and_practitioner_manual.doc
- Live_online_exercise_session.docx

Data are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgements

We thank the wonderful group of people with long-COVID who participated fully in the development of the intervention at a time when their health and wellbeing was significantly impacted by COVID-19 – Elaine Fairbrother, Zoe Noakes, Bogdan Zutic, Helen Atherton, Lee Austin, Tom Forsyth, Anne Bush, Alan Bush, Patrick Sawdon. Also, our intervention team at Atrium Health – Jonathon Prosser, Lee Tompkins, Scott McGuire, Grace Lobley, Francesca Denton, David Montgomery.

References

- Office for National Statistics: **Prevalence of ongoing symptoms following coronavirus (COVID-19) infection in the UK 5**. 2023.
[Reference Source](#)
- Higgins V, Sohaei D, Diamandis EP, et al.: **COVID-19: from an acute to chronic disease? Potential long-term health consequences**. *Crit Rev Clin Lab Sci*. 2021; **58**(5): 297–310.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Ladds E, Rushforth A, Wieringa S, et al.: **Persistent symptoms after Covid-19: qualitative study of 114 “long Covid” patients and draft quality principles for services**. *BMC Health Serv Res*. 2020; **20**(1): 1144.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mahase E: **Covid-19: What do we know about “long covid”?** *BMJ*. 2020; **370**: m2815.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Evans RA, McAuley H, Harrison EM, et al.: **Physical, cognitive, and mental health impacts of COVID-19 after hospitalisation (PHOSP-COVID): a UK multicentre, prospective cohort study**. *Lancet Respir Med*. 2021; **9**(11): 1275–1287.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- NIHR: **Themed Review: Living with Covid19 - Second review**. NIHR; 2021.
[Reference Source](#)
- National Institute for Health and Care Excellence (NICE) SIGNS, Royal College of General Practitioners (RCGP): **COVID-19 rapid guideline: managing the long-term effects of COVID-19**.
[PubMed Abstract](#)
- NHS England: **Your COVID Recovery**.
[Reference Source](#)
- Fugazzaro S, Contri A, Esseroukh O, et al.: **Rehabilitation Interventions for Post-Acute COVID-19 Syndrome: A Systematic Review**. *Int J Environ Res Public Health*. 2022; **19**(9): 5185.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- McCarthy B, Casey D, Devane D, et al.: **Pulmonary rehabilitation for chronic obstructive pulmonary disease**. *Cochrane Database Syst Rev*. 2015; **2015**(2): CD003793.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Lau HMC, Ng GYF, Jones AYM, et al.: **A randomised controlled trial of the effectiveness of an exercise training program in patients recovering from severe acute respiratory syndrome**. *Aust J Physiother*. 2005; **51**(4): 213–9.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- McGregor G, Sandhu H, Bruce J, et al.: **Rehabilitation Exercise and psychological support After covid-19 InfectioN’ (REGAIN): a structured summary of a study protocol for a randomised controlled trial**. *Trials*. 2021; **22**(1): 8.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Skivington K, Matthews L, Simpson SA, et al.: **A new framework for developing and evaluating complex interventions: update of Medical Research Council guidance**. *BMJ*. 2021; **374**: n2061.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Rhon DI, Fritz JM, Kerns RD, et al.: **TIDieR-telehealth: precision in reporting of telehealth interventions used in clinical trials - unique considerations for the Template for the Intervention Description and Replication (TIDieR) checklist**. *BMC Med Res Methodol*. 2022; **22**(1): 161.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Liu K, Zhang W, Yang Y, et al.: **Respiratory rehabilitation in elderly patients with COVID-19: A randomized controlled study**. *Complement Ther Clin Pract*. 2020; **39**: 101166.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Larun L, Brurberg KG, Odgaard-Jensen J, et al.: **Exercise therapy for chronic fatigue syndrome**. *Cochrane Database Syst Rev*. 2017; **4**(4): CD003200.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Ngai JC, Ko FW, Ng SS, et al.: **The long-term impact of severe acute respiratory syndrome on pulmonary function, exercise capacity and health status**. *Respirology*. 2010; **15**(3): 543–50.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Stussman B, Williams A, Snow J, et al.: **Characterization of Post-exertional Malaise in Patients With Myalgic Encephalomyelitis/Chronic Fatigue Syndrome**. *Front Neurol*. 2020; **11**: 1025.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Kop WJ: **Biopsychosocial Processes of Health and Disease During the COVID-19 Pandemic**. *Psychosom Med*. 2021; **83**(4): 304–308.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Chakrawarty A, Ranjan P, Klanidhi KB, et al.: **Psycho-social and behavioral impact of COVID-19 on middle-aged and elderly individuals: A qualitative study**. *J Educ Health Promot*. 2021; **10**: 269.
[PubMed Abstract](#) | [Free Full Text](#)
- Michie S, van Stralen MM, West R: **The behaviour change wheel: a new method for characterising and designing behaviour change interventions**. *Implement Sci*. 2011; **6**: 42.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Bandura A: **Self-efficacy: toward a unifying theory of behavioral change**. *Psychol Rev*. 1977; **84**(2): 191–215.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Copeland L, McNamara R, Kelson M, et al.: **Mechanisms of change within motivational interviewing in relation to health behaviors outcomes: A systematic review**. *Patient Educ Couns*. 2015; **98**(4): 401–11.
[PubMed Abstract](#) | [Publisher Full Text](#)
- Borek AJ, Abraham C: **How do Small Groups Promote Behaviour Change? An Integrative Conceptual Review of Explanatory Mechanisms**. *Appl Psychol Health Well Being*. 2018; **10**(1): 30–61.
[PubMed Abstract](#) | [Publisher Full Text](#)
- British Psychological Society: **Guidance for the public on the mental health and wellbeing aspects of coronavirus (COVID-19)**.
- McGregor G: **REGAIN trial intervention development process evaluation data**. Dryad. [Dataset], 2023.
<http://www.doi.org/10.5061/dryad.n8pk0p30r>

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 28 July 2023

<https://doi.org/10.3310/nihropenres.14584.r29899>

© 2023 Kulnik S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Stefan Tino Kulnik 

Ludwig Boltzmann Institute for Digital Health and Prevention, Ludwig Boltzmann Gesellschaft, Salzburg, Austria

I thank the authors for addressing my comments.

I note that the members of the PPI group have now been named in the acknowledgements, which is a lovely touch.

(One small correction can be incorporated in the typesetting: there is an incomplete part-sentence in the second paragraph of the discussion which reads "that the opposite was also true" and which can be deleted.)

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Digital health, rehabilitation, physiotherapy, co-design, patient and public involvement, complex interventions, behaviour change

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 28 July 2023

<https://doi.org/10.3310/nihropenres.14584.r29898>

© 2023 Tew G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Garry A. Tew 

Institute for Health and Care Improvement, York St John University, York, UK

Thank you for updating the manuscript.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Professor of Clinical Exercise Science

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 02 June 2023

<https://doi.org/10.3310/nihropenres.14502.r29251>

© 2023 Kulnik S. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Stefan Tino Kulnik 

Ludwig Boltzmann Institute for Digital Health and Prevention, Ludwig Boltzmann Gesellschaft, Salzburg, Austria

Thank you for inviting my peer review of this manuscript entitled „Development of an online intervention for the Rehabilitation Exercise and psycholoGical support After covid-19 InfectioN (REGAIN) trial“. The manuscript describes the stepwise design of a tele-rehabilitation intervention for individuals with persisting symptoms of COVID-19 more than 3 months after infection. Intervention design was guided by the Medical Research Council framework for the development of complex interventions. The reporting followed the TIDieR checklist, including the recent extension for telehealth interventions.

I appreciate the detailed description of the intervention development process in this well-presented manuscript, which demonstrates good practice for transparency and scientific rigour and can be helpful for other intervention developers. I have the following comments for the authors' consideration, which I suggest to further strengthen the manuscript:

In the plain English summary, you could explicitly state that this is an online/remote intervention.

In the Methods:

Figure 1 is very helpful. I noticed that there are 4 components described in Stage 1 in Figure 1 (patient experience, evidence base, expert opinion, and current NHS practice), but in the main text under Stage 1 there is description of the review of evidence only. For consistency, I suggest to also briefly describe the other 3 components in the main text under Stage 1.

It would also be helpful for readers to be provided more information on the make-up of the PPI group (number of participants, gender distribution, age groups, months of living with long COVID), and the number and clinical roles of the individuals who were consulted for their expert opinions.

Stage 3 of the development process is described as 'pre-pilot', which could be read to imply that there follows a formal 'full' pilot study – but I understand that the intervention went directly to a definitive trial. Perhaps you can clarify this in the manuscript.

Please add a description of the timeline (month/year to month/year), ideally for each of the Stages.

In the Results:

I enjoyed reading about the many good design decisions that were made during the development process, such as the short introductory videos to stimulate support group discussion, and the thorough approach for screening for adverse events.

Tables 1, 2, and 3 and Figure 2 provide a good amount of relevant and helpful information. Some sentences seem abbreviated with missing articles, which I feel could be written out for better readability. Acronyms used in the tables could be explained in a legend. In Figure 2, I noticed that the wording of the online group support sessions 1-6 is not completely consistent with Table 3 and the main text. The image to illustrate the exercise programme in Figure 2 could better be an image without exercise equipment, as I understand it was a deliberate design decision to select exercises without need for equipment. In table 3 (row 'Group session 2', last column) the first behaviour change taxonomy seems to be incomplete ('Information about').

It was not quite clear to me whether all intervention components are intended to be delivered by clinical exercise physiologists or physiotherapists (i.e., the group support sessions also). You could state this more clearly.

Regular quality control reviews and recording of online sessions for independent review are described. Here it would be helpful to have more detail on who conducts these reviews, whether any particular assessment tools are used, and whether this is intended solely for trial purposes or also as an ongoing quality assurance measure if the intervention would be rolled out in practice.

I appreciated the design decisions for patient safety. In table 1, there is mention of aiming at a rate of perceived exertion (RPE) of 11-14. You could elaborate on this in the main text also and add a reference to the Borg RPE scale.

Table 3 provides a helpful linkage of intervention components with behaviour change theory. I would suggest that adding a figure with an explicit logic model is also required, to give readers an overview of the proposed mechanisms of action, linking through from the context and problem to the anticipated behavioural and clinical outcomes. This would be in line with the MRC framework recommendations.

Although the TIDieR checklist is not provided as a separate document, I think the manuscript covers most items of the checklist. Specifically with regard to the tele-rehabilitation/telehealth

aspect, you could add details of the technical and data regulatory requirements of the intervention (e.g., is it necessary to use certain video-conferencing tools to meet data protection regulations, etc.).

In the Discussion, 2nd paragraph, the meaning of the statement, but that the opposite was also true was unclear to me.

I note the two supplemental data files which are available for download on Dryad. These are helpful, but you could augment the files by adding more contextual detail for the reader, such as dates, number and characteristics of persons involved, who moderated/facilitated the activities, etc.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Digital health, rehabilitation, physiotherapy, co-design, patient and public involvement, complex interventions, behaviour change

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 23 Jul 2023

Gordon McGregor

Reviewer 2

Thank you for inviting my peer review of this manuscript entitled „Development of an online intervention for the Rehabilitation Exercise and psycholoGical support After covid-19 InfectioN

(REGAIN) trial". The manuscript describes the stepwise design of a tele-rehabilitation intervention for individuals with persisting symptoms of COVID-19 more than 3 months after infection. Intervention design was guided by the Medical Research Council framework for the development of complex interventions. The reporting followed the TIDieR checklist, including the recent extension for telehealth interventions.

I appreciate the detailed description of the intervention development process in this well-presented manuscript, which demonstrates good practice for transparency and scientific rigour and can be helpful for other intervention developers. I have the following comments for the authors' consideration, which I suggest to further strengthen the manuscript:

Thank you for your thorough and insightful review. We have amended the manuscript, figures, and tables accordingly for detail, clarity, consistency and readability (unless otherwise stated) in response to each of your suggestions below.

In the plain English summary, you could explicitly state that this is an online/remote intervention.

In the Methods:

Figure 1 is very helpful. I noticed that there are 4 components described in Stage 1 in Figure 1 (patient experience, evidence base, expert opinion, and current NHS practice), but in the main text under Stage 1 there is description of the review of evidence only. For consistency, I suggest to also briefly describe the other 3 components in the main text under Stage 1.

It would also be helpful for readers to be provided more information on the make-up of the PPI group (number of participants, gender distribution, age groups, months of living with long COVID), and the number and clinical roles of the individuals who were consulted for their expert opinions.

Stage 3 of the development process is described as 'pre-pilot', which could be read to imply that there follows a formal 'full' pilot study – but I understand that the intervention went directly to a definitive trial. Perhaps you can clarify this in the manuscript.

The pre-pilot phase preceded a 3-month internal pilot which continued seamlessly into the full trial.

Please add a description of the timeline (month/year to month/year), ideally for each of the Stages.

In the Results:

I enjoyed reading about the many good design decision that were made during the development process, such as the short introductory videos to stimulate support group discussion, and the thorough approach for screening for adverse events.

Tables 1, 2, and 3 and Figure 2 provide a good amount of relevant and helpful information. Some sentences seem abbreviated with missing articles, which I feel could be written out for better readability.

Acronyms used in the tables could be explained in a legend.

In Figure 2, I noticed that the wording of the online group support sessions 1-6 is not completely consistent with Table 3 and the main text.

The image to illustrate the exercise programme in Figure 2 could better be an image without exercise equipment, as I understand it was a deliberate design decision to select exercises without need for equipment.

In table 3 (row ,Group session 2', last column) the first behaviour change taxonomy seems to be incomplete (,Information about').

It was not quite clear to me whether all intervention components are intended to be delivered by clinical exercise physiologists or physiotherapists (i.e., the group support sessions also). You could state this more clearly.

Regular quality control reviews and recording of online sessions for independent review are described. Here it would be helpful to have more detail on who conducts these reviews, whether any particular assessment tools are used, and whether this is intended solely for trial purposes or also as an ongoing quality assurance measure if the intervention would be rolled out in practice.

Determination of delivery practices should the intervention be rolled out in clinical practice are out with the scope of this manuscript.

I appreciated the design decisions for patient safety. In table 1, there is mention of aiming at a rate of perceived exertion (RPE) of 11-14. You could elaborate on this in the main text also and add a reference to the Borg RPE scale.

Reference added.

Table 3 provides a helpful linkage of intervention components with behaviour change theory. I would suggest that adding a figure with an explicit logic model is also required, to give readers an overview of the proposed mechanisms of action, linking through from the context and problem to the anticipated behavioural and clinical outcomes. This would be in line with the MRC framework recommendations.

Logic model added (figure 3).

Although the TIDieR checklist is not provided as a separate document, I think the manuscript covers most items of the checklist. Specifically with regard to the tele-rehabilitation/ telehealth aspect, you could add details of the technical and data regulatory requirements of the intervention (e.g., is it necessary to use certain video-conferencing tools to meet data protection regulations, etc.).

Regulatory compliance section added.

In the Discussion, 2nd paragraph, the meaning of the statement, but that the opposite was also true was unclear to me.

I note the two supplemental data files which are available for download on Dryad. These are helpful, but you could augment the files by adding more contextual detail for the reader, such as dates, number and characteristics of persons involved, who moderated/facilitated the activities, etc.

These are raw data, thus cannot be altered. Information regarding data collection is outlined in the manuscript.

Competing Interests: none

Reviewer Report 18 May 2023

<https://doi.org/10.3310/nihropenres.14502.r29247>

© 2023 Tew G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Garry A. Tew 

Institute for Health and Care Improvement, York St John University, York, UK

The manuscript by Ennis and colleagues describes the development of the REGAIN intervention; an online group exercise and behavioural support intervention for people with long-COVID.

A multi-stage development process was undertaken, including: (1) background research, (2) co-production meetings to draft the intervention, (3) pre-piloting of the intervention with people suffering from long-Covid, and (4) refinement of the intervention based on initial experiences and feedback.

The final intervention included four main components: (1) an online 1:1 consultation, (2) online group exercise sessions, (3) online group behavioural support sessions, and (4) on-demand exercise and educational videos.

The manuscript is clearly written and includes everything that I would expect to see for this type of paper. I will be interested to see the results of the trial. I only have two relatively minor suggestions:

1. In figure 2, consider adding something about the on-demand exercise sessions and consider repositioning the information on the behavioural support sessions so that it doesn't look like it happens after the exercise programme.

2. I was left wanting a bit more information about the content of the group exercise sessions and on-demand exercise videos. Consider providing access to at least one of the 12 exercise session templates via the manuscript. And a link to at least one of the 14 videos.

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Yes

If applicable, is the statistical analysis and its interpretation appropriate?

Not applicable

Are all the source data underlying the results available to ensure full reproducibility?

Yes

Are the conclusions drawn adequately supported by the results?

Yes

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Professor of Clinical Exercise Science

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Author Response 23 Jul 2023

Gordon McGregor

Thank you for these helpful suggestions. We have amended figure 2 accordingly and have added an example exercise session template.

Competing Interests: none