



Cochrane
Library

Cochrane Database of Systematic Reviews

Chemotherapy for non-small cell lung cancer (Review)

for the Non-Small Cell Lung Cancer Collaborative Group

for the Non-Small Cell Lung Cancer Collaborative Group.
Chemotherapy for non-small cell lung cancer.
Cochrane Database of Systematic Reviews 2000, Issue 2. Art. No.: CD002139.
DOI: [10.1002/14651858.CD002139](https://doi.org/10.1002/14651858.CD002139).

www.cochranelibrary.com

Chemotherapy for non-small cell lung cancer (Review)
Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

WILEY

TABLE OF CONTENTS

ABSTRACT	1
PLAIN LANGUAGE SUMMARY	2
BACKGROUND	3
OBJECTIVES	3
METHODS	3
RESULTS	4
DISCUSSION	6
AUTHORS' CONCLUSIONS	7
ACKNOWLEDGEMENTS	8
REFERENCES	9
CHARACTERISTICS OF STUDIES	17
DATA AND ANALYSES	70
Analysis 1.1. Comparison 1 surgery vs surgery + chemotherapy, Outcome 1 survival.	71
Analysis 1.2. Comparison 1 surgery vs surgery + chemotherapy, Outcome 2 subgroup for survival - age.	72
Analysis 1.3. Comparison 1 surgery vs surgery + chemotherapy, Outcome 3 subgroup for survival - sex.	72
Analysis 1.4. Comparison 1 surgery vs surgery + chemotherapy, Outcome 4 subgroup for survival - performance status.	72
Analysis 1.5. Comparison 1 surgery vs surgery + chemotherapy, Outcome 5 subgroup for survival - histology.	72
Analysis 1.6. Comparison 1 surgery vs surgery + chemotherapy, Outcome 6 subgroup for survival - stage.	73
Analysis 2.1. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 1 survival.	73
Analysis 2.2. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 2 subgroup for survival - age.	74
Analysis 2.3. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 3 subgroup for survival - sex.	74
Analysis 2.4. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 4 subgroup for survival - performance status.	74
Analysis 2.5. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 5 subgroup for survival - histology.	75
Analysis 2.6. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 6 subgroup for survival - stage.	75
Analysis 3.1. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 1 survival.	76
Analysis 3.2. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 2 subgroup for survival - age.	77
Analysis 3.3. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 3 subgroup for survival - sex.	77
Analysis 3.4. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 4 subgroup for survival - performance status.	77
Analysis 3.5. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 5 subgroup for survival - histology.	77
Analysis 3.6. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 6 subgroup for survival - stage.	78
Analysis 4.1. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 1 survival.	78
Analysis 4.2. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 2 subgroup for survival - age.	79
Analysis 4.3. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 3 subgroup for survival - sex.	80
Analysis 4.4. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 4 subgroup for survival - performance status.	80
Analysis 4.5. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 5 subgroup for survival - histology. ..	80
Analysis 4.6. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 6 subgroup for survival - stage.	80
Analysis 5.1. Comparison 5 RT + C vs Rt, Outcome 1 Survival.	81
ADDITIONAL TABLES	82
WHAT'S NEW	82
HISTORY	82
DECLARATIONS OF INTEREST	83
SOURCES OF SUPPORT	83

NOTES	83
INDEX TERMS	84

[Intervention Review]

Chemotherapy for non-small cell lung cancer

for the Non-Small Cell Lung Cancer Collaborative Group¹

¹Meta-analysis Group, MRC Clinical Trials Unit at UCL, London, UK

Contact: Sarah Burdett, Meta-analysis Group, MRC Clinical Trials Unit at UCL, London, UK. sarah.burdett@ucl.ac.uk. Sarah Burdett, Meta-analysis Group, MRC Clinical Trials Unit at UCL, London, UK. .

Editorial group: Cochrane Lung Cancer Group.

Publication status and date: Stable (no update expected for reasons given in 'What's new'), published in Issue 3, 2015.

Citation: for the Non-Small Cell Lung Cancer Collaborative Group. Chemotherapy for non-small cell lung cancer. *Cochrane Database of Systematic Reviews* 2000, Issue 2. Art. No.: CD002139. DOI: [10.1002/14651858.CD002139](https://doi.org/10.1002/14651858.CD002139).

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

ABSTRACT

Background

The role of chemotherapy in the treatment of patients with non-small cell lung cancer was not clear. A systematic review and quantitative meta-analysis was therefore undertaken to evaluate the available evidence from all relevant randomised trials.

Objectives

To evaluate the effect of cytotoxic chemotherapy on survival in patients with non-small cell lung cancer. To investigate whether or not pre-defined patient sub-groups benefit more or less from chemotherapy.

Search methods

MEDLINE and CANCELIT searches (1963-june 1992) were supplemented by information from trial registers and by hand searching relevant meeting proceedings and by discussion with relevant trialists and organisations.

Selection criteria

Trials comparing primary treatments of surgery, surgery + radiotherapy, radical radiotherapy or supportive care versus the same primary treatment, plus chemotherapy were eligible for inclusion provided that they randomised non-small cell lung cancer patients using a method which precluded prior knowledge of treatment assignment.

Data collection and analysis

A quantitative meta-analysis using updated information from individual patients from all available randomised trials was carried out. Data from all patients randomised in all eligible trials were sought directly from those responsible. Updated information on survival, and date of last follow up were obtained, as were details of treatment allocated, date of randomisation, age, sex, histological cell type, stage and performance status. To avoid potential bias, information was requested for all randomised patients including those who had been excluded from the investigators' original analyses. All analyses were done on intention to treat on the endpoint of survival. For trials using cisplatin-based regimens, subgroup analyses by age, sex, histological cell type, tumour stage and performance status were also done.

Main results

Data from 52 trials and 9387 patients were included. The results for modern regimens containing cisplatin favoured chemotherapy in all comparisons and reached conventional levels of significance when used with radical radiotherapy and with supportive care. Trials comparing surgery with surgery plus chemotherapy gave a hazard ratio of 0.87 (13% reduction in the risk of death, equivalent to an absolute benefit of 5% at 5 years). Trials comparing radical radiotherapy with radical radiotherapy plus chemotherapy gave a hazard ratio 0.87 (13% reduction in the risk of death equivalent to an absolute benefit of 4% at 2 years), and trials comparing supportive care with supportive care plus chemotherapy gave a hazard ratio of 0.73 (27% reduction in the risk of death equivalent to a 10% improvement in survival at one year). The essential drugs needed to achieve these effects were not identified. No difference in the size of effect was seen in any subgroup

Chemotherapy for non-small cell lung cancer (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

of patients. In all but the radical radiotherapy setting, older trials using long term alkylating agents tended to show a detrimental effect of chemotherapy. This effect reached conventional significance in the adjuvant surgical comparison.

Authors' conclusions

At the outset of this meta-analysis there was considerable pessimism about the role of chemotherapy in the treatment of non-small cell lung cancer. These results offer hope of progress and suggest that chemotherapy may have a role in treating this disease.

PLAIN LANGUAGE SUMMARY

Chemotherapy can improve survival rates for non-small cell lung cancer

Non-small cell lung cancer is the most common type of lung cancer. The standard treatment for small tumours is surgery (operation to remove the tumour) or surgery and radiotherapy (x-ray treatment). Where the tumour has spread within the chest, standard treatment is radiotherapy. Where the tumour has spread beyond the chest supportive (treatment to relieve symptoms) is given. Trials have tried giving chemotherapy (drugs) after these standard treatments to find out whether it can help people to live longer. This review found that giving chemotherapy after either radiotherapy or supportive care did seem to help patients live longer. Giving chemotherapy after radiotherapy to 1000 patients would mean that an extra 40 patients would be expected to be alive 2 years later, than if the chemotherapy was not given. Giving chemotherapy after supportive care to 1000 patients would mean that 100 more would be expected to be alive 2 years later, than if the chemotherapy was not given. Chemotherapy after surgery may also help patients live longer although the evidence to support this is less clear.

BACKGROUND

More than half a million new cases of lung cancer are diagnosed each year (Parkin 1993). About 80% of these tumours are of non-small cell histological type (Rankin 1986), including adenocarcinomas and squamous cell and large cell carcinomas. Non-small cell lung cancer is the main cause of deaths related to cancer (Silverberg 1990), and five year survival across all stages of disease is about 12% (Boring 1993). Surgery is generally regarded as the best treatment option, but only about 20% of tumours are suitable for potentially curative resection (Rudd 1991). A further, small proportion of patients, usually those presenting with locally advanced disease, undergo radical thoracic radiotherapy. Most patients with late stage or metastatic disease are treated palliatively.

Although cytotoxic chemotherapy is used routinely in treating small cell lung cancer, its role in non-small cell lung cancer remained controversial. This was despite over thirty years of research involving more than 10,000 patients in over 50 randomised clinical trials examining the efficacy of chemotherapy when combined with local treatment or best supportive care. With few exceptions, most trials were too small to reliably detect moderate treatment effects. Consequently, although a few trials reported significant results, both for and against chemotherapy, most trials were inconclusive. In 1991, an international consensus report concluded that post-operative chemotherapy was of unproved benefit and should be considered experimental (Holmes 1991). In the same year, the British Medical Research Council's Cancer Trials Office, Cambridge; the Institut Gustave Roussy, Villejuif, France; and the Istituto Mario Negri, Milan, Italy initiated an individual patient data meta-analysis to assess the role of chemotherapy in the treatment of non-small cell lung cancer. This approach to meta-analysis and systematic review involves the central collection, validation and analysis of the original trial data. It does not rely on data extracted from publications. At the outset, the secretariat contacted the investigators responsible for each trial and established the Non-small Cell Lung Cancer Collaborative Group on whose behalf the meta-analysis was carried out and published in the British Medical Journal in 1995 (NSCLCCG 1995). Since that time a number of further trials have been completed. However, the majority of these are not yet published and several large trials are still ongoing. Members of the secretariat met in 1999 and decided that the meta-analysis should be updated when the results of these further trials become available. This is likely to be in 2000/2001.

OBJECTIVES

To compare, in terms of overall survival:

1. Surgery versus surgery plus adjuvant chemotherapy
2. Surgery plus radiotherapy versus surgery plus radiotherapy plus chemotherapy
3. Radical radiotherapy versus radical radiotherapy plus chemotherapy
4. Supportive care versus supportive care plus chemotherapy

in patients with histologically diagnosed non-small cell lung cancer.

Trials where chemotherapy was given before surgery (neo-adjuvant) were not included.

To investigate whether or not pre-defined patient sub-groups benefit more or less from cisplatin-based chemotherapy in terms of survival. Quality of life was measured in only a few trials and so could not be reviewed in the meta-analysis.

METHODS

Criteria for considering studies for this review

Types of studies

Both published and unpublished trials were eligible for inclusion provided they randomised patients with non-small cell lung cancer between one of the above four primary treatments and the same treatment plus an established form of cytotoxic chemotherapy. Each trial had to be unconfounded (i.e. differ only by the addition of chemotherapy to the treatment arm) and properly randomised. Trials allocating treatment by quasi random methods, e.g. by date of birth were not included. Trials were eligible if they started recruitment after 1 January 1965 and completed recruitment by 31 December 1991 (This upper date limit will be revised for the forthcoming update). Trials allowing patients to have received chemotherapy before randomisation were excluded. Trials in the early and locally advanced setting should not have permitted previous treatment for any other malignancy. Surgical trials were eligible only if they had randomised patients who had undergone a potentially curative resection and trials of neo-adjuvant treatment were not included in this comparison as it was considered too early to evaluate the neo-adjuvant approach. Trials of radical radiotherapy using orthovoltage radiotherapy or a total radiation dose of <30 Gy were excluded, as were trials in which drugs were used with the primary aim of sensitisation to radiation.

Types of participants

Eligible trials included individuals with histologically confirmed non-small cell lung cancer. Individual data from all randomised patients were included in the meta-analysis and where possible data were obtained for individuals who had been excluded from the original trial analyses. These individuals were included in the meta-analysis. Patients with small cell lung cancer that were included in early trials that randomised all types of lung cancer were excluded from the meta-analysis.

Types of interventions

1. Surgery vs surgery + adjuvant chemotherapy
2. Surgery + radiotherapy vs surgery + radiotherapy + chemotherapy
3. Radical radiotherapy vs radical radiotherapy + chemotherapy
4. Supportive care vs supportive care + chemotherapy

Trials investigating neo-adjuvant chemotherapy, that is chemotherapy given before surgery were not included.

Trials were classified as belonging to one of four categories of chemotherapy:

- a) Regimens containing cisplatin
- b) Regimens using long-term alkylating agents but not cisplatin
- c) Regimens containing etoposide or vinca alkaloids but not cisplatin
- d) Other regimens

Types of outcome measures

Mortality, death by any cause

As is usual with cancer trials and meta-analyses, results are discussed in terms of survival

Search methods for identification of studies

MEDLINE and CANCELIT searches were carried out for the period 1963-1990, and were updated to cover up to June 1992. Meetings abstracts of ASCO and the World Lung Cancer Conferences were hand searched as were bibliographies of books, reviews and specialist journals. Trial registers managed by the National Cancer Institute (PDQ, ClinProt), United Kingdom Coordinating Committee for Cancer Research and the Union Internationale Contre le Cancer were also consulted. All trialists who took part in the meta-analysis were also asked to help identify trials.

At the time of the initial literature searches, the Cochrane Collaboration optimal search strategy was not yet established. Since the publication of the meta-analysis (NSCLCCG 1995), all searches have been repeated using a modified version of the Cochrane Collaboration optimal search strategy.

Data collection and analysis

This review is based on individual patient data obtained directly from the responsible trialist or data centre. It does not use information extracted from published papers. All data were collected, checked and analysed centrally.

Trials were classified as belonging to one of four categories of chemotherapy:

- a) Regimens containing cisplatin
- b) Regimens using long-term alkylating agents but not cisplatin
- c) Regimens containing etoposide or vinca alkaloids but not cisplatin
- d) Other regimens

Note that the trials using oral alkylating agents (b) gave treatments for extended periods of more than one year as was considered best practice at the time, and that duration of treatment was not an exclusion criterium, but is rather purely descriptive.

Data were sought for all patients randomised in all eligible randomised trials (published or unpublished) and updated follow-up requested. For all comparisons the following data were collected: patient identifier, treatment allocated, date of randomisation, survival status, date of last follow up or death and whether the individual was excluded from the original analyses. Data on age, sex, stage, histology and performance status were also collected. Collection and validation of data were carried out in two centres (Cancer Trials Office and Institut Gustave Roussy).

All data were checked thoroughly and a common database was agreed. The final database entries for each trial were verified by the responsible trialist or data centre. As stage was recorded using different classification systems, for the purposes of this meta-analysis, all stage data were translated to a common staging system. [Table 1](#)

All analyses were based on intention to treat. Survival analyses were stratified by trial, and the log rank expected number of deaths and variance used to calculate individual and pooled hazard

ratios (HRs) using the fixed effect model (Yusuf 1985). Thus, the times to death for individual patients were used within trials to calculate HRs representing the overall chance of dying when receiving chemotherapy in addition to primary treatment alone. HRs were also calculated for pre-specified sub-groups of patients using similar stratified methodology. Analyses were performed for each pre-specified category, for example, for males and for females within each individual trial. These trial results were then combined to give overall HRs for males and females. Results are also presented as absolute differences at 2 years calculated using the HR and baseline event rate on the treatment alone arm; proportional hazards are assumed. Confidence intervals for absolute differences were similarly calculated from the baseline event rate and the HR at the 95% confidence interval boundary values. Chi-squared tests were used to test for gross statistical heterogeneity over all trials in a comparison, between sub-sets of trials, and between subgroups using the test for interaction or trend as appropriate (EBCTCG 1990). These tests are aimed primarily at detecting differences in effect size rather than direction and were chosen because qualitative differences were not anticipated.

Analyses of the "raw" individual patient data were done using an in-house program (SCHARP). For transfer to the Cochrane Library, the log rank summary statistics of these analyses (odds and variance) were entered into RevMan under the individual patient data category. Survival curves were drawn as simple (non-stratified) Kaplan Meier (Kaplan 1958) curves. These are not currently reproducible in the Cochrane Library but can be found in the original meta-analysis publication (NSCLCCG 1995). All P values quoted are two-sided.

RESULTS

Description of studies

In total, 91 trials were identified as potentially eligible for the meta-analysis. Thirty three of these were found to be ineligible and therefore excluded. Reasons for exclusions are listed in the table of excluded studies. Of the 58 eligible trials, data were not available from six as they had been lost, destroyed or were untraceable. These trials are also listed in the table of excluded studies. Data from 52 randomised trials and 9387 patients were therefore included in this meta-analysis.

Risk of bias in included studies

Only trials with adequate methods of randomisation were included. Trials using quasi random methods such as birthdate were not included. All "raw" data received on individual patients were checked thoroughly to ensure both the accuracy of the meta-analysis database and the quality of randomisation and follow-up. Any queries were resolved and the final database entries verified by discussion with the responsible trial investigator or statistician.

Effects of interventions

EARLY DISEASE

1. Surgery vs surgery plus adjuvant chemotherapy

Data were available from 14 trials (4357 patients, 2574 deaths). Five early trials used long term alkylating agents, mainly cyclophosphamide and nitrosourea. Eight more recent trials used cisplatin based combination chemotherapy. Three of these used the regimen of cisplatin, doxorubicin and cyclophosphamide (CAP) and three used cisplatin with vindesine. The intended dose of

cisplatin ranged from 40 mg/m² to 80 mg/m² per cycle and total dose from 50 mg/m² to 240 mg/m². A further three trials used other drug regimens, all of which included tegafur or UFT (tegafur plus uracil), a drug similar to fluorouracil. In all the trials chemotherapy was scheduled to start no later than six weeks after surgery.

There is considerable diversity of results across all trials and clear evidence of a difference in the direction of effect between the predefined categories of chemotherapy. The test for overall statistical heterogeneity is conventionally significant ($P=0.02$), as is the test for interaction ($P=0.004$). There is no evidence, however, of heterogeneity within each category of drugs ($P=0.21$). The results for each of the predefined chemotherapy categories should therefore be considered independently.

Trials using long-term alkylating agents

The results for trials using long term alkylating agents are consistent, all favour surgery alone. The combined hazard ratio is 1.15 ($P=0.005$) in favour of surgery alone. This 15% increase in the relative risk of death is equivalent to absolute detriments of chemotherapy of 4% at 2 years reducing survival from 70% to 66% and 5% at 5 years reducing survival from 50% to 45%.

Trials using cisplatin-based regimens

For regimens containing cisplatin, the results of most trials favour chemotherapy. There is no obvious statistical heterogeneity between the results of these trials ($P=0.55$). The overall hazard ratio of 0.87 ($P=0.08$), or 13% reduction in the risk of death, suggests an absolute benefit from chemotherapy of 3% at 2 years, improving survival from 70% to 73% and 5% at 5 years, improving survival from 50 to 55%. On their own these results are not conclusive such that the 95% confidence intervals for absolute difference in survival are consistent with a 0.5% detriment to a 7% benefit of chemotherapy at 2 years and similarly consistent with a 1% detriment to a 10% benefit at 5 years.

Other trials

The trials that were classified as using other regimens give an estimated hazard ratio of 0.89 in favour of chemotherapy ($P=0.30$), but there was insufficient information to draw any reliable conclusions.

2. Surgery + radiotherapy vs surgery + radiotherapy + chemotherapy
 Data were available from all seven eligible trials (807 patients and 619 deaths), six of which used a cisplatin based regimen. Intended doses of cisplatin ranged from 40 mg/m² to 100 mg/m² per cycle and total dose from 80 mg/m² to 400 mg/m². Total planned doses of radiotherapy ranged from 40 Gy in 10 fractions to 65 Gy in 33 fractions. The delay between surgery and the first adjuvant treatment was scheduled to be no longer than seven weeks.

The overall hazard ratio of 0.98 ($P=0.76$) is marginally in favour of chemotherapy. There is no gross statistical heterogeneity between the trials ($P=0.73$). For the cisplatin based trials the hazard ratio of 0.94 ($P=0.46$), or 6% reduction in the risk of death, favours chemotherapy and is equivalent to a 2% absolute benefit at both 2 and 5 years improving from 50% to 52% and from 15% to 17% respectively. The results are however not conventionally significant, the 95% confidence intervals range from a 4% detriment to an 8% benefit at 2 years and from a 3% detriment to an 8% benefit at 5 years.

LOCALLY ADVANCED DISEASE

3. Radical radiotherapy vs radical radiotherapy plus chemotherapy
 Data were available from 22 trials (3033 patients and 2814 deaths). Five trials used long term alkylating agents, mainly cyclophosphamide or nitrosourea in combination with methotrexate. Three used vinca-alkaloids or etoposide, and three used 'other' regimens, which in this comparison were mostly based on doxorubicin. Eleven trials used chemotherapy regimens containing cisplatin. Two used the regimen of cisplatin, doxorubicin and cyclophosphamide and seven used a combination of cisplatin plus a vinca-alkaloid or etoposide. Intended doses of cisplatin ranged from 40 mg/m² to 120 mg/m² per cycle and total doses from 120 mg/m² to 800 mg/m². The intended radiation dose for cisplatin based trials ranged from 50 Gy in 20 fractions to 65 Gy in 30 fractions. Ten of these trials started chemotherapy before radiotherapy.

In this comparison there was no gross statistical heterogeneity between trials ($P=0.56$), nor such strong evidence of a difference between chemotherapy categories, reflected in the non-significant test for interaction ($P=0.59$). The overall results show a significant overall benefit of chemotherapy. The hazard ratio of 0.90 ($P=0.006$), or 10% reduction in the risk of death, corresponds to absolute benefits of 3% at 2 years and 2% at 5 years. However it is useful to also consider each of the chemotherapies independently.

Trials using cisplatin-based regimens

Trials using cisplatin based chemotherapy provided the most information (more than 50%) and the strongest evidence for an effect in favour of chemotherapy. The hazard ratio of 0.87 ($P=0.005$), or 13% reduction in the risk of death, is equivalent to absolute benefits of 4% (95% confidence interval 1% to 7%) at 2 years improving survival from 15% to 19% and 2% (95% confidence interval 1% to 4%) at 5 years improving survival from 5% to 7%.

Other trials

Trials using long term alkylating agents and 'other' regimens both give a hazard ratio of 0.98 ($P=0.81$ and $P=0.88$ respectively), both marginally in favour of chemotherapy, but inconclusive. Trials using regimens containing vinca alkaloids or etoposide also favour chemotherapy, with a hazard ratio of 0.87 ($P=0.23$) or 13% reduction in the risk of death, but no firm conclusions can be drawn. Furthermore, there was no firm evidence that the results of the trials using regimens containing vinca alkaloids or etoposide or of those using other regimens of modern drugs are any different from those using cisplatin based chemotherapy.

ADVANCED DISEASE

4. Supportive care vs supportive care plus chemotherapy
 Data were available from all 11 eligible trials (1190 patients and 1144 deaths). Two trials used long term alkylating agents and one used etoposide as a single agent. The remaining eight trials used cisplatin based chemotherapy, seven of which used a combination of cisplatin and vinca alkaloids or etoposide. The intended dose of cisplatin ranged from 40 mg/m² to 120 mg/m² per cycle, with total doses of 280 mg/m² upwards. This included several trials in which chemotherapy was given until the disease progressed or the toxicity was unacceptable. In this advanced disease setting, however, many patients would not have received the planned number of treatment cycles. One trial allowed entry of only patients with metastatic disease, the rest included patients with both locally advanced and advanced disease.

There is considerable overall statistical heterogeneity ($P < 0.0001$) and a pronounced difference in the results for the different chemotherapy categories, ($P = 0.003$) and again it is wise to focus on the results within each of the pre-defined chemotherapy categories.

Trials using long-term alkylating agents

The result for trials using long term alkylating agents suggested a detriment of chemotherapy with a hazard ratio of 1.26 or 26% increase in the relative risk of death. However, with only two such trials, the confidence intervals are wide (0.96 to 1.66) and the result does not reach conventional levels of significance ($P = 0.095$).

Trials using cisplatin-based chemotherapy

The cisplatin based trials show a clear benefit of chemotherapy with a hazard ratio of 0.73 ($P < 0.0001$) or 27% reduction in the risk of death. This is equivalent to an absolute improvement in survival of 10% at one year, improving survival from 15% to 25%, or an increased median survival of 1.5 months, improving median survival from 4 months to 5.5 months. One trial (CEP-85) showed an extreme result in favour of chemotherapy. When this trial is excluded from the analysis, the results are still significantly in favour of chemotherapy (hazard ratio 0.77 (0.63 to 0.85, $P = 0.001$). When this trial is removed, there is no gross statistical heterogeneity within the cisplatin based category ($P = 0.09$).

TREATMENT EFFECT IN PATIENT SUBGROUPS

Predefined subgroups of patients were analysed to determine whether we could identify particular types of patient or tumour that benefited more (or less) from chemotherapy. To minimise heterogeneity, only cisplatin based regimens were included in this analysis. Data on stage were available for 92% of patients, performance status for 94% of patients and age, sex and histological cell type for more than 99% of patients. There is no evidence that any group of patients specified by age, sex, histological cell type, performance status or stage benefit more or less from chemotherapy. This means, for example, that patients of all ages appear to gain the same relative benefit from chemotherapy. Note that these analyses do not compare the underlying survival of patients. It does not imply that old and young patients live for the same amount of time.

UPDATING RESULTS

As this systematic review is based on the original data from trials (not data taken from publications), updates are major projects taking many months of full time work and requiring the input of numerous individuals and groups. Standard practice for IPD meta-analyses are to undertake full updates, when appropriate, depending on the maturity of data and the rate at which further trials are completed and published.

This systematic review has now been updated to include more patients from more randomised controlled trials published since 1995. Details of current numbers of patients and trials included and the most up to date citations are listed below. Several new Cochrane Reviews will be submitted.

1. Surgery vs. surgery + adjuvant chemotherapy

Updated to include: 8447 patients, 34 trial comparisons

Most recent citation: NSCLC Meta-analysis Collaborative Group. Adjuvant chemotherapy, with or without postoperative

radiotherapy, in operable non-small-cell lung cancer: two meta-analyses of individual patient data. *Lancet*. 2010;375:1267-77.

2. Surgery + radiotherapy vs. surgery + radiotherapy + chemotherapy

Updated to include: 2660 patients, 13 trial comparisons

Most recent citation: NSCLC Meta-analysis Collaborative Group.

Adjuvant chemotherapy, with or without postoperative radiotherapy, in operable non-small-cell lung cancer: two meta-analyses of individual patient data. *Lancet*. 2010;375:1267-77.

3. Radiotherapy vs. radiotherapy + sequential chemotherapy

Updated to include: 3839 patients, 22 RCTs

Most recent citation: Le Pechoux C, Burdett S, Auperin A. Individual patient data (IPD) meta-analysis (MA) of chemotherapy (CT) in locally advanced non-small cell lung cancer (NSCLC). *Journal of Thoracic Oncology*. 2008; 3(Supplement 1):S20, 35IN.

4. Supportive care vs. supportive care + chemotherapy

Updated to include: 2714 patients, 16 RCTs

Most recent citations:

Non-Small Cell Lung Cancer Collaborative Group. Chemotherapy and supportive care versus supportive care alone for advanced non-small cell lung cancer. *Cochrane Database of Systematic Reviews* 2010, Issue 5. Art. No.: CD007309. DOI: 10.1002/14651858.CD007309.pub2

NSCLC Meta-Analyses Collaborative Group. Chemotherapy in addition to supportive care improves survival in advanced non-small cell lung cancer: a systematic review and meta-analysis of individual patient data from 16 randomized trials. *J Clin Oncol*. 2008;26:4617-25.

Three other comparisons of timing of chemotherapy have also been explored:

a) Radiotherapy vs. radiotherapy + concomitant chemotherapy

Includes: 2910 patients, 16 RCTs

Most recent citation: Le Pechoux C, Burdett S, Auperin A. Individual patient data (IPD) meta-analysis (MA) of chemotherapy (CT) in locally advanced non-small cell lung cancer (NSCLC). *Journal of Thoracic Oncology*. 2008; 3(4, Supplement 1):S20, 35IN.

b) Radiotherapy + sequential chemotherapy vs. radiotherapy + concomitant chemotherapy

Includes: 1205 patients, 6 RCTs

Most recent citation: Aupérin A, Le Pécoux C, Rolland E, et al on behalf of the NSCLC Collaborative Group. Concomitant versus sequential radiochemotherapy in locally advanced non-small cell lung cancer: A meta-analysis of individual data of 1205 patients. *J Clin Oncol*. 2010;28:2181-90.

c) Surgery vs. surgery + neoadjuvant chemotherapy

Status: ongoing

DISCUSSION

This meta-analysis is based on an extensive dataset comprising individual data on 9387 patients from 52 randomised controlled

trials that compared local surgical or radiotherapy treatment or best supportive care with the same treatment plus chemotherapy in non-small cell lung cancer. A number of methods were employed to try to identify all trials and both published and unpublished data were included thereby minimising the influence of publication bias. Furthermore, at the time the meta-analysis was completed only six other eligible trials were found, for which data were not available. These were mostly older trials using chemotherapy regimens based on the long term administration of oral alkylating agents, regimens that are no longer used. Only one of the unavailable trials used a cisplatin based regimen and data from approximately 99% of all patients ever entered into all known relevant trials of modern chemotherapies were analysed. Thus it is unlikely that the observed results would be changed by the unavailable data. Furthermore, for almost all trials the data on individual patient had been updated to the point of data collection, which was often many years after the publication of the trial's results. Although a number of trials have been completed since the meta-analysis was published, most of these are not yet in the public domain. This meta-analysis therefore currently provides a comprehensive and reliable assessment of the average treatment effect of broad categories of chemotherapy regimens among broad classes of patients with non-small cell lung cancer. We aim to obtain data from the new trials including data from further large trials due to complete soon in the next update of the individual patient data meta-analysis.

One of the most striking aspects of the results is the consistency in the direction, and indeed in the estimated hazard ratios, of the various chemotherapy categories among the different primary treatment comparisons. This consistency allows stronger conclusions to be drawn than perhaps could be inferred from each of the individual results.

In the early and advanced settings, older trials using long term alkylating agents tended to show a detrimental effect of chemotherapy. This effect was conventionally significant for the adjuvant surgical trials. Chemotherapies of the type used in the early 1970s based on long term administration of oral alkylating agents are therefore likely to be detrimental to patients with non-small cell lung cancer. The mechanism for this is unknown, although some occurrences of leukaemia after treatment with busulphan have been described for non-small cell lung cancer (Stott 1977), and a possible model for an observed detrimental effect of cyclophosphamide and other alkylating agents in non-small cell lung cancer has been proposed (Stewart 1992). Clearly, such regimens are not used today, but the result could have implications for other disease sites, albeit that the administration of chemotherapy and the drugs used have changed considerably over the past twenty years.

In all comparisons, results for modern cisplatin-containing regimens favour chemotherapy. These are conventionally significant in the locally advanced and supportive care settings. However, it should be stressed that this categorisation of drug regimens was chosen mainly as an objective way of classifying modern chemotherapy. Furthermore, several cisplatin based regimens were used and it is not possible to deduce to what extent the observed effects are due to the cisplatin or to the other drugs, in the combinations studied. Indeed cisplatin was used in combination with vinca alkaloids or etoposide in two thirds of trials.

It is therefore not possible to recommend a particular regimen over another.

Trials using regimens containing vinca alkaloids or etoposide and those in the "other drug" category also tend to favour chemotherapy. For these categories the confidence intervals are relatively wide and no reliable conclusions can be drawn.

The meta-analysis provides no evidence that modern cisplatin based chemotherapy is more or less effective in any particular subgroup of patients. Thus, no good evidence exists that the relative effect of chemotherapy is any smaller or larger for any particular type of patient. Nevertheless, as certain types of patient may have intrinsically different prognoses and consequently differing baseline survivals, the same hazard ratio or relative effect may provide different absolute differences in survival. For example, in the surgical setting, the hazard ratio of 0.87 would increase the 2-year survival of patients with a good prognosis from a baseline of 80%, to 82%. For patients with a poor prognosis this same HR would improve survival from a 2-year baseline of 40% to 45%. Similarly, the same observed hazard ratio of 0.87 in the locally advanced setting would increase the survival of patients with a good prognosis from a baseline of 30%, to 35% and patients with a poor prognosis from 5% to 7%. Thus the absolute benefit derived from the same relative risk and consequently the clinical decisions reached may be different for older or younger patients for example. It is also worth noting that the patients included in these trials are generally of better prognosis than those in the lung cancer population at large. For example few patients older than 75 years or with poor performance status were included in the trials.

The meta-analysis suggests that modern chemotherapy regimens may provide absolute benefits of about 5% in the surgical and 2% in the radical radiotherapy setting at 5 years and 10% at one year in the supportive care setting. The confidence intervals are such, however, that the results are consistent with benefits of as much as 10%, 4% and 15% respectively or with as little as a 1% detriment and 1% and 5% benefits respectively. Although modest, such improvements could, given the high incidence of lung cancer, be important in public health terms, and studies of patients' opinions of treatments for cancer have shown that many patients accept considerable toxicity in return for small improvements in survival (Slevin 1990). However, patients are not uniform in their preferences, and the trade offs involved in choosing between more and less intensive therapy are not necessarily straightforward and warrant further study (Till 1992).

An important consideration when making such choices is the effect that chemotherapy may have on quality of life. Unfortunately, because few trials reported it, quality of life could not be addressed in the meta-analysis.

AUTHORS' CONCLUSIONS

Implications for practice

Although, inevitably, meta-analyses give only average estimates of treatment effects, these are probably the best estimates on which to base treatment policy. At the outset of this meta-analysis there was considerable pessimism about the treatment of non-small cell lung cancer. Although the suggested benefits of modern chemotherapies are modest, these results offer hope of progress and show that chemotherapy may have a role in treating this

disease. Some patients and clinicians would need to observe larger treatment effects than others before being convinced that chemotherapy is worthwhile, and undoubtedly these results will be applied differently by individual clinicians and patients around the world. Some groups may consider these results to be good enough evidence to use cisplatin based chemotherapy for certain patients. As essential drugs were not determined by this meta-analysis, however, others may need further evidence to decide whether to use chemotherapy routinely in the treatment of non-small cell lung cancer.

Implications for research

Extended follow up on existing trials and the inclusion of further randomised trials will add to the evidence in the next update of this meta-analysis. Continued research into screening new drugs and improving chemotherapy regimens is required as is measurement of quality of life.

ACKNOWLEDGEMENTS

The following investigators, groups and secretariat were members of the Non-small Cell Lung Cancer Collaborative Group and participated in this meta-analysis:

The investigators were:

W Alberti, G Anderson, A Bartolucci, D Bell, J Blanco Villalba, O Brodin, C Cardiello, F Cartei, G Cartei, R Cellerino, C Chastang, Y Cormier, J Cox, L Crino, J Crowley, B Dautzenberg, A Depierre, A Dietemann, R Dillman, O Doi, R Feld, R Figlin, P Ganz, M Green, A Gregor, P Helle, J Herndon, S Hitomi, H Host, M Imaizumi, J Jett, D Johnson, S Kaasa, H Kimura, J Klastersky, H Kondo, H Kreisman, M Kris, K Kunishima, O Kuwahara, T Lad, A Laing, F Macbeth, A Masaoka, K Mattson, E Minatel, J Mira, T Mori, C Mountain, N Neiderle, A Niiranen, E Nou, W Page, J Pater, S Piantadosi, K Pisters, S Pyrhonen, E Quoix, E Rapp, N Rowell, T Sahnoud, K Sawamura, D Schallier, C Scott, J Simpson, M Stagg, T Teramatsu, M Trovo, R Tsuchiya, D Tummarello, P van Houtte, N Van Zandwijk, R Vincent, H Wada, J White, C Williams, R Wood, Y Yamaguchi

The organisations and groups that collaborated in the meta-analysis were: Cancer and Leukemia Group B, Cercle d'Etudes Pneumologiques (Strasbourg), European Organisation for Research and Treatment of Cancer, Fédération Nationale des Centres de Lutte Contre le Cancer, Finnish Lung Cancer Study, Groupe d'Etude et de Traitement des Cancers Bronchiques, Japan Lung Cancer Surgical Study Group, Lung Cancer Study Group, Medical Research Council, National Cancer Institute of Canada Clinical Trials Group, North Central Cancer Treatment Group, Osaka Lung Cancer Study Group, Radiation Therapy Oncology Group, Southeastern Cancer Study Group, South West Oncology Group, Study Group of Adjuvant Chemotherapy for Lung Cancer (Chibu, Japan), Swedish Lung Cancer Study Group, Veterans Administration Surgical Adjuvant Group, Veterans Administration Surgical Oncology Group, West of Scotland Lung Cancer Research Group, West Japan Study Group for Lung Cancer Surgery.

The members of the secretariat were:

R Arriagada, J Cartmell-Davies, D Girling, T Le Chevalier, S Marsoni, M Parmar, JP Pignon, C Rekeawicz, R Souhami, L Stewart, M Tarayre, A Tinazzi, V Torri

We thank all the patients who took part in all the trials included in the meta-analysis. We also thank the following institutions that funded the Non-small Cell Lung Cancer Collaborative Group Meeting in September 1993, at which preliminary results of this meta-analysis were presented and discussed: Bristol-Myers Oncology, Bristol-Myers Squibb, Amgen, Asta-Medica Ltd, Asta-Sarget, Beecham Sevigne, Roger Bellon, Cancer Research Campaign, Chugai, Ciba-Geigy, Eurocetus, Farmitalia Carlo Erba, Glaxo France, Harold Hyam Wingate Foundation, Institut National de la Santé et de la Recherche Médicale, Ipsen Biotech, Lederle, Lilly France, Lilly Industries, Medical Research Council, Pfizer, Rhône-Poulenc and Servier Medical. The Cancer Trials Office's secretariat thanks Richard Wood, Sarah Walker and John Machin, all the computing, data management and secretarial staff for their support throughout this project. The Institute Gustave Roussy's secretariat thanks Ben Affaied for programming support and Sylviane Iacobelli for secretarial help.

REFERENCES

References to studies included in this review

a01 MRC LU02 {unpublished data only}

* Girling DJ, Stott H, Stephens RJ, Fox W. Fifteen-year follow-up of all patients in a study of post-operative chemotherapy for bronchial carcinoma. *British Journal of Cancer* 1985;**52**:867-873.

a02 VASAG {unpublished data only}

* Shields TW, Humphrey EW, Eastridge CE, Keehn RJ. Adjuvant cancer chemotherapy after resection of carcinoma of the lung. *Cancer* 1977;**40**:2057-2062.

a03 EORTC 08741a {unpublished data only}

* Israel L, Bonadonna G, Sylvester R. Controlled study with adjuvant radiotherapy, chemotherapy, immunotherapy and chemoimmunotherapy in operable squamous carcinoma of the lung. In: Muggia FM, Rozenzweig M editor(s). *Lung Cancer: Progress in Therapeutic Research*. New York: Raven Press, 1979:443-452.

a04 VASOG 5 {unpublished data only}

* Shields TW, Higgins GA, Humphrey EW, Matthews MJ, Keehn RJ. Prolonged intermittent adjuvant chemotherapy with CCNU and hydroxyurea after resection of carcinoma of the lung. *Cancer* 1982;**50**:1713-1721.

a05 WPL 7351 {unpublished data only}

* Mountain CF, Vincent RG, Sealy R, et al. A clinical trial of CCNU as surgical adjuvant treatment for patients with surgical stage I and stage II non-small cell lung cancer: Preliminary findings.. In: Muggia FM, Rozenzweig M editor(s). *Lung Cancer: Progress in Therapeutic Research..* New York: Raven Press, 1979:421-431.

a06 OLCSG 1a {unpublished data only}

* Sawamura K, Mori T, Doi O, Yasumitsu T, Kuwahara O, Kuwabara M, Nakahara K, Kurata M, Sagara N. A prospective randomized controlled study of the postoperative adjuvant therapy for non-small cell lung cancer. *Lung Cancer* 1988;**4**:A166.

a07 OLCSG 1b {unpublished data only}

* Sawamura K, Mori T, Doi O, Yasumitsu T, Kuwahara O, Kuwabara M, Nakahara K, Kurata M, Sagara N. A prospective randomized controlled study of the postoperative adjuvant therapy for non-small cell lung cancer. *Lung Cancer* 1988;**4**:A166.

a08 SGACLC 1 {unpublished data only}

* Study Group for Adjuvant Chemotherapy for Lung Cancer. A randomised controlled trial of postoperative adjuvant chemotherapy in non-small cell lung cancer (in Japanese.). *Hai-gan* 1992;**32**:481-486.

a09 WJSG 2 {unpublished data only}

Teramatsu T on behalf of the Society of adjuvant chemotherapy for lung cancer surgery in West Japan. Assessment of postoperative adjuvant chemotherapy on non-small cell lung cancer (abstract). *Lung Cancer* 1991;**7**(Suppl):124.

* Wada H, Hitomi S, Takashi T, and the West Japan Study Group for Lung Cancer Surgery. Adjuvant chemotherapy after complete resection in non-small cell lung cancer (full publication). *Journal of Clinical Oncology* 1996;**14**:1048-1054.

a10 LCSG 801 {unpublished data only}

* Feld R, Rubinstein L, Thomas PA. Adjuvant chemotherapy with cyclophosphamide, doxorubicin and cisplatin in patients with completely resected stage I non-small-cell lung cancer. *Journal of the National Cancer Institute* 1993;**85**(4):299-306.

a11 OLCSG 1c {unpublished data only}

* Sawamura K, Mori T, Doi O, Yasumitsu T, Kawahara O, Kuwabara M, Nakahara K, Kurata M, Sagara N. A prospective randomized controlled study of the postoperative adjuvant therapy for non-small cell lung cancer. *Lung Cancer* 1988;**4**:A166.

a12 FLCSG 1 {unpublished data only}

* Niiranen A, Niitamo-Korhonen S, Kouri M, Assendelft A, Mattson K, Pyrhönen S. Adjuvant chemotherapy after radical surgery for non-small cell lung cancer: A randomized study. *Journal of Clinical Oncology* 1992;**10**(12):1927-1932.

a13 SGACLC 2 {unpublished data only}

* Study Group for Adjuvant Chemotherapy for Lung Cancer. A randomized trial of postoperative adjuvant chemotherapy in non-small cell lung cancer (the second cooperative study). *European Journal of Surgical Oncology* 1995;**21**:69-77.

a14 IPCR, Chiba {unpublished data only}

* Kimura H, Yamaguchi Y, Fujisawa T, Baba M, Shiba M. A randomized controlled study of postoperative adjuvant chemoimmunotherapy of resected non-small cell lung cancer with IL2 and LAK cells. *Lung Cancer* 1991;**7**(Suppl):113.

a15 LCSG 853 {unpublished data only}

* Lung Cancer Study Group. A clinical trial in patients with stage II and III completely resected non-small cell cancer of the lung comparing chemotherapy (CAP) versus no therapy following surgery.

a16 JLCSSG {unpublished data only}

* Ohta M, Tsuchiya R, Shimoyama M, Sawamura K, Mori T, Miyazawa N, Suemasu K, Watanabe Y, Tomita M, Terashima M. Adjuvant chemotherapy for completely resected stage III non-small cell lung cancer. *Journal of Thoracic and Cardiovascular Surgery* 1993;**106**:703-708.

b01 EORTC 08741b {unpublished data only}

* Israel L, Bonadonna G, Sylvester R and members of the EORTC Lung Cancer Group. Controlled study with adjuvant radiotherapy, chemotherapy, immunotherapy and chemoimmunotherapy in operable squamous carcinoma of the lung. In: Muggia FM, Rozenzweig M editor(s). *Lung Cancer: Progress in Therapeutic Research*. New York: Raven Press, 1979:443-452.

b02 LCSG 791 {unpublished data only}

Lad T. The comparison of CAP chemotherapy and radiotherapy to radiotherapy alone for resected lung cancer with positive margin or involved highest sampled paratracheal node (stage IIIa). *Chest* 1994;**106**(6):303S.

* Lad T, Rubinstein L, Sadeghi A. The benefit of adjuvant treatment for resected locally advanced non-small cell lung cancer. *Journal of Clinical Oncology* 1988;**6**:9-17.

b03 MSKCC 80-53 {unpublished data only}

* Pisters KMW, Kris MG, Gralla RT, Hilaris B, McCormack PM, Bains MS. Randomized trial comparing post-operative chemotherapy with vindesine and cisplatin plus thoracic irradiation with irradiation alone in stage III (N2) non-small cell lung cancer. *Journal of Surgical Oncology* 1994;**56**:236-241.

b04 FLCSG 3 {unpublished data only}

* Niiranen, Kouri M, Pyrhonen S, Mattson K. Postsurgical radiotherapy versus postsurgical radiotherapy plus chemotherapy for non-small cell lung cancer.

b05 GETCB 01CB82 {unpublished data only}

* Chastang CI, Dautzenberg B, Arriagada R, Le Chevalier T, Belpomme D, Hurdebourg J, the GETCB. Results of a randomized clinical trial of a postoperative chemotherapy (COPAC) in stage II and III resected bronchogenic carcinoma. *European Journal of Cancer* 1991;**27**(Suppl 2):A1017.

b06 OLCSG 1d {unpublished data only}

* Sawamura K, Mori T, Doi O, Yasumitsu T, Kawahara O, Kuwabara M, Nakahara K, Kurata M, Sagara N. A prospective randomized controlled study of the postoperative adjuvant therapy for non-small cell lung cancer. *Lung Cancer* 1988;**4**:A166.

b07 EORTC 08861 {unpublished data only}

* EORTC Lung Cancer Cooperative Group. Phase III randomized trial of adjuvant radiotherapy vs radiotherapy plus chemotherapy with DDP/VDS vs no adjuvant therapy in patients with completely resected non-small cell lung cancer.

c01 NRH NSC 26271 {unpublished data only}

* Høst H. Cyclophosphamide as adjuvant to radiotherapy in the treatment of unresectable bronchogenic carcinoma. *Cancer Chemotherapy Reports* 1973;**4**(Part 3):161-164.

c02 EORTC 08742 {unpublished data only}

* Israel L, Depierre A, Dalesio O. Interim results of EORTC protocol 08742: comparison, after irradiation of locally advanced squamous cell bronchial carcinoma, of abstention, immunotherapy, combination chemotherapy or chemoimmunotherapy. *Recent Results on Cancer Research* 1982;**80**:214-218.

c03 RTOG 7302 a {unpublished data only}

* Simpson JR, Francis ME, Perez-Tamayo R, Mark RD, Rao DV. Palliative radiotherapy for inoperable carcinoma of the lung: Final report of a RTOG multi-institutional trial. *International Journal of Radiation, Oncology, Biology and Physics* 1985;**11**(4):751-758.

c04 RTOG 7302 b {unpublished data only}

* Simpson JR, Francis ME, Perez-Tamayo R, Mark RD, Rao DV. Palliative radiotherapy for inoperable carcinoma of the lung: Final report of a RTOG multi-institutional trial. *International Journal of Radiation, Oncology, Biology and Physics* 1985;**11**(4):751-758.

c05 RTOG 7302 c {unpublished data only}

* Simpson JR, Francis ME, Perez-Tamayo R, Mark RD, Rao DV. Palliative radiotherapy for inoperable carcinoma of the lung: Final report of a RTOG multi-institutional trial. *International Journal of Radiation, Oncology, Biology and Physics* 1985;**11**(4):751-758.

c06 MCL-1 {unpublished data only}

* Sidorowicz E, Hirsch V, Hand R, Guerra J, Margolese R, Thirlwell M, Colman N, Frank H, Kreisman H. Comparison of radiotherapy (RT) and RT-MACC chemotherapy in limited non-small cell lung cancer (NSCLC). *Proceedings of the IVth world conference on lung cancer* 1985:A277.

c07 Aviano {unpublished data only}

* Trovo MG, Minatel E, Veronesi A, Roncadin M, De Paoli A, Franchin G, Magri DM, Tirelli U, Carboma A, Grigoletto E. Combined radiotherapy and chemotherapy versus radiotherapy alone in locally advanced epidermoid bronchogenic carcinoma. *Cancer* 1990;**65**:400-404.

c08 AZ-OC-1-80 {unpublished data only}

* Schallier DC, de Neve WJ, de Greve JL, van Belle SP, de Wasch GJ, Dotremont G, Storme GA. Is adjuvant treatment with vinblastine effective in reducing the occurrence of distant metastasis in limited squamous cell lung cancer? A randomized study. *Clinical and Experimental Metastasis* 1988;**6**(1):39-48.

c09 Gwent 3 {unpublished data only}

* Anderson G. Study of radiotherapy versus radiotherapy + VP-16-213 in carcinoma of the bronchus.

c10 SECSG 81 LUN375 {unpublished data only}

* Johnson DH, Einhorn LH, Bartolucci A, Birch R, Omura G, Perez CA, Greco FA. Thoracic radiotherapy does not prolong survival in patients with locally advanced unresectable non-small cell lung cancer. *Annals of Internal Medicine* 1990;**113**:33-38.

c11 Gwent 1 {unpublished data only}

* Anderson G, Deeley TJ, Smith C, Jani J. Comparison of radiotherapy alone and radiotherapy with chemotherapy using adriamycin and 5-fluorouracil in bronchogenic carcinoma. *Thorax* 1981;**36**:190-193.

c12 SWOG 7635 {unpublished data only}

* White JE, Chen T, Reed R, Mira J, Stuckey WJ, Weatherall T, O'Bryan R, Samson MK, Seydel HG. Limited squamous cell carcinoma of the lung: A Southwest Oncology Group randomised study of radiation with or without doxorubicin chemotherapy and with or without levasmisole immunotherapy. *Cancer Treatment Reports* 1982;**66**(5):1113-1120.

c13 NCCTG 822451 {unpublished data only}

* Morton RF, Jett JR, McGinnis WL, Earle JD, Thernea TM, Krook JE, Elliott TE, Mailliard JA, Nelimark RA, Maksymiuk AW, et al. Thoracic radiation therapy alone compared with combined chemoradiotherapy for locally unresectable non-small cell lung cancer. *Annals of Internal Medicine* 1991;**115**:681-686.

c14 Buenos Aires {unpublished data only}

* Cardiello C, Blanco Villalba J, Anac S, Francheri Wilson C, Trodler C, Hunis A, Alvarez Bermudez C. Combined radiochemotherapy (RtCt) versus radiotherapy (Rt) in limited inoperable non-small cell carcinoma of the lung (NSCLC). *Proceedings of the American Society of Clinical Oncology* 1985;**4**:117.

c15 Brussels {unpublished data only}

* van Houtte P, Klastersky J, Renauld A, Michel J, Vandermoten G, Nguyen H, Sculier JP, Devriendt J, Mommen P. Induction chemotherapy with cisplatin etoposide and vindesine before radiation therapy for non-small cell lung cancer. *Antibiotic Chemotherapy* 1988;**41**:131-137.

c16 FLCSG 2 {unpublished data only}

* Mattson K, Holsti LR, Holsti P, Jakobsson M, Kajanti M, Liippo K, Mäntylä M, Niitamo-Korhonen S, Nikkanen V, Nordman E, et al. Inoperable non-small cell lung cancer: radiation with or without chemotherapy. *European Journal of Clinical Oncology* 1988;**24**:477-482.

c17 Essen {unpublished data only}

* Alberti W, Niederle N, Budach V, Konietzko N, Sack H. Prospective randomized study comparing immediate radiotherapy, chemo- plus radiotherapy or delayed radiotherapy in non-small cell lung cancer. *Journal of Cancer Research and Clinical Oncology* 1990;**116**(part 1):503.

c18 SLCSG {unpublished data only}

* Brodin O, Nou E, on behalf of the Swedish Multicentric Group for the study of squamous cell carcinoma of the lung. Patients with non-resectable squamous cell carcinoma of the lung. A prospective randomised study. *Lung Cancer* 1991;**7**:A615.

c19 CEBI 138 {unpublished data only}

* Le Chevalier T, Arriagada R, Quoix E, Ruffie P, Martin M, Tarayre M, Lacombe-Terrier M-J, Douillard J-Y, Laplanche A. Radiotherapy alone versus combined chemotherapy and radiotherapy in nonresectable non-small cell lung cancer: First analysis of a randomized trial in 353 patients. *Journal of the National Cancer Institute* 1991;**83**(6):417-423.

c20 WSLCRG/FI {unpublished data only}

* Gregor A, Macbeth FR, Paul J, Cram L, Hansen HH. Radical radiotherapy in localized inoperable non-small cell lung cancer: A randomized trial. *Journal of the National Cancer Institute* 1993;**85**(12):997-999.

c21 Perugia {unpublished data only}

* Crinò L, Latini P, Meacci M, Corgna E, Maranzano E, Darwish S, Minotti V, Santucci A, Tonato M. Induction chemotherapy plus high-dose radiotherapy versus radiotherapy alone in locally

advanced unresectable non-small cell lung cancer. *Annals of Oncology* 1993;**4**:847-851.

c22 CALGB 8433 {unpublished data only}

* Dillman RO, Seagren SL, Propert KJ, Guerra J, Eaton WL, Perry MC, Carey R, Frei EF, Green MR. A randomized trial of induction chemotherapy plus high-dose radiation versus radiation alone in stage III non-small cell lung cancer. *New England Journal of Medicine* 1990;**323**:940-945.

c23 EORTC 08842 {unpublished data only}

* Planting A, Helle P, Drings P, Dalesio O, Kirkpatrick A, McVie G, Giaccone G. A randomized study of high-dose split course radiotherapy preceded by high-dose chemotherapy versus high-dose radiotherapy only in locally advanced non-small cell lung cancer. *Annals of Oncology* 1996;**7**:139-144.

c24 SWOG 8300 a {unpublished data only}

* Miller TP, Mira JG, Horn B. Treatment of limited non-small cell lung cancer: Radiation versus radiation plus chemotherapy.

c25 SWOG 8300 b {unpublished data only}

* Miller TP, Mira JG, Horn B. Treatment of non-small cell lung cancer: Radiation versus radiation plus chemotherapy.

d01 Oxford {unpublished data only}

* Laing AH, Berry RJ, Newman CR, Peto J. Treatment of inoperable carcinoma of bronchus. *The Lancet* 1975;**1161-1164**.

d02 Quebec {unpublished data only}

* Cormier Y, Bergeron D, La Forge J, Lavindier M, Fournier M, Chenard J, Desmeules M. Benefits of polychemotherapy in advanced non-small cell bronchogenic carcinoma. *Cancer* 1982;**50**:845-849.

d03 Gwent 2 {unpublished data only}

* Anderson G, Payne H. Response rate and toxicity of etoposide (VP-16) in squamous carcinoma of the lung: Report from the Lung Cancer Treatment Study Group. *Seminars of Oncology* 1985;**12**(1 Suppl 2):21-22.

d04 RLW 8351 {unpublished data only}

* Woods RL, Williams CJ, Levi J, Page J, Bell D, Byrne M, Kerestes ZL. A randomised trial of cisplatin and vindesine versus supportive care only in advanced non-small cell lung cancer. *British Journal of Cancer* 1990;**61**:608-611.

d05 NCIC CTG BR5 {unpublished data only}

* Rapp E, Pater JL, Willan A, Cormier Y, Murray N, Evans WK, Hodson DI, Clark DA, Feld R, Arnold AM, et al. Chemotherapy can prolong survival in patients with advanced non-small cell lung cancer-Report of a Canadian multicenter randomized trial. *Journal of Clinical Oncology* 1988;**6**(4):633-641.

d06 Southampton {unpublished data only}

* Woods RL, Williams CJ, Levi J, Page J, Bell D, Byrne M, Kerestes ZL. A randomised trial of cisplatin and vindesine versus supportive care only in advanced non-small cell lung cancer. *British Journal of Cancer* 1990;**61**:608-611.

d07 NRH {unpublished data only}

* Kaasa S, Lund E, Thorud E, Hatlevoll R, Høst H. Symptomatic treatment versus combination chemotherapy for patients with extensive non-small cell lung cancer. *Cancer* 1991;**67**:2443-2447.

d08 UCLA {unpublished data only}

* Ganz PA, Figlin RA, Haskell CM, La Soto N, Siau J. Supportive care versus supportive care and combination chemotherapy in metastatic non-small cell lung cancer. Does chemotherapy make a difference?. *Cancer* 1989;**63**:1271-1278.

d09 Ancona 1 {unpublished data only}

* Cellerino R, Tummarello D, Guidi F, Isidori P, Raspugli M, Biscottini B, Fatati G. A randomized trial of alternating chemotherapy versus best supportive care in advanced non-small cell lung cancer. *Journal of Clinical Oncology* 1991;**9**(8):1453-1461.

d10 AOI-Udine {unpublished data only}

* Cartei G, Cartei F, Cantone A, Causarano D, Genco G, Tobaldin A, Interlandi G, Giraldo T. Cisplatin-cyclophosphamide-mitomycin combination chemotherapy with supportive care versus supportive care alone for treatment of metastatic non-small cell lung cancer. *Journal of the National Cancer Institute* 1993;**85**(10):794-800.

d11 CEP-85 {unpublished data only}

* Quoix É, Dietemann A, Charbonneau J, Boutin C, Meurice JC, Orlando JP, Ducolone A, Pauli G, Roegel É. Is cisplatin-based chemotherapy useful in disseminated non small cell lung cancer? Report of a French multicenter randomized trial [La chimiothérapie comportant du cisplatine est-elle utile dans le cancer bronchique non microcellulaire au stade IV? Résultats d'une étude randomisée]. *Bulletin du Cancer (Paris)* 1991;**78**:341-346.

ya(surg) <=54 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the surgical setting.

yb(surg) 55-59 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the surgical setting.

yc(surg) 60-64 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the surgical setting.

yd(surg) >=65 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the surgical setting.

ye(surg) male {unpublished data only}

Subgroup analysis by sex for cisplatin-based trials in the surgical setting.

yf(surg) female {unpublished data only}

Subgroup analysis by sex for cisplatin-based trials in the surgical setting.

yg(surg) good PS {unpublished data only}

Subgroup analysis by performance status for cisplatin-based trials in the surgical setting.

yh(surg) poor PS {unpublished data only}

Subgroup analysis by performance status for cisplatin-based trials in the surgical setting.

yi(surg) adeno {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the surgical setting.

yj(surg) squamous {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the surgical setting.

yk(surg) other {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the surgical setting.

yl(surg) stage I,II {unpublished data only}

Subgroup analysis by stage for cisplatin-based trials in the surgical setting.

ym(surg) stage III {unpublished data only}

Subgroup analysis by stage for cisplatin-based trials in the surgical setting.

yn(s+rt) <=54 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the surgery + RT setting.

yo(s+rt) 55-59 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the surgery + RT setting.

yp(s+rt) 60-64 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the surgery + RT setting.

yq(s+rt) >=65 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the surgery + RT setting.

yr(s+rt) male {unpublished data only}

Subgroup analysis by sex for cisplatin-based trials in the surgery + RT setting.

ys(s+rt) female {unpublished data only}

Subgroup analysis by sex for cisplatin-based trials in the surgery + RT setting.

yt(s+rt) good PS {unpublished data only}

Subgroup analysis by performance status for cisplatin-based trials in the surgery + RT setting.

yu(s+rt) poor PS {unpublished data only}

Subgroup analysis by performance status for cisplatin-based trials in the surgery + RT setting.

yv(s+rt) adeno {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the surgery + RT setting.

yw(s+rt) squamous {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the surgery + RT setting.

yx(s+rt) other {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the surgery + RT setting.

yy(s+rt) stage I,II {unpublished data only}

Subgroup analysis by stage for cisplatin-based trials in the surgery + RT setting.

yz(s+rt) stage III {unpublished data only}

Subgroup analysis by stage for cisplatin-based trials in the surgery + RT setting.

za(radrt) <=54 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the radical RT setting.

zb(radrt) 55-59years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the radical RT setting.

zc(radrt) 60-64years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the radical RT setting.

zd(radrt) >=65 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the radical RT setting.

ze(radrt) male {unpublished data only}

Subgroup analysis by sex for cisplatin-based trials in the radical RT setting.

zf(radrt) female {unpublished data only}

Subgroup analysis by sex for cisplatin-based trials in the radical RT setting.

zg(radrt) good PS {unpublished data only}

Subgroup analysis by performance status for cisplatin-based trials in the radical RT setting.

zh(radrt) poor PS {unpublished data only}

Subgroup analysis by performance status for cisplatin-based trials in the radical RT setting.

zi(radrt) adeno {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the radical RT setting.

zj(radrt) squamous {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the radical RT setting.

zk(radrt) other {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the radical RT setting.

zl(radrt) stage I,II {unpublished data only}

Subgroup analysis by stage for cisplatin-based trials in the radical RT setting.

zm(radrt) stage III {unpublished data only}

Subgroup analysis by stage for cisplatin-based trials in the radical RT setting.

zn(sc) <=54 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the supportive care setting.

zo(sc) 55-59 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the supportive care setting.

zp(sc) 60-64 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the supportive care setting.

zq(sc) >=65 years {unpublished data only}

Subgroup analysis by age for cisplatin-based trials in the supportive care setting.

zr(sc) male {unpublished data only}

Subgroup analysis by sex for cisplatin-based trials in the supportive care setting.

zs(sc) female {unpublished data only}

Subgroup analysis by sex for cisplatin-based trials in the supportive care setting.

zt(sc) good PS {unpublished data only}

Subgroup analysis by performance status for cisplatin-based trials in the supportive care setting.

zu(sc) poor PS {unpublished data only}

Subgroup analysis by performance status for cisplatin-based trials in the supportive care setting.

zv(sc) adeno {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the supportive care setting.

zw(sc) squamous {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the supportive care setting.

zx(sc) other {unpublished data only}

Subgroup analysis by histology for cisplatin-based trials in the supportive care setting.

zy(sc) non-metastatic {unpublished data only}

Subgroup analysis by stage for cisplatin-based trials in the supportive care setting.

zz(sc)metastatic {unpublished data only}

Subgroup analysis by stage for cisplatin-based trials in the supportive care setting.

References to studies excluded from this review

Aliev 1985 {published data only}

* Aliev BM, Starichkov MS, Mikhina ZP, Petrukhin OD, Bychkov MB. Radiotherapy and combined chemotherapy and radiotherapy of patients with squamous-cell lung cancer (a randomized study). *Voprosy Onkologii* 1985;**31**(10):71-6.

Bergsagel 1972 {published data only}

* Bergsagel DE, Jenkin RDT, Pringle JF, White DM, Fetterly JCM, Klaassen DJ. [Lung Cancer: clinical trial of radiotherapy alone vs radiotherapy plus cyclophosphamide]. *Cancer* 1972;**30**:621-627.

Blaha 1973 {published data only}

* Blaha H, Dierkesmann R, Feuerer W, et al. Adjuvant therapy of non-small cell bronchial cancer with mopidamol [Adjuvante therapie des nicht-kleinzelligen bronchialkarzinoms mit mopidamol]. *Pneumologie* 1973;**43**:299-304.

Brunner 1971 {published data only}

* Brunner KW, Marthaler TH, Muller W. Unfavourable effects of long-term adjuvant chemotherapy with endoxan in radically operated bronchogenic carcinoma. *European Journal of Cancer* 1971;**7**:285-294.

Buyze 1973 {published data only}

* Buyze EAC, Nelemans FA. A study of post-operative cytostatic medication in patients with operable carcinoma of the lung. *Arzneim. Forsh (Drug Research)* 1973;**23**(6):860-862.

Byar 1978 {published data only}

* Byar D, Kenis Y, van Andel JG, et al. Results of a E.O.R.T.C. randomized trial of cyclophosphamide and radiotherapy in inoperable lung cancer: Prognostic factors and treatment results. *European Journal of Cancer* 1978;**14**:919-930.

Carr 1972 {published data only}

* Carr DT, Childs DS, Lee RE. Radiotherapy plus 5-FU compared to radiotherapy alone for inoperable and unresectable bronchogenic carcinoma. *Cancer* 1972;**29**(2):375-380.

Castberg 1976 {published data only}

* Castberg T, Kruse-Blinkenberg HO. Cancer chemotherapy as a supplement to operation in bronchogenic carcinoma [Cancerkemoterapi som supplement til operation ved bronkogent karcinom]. *Ugestr Laeg* 1976;**138**:1449-1455.

Crosbie 1966 {published data only}

* Crosbie WA, Kamdar HH, Belcher JR. Controlled trial of vinblastine sulphate in the treatment of cancer of the lung. *British Journal of Diseases of the Chest* 1966;**60**:28.

Dolton 1970 {published data only}

* Dolton EG. Combined surgery and chemotherapy for carcinoma of bronchus. *Lancet* 1970;**Jan 3**:40-41.

Durrant 1971 {published data only}

* Durrant KR, Ellis F, Black JM, Berry RJ, Ridehalgh FR, Hamilton WS. Comparison of treatment policies in inoperable bronchial carcinoma. *Lancet* 1971;**1**(7702):715-719.

Hall 1967 {published data only}

* Hall TC, Dederick MM, Chalmers TC, et al. A clinical pharmacologic study of chemotherapy and x-ray therapy in lung cancer. *American Journal of Medicine* 1943;**2**:186-193.

Hansen 1973 {published data only}

* Hansen JL. Clinical evaluation of an innocuous antineoplastic agent: Preliminary study of 106 patients treated with or without RSV after pulmonary resection for bronchogenic carcinoma. *Cancer Chemotherapy Reports* 1973;**4**(Part 3):261-264.

Helsper 1962 {published data only}

Helsper JT, Sharpe GS. Combination therapy with 5-fluorouracil and colbalt-60 for inoperable carcinoma of the lungs. *Cancer Chemotherapy Reports* 1962;**20**:103-105.

Holsti 1971 {published data only}

* Holsti LR. Adjuvant Chemotherapy. In: Deeley editor(s). *Modern Radiotherapy*. 222-240.

Hosley 1962 {published data only}

* Hosley HF, Marangoudakis S, Ross CA, Murray WT, Holland JF. Combination radiation-chemotherapy for bronchogenic carcinoma - pilot study. *Cancer Chemotherapy Reports* 1962;**16**:467-471.

Karageorgis 1991 {published data only}

* Karageorgis P, Koliarakis N, Paraskevaidis M. NSCLC st. IIIb patients survival after chemotherapy, radiotherapy and combination of the above modalities. Preliminary report. *Proceedings of ECCO* 1991;**6**:S180.

Karp {published data only}

* Karp DD, Daly BDT, Diehl JT, et al. A pilot study of radiation therapy with or without concomitant cisplatin and etoposide as adjuvant therapy in non-small cell lung cancer. In: Bernal SD, Hesketh PJ editor(s). *Lung Biology in Health and Disease*. 58. 1992:325-334.

Karrer 1978 {published data only}

* Karrer K, Prindun N, Denck H. Chemotherapy as an adjuvant to surgery in lung cancer. *Cancer Chemotherapy and Pharmacology* 1978;**1**:145-159.

Kaung 1974 {published data only}

* Kaung DT, Wolf J, Hyde L, Zelen M. [Preliminary report on the treatment of nonresectable cancer of the lung]. *Cancer Chemotherapy Reports* 1974;**58**(Part 1):359-364.

NCCTG852451 {unpublished data only}

NCCTG. Phase III randomized comparison of surgical adjuvant combination chemotherapy with CAP (CTX/ADR/CDDR) vs observation in patients with totally resected stage I, II or III.

Newman 1985 {published data only}

Newman CE, Cox R, Ford CHJ, Johnson JR, Jones DR, Wheaton M. Reduced survival with radiotherapy and razoxane compared with radiotherapy alone for inoperable lung cancer in a randomised double-blind trial. *British Journal of Cancer* 1985;**51**:731-732.

Okawa 1992 {published data only}

* Okawa T, Hashimoto S, Niibe H, et al. Effect of sizofiran (SPG) combined with radiotherapy on non-small cell lung cancer (NSCLC), results of a prospective randomized controlled study, cooperative study on SPG lung cancer. *Proceedings of the American Society of Clinical Oncology* 1992;**11**:288.

Osterlind 1985 {published data only}

* Osterlind K, Hansen M, Hansen HH, Dombernowsky P. Controlled trial of RSV, herbs or placebo as adjuvants to complete resection of squamous cell lung cancer. *European Journal of Surgical Oncology* 1985;**11**(4):349-351.

Petrovic 1978 {published data only}

* Petrovich Z, Ohanian M, Cox J. [Clinical research on the treatment of locally advanced lung cancer. Final report of VALG protocol 13 limited]. *Cancer* 1978;**42**:1129-1134.

Pirogov 1976 {published data only}

* Pirogov AI, Traktenburg AKh. Results and prospects for combined surgery and antitumour chemotherapy for lung cancer. *Cancer Treatment Reports* 1976;**60**(10):1489-1491.

Privitera 1987 {published data only}

Privitera G, Ciottoli GB, Patanè C, et al. Phase II double-blind randomized study of lonidamine and radiotherapy in epidermoid carcinoma of the lung. *Radiotherapy and Oncology* 1987;**10**:285-90.

Scheer 1974 {published data only}

* Scheer AC, Wilson RF, Kalisher L. Combined radiotherapy and hydroxyurea in the management of lung cancer. *Clinical Radiology* 1974;**25**:415-418.

Selawry 1977 {published data only}

* Selawry O, Krant M, Scotto J, et al. Methotrexate compared with placebo in lung cancer. *Cancer* 1977;**40**:4-8.

Slack 1970 {published data only}

* Slack NH. Bronchogenic carcinoma: nitrogen mustard as a surgical adjuvant and factors influencing survival. University surgical adjuvant lung project. *Cancer* 25;**5**:987-1002.

Spatti 1985 {published data only}

* Spatti G, Regazzoni M, Koronel R, Zecchini AM, de Palo G. Adjuvant treatment with melphalan in ovarian carcinoma with no residual disease following surgery. *Tumori* 1987;**73**:157-162.

Spittle 1980 {published data only}

* Spittle M, Newton KA, White W, et al. Razoxane as a radiopotentiator for lung cancer.. Second World Conference on Lung Cancer, Copenhagen. 1980; Vol. 220.

Vincent 1975 {published data only}

* Vincent RG, Pickren JW, Fergen TB, Takita H. Evaluation of methotrexate in the treatment of bronchogenic carcinoma. *Cancer* 1975;**36**:873-880.

Wils 1984 {published data only}

* Wils JA, Utama I, Naus A, Verschuieren TA. Phase II randomized trial of radiotherapy alone vs the sequential use of chemotherapy and radiotherapy in stage II non-small cell lung cancer. Phase II trial of chemotherapy alone in stage IV non-small cell lung cancer. *European Journal of Clinical Oncology* 1984;**20**(7):911-914.

Wingfield 1970 {published data only}

* Wingfield HV. Combined surgery and chemotherapy for carcinoma of the bronchus. *Lancet* 1970;**1**(7644):470-471.

Wolf 1991 {published data only}

* Wolf K, Havemann K, Hans K, et al. Radiotherapy versus chemotherapy followed by radiotherapy in inoperable limited stage non-small cell lung cancer (NSCLC). *Lung Cancer* 1991;**7s**:A611.

References to ongoing studies
Ancona {unpublished data only}

Gasparini S, Zucchetto L, Massei V. Lonidamine and chemotherapy plus lonidamine vs the best supportive care in the treatment of advanced non-small cell lung cancer.

ANITA {unpublished data only}

Adjuvant Navelbine International Trialist Association (ANITA). Phase III randomized trial of adjuvant chemotherapy following curative resection for non-small cell lung cancer.

BLT {published and unpublished data}

* MRC CTU. Phase III randomized study of cisplatin-based chemotherapy in patients with non-small cell lung cancer.

CAN NCIC BR10 {published and unpublished data}

* NCIC Clinical Trials Group. Phase III randomized study of adjuvant chemotherapy with VNB/CDDP vs no adjuvant chemotherapy for completely resected non-small cell lung cancer.

CLB 9633 {published and unpublished data}

* Cancer and Leukemia Group B. Phase III randomized study of carboplatin / paclitaxel vs no adjuvant chemotherapy after resection of stage IB non-small cell lung cancer.

CNR ALPI {published and unpublished data}

* Consiglio Nazionale Ricerche, Italy / EORTC. Phase III randomized study of adjuvant MVP vs no adjuvant chemotherapy (with or without adjuvant radiotherapy) for radically resected stage I/II/IIIa non-small cell lung cancer.

CRC-TU-LU3001 {published and unpublished data}

* Cullen MH, Billingham LJ, Woodroffe CM, et al. Mitomycin, Ifosfamide, and Cisplatin in unresectable non-small cell lung

cancer: effect on survival and quality of life. *Journal of Clinical Oncology* 1999;**17**:3188-3194.

CRC-TU-LU3002 {published data only}

* Cullen MH, Billingham LJ, Woodroffe CM, et al. Mitomycin, Ifosphamide, and Cisplatin in unresectable non-small cell lung cancer: effect on survival and quality of life. *Journal of Clinical Oncology* 1999;**17**:3188-3194.

EORTC 08922 {published and unpublished data}

* EORTC. Phase III randomised trial of adjuvant MICE (Mesna/ IFF/CBDCA/VP-16) or EP (VP-16/CDDP) vs no adjuvant therapy in patients with completely resected stage I/II/IIIa non-small cell lung cancer containing the activated oncogene K-ras.

FRE IALT {published and unpublished data}

* Institut Gustave Roussy. Phase III randomized study of adjuvant chemotherapy following curative resection for non-small cell lung cancer.

GAP {published data only}

Clerici M, Barni S, Cantaluppi G, et al. Adjuvant chemotherapy in non-small cell lung cancer: a randomised trial. *European Journal of Cancer* 1991;**27**(Suppl 2):S173.

Helsing {published and unpublished data}

* Helsing M, Bergman B, Thaning L, Hero U. Quality of life and survival in patients with advanced non-small cell lung cancer receiving supportive care plus chemotherapy with carboplatin and etoposide or supportive care only. A multicentre randomised phase III trial. *European Journal of Cancer* 1998;**34**(7):1036-1044.

Ichinose 1991 {published data only}

* Ichinose Y, Hara N, Ohta M, Motohiro A, Kuda T, Aso H. Postoperative adjuvant chemotherapy in non-small cell lung cancer: Prognostic value of DNA ploidy and postrecurrent survival. *Journal of Surgical Oncology* 1991;**46**:15-20.

Kim 1993 {published data only}

Kim NK, Yang SH, Im YH, et al. A phase III randomized comparison of neoadjuvant chemotherapy with cisplatin, etoposide and vinblastine plus radiation vs. radiation alone in stage III non-small cell lung cancer. *Proceeding of the American Society for Clinical Oncology* 1993;**12**:330.

NCCTG-822451 {published data only}

* North Central Cancer Treatment Group. Phase III randomized comparison of surgical adjuvant combination chemotherapy with CAP vs observation in patients with totally resected stage I, II or III non-small cell lung cancer.

RICUM {published data only}

Ayoub J, Duranceau A, Lorange G, et al. Effectiveness of adjuvant chemotherapy in operable non-small cell lung cancer. In: Salmon SE editor(s). *Adjuvant therapy of cancer*. New York: and Stratton, 1987:157-164.

RTOG 8808 {unpublished data only}

Sause WT, Scott C, Taylor S, et al. Radiation Therapy Oncology Group (RTOG) 88-08 and Eastern Cooperative Oncology Group

(ECOG) 4588: Preliminary results of a phase III trial in regionally advanced unresectable non-small cell lung cancer. *Journal of the National Cancer Institute* 1995;**87**:198-205.

TELCVIS 1999 {published data only}

* The Elderly Lung Cancer Vinorelbine Italian Study. Effects of Vinorelbine on quality of life and survival of elderly patients with advanced non-small cell lung cancer. *Journal of the National Cancer Institute* 1999;**91**:66-72.

Thongprasert 1999 {published and unpublished data}

Thongprasert S, Sanguanmitra P, Juthapan W, Clinch J. Relationship between quality of life and clinical outcomes in advanced non-small cell lung cancer: best supportive care (BSC) versus BSC plus chemotherapy. *Lung Cancer* 1999;**24**:17-24.

Additional references

Asola 1991

Asola R, Suovuori A, Karnani P. Peroral tegafur and calcium folinate in the treatment of advanced colorectal cancer: FINN-SIC phase I trial. *European Journal of Cancer* 1991;**2**(suppl):S91, A529.

Boring 1993

Boring CC, Squires TS, Tong T. Cancer statistics. *Ca: a Cancer Journal for Clinicians* 1993;**43**:7-26.

DerSimonian 1986

DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials* 1986;**7**:177-88.

EBCTCG 1990

Early Breast Cancer Trialists Collaborative Group. *Treatment of Early Breast Cancer Vol 1. Worldwide evidence 1985-1990*. Oxford University Press, 1990.

Holmes 1991

Holmes EC, Bleeher NM, Le Chevalier T, Ettinger D, Jett J, Johnson D. Post-operative adjuvant treatments for non-small cell lung cancers: a consensus report. *Lung Cancer* 1991;**7**:11-3.

Kaplan 1958

Kaplan EL, Meier P. Nonparametric estimation from incomplete observation. *Journal of the American Statistical Association* 1958;**53**:457-81.

Parkin 1993

Parkin DM, Sasco AJ. Lung cancer: Worldwide variation in occurrence and proportion attributable to tobacco use. *Lung Cancer* 1993;**9**:1-16.

Rankin 1986

Rankin EM. Non-small cell lung cancer. In: Slevin ML, Staquet MJ editor(s). *Randomised clinical trials in cancer: A critical review by site*. New York: Raven Press, 1986:447-92.

Rudd 1991

Rudd R. Chemotherapy in the treatment of non-small cell lung cancer. *Respiratory Disease in Practice* 1991;**7**(6):12-4.

Silverberg 1990

Silverberg E, Boring CC, Squires TS. Cancer Statistics. *Ca: a Cancer Journal for Clinicians* 1990;**40**:9-26.

Slevin 1990

Slevin ML, Stubbs L, Plant WJ, Wilson P, Gregory WP, Armes PJ. Attitudes to chemotherapy: comparing views of patients with cancer with those of doctors, nurses and the general public. *British Medical Journal* 1990;**300**:1458-60.

Squires 1993

Squires TS, Tong T. Cancer Statistics. *Ca: a Cancer Journal for Clinicians* 1993;**43**:7-26.

Stewart 1992

Stewart THM, Raman S. Adjuvant treatment of non-small cell lung cancer, a meta-analysis. In: Stewart THM, Wheelock EF editor(s). Cellular immune mechanisms and tumour dormancy. Boca Raton, Florida: CRC Press, 213-38.

Stott 1977

Stott H, Fox W, Girling DJ, Stephens RJ, Dalton DAG. Acute leukaemia after busulphan. *British Medical Journal* 1977;**2**:1513-7.

Till 1992

Till JE, Sutherland HJ, Meslin EM. Is there a role for preference assessments in research on quality of life in oncology?. *Quality of Life Research* 1992;**1**:31-40.

Yusuf 1985

Yusuf S, Peto R, Lewis J, Collins R, Sleight P. Beta-blockade during and after myocardial infarction: An overview of the randomised trials. *Progress in Cardiovascular Diseases* 1985;**27**(5):335-71.

References to other published versions of this review
NSCLCCG 1995

Non-small Cell Lung Cancer Collaborative Group. Chemotherapy in non-small cell lung cancer: a meta-analysis using updated data on individual patients from 52 randomised clinical trials. *British Medical Journal* 1995;**311**:899-909.

* Indicates the major publication for the study

CHARACTERISTICS OF STUDIES
Characteristics of included studies [ordered by study ID]
a01 MRC LU02

Methods	RCT - 1965-68
Participants	643 patients
Interventions	surgery vs surgery + chemotherapy (i) cyclophosphamide* 200/75† (ii) busulphan 4*/1.5
Outcomes	survival
Notes	daily treatment 2 years

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a02 VASAG

Methods	RCT - 1968-73
Participants	443 patients
Interventions	surgery vs surgery + chemotherapy

Chemotherapy for non-small cell lung cancer (Review)

a02 VASAG (Continued)

 (i) cyclophosphamide 40‡
 (ii) cyclophosphamide 40‡
 methotrexate 50§

Outcomes	survival
Notes	15 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a03 EORTC 08741a

Methods	RCT - 1973-79
Participants	146 patients
Interventions	surgery vs surgery + chemotherapy lomustine* 70 cyclophosphamide 1000 methotrexate 40
Outcomes	survival
Notes	13 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a04 VASOG 5

Methods	RCT - 1973-79
Participants	841 patients
Interventions	surgery vs surgery + chemotherapy lomustine* 70 nitrogen mustard* 2000
Outcomes	survival
Notes	lomustine in 9 cycles nitrogen mustard in 52 cycles

Risk of bias
Chemotherapy for non-small cell lung cancer (Review)

a04 VASOG 5 (Continued)

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a05 WPL 7351

Methods	RCT - 1974-76
Participants	72 patients
Interventions	surgery vs surgery + chemotherapy lomustine* 130
Outcomes	survival
Notes	17 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a06 OLCSG 1a

Methods	RCT - 1982-87
Participants	321 patients
Interventions	surgery vs surgery + chemotherapy tegafur* 600-800\$
Outcomes	survival
Notes	daily treatment >1 year

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a07 OLCSG 1b

Methods	RCT - 1982-86
Participants	83 patients

Chemotherapy for non-small cell lung cancer (Review)

a07 OLCSG 1b (Continued)

Interventions	surgery vs surgery + chemotherapy doxorubicin 100\$ mitomycin C 20\$ tegafur(a)* 600-800\$ tegafur(b)* 600-800\$
Outcomes	survival
Notes	doxorubicin given in 3 cycles tegafur (a) daily treatment tegafur (b) daily treatment > 1 year

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a08 SGACLC 1

Methods	RCT - 1982-85
Participants	306 patients
Interventions	surgery vs surgery + chemotherapy mitomycin C 0.08‡ cyclophosphamide 2‡ tegafur* 12‡
Outcomes	survival
Notes	mitomycin C given in 10 cycles tegafur daily treatment > 6 months

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a09 WJSG 2

Methods	RCT - 1985-88
Participants	323 patients
Interventions	surgery vs surgery + chemotherapy (i) cisplatin 50 vindesine 6-9\$ UFT* 400\$ (ii) UFT* 400\$

Chemotherapy for non-small cell lung cancer (Review)

a09 WJSG 2 (Continued)

Outcomes	survival
Notes	1 cycle of cisplatin / vindesine given UFT daily treatment 1 year

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a10 LCSG 801

Methods	RCT - 1980-86
Participants	283 patients
Interventions	surgery vs surgery + chemotherapy cisplatin 60 doxorubicin 40 cyclophosphamide 400
Outcomes	survival
Notes	4 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a11 OLCSG 1c

Methods	RCT - 1982-87
Participants	28 patients
Interventions	surgery vs surgery + chemotherapy cisplatin 80 tegafur* 600-800\$
Outcomes	survival
Notes	cisplatin given in 1 cycle tegafur daily treatment > 1 year

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

a11 OLCSG 1c *(Continued)*

Allocation concealment (selection bias)	Low risk	A - Adequate
--	----------	--------------

a12 FLCSG 1

Methods	RCT - 1982-87
Participants	110 patients
Interventions	surgery vs surgery + chemotherapy cisplatin 40 doxorubicin 40 cyclophosphamide 400
Outcomes	survival
Notes	6 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a13 SGACLC 2

Methods	RCT - 1985-87
Participants	332 patients
Interventions	surgery vs surgery + chemotherapy cisplatin 66 doxorubicin 26 UFT* 8‡
Outcomes	survival
Notes	Unpublished UFT daily treatment > 6 months

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a14 IPCR, Chiba

Methods	RCT - 1985-91
Participants	29 patients
Interventions	surgery vs surgery + chemotherapy cisplatin 80 vindesine 3 mitomycin c¶ 8
Outcomes	survival
Notes	>2 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a15 LCSG 853

Methods	RCT - 1985-89
Participants	188 patients
Interventions	surgery vs syrgery + chemotherapy cisplatin 60 doxorubicin 40 cyclophosphamide 400
Outcomes	survival
Notes	Unpublished 4 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

a16 JLCSSG

Methods	RCT - 1986-88
Participants	209 patients
Interventions	surgery vs surgery + chemotherapy cisplatin 80 vindesine 6

Chemotherapy for non-small cell lung cancer (Review)

a16 JLCSSG (Continued)

Outcomes	survival
Notes	2-3 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

b01 EORTC 08741b

Methods	RCT - 1973-79
Participants	139 patients
Interventions	surgery + radiotherapy vs surgery + radiotherapy + chemotherapy lomustine* 70 cyclophosphamide 1000 methotrexate 40
Outcomes	survival
Notes	13 cycles of chemotherapy radiotherapy 45 Gy in 14-25 fractions complete resection

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

b02 LCSG 791

Methods	RCT - 1979-85
Participants	172 patients
Interventions	surgery + radiotherapy vs surgery + radiotherapy + chemotherapy cyclophosphamide 400 doxorubicin 40 cisplatin 40
Outcomes	survival
Notes	6 cycles of chemotherapy radiotherapy 40 Gy in 10 fractions** incomplete resection

b02 LCSG 791 (Continued)

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

b03 MSKCC 80-53

Methods	RCT - 1981-87
Participants	72 patients
Interventions	surgery + radiotherapy vs surgery + radiotherapy + chemotherapy cisplatin 120 vindesine 9
Outcomes	survival
Notes	4 cycles of chemotherapy radiotherapy 46 Gy complete and incomplete resections

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

b04 FLCSG 3

Methods	RCT - 1982-87
Participants	86 patients
Interventions	surgery + radiotherapy vs surgery + radiotherapy + chemotherapy cyclophosphamide 400 doxorubicin 40 cisplatin 40
Outcomes	survival
Notes	unpublished 8 cycles of chemotherapy, 2 given before radiotherapy radiotherapy 55 Gy in 20 fractions** incomplete resection

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

b04 FLCSG 3 (Continued)

Allocation concealment (selection bias)	Low risk	A - Adequate
--	----------	--------------

b05 GETCB 01CB82

Methods	RCT - 1982-86
Participants	267 patients
Interventions	surgery + radiotherapy vs surgery + radiotherapy + chemotherapy doxorubicin 40 vincristine 1.2 cisplatin 75 lomustine 80\$ alternating with cyclophosphamide 600
Outcomes	survival
Notes	3 cycles of chemotherapy given before radiotherapy radiotherapy 60-65 Gy in 30-33 fractions complete and incomplete resection

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

b06 OLCSG 1d

Methods	RCT - 1983-87
Participants	49 patients
Interventions	surgery + radiotherapy vs surgery + radiotherapy + chemotherapy cisplatin 80 tegafur* 600-800
Outcomes	survival
Notes	cisplatin given once tegafur daily treatment radiotherapy 40 Gy in 20 fractions complete resection

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

b07 EORTC 08861

Methods	RCT - 1986-90
Participants	22 patients
Interventions	surgery + radiotherapy vs surgery + radiotherapy + chemotherapy cisplatin 100 vindesine 6
Outcomes	survival
Notes	unpublished 4 cycles of chemotherapy, 2 given before radiotherapy radiotherapy 56 Gy in 28 fractions complete resection

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c01 NRH NSC 26271

Methods	RCT - 1968-71
Participants	74 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cyclophosphamide 400†† cyclophosphamide* 100
Outcomes	survival
Notes	oral cyclophosphamide given as daily treatment until tumour progression or toxicity radiotherapy 50 Gy in 25-31 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c02 EORTC 08742

Methods	RCT - 1973-80
Participants	117 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy

Chemotherapy for non-small cell lung cancer (Review)

c02 EORTC 08742 (Continued)

cyclophosphamide 1000
 lomustine* 100
 methotrexate 40

Outcomes survival

Notes 12 cycles of chemotherapy
 radiotherapy 50-60 Gy

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

c03 RTOG 7302 a

Methods RCT - 1973-78

Participants 111 patients

Interventions radical radiotherapy vs radical radiotherapy + chemotherapy
 cyclophosphamide 1000

Outcomes survival

Notes chemotherapy given until tumour progression or toxicity
 radiotherapy 40 Gy in 10 fractions**

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

c04 RTOG 7302 b

Methods RCT - 1973-78

Participants 96 patients

Interventions radical radiotherapy vs radical radiotherapy + chemotherapy
 cyclophosphamide 1000

Outcomes survival

Notes chemotherapy given until tumour progression or toxicity
 radiotherapy 30 Gy in 10 fractions

Risk of bias
Chemotherapy for non-small cell lung cancer (Review)

c04 RTOG 7302 b (Continued)

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c05 RTOG 7302 c

Methods	RCT - 1973-78
Participants	104 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cyclophosphamide 1000
Outcomes	survival
Notes	chemotherapy given until tumour progression or toxicity radiotherapy 40 Gy in 20 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c06 MCL-1

Methods	RCT - 1980-84
Participants	52 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy doxorubicin 40 lomustine* 30 cyclophosphamide 400 methotrexate 30
Outcomes	survival
Notes	chemotherapy given until tumour progression or toxicity radiotherapy 55 Gy in 25 fractions**

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c07 Aviano

Methods	RCT - 1980-84
Participants	111 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy doxorubicin 40 cyclophosphamide 600 methotrexate 30 procarbazine* 1000
Outcomes	survival
Notes	12 cycles of chemotherapy radiotherapy 45 Gy in 15 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c08 AZ-OC-1-80

Methods	RCT - 1981-85
Participants	52 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy vinblastine 6
Outcomes	survival
Notes	chemotherapy given until tumour progression or toxicity radiotherapy 55 Gy in 28 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c09 Gwent 3

Methods	RCT - 1981-85
Participants	85 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy etoposide* 1000

c09 Gwent 3 (Continued)

Outcomes	survival	
Notes	unpublished 7 cycles of chemotherapy given radiotherapy 32 Gy in 8 fractions	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c10 SECSG 81 LUN375

Methods	RCT - 1981-85	
Participants	212 patients	
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy vindesine(a) 3 vindesine(b) 6	
Outcomes	survival	
Notes	5 cycles of vindesine(a) then 10 cycles of vindesine(b) radiotherapy 60 Gy in 33 fractions	
Risk of bias		
Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c11 Gwent 1

Methods	RCT - 1974-76	
Participants	56 patients	
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy doxorubicin 50 fluorouracil 1200\$	
Outcomes	survival	
Notes	4 cycles of chemotherapy radiotherapy 32 Gy in 8 fractions	
Risk of bias		

c11 Gwent 1 (Continued)

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c12 SWOG 7635

Methods	RCT - 1977-79
Participants	62 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy doxorubicin 50
Outcomes	survival
Notes	8 cycles of chemotherapy radiotherapy 60 Gy in 20 fractions**

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c13 NCCTG 822451

Methods	RCT - 1983-87
Participants	121 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy doxorubicin 40 cyclophosphamide 400 methotrexate 40 lomustine* 30
Outcomes	survival
Notes	4 cycles of chemotherapy, 2 given before radiotherapy radiotherapy 60 Gy in 30 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c14 Buenos Aires

Methods	RCT - 1981-85
Participants	81 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cisplatin 40 doxorubicin 40 cyclophosphamide 400
Outcomes	survival
Notes	6 cycles of chemotherapy radiotherapy 55 Gy in 22 fractions**

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c15 Brussels

Methods	RCT - 1981-84
Participants	65 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cisplatin 60 etoposide 360 vindesine 3
Outcomes	survival
Notes	3 cycles of chemotherapy given before radiotherapy radiotherapy 55 Gy in 28 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c16 FLCSG 2

Methods	RCT - 1982-84
Participants	252 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cisplatin 40 doxorubicin 40

Chemotherapy for non-small cell lung cancer (Review)

c16 FLCSG 2 (Continued)

cyclophosphamide 400

Outcomes survival

 Notes 6 cycles of chemotherapy, 3 given before radiotherapy
 radiotherapy 55 Gy in 20 fractions**

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

c17 Essen

Methods RCT - 1983-87

Participants 48 patients

 Interventions radical radiotherapy vs radical radiotherapy + chemotherapy
 cisplatin 80
 vindesine 6

Outcomes survival

 Notes 3 cycles of chemotherapy given before radiotherapy
 radiotherapy 52-56 Gy in 13-14 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

c18 SLCSG

Methods RCT - 1983-89

Participants 327 patients

 Interventions radical radiotherapy vs radical radiotherapy vs chemotherapy
 cisplatin 120
 etoposide 300

Outcomes survival

 Notes 3 cycles of chemotherapy given before radiotherapy
 radiotherapy 56 Gy in 28 fractions

Risk of bias

c18 SLCSG (Continued)

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c19 CEBI 138

Methods	RCT - 1983-89
Participants	353 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cisplatin 100 cyclophosphamide 600 vindesine 3 lomustine* 75
Outcomes	survival
Notes	6 cycles of chemotherapy, 3 given before radiotherapy radiotherapy 65 Gy in 26 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c20 WSLCRG/FI

Methods	RCT - 1984-89
Participants	79 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cisplatin 100 vindesine 6
Outcomes	survival
Notes	8 cycles of chemotherapy, 2 given before radiotherapy radiotherapy 50 Gy in 20 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c21 Perugia

Methods	RCT - 1984-88
Participants	66 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cisplatin 100 etoposide 360
Outcomes	survival
Notes	3 cycles of chemotherapy given before radiotherapy radiotherapy 56 Gy in 30 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c22 CALGB 8433

Methods	RCT - 1984-87
Participants	180 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cisplatin 100 vinblastine 5
Outcomes	survival
Notes	2 cycles of cisplatin given before radiotherapy 5 cycles of vinblastine given before radiotherapy radiotherapy 60 Gy in 30 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c23 EORTC 08842

Methods	RCT - 1984-89
Participants	75 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy cisplatin 100

c23 EORTC 08842 (Continued)

vindesine 6

Outcomes	survival
Notes	unpublished 3 cycles of chemotherapy, 2 given before radiotherapy radiotherapy 55 Gy in 20 fractions**

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c24 SWOG 8300 a

Methods	RCT - 1984-88
Participants	128 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy fluorouracil 1200 vincristine 2 mitomycin C 10 alternating with cisplatin 40 doxorubicin 40 cyclophosphamide 400
Outcomes	survival
Notes	unpublished 6 cycles of chemotherapy, 2 given before radiotherapy radiotherapy 58 Gy in 29 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

c25 SWOG 8300 b

Methods	RCT - 1984-88
Participants	126 patients
Interventions	radical radiotherapy vs radical radiotherapy + chemotherapy fluorouracil 1200 vincristine 2 mitomycin C 10 cisplatin 40

Chemotherapy for non-small cell lung cancer (Review)

c25 SWOG 8300 b *(Continued)*

 doxorubicin 40
 cyclophosphamide 400

Outcomes survival

 Notes unpublished
 6 cycles of chemotherapy, 2 given before radiotherapy
 radiotherapy 58 Gy in 29 fractions

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

d01 Oxford

Methods RCT - 1970-73

Participants 188 patients

 Interventions supportive care vs supportive care + chemotherapy
 (i) procarbazine* 2.5† ††
 (ii) nitrogen mustard 0.3†
 vinblastine 0.5†
 procarbazine* 35†
 prednisolone 560†

Outcomes survival

 Notes (i) daily treatment for 1 year
 (ii) 11 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

d02 Quebec

Methods RCT - 1978-79

Participants 38 patients

 Interventions supportive care vs supportive care + chemotherapy
 methotrexate 40
 doxorubicin 40§§
 cyclophosphamide 400
 lomustine* 30

d02 Quebec (Continued)

Outcomes	survival
Notes	chemotherapy given until tumour progression or toxicity

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

d03 Gwent 2

Methods	RCT - 1982-84
Participants	186 patients
Interventions	supportive care vs supportive care + chemotherapy etoposide* 600
Outcomes	survival
Notes	6 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

d04 RLW 8351

Methods	RCT - 1982-86
Participants	167 patients
Interventions	supportive care vs supportive care + chemotherapy cisplatin 120 videsine 3
Outcomes	survival
Notes	chemotherapy given until tumour progression or toxicity

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

d05 NCIC CTG BR5

Methods	RCT - 1983-86
Participants	150 patients
Interventions	supportive care vs supportive care + chemotherapy (i) cisplatin 120 vindesine 3 (ii) cisplatin 40 doxorubicin 40 cyclophosphamide 400
Outcomes	survival
Notes	chemotherapy given until tumour progression or toxicity

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

d06 Southampton

Methods	RCT - 1983-86
Participants	32 patients
Interventions	supportive care vs supportive care + chemotherapy cisplatin 120 vindesine 3
Outcomes	survival
Notes	Same ref as RLW 8351 cisplatin given in 6 cycles vindesine given in 15 cycles

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

d07 NRH

Methods	RCT - 1983-87
Participants	87 patients

d07 NRH (Continued)

Interventions	supportive care vs supportive care + chemotherapy cisplatin 70 etoposide 100 etoposide* 400
Outcomes	survival
Notes	4 cycles of chemotherapy

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

d08 UCLA

Methods	RCT - 1984-86
Participants	63 patients
Interventions	supportive care vs supportive care + chemotherapy cisplatin 120 vinblastine 6
Outcomes	survival
Notes	chemotherapy given until tumour progression or toxicity

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

d09 Ancona 1

Methods	RCT - 1985-88
Participants	128 patients
Interventions	supportive care vs supportive care + chemotherapy cisplatin 80 cyclophosphamide 500 epirubicin 50 methotrexate 30 etoposide 200 lomustine* 70
Outcomes	survival

d09 Ancona 1 (Continued)

Notes chemotherapy given until tumour progression or toxicity

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

d10 AOI-Udine

Methods	RCT - 1984-86
Participants	102 patients
Interventions	supportive care vs supportive care + chemotherapy cisplatin 75 cyclophosphamide 400 mitomycin C 10
Outcomes	survival
Notes	6 cycles of chemotherapy given

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

d11 CEP-85

Methods	RCT - 1985-88
Participants	49 patients
Interventions	supportive care vs supportive care + chemotherapy cisplatin 120 vindesine 3
Outcomes	survival
Notes	cisplatin given in 8 cycles vindesine given in 18 cycles

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

Chemotherapy for non-small cell lung cancer (Review)

ya(surg) <=54 years

Methods	Subgroup analysis for age
Participants	
Interventions	
Outcomes	
Notes	Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yb(surg) 55-59 years

Methods	
Participants	
Interventions	
Outcomes	
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yc(surg) 60-64 years

Methods	
Participants	
Interventions	
Outcomes	
Notes	

Risk of bias

--	--

yc(surg) 60-64 years *(Continued)*

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yd(surg) >=65 years

Methods
Participants
Interventions
Outcomes
Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

ye(surg) male

Methods	Subgroup analysis for sex
Participants	
Interventions	
Outcomes	
Notes	Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yf(surg) female

Methods
Participants
Interventions

Chemotherapy for non-small cell lung cancer (Review)

yf(surg) female *(Continued)*

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yg(surg) good PS

Methods Subgroup analysis for performance status

Participants

Interventions

Outcomes

Notes Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yh(surg) poor PS

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yi(surg) adeno

Methods	Subgroup analysis for histology
Participants	
Interventions	
Outcomes	
Notes	Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yj(surg) squamous

Methods	
Participants	
Interventions	
Outcomes	
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yk(surg) other

Methods	
Participants	
Interventions	
Outcomes	
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

yk(surg) other (Continued)

Allocation concealment (selection bias)	Low risk	A - Adequate
--	----------	--------------

yl(surg) stage I,II

Methods	Subgroup analysis for stage
Participants	
Interventions	
Outcomes	
Notes	Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

ym(surg) stage III

Methods	
Participants	
Interventions	
Outcomes	
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yn(s+rt) <=54 years

Methods	Subgroup analysis for age
Participants	
Interventions	
Outcomes	

Chemotherapy for non-small cell lung cancer (Review)

yn(s+rt) <=54 years (Continued)

Notes Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yo(s+rt) 55-59 years

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yp(s+rt) 60-64 years

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yq(s+rt) >=65 years

Methods

Chemotherapy for non-small cell lung cancer (Review)

yq(s+rt) >=65 years (Continued)

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

yr(s+rt) male

Methods	Subgroup analysis for sex
---------	---------------------------

Participants

Interventions

Outcomes

Notes	Stratified analysis of data from a number of trials
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

ys(s+rt) female

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

yt(s+rt) good PS

Methods	Subgroup analysis for performance status
Participants	
Interventions	
Outcomes	
Notes	Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yu(s+rt) poor PS

Methods	
Participants	
Interventions	
Outcomes	
Notes	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yv(s+rt) adeno

Methods	Subgroup analysis for histology
Participants	
Interventions	
Outcomes	
Notes	Stratified analysis of data from a number of trials

Risk of bias

--	--

yv(s+rt) adeno *(Continued)*

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yw(s+rt) squamous

Methods
Participants
Interventions
Outcomes
Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yx(s+rt) other

Methods
Participants
Interventions
Outcomes
Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yy(s+rt) stage I,II

Methods	Subgroup analysis for stage
Participants	
Interventions	

Chemotherapy for non-small cell lung cancer (Review)

yy(s+rt) stage I,II *(Continued)*

Outcomes

Notes Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

yz(s+rt) stage III

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

za(radrt) <=54 years

Methods Subgroup analysis for age

Participants

Interventions

Outcomes

Notes Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zb(radrt) 55-59years

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zc(radrt) 60-64years

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zd(radrt) >=65 years

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

zd(radrt) >=65 years *(Continued)*

Allocation concealment (selection bias)	Low risk	A - Adequate
--	----------	--------------

ze(radrt) male

Methods	Subgroup analysis for sex	
Participants		
Interventions		
Outcomes		
Notes	Stratified analysis of data from a number of trials	

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zf(radrt) female

Methods		
Participants		
Interventions		
Outcomes		
Notes		

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zg(radrt) good PS

Methods	Subgroup analysis for performance status	
Participants		
Interventions		
Outcomes		

Chemotherapy for non-small cell lung cancer (Review)

zg(radrt) good PS *(Continued)*

Notes Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zh(radrt) poor PS

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zi(radrt) adeno

Methods Subgroup analysis for histology

Participants

Interventions

Outcomes

Notes Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zj(radrt) squamous

Methods

Chemotherapy for non-small cell lung cancer (Review)

zj(radrt) squamous *(Continued)*

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
--	----------	--------------

zk(radrt) other

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
--	----------	--------------

zl(radrt) stage I,II

Methods	Subgroup analysis for stage
---------	-----------------------------

Participants

Interventions

Outcomes

Notes	Stratified analysis of data from a number of trials
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
--	----------	--------------

Chemotherapy for non-small cell lung cancer (Review)

zm(radrt) stage III

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

zn(sc) <=54 years

Methods	Subgroup analysis for age
---------	---------------------------

Participants

Interventions

Outcomes

Notes	Stratified analysis of data from a number of trials
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

Allocation concealment (selection bias)	Low risk	A - Adequate
---	----------	--------------

zo(sc) 55-59 years

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias
Chemotherapy for non-small cell lung cancer (Review)

zo(sc) 55-59 years (Continued)

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zp(sc) 60-64 years

Methods
Participants
Interventions
Outcomes
Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zq(sc) >=65 years

Methods
Participants
Interventions
Outcomes
Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zr(sc) male

Methods	Subgroup analysis for sex
Participants	
Interventions	

Chemotherapy for non-small cell lung cancer (Review)

zr(sc) male (Continued)

Outcomes

Notes Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zs(sc) female

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zt(sc) good PS

Methods Subgroup analysis for performance status

Participants

Interventions

Outcomes

Notes Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zu(sc) poor PS

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zv(sc) adeno

Methods Subgroup analysis for histology

Participants

Interventions

Outcomes

Notes Stratified analysis of data from a number of trials

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zw(sc) squamous

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
------	--------------------	-----------------------

zw(sc) squamous *(Continued)*

Allocation concealment (selection bias)	Low risk	A - Adequate
--	----------	--------------

zx(sc) other

Methods

Participants

Interventions

Outcomes

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zy(sc)non-metastatic

Methods	Subgroup analysis for stage
---------	-----------------------------

Participants

Interventions

Outcomes

Notes	Stratified analysis of data from a number of trials
-------	---

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

zz(sc)metastatic

Methods

Participants

Interventions

Outcomes

Chemotherapy for non-small cell lung cancer (Review)

zz(sc)metastatic (Continued)

Notes

Risk of bias

Bias	Authors' judgement	Support for judgement
Allocation concealment (selection bias)	Low risk	A - Adequate

 drug doses given in mg/m² unless specified

* given orally

† after 10 days patients switched to maintenance chemotherapy. For first year only, drug doses were cyclophosphamide 150mg and busulphan 3mg

‡ dose in mg/kg

§ total dose

¶ mitomycin c was added to regimen from 1990

** split course of radiotherapy

†† daily during radiotherapy

†† dose escalating to 5mg/kg during weeks 3-6 then reduced to starting dose

 §§ stopped at total dose 450 mg/m²
Characteristics of excluded studies [ordered by study ID]

Study	Reason for exclusion
Aliev 1985	non-random use of radiotherapy
Bergsagel 1972	ELIGIBLE - IPD not available from trialist
Blaha 1973	mopidamol not established cytotoxic chemotherapy
Brunner 1971	ELIGIBLE - IPD not available from trialist
Buyze 1973	pre 1965 and pseudo-random (birthdate)
Byar 1978	prior chemotherapy allowed
Carr 1972	pre 1965, includes small cell lung cancer
Castberg 1976	pseudo-random (birthdate)
Crosbie 1966	pre 1965, pseudo-random
Dolton 1970	pre 1965, historical control
Durrant 1971	pre 1965
Hall 1967	pre 1965, prior chemotherapy allowed
Hansen 1973	RSV (1,2-diphenyl-ab-dicetone) not established cytotoxic chemotherapy
Helsper 1962	pre 1965, historical control
Holsti 1971	pre 1965, pseudo-random (birthdate)

Study	Reason for exclusion
Hosley 1962	pre 1965, used orthovoltage RT
Karageorgis 1991	confounded, RT dose not equal on each arm
Karp	confounded, RT dose not equal on each arm
Karrer 1978	ELIGIBLE - IPD not available from trialist
Kaung 1974	ELIGIBLE - IPD not available from trialist
NCCTG852451	Trial closed with no patients recruited
Newman 1985	razoxane not established cytotoxic chemotherapy
Okawa 1992	SPG (sizofiran) not established cytotoxic chemotherapy
Osterlind 1985	RSV (1,2-diphenyl-ab-dicetone) not established cytotoxic chemotherapy
Petrovic 1978	ELIGIBLE - IPD not available from trialist
Pirogov 1976	pseudo randomised
Privitera 1987	lonidamine not established cytotoxic chemotherapy
Scheer 1974	randomised all lung cancer, no histology
Selawry 1977	prior chemotherapy allowed
Slack 1970	pre 1965
Spatti 1985	used histological control
Spittle 1980	razoxane not established cytotoxic chemotherapy
Vincent 1975	pre 1965, only partly randomised
Wils 1984	confounded, RT dose different on each arm
Wingfield 1970	non-randomised
Wolf 1991	confounded, cisplatin used as a sensitiser on each arm

Characteristics of ongoing studies [ordered by study ID]

Ancona

Trial name or title	
Methods	
Participants	47
Interventions	supportive care vs supportive care + chemotherapy

Chemotherapy for non-small cell lung cancer (Review)

Ancona *(Continued)*

Outcomes

Starting date

Contact information

 Notes Ongoing at time of review to be included in update
ANITA

Trial name or title

Methods

 Participants 800 projected

 Interventions surgery versus surgery + chemotherapy

Outcomes

Starting date

Contact information

 Notes Ongoing, to be included in update
BLT

Trial name or title

Methods

 Participants 3300 projected

 Interventions
 (1)surgery + radiotherapy vs surgery + radiotherapy + chemotherapy
 (2) radiotherapy vs radiotherapy + chemotherapy
 (3)supportive care vs supportive care + chemotherapy

Outcomes

Starting date

Contact information

 Notes Ongoing, to be included in update
CAN NCIC BR10

Trial name or title

Chemotherapy for non-small cell lung cancer (Review)

CAN NCIC BR10 (Continued)

Methods	
Participants	600 projected
Interventions	surgery vs surgery + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Ongoing, to be included in update

CLB 9633

Trial name or title	
Methods	
Participants	500 projected
Interventions	surgery vs surgery + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Ongoing, to be included in update

CNR ALPI

Trial name or title	
Methods	
Participants	1200
Interventions	surgery vs surgery + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Closed, to be included in update

CRC-TU-LU3001

Trial name or title	
Methods	
Participants	466 projected
Interventions	radiotherapy vs radiotherapy + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Ongoing at time of review to be included in update

CRC-TU-LU3002

Trial name or title	
Methods	
Participants	359 projected
Interventions	supportive care vs supportive care + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Ongoing at time of review to be included in update

EORTC 08922

Trial name or title	
Methods	
Participants	34
Interventions	surgery vs surgery + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Ongoing at time of review, now closed, to be included in update

Chemotherapy for non-small cell lung cancer (Review)

FRE IALT

Trial name or title	
Methods	
Participants	3300 projected
Interventions	surgery vs surgery + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Ongoing, to be included in update

GAP

Trial name or title	
Methods	
Participants	200
Interventions	surgery vs surgery + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Ongoing at time of review, closed 92 to be included in update

Helsing

Trial name or title	
Methods	
Participants	48
Interventions	supportive care vs supportive care + chemotherapy
Outcomes	
Starting date	
Contact information	

Helsing (Continued)

Notes	Reported in 1998 after review published. To be included in update.
-------	--

Ichinose 1991

Trial name or title	
Methods	
Participants	86 patients
Interventions	surgery vs surgery + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Identified after meta-analysis published. To be included in update

Kim 1993

Trial name or title	
Methods	
Participants	101
Interventions	Radical RT versus radical RT + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Ongoing at the time of the review. To be included in update

NCCTG-822451

Trial name or title	
Methods	
Participants	120 projected
Interventions	radiotherapy vs radiotherapy + chemotherapy
Outcomes	

NCCTG-822451 *(Continued)*

Starting date

Contact information

Notes	Completed in 1993 (outside time boundary of review). To be included in update.
-------	--

RICUM

Trial name or title

Methods

Participants	242 173
--------------	------------

Interventions	(1)surgery vs surgery + chemotherapy (2)surgery + radiotherapy vs surgery + radiotherapy + chemotherapy
---------------	--

Outcomes

Starting date

Contact information

Notes	Ongoing at time of review,closed 92 to be included in update
-------	--

RTOG 8808

Trial name or title

Methods

Participants	327
--------------	-----

Interventions	radiotherapy vs radiotherapy + chemotherapy
---------------	---

Outcomes

Starting date

Contact information

Notes	Ongoing at time of review,closed 92 to be included in update
-------	--

TELCVIS 1999

Trial name or title

Methods

Chemotherapy for non-small cell lung cancer (Review)

TELCVIS 1999 (Continued)

Participants	191
Interventions	supportive care versus supportive care + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Reported in 1999 after review published. To be included in update

Thongprasert 1999


Trial name or title	
Methods	
Participants	288
Interventions	supportive care vs supportive care + chemotherapy
Outcomes	
Starting date	
Contact information	
Notes	Reported in 1996 after review published. To be included in update.


DATA AND ANALYSES
Comparison 1. surgery vs surgery + chemotherapy

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 survival	16	4457	Peto Odds Ratio (95% CI)	1.04 [0.96, 1.12]
1.1 long term alkylating agents	5	2145	Peto Odds Ratio (95% CI)	1.15 [1.04, 1.27]
1.2 other drugs	4	918	Peto Odds Ratio (95% CI)	0.89 [0.72, 1.11]
1.3 cisplatin based	8	1394	Peto Odds Ratio (95% CI)	0.87 [0.74, 1.02]
2 subgroup for survival - age	4		Peto Odds Ratio (95% CI)	Totals not selected
3 subgroup for survival - sex	2		Peto Odds Ratio (95% CI)	Totals not selected


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
4 subgroup for survival - performance status	2		Peto Odds Ratio (95% CI)	Totals not selected
5 subgroup for survival - histology	3		Peto Odds Ratio (95% CI)	Totals not selected
6 subgroup for survival - stage	2		Peto Odds Ratio (95% CI)	Totals not selected

Analysis 1.1. Comparison 1 surgery vs surgery + chemotherapy, Outcome 1 survival.







Analysis 1.2. Comparison 1 surgery vs surgery + chemotherapy, Outcome 2 subgroup for survival - age.




Analysis 1.3. Comparison 1 surgery vs surgery + chemotherapy, Outcome 3 subgroup for survival - sex.



Analysis 1.4. Comparison 1 surgery vs surgery + chemotherapy, Outcome 4 subgroup for survival - performance status.



Analysis 1.5. Comparison 1 surgery vs surgery + chemotherapy, Outcome 5 subgroup for survival - histology.



Analysis 1.6. Comparison 1 surgery vs surgery + chemotherapy, Outcome 6 subgroup for survival - stage.

Study or subgroup	Treatment n/N	Control n/N	Peto Odds Ratio 95% CI	Peto Odds Ratio 95% CI
yl(surg) stage I,II	180/485	222/488		0.78[0.64,0.94]
ym(surg) stage III	117/216	93/196		1.01[0.77,1.33]

Favours Treatment 0.1 0.2 0.5 1 2 5 10 Favours Control


Comparison 2. surgery + radiotherapy vs surgery + radiotherapy + chemotherapy

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 survival	7	807	Peto Odds Ratio (95% CI)	0.98 [0.83, 1.14]
1.1 long term alkylating agents	1	139	Peto Odds Ratio (95% CI)	1.35 [0.83, 2.20]
1.2 cisplatin based	6	668	Peto Odds Ratio (95% CI)	0.94 [0.79, 1.11]
2 subgroup for survival - age	4		Peto Odds Ratio (95% CI)	Totals not selected
3 subgroup for survival - sex	2		Peto Odds Ratio (95% CI)	Totals not selected
4 subgroup for survival - performance status	2		Peto Odds Ratio (95% CI)	Totals not selected
5 subgroup for survival - histology	3		Peto Odds Ratio (95% CI)	Totals not selected
6 subgroup for survival - stage	2		Peto Odds Ratio (95% CI)	Totals not selected


Analysis 2.1. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 1 survival.

Study or subgroup	Treatment n/N	Control n/N	Peto Odds Ratio 95% CI	Weight	Peto Odds Ratio 95% CI
2.1.1 long term alkylating agents					
a03 EORTC 08741a	33/66	40/73		10.72%	1.35[0.83,2.2]
Subtotal (95% CI)	66	73		10.72%	1.35[0.83,2.2]
Total events: 33 (Treatment), 40 (Control)					
Heterogeneity: Not applicable					
Test for overall effect: Z=1.21(P=0.23)					
2.1.2 cisplatin based					
b02 LCSG 791	68/82	75/90		23.42%	0.85[0.61,1.19]
b03 MSKCC 80-53	32/36	27/36		9.6%	1.23[0.74,2.06]
b04 FLCSG 3	34/40	42/46		12.48%	0.84[0.54,1.33]
b05 GETCB 01CB82	120/138	113/129		38.31%	0.95[0.73,1.23]
b06 OLCSG 1d	13/26	10/23		3.72%	1.02[0.45,2.34]
b07 EORTC 08861	5/10	7/12		1.77%	1.06[0.32,3.52]
Subtotal (95% CI)	332	336		89.28%	0.94[0.79,1.11]
Total events: 272 (Treatment), 274 (Control)					


Favours Treatment 0.1 0.2 0.5 1 2 5 10 Favours Control




Analysis 2.2. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 2 subgroup for survival - age.




Analysis 2.3. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 3 subgroup for survival - sex.




Analysis 2.4. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 4 subgroup for survival - performance status.



Analysis 2.5. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 5 subgroup for survival - histology.




Analysis 2.6. Comparison 2 surgery + radiotherapy vs surgery + radiotherapy + chemotherapy, Outcome 6 subgroup for survival - stage.




Comparison 3. radical radiotherapy vs radical radiotherapy + chemotherapy


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 survival	25	3033	Peto Odds Ratio (95% CI)	0.90 [0.83, 0.97]
1.1 long term alkylating agent	7	665	Peto Odds Ratio (95% CI)	0.98 [0.83, 1.16]
1.2 vinca alkaloids / etoposide	3	349	Peto Odds Ratio (95% CI)	0.87 [0.70, 1.09]
1.3 other drugs	3	239	Peto Odds Ratio (95% CI)	0.98 [0.74, 1.29]
1.4 cisplatin based	12	1780	Peto Odds Ratio (95% CI)	0.87 [0.79, 0.96]
2 subgroup for survival - age	4		Peto Odds Ratio (95% CI)	Totals not selected
3 subgroup for survival - sex	2		Peto Odds Ratio (95% CI)	Totals not selected
4 subgroup for survival - performance status	2		Peto Odds Ratio (95% CI)	Totals not selected
5 subgroup for survival - histology	3		Peto Odds Ratio (95% CI)	Totals not selected
6 subgroup for survival - stage	2		Peto Odds Ratio (95% CI)	Totals not selected

Analysis 3.1. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 1 survival.







Analysis 3.2. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 2 subgroup for survival - age.




Analysis 3.3. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 3 subgroup for survival - sex.




Analysis 3.4. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 4 subgroup for survival - performance status.




Analysis 3.5. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 5 subgroup for survival - histology.






Analysis 3.6. Comparison 3 radical radiotherapy vs radical radiotherapy + chemotherapy, Outcome 6 subgroup for survival - stage.




Comparison 4. supportive care vs supportive care + chemotherapy


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 survival	11	1190	Peto Odds Ratio (95% CI)	0.84 [0.74, 0.95]
1.1 long term alkylating agents	2	226	Peto Odds Ratio (95% CI)	1.26 [0.96, 1.66]
1.2 vinca alkaloids / etoposide	1	186	Peto Odds Ratio (95% CI)	0.87 [0.64, 1.20]
1.3 cisplatin based	8	778	Peto Odds Ratio (95% CI)	0.73 [0.63, 0.85]
2 subgroup for survival - age	4		Peto Odds Ratio (95% CI)	Totals not selected
3 subgroup for survival - sex	2		Peto Odds Ratio (95% CI)	Totals not selected
4 subgroup for survival - performance status	2		Peto Odds Ratio (95% CI)	Totals not selected
5 subgroup for survival - histology	3		Peto Odds Ratio (95% CI)	Totals not selected
6 subgroup for survival - stage	2		Peto Odds Ratio (95% CI)	Totals not selected

Analysis 4.1. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 1 survival.







Analysis 4.2. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 2 subgroup for survival - age.




Analysis 4.3. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 3 subgroup for survival - sex.




Analysis 4.4. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 4 subgroup for survival - performance status.



Analysis 4.5. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 5 subgroup for survival - histology.



Analysis 4.6. Comparison 4 supportive care vs supportive care + chemotherapy, Outcome 6 subgroup for survival - stage.





Comparison 5. RT + C vs Rt

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1 Survival	25	3055	Peto Odds Ratio (99% CI)	0.90 [0.83, 0.97]
1.1 Adjuvant Chemo	24	1235	Peto Odds Ratio (99% CI)	0.94 [0.83, 1.06]

Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
1.2 Neoadjuvant chemo	12	1820	Peto Odds Ratio (99% CI)	0.88 [0.80, 0.97]

Analysis 5.1. Comparison 5 RT + C vs Rt, Outcome 1 Survival.





ADDITIONAL TABLES

Table 1. Common meta-analysis stage scale

T	N	M	Meta-analysis Stage	AJC Stage
0,1,2,X,S	0	0	I	I
0,1,2,X,S	1	0	II	II
Any	2,3	0	III	III non-metastatic
3,4	Any	0	III	III non-metastatic
Any	Any	1	IV	Any metastatic

WHAT'S NEW

Date	Event	Description
10 March 2015	Review declared as stable	This review has been superseded by a new review "Adjuvant chemotherapy for resected early-stage NSCLC" (DOI: 10.1002/14651858.CD011430). It will no longer be updated.

HISTORY

Protocol first published: Issue 2, 2000
Review first published: Issue 2, 2000

Date	Event	Description
14 October 2010	Amended	Some information about the update of this review has been included in the results section.
7 September 2010	Amended	Some information about the update of this review has been included. Contact person has changed.
18 September 2008	Amended	Converted to new review format.
1 January 2000	New citation required and conclusions have changed	Substantive amendment

DECLARATIONS OF INTEREST

There is no known conflict of interest

SOURCES OF SUPPORT

Internal sources

- Medical Research Council, UK.
- Institute Gustave Roussy, France.

External sources

- INSERM 921204, France.
- ARC 2025, France.

NOTES

This review has been superseded by a new review from the same authors "Adjuvant chemotherapy for resected early-stage NSCLC" (DOI: 10.1002/14651858.CD011430). It will therefore no longer be updated. Please go to that review for the most up to date information.

Updated Results

As this systematic review is based on the original data from trials (not data taken from publications), updates are major projects taking many months of full time work and requiring the input of numerous individuals and groups. Standard practice for IPD meta-analyses are to undertake full updates, when appropriate, depending on the maturity of data and the rate at which further trials are completed and published.

This systematic review has now been updated to include more patients from more randomised controlled trials published since 1995. Details of current numbers of patients and trials included and the most up to date citations are listed below. Several new Cochrane Reviews will be submitted.

1. Surgery vs. surgery + adjuvant chemotherapy
 Updated to include: 8447 patients, 34 trial comparisons

Most recent citation: NSCLC Meta-analysis Collaborative Group. Adjuvant chemotherapy, with or without postoperative radiotherapy, in operable non-small-cell lung cancer: two meta-analyses of individual patient data. *Lancet*. 2010;375:1267-77.

2. Surgery + radiotherapy vs. surgery + radiotherapy + chemotherapy
 Updated to include: 2660 patients, 13 trial comparisons

Most recent citation: NSCLC Meta-analysis Collaborative Group. Adjuvant chemotherapy, with or without postoperative radiotherapy, in operable non-small-cell lung cancer: two meta-analyses of individual patient data. *Lancet*. 2010;375:1267-77.

3. Radiotherapy vs. radiotherapy + sequential chemotherapy

Updated to include: 3839 patients, 22 RCTs

Most recent citation: Le Pechoux C, Burdett S, Auperin A. Individual patient data (IPD) meta-analysis (MA) of chemotherapy (CT) in locally advanced non-small cell lung cancer (NSCLC). *Journal of Thoracic Oncology*. 2008; 3(Supplement 1):S20, 35IN.

Chemotherapy for non-small cell lung cancer (Review)

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

4. Supportive care vs. supportive care + chemotherapy

Updated to include: 2714 patients, 16 RCTs

Most recent citations:

Non-Small Cell Lung Cancer Collaborative Group. Chemotherapy and supportive care versus supportive care alone for advanced non-small cell lung cancer. *Cochrane Database of Systematic Reviews* 2010, Issue 5. Art. No.: CD007309. DOI: 10.1002/14651858.CD007309.pub2

NSCLC Meta-Analyses Collaborative Group. Chemotherapy in addition to supportive care improves survival in advanced non-small cell lung cancer: a systematic review and meta-analysis of individual patient data from 16 randomized trials. *J Clin Oncol.* 2008;26:4617-25.

Three other comparisons of timing of chemotherapy have also been explored:

a) Radiotherapy vs. radiotherapy + concomitant chemotherapy

Includes: 2910 patients, 16 RCTs

Most recent citation: Le Pechoux C, Burdett S, Auperin A. Individual patient data (IPD) meta-analysis (MA) of chemotherapy (CT) in locally advanced non-small cell lung cancer (NSCLC). *Journal of Thoracic Oncology.* 2008; 3(4, Supplement 1):S20, 35IN.

b) Radiotherapy + sequential chemotherapy vs. radiotherapy + concomitant chemotherapy

Includes: 1205 patients, 6 RCTs

Most recent citation: Aupérin A, Le Péchoux C, Rolland E, et al on behalf of the NSCLC Collaborative Group. Concomitant versus sequential radiochemotherapy in locally advanced non-small cell lung cancer: A meta-analysis of individual data of 1205 patients. *J Clin Oncol.* 2010;28:2181-90.

c) Surgery vs. surgery + neoadjuvant chemotherapy

Status: ongoing

INDEX TERMS

Medical Subject Headings (MeSH)

Antineoplastic Agents [*therapeutic use]; Carcinoma, Non-Small-Cell Lung [*drug therapy]; Lung Neoplasms [*drug therapy]; Meta-Analysis as Topic; Randomized Controlled Trials as Topic

MeSH check words

Humans