

Mycobacterium avium Bacilli Grow Saprozoically in Coculture with *Acanthamoeba polyphaga* and Survive within Cyst Walls

MICHAEL STEINERT,^{1*} KRISTIN BIRKNESS,² ELIZABETH WHITE,³
BARRY FIELDS,¹ AND FREDERICK QUINN²

Respiratory Diseases Branch, Division of Bacterial and Mycotic Diseases,¹ Tuberculosis/Mycobacteriology Branch,
Division AIDS, STD and TB Laboratory Research,² and Molecular Pathology and Ultrastructure Activity,
Division of Viral and Rickettsial Diseases,³ National Center for Infectious Diseases,
Centers for Disease Control and Prevention, Atlanta, Georgia 30333

Received 1 December 1997/Accepted 17 February 1998

Protozoans are gaining recognition as environmental hosts for a variety of waterborne pathogens. We compared the growth of *Mycobacterium avium*, a human pathogen associated with domestic water supplies, in coculture with the free-living amoeba *Acanthamoeba polyphaga* with the growth of *M. avium* when it was separated from amoebae by a 0.1- μ m-pore-size polycarbonate membrane (in a parachamber). Although viable mycobacteria were observed within amoebal vacuoles, there was no significant difference between bacterial growth in coculture and bacterial growth in the parachamber. This suggests that *M. avium* is able to grow saprozoically on products secreted by the amoebae. In contrast, *Legionella pneumophila*, a well-studied intracellular parasite of amoebae, multiplied only in coculture. A comparison of amoebae infected with *L. pneumophila* and amoebae infected with *M. avium* by electron microscopy demonstrated that there were striking differences in the locations of the bacteria within amoebal cysts. While *L. pneumophila* resided within the cysts, *M. avium* was found within the outer walls of the double-walled cysts of *A. polyphaga*. These locations may provide a reservoir for the bacteria when environmental conditions become unfavorable.

The facultative intracellular pathogen *Mycobacterium avium* is one of the primary health threats to patients with AIDS. It can cause bacteremia and disseminated multiorgan bacterial disease, including pulmonary infections of immunocompetent individuals (10, 15). The interaction of mycobacteria with host phagocytic cells likely is central to mycobacterial pathogenesis. Potential virulence mechanisms against phagocytic cells include prevention of the acidification of phagocytic vesicles and limited fusion of the phagosomes with the endosomal and lysosomal compartments that can lead to bacterial replication within macrophages (12, 13, 18). Within phagocytic cells *M. avium* bacilli tend to occupy individual vacuoles and to release superoxide dismutase (15) and the cell wall constituent liparabinomannan into the cytoplasm and into other lysosomal vesicles (34). Immunoelectron microscopy of macrophages infected with *M. avium* indicates that the parasitophoric vacuolar membrane possesses the late endosomal marker lysosome-associated membrane protein 1 but lacks the vesicular proton-ATPase (34).

The epidemiology of *M. avium* has been compared to that of *Legionella pneumophila* (8) because both organisms are widespread in aquatic environments, including municipal drinking water systems (9, 14, 16, 27, 32). Previous studies have proposed that amoebae might be environmental hosts of certain mycobacteria (4, 24). More recently, it has been shown that *M. avium* survives intracellularly within *Acanthamoeba castellanii* and interferes with the fusion of the lysosomal and parasitophoric vacuoles. In addition, the growth of *M. avium* in environmental amoebae results in increased virulence in the beige mouse model (8). These observations are reminiscent of the intracellular parasitism of *L. pneumophila* in the trophozoites

of a variety of free-living amoebae. Meanwhile, it is well-documented that *L. pneumophila* multiplies within amoebae and that *Acanthamoeba* cysts are able to protect legionellae from certain disinfection procedures (17, 22).

Since the biology of mycobacteria and the biology of legionellae have some significant similarities, we compared these two pathogens with regard to their interactions with amoebae. We confirmed that *M. avium* survives intracellularly and that growth occurs in coculture with amoebae. In contrast to *L. pneumophila*, *M. avium* exhibits saprophytic growth on products secreted by *Acanthamoeba* cells. However, since the mycobacteria are viable in trophozoites and cysts of *Acanthamoeba*, they might benefit from this interaction under adverse conditions.

MATERIALS AND METHODS

Bacterial strains and growth conditions. The *M. avium* serotype 4 strain used was a clinical isolate obtained from an AIDS patient (19). The strain was cultured on Middlebrook 7H11 agar (Difco) supplemented with oleic acid, albumin, dextrose, and catalase (BBL) for 10 days at 37°C. For amoeba infection assays, *M. avium* was grown in Middlebrook 7H9 broth (Difco) supplemented with albumin, dextrose, and catalase (BBL) to the late logarithmic phase with slow shaking (50 rpm) for 8 days at 37°C in the presence of 5% CO₂. *L. pneumophila* Philadelphia I JR32 was grown on buffered charcoal-yeast extract agar. The plates were incubated at 37°C in the presence of 5% CO₂ for 3 days. *Escherichia coli* HB101 was maintained on Luria-Bertani agar at 37°C.

***Acanthamoeba polyphaga*.** *Acanthamoeba polyphaga* ATCC 30872 was obtained from the American Type Culture Collection and was maintained axenically at room temperature in PYG 712 broth [2% proteose peptone, 0.1% yeast extract, 0.1 M glucose, 4 mM MgSO₄, 0.4 M CaCl₂, 0.1% sodium citrate dihydrate, 0.05 mM Fe(NH₄)₂(SO₄)₂ · 6H₂O, 2.5 mM NaH₂PO₃, 2.5 mM K₂HPO₃] as monolayers in 75-cm² tissue culture flasks. Amoebae were suspended by tapping the flasks. Cell counts were determined with a modified Fuchs-Rosenthal chamber. Amoebae were subcultured at intervals of 10 days.

Infection of *A. polyphaga* monolayers and parachamber experiments. *L. pneumophila* and *E. coli* were suspended in *Acanthamoeba* buffer (see below) and centrifuged at the maximum speed in an Eppendorf microcentrifuge, and the pellets were washed twice in *Acanthamoeba* buffer. Broth cultures of *M. avium* were likewise centrifuged and washed twice in buffer. After washing, *M. avium* and *L. pneumophila* were incubated in *Acanthamoeba* buffer for 10 days (33°C, 5% CO₂) without shaking to deplete the stored nutrients in the bacteria. This

* Corresponding author. Mailing address: Respiratory Diseases Branch, Division of Bacterial and Mycotic Diseases, Centers for Disease Control and Prevention, 1600 Clifton Road, Atlanta, GA 30333. Phone: (404) 639-0855. Fax: (404) 639-4215. E-mail: zma7@cdc.gov.

10-day starvation procedure was necessary since the residual growth rate of *M. avium* in buffer was very high. After this step the bacterial suspensions were vortexed for 1 min and adjusted in *Acanthamoeba* buffer to a concentration of 10^8 cells/ml, as determined by optical density at 600 nm.

Axenic cultures of *A. polyphaga* in the logarithmic phase were centrifuged ($200 \times g$, 15 min), and the pellets were resuspended in *Acanthamoeba* buffer (PYG 712 medium without proteose peptone, yeast extract, sodium citrate, and glucose). After two repetitions of this washing step, the cells were adjusted to a titer of 10^5 cells per ml in this buffer. One milliliter of this amoebal suspension was pipetted into each well of a 24-well microtiter plate (Costar, Cambridge, Mass.). Following 2 h of incubation at 33°C, the amoeba monolayer was inoculated with $10 \mu\text{l}$ of a bacterial suspension (10^5 bacteria/ml), which resulted in a preparation containing 10^3 bacteria/ml (multiplicity of infection, 10^{-2}). The numbers of CFU were determined at time zero and after 1, 7, and 14 days. In a control experiment the multiplication of the bacteria in buffer without amoebae was determined. To minimize evaporation, the plates were sealed in plastic bags.

In the parachamber experiment the bacteria were washed as described above, and 1 ml of a suspension containing 10^3 bacteria/ml in *Acanthamoeba* buffer was pipetted into each well of a 24-well microtiter plate (Costar). After insertion of a parachamber (diameter, 6.5 mm; transwell pore size, 0.1 μm ; Costar) into each well, $200 \mu\text{l}$ of an amoebal suspension containing 5×10^5 cells/ml in *Acanthamoeba* buffer was pipetted into each transwell. Colony counts were determined at time zero and after 1, 7, and 14 days. As described above for the *Acanthamoeba* infection, control experiments in buffer were performed and precautions to prevent evaporation were taken. In addition, the separation of bacteria and amoebae by the 0.1- μm -pore-size polycarbonate membrane filter was confirmed by viable cell counting and by light microscopy of the contents of the lower chamber.

Localization of intracellular bacteria and viability testing of *M. avium*. One milliliter of an *Acanthamoeba* cell suspension (10^5 cells/ml in *Acanthamoeba* buffer) was inoculated with a minimal volume containing 10^7 bacteria (in *Acanthamoeba* buffer) to yield a multiplicity of infection of 100. At 1, 6, 24, and 48 h the amoeba monolayer was washed twice with 1 ml of phosphate-buffered saline to eliminate noningested bacteria. To localize intracellular bacteria, the cells were processed for acid-fast staining and electron microscopy. For electron microscopy the cells were fixed in 2.5% cacodylate-buffered glutaraldehyde, postfixed with 1% osmium tetroxide in 0.1 M cacodylate buffer (1 h), and embedded in Epon resin. Thin sections were stained with 2% uranyl acetate and lead citrate and examined with a transmission electron microscope (Phillips Electronic Instruments, Mahwah, N.J.) at 40 kV.

To determine the viability of intracellular bacteria, a Baclight Live/Dead kit (Molecular Probes, Junction City, Oreg.) was used as described by the manufacturer. Briefly, after 2 days of incubation at 33°C, infected acanthamoebae (see above) were mounted on glass slides. Saline ($100 \mu\text{l}$) was placed on the air-dried cells, and $1 \mu\text{l}$ of SYTO 9 nucleic acid stain and $1 \mu\text{l}$ of a propidium iodide solution were suspended in the saline. SYTO 9 is a nucleic acid stain that labels bacterial cells with green fluorescence; propidium iodide, a red fluorescent nucleic acid stain, penetrates only bacteria with damaged membranes and effectively competes with SYTO 9 for nucleic acid binding sites. Thus, damaged cells were identified by red fluorescence, while live cells were identified by green fluorescence. The preparation was incubated in the dark for 15 min and then examined by fluorescence microscopy.

Synchronous encystment of infected *A. polyphaga* cells. The experiment to study infection of *A. polyphaga* monolayers was set up as described above for the coculture experiment except that the mycobacterial inoculum used was larger (final concentration, 10^7 bacteria/ml). After incubation for 24 h at 33°C, the supernatant was discarded, and the amoeba monolayer was rinsed with encystment buffer [0.1 M KCl, 0.02 M tris(2-amino-2-hydroxymethyl)-1,3-propanediol, 8 mM MgSO_4 , 0.4 mM CaCl_2 , 1 mM NaHCO_3] and then incubated in fresh encystment buffer at 33°C. After 3 days, the cell suspension was centrifuged ($1,000 \times g$, 20 min), and the pellet was resuspended in 3% (vol/vol) hydrochloric acid. This acid treatment was sufficient to kill the remaining trophozoites, immature cysts, and extracellular bacteria after 36 h. During the treatment the percentages of viable amoebae and bacteria were determined by Trypan blue exclusion and plating on Middlebrook 7H11 agar, respectively. After the acid treatment the cysts were washed three times with *Acanthamoeba* buffer. One-half of the sample was processed for electron microscopy (see above), and the other half was incubated in PYG medium at 33°C for 7 days. The excystment of the cysts was examined by light microscopy, and the presence of viable bacteria was determined by viable counting on Middlebrook 7H11 agar plates.

RESULTS

Growth of bacteria in *Acanthamoeba* buffer. Despite intensive washing with buffer, unstarved mycobacteria showed significant residual growth in buffer after 10 days of incubation (Table 1). The increases in CFU per milliliter were 72-fold for the unstarved bacilli and 6-fold for the starved mycobacteria. In contrast, the number of CFU of *L. pneumophila* was almost unaffected by starvation (0.3-fold decrease for unstarved legio-

TABLE 1. Growth of unstarved and starved bacteria in *Acanthamoeba* buffer

Strain	Growth increment in <i>Acanthamoeba</i> buffer ^a	
	Unstarved bacteria	Starved bacteria ^b
<i>M. avium</i> serotype 4	72.2 ± 59.6	6.1 ± 6.3
<i>L. pneumophila</i> JR32	0.31 ± 0.13	0.72 ± 0.24
<i>E. coli</i> HB101	0.049 ± 0.067	ND ^c

^a Number of CFU per milliliter after 10 days divided by initial number of CFU per milliliter (mean ± standard deviation from four independent experiments). The preparations were incubated at 33°C.

^b Starved *M. avium* and *L. pneumophila* cells were incubated in nutrient-free buffer for 10 days to deplete the stored nutrients in the bacteria.

^c ND, not determined.

nellae and 0.7-fold decrease for starved legionellae). The viability of unstarved *E. coli* decreased 0.049-fold. Although media and growth conditions are known to influence virulence, there was no obvious difference in the initial uptake by amoebae between starved and unstarved bacteria (data not shown).

Direct coculture and parachamber culture. *E. coli* and starved cells of *M. avium* and *L. pneumophila* were tested for the ability to grow in coculture with *A. polyphaga*. The results of this growth assay are shown in Fig. 1. For *M. avium* the number of bacteria increased from 1.5×10^3 CFU in the inoculum to 3×10^5 CFU after 7 days and to 4×10^6 CFU after 14 days. In monolayers infected with *L. pneumophila* the number of bacteria increased from 5×10^2 CFU in the inoculum to 3×10^5 CFU after 2 days to 2×10^7 CFU after 14 days. *E. coli* also multiplied in coculture; the number of bacteria increased from 866 CFU to 1.6×10^6 CFU within 1 day.

Parachamber experiments were used to determine if the amoeba-associated growth of the bacteria was due to intracellular parasitism of *A. polyphaga* (Fig. 2). *M. avium* and *E. coli* showed approximately the same growth kinetics as those observed in the direct coculture experiment. In contrast, *L. pneumophila* showed no increase in CFU when it was separated from the amoebae by the 0.1- μm -pore-size polycarbonate membrane. Due to the low-nutrient-content environment the number of amoebal cysts increased during the course of the experiment.




FIG. 1. Growth of the *M. avium* serotype 4 strain, *L. pneumophila* JR32, and *E. coli* HB101 in cocultures with *A. polyphaga*. Each coculture experiment was performed three times in *Acanthamoeba* buffer at 33°C. The values shown are the mean numbers of CFU (± standard deviations) determined at zero time and at 1, 7, and 14 days after coincubation was started.




FIG. 2. Growth of the *M. avium* serotype 4 strain, *L. pneumophila* JR32, and *E. coli* HB101 when they were separated from *A. polyphaga* by a 0.1- μ m-pore-size polycarbonate membrane (parachamber). Each parachamber experiment was performed three times in *Acanthamoeba* buffer at 33°C. The values shown are the mean numbers of CFU (\pm standard deviations) determined at zero time and at 1, 7, and 14 days after inoculation.

Intracellular survival. Acid-fast staining, Gimenez staining, and transmission electron microscopy were used to demonstrate bacterial ingestion by *A. polyphaga* during coculture. These studies confirmed previous observations that great num-

bers of legionellae are found intracellularly inside a single vacuole. The numbers of legionellae increase over time, and finally the bacteria fill the entire host cell. In contrast, the numbers of intracellular *M. avium* cells were much lower (1 to 20 bacteria per vacuole), and the bacteria were found in several vacuoles (one to six vacuoles per cell) (Fig. 3). The viabilities of intracellular bacteria could be confirmed by using differential live-dead fluorescence staining (Fig. 4). The percentage of infected amoebae increased over time, while the number of intracellular mycobacteria within a single host cell remained constant after 2 days of cocultivation. This observation suggests that the primary mechanism of mycobacterial growth in coculture is not intracellular. The very few amoebae that contained *E. coli* cells contained one to three bacteria in a single vacuole.

***M. avium* within *Acanthamoeba* cysts.** The ability of *L. pneumophila* to survive within the cysts of *A. polyphaga* was previously demonstrated by Kilvington and Price (22). To determine whether *M. avium* and *E. coli* have similar capabilities, the encystment of amoebae after coculture with these species was induced by incubation in encystment buffer. Six hours after cyst induction the trophozoites rounded up, and encystment of a trophozoite appeared to be complete after 18 h. After 3 days about 92% of the trophozoites produced mature cysts (as determined by the presence of a double wall). The upper temperature limit that permits encystment is 40°C, but optimal numbers of infected cysts were found at 33°C. In order to kill




FIG. 3. Transmission electron micrograph of *M. avium* serotype 4 bacilli within cytoplasmic vesicles of an *A. polyphaga* trophozoite after 2 days of cocultivation. Bar = 1 μ m.




FIG. 4. Differential live-dead fluorescence staining of *M. avium* serotype 4 bacilli within vesicles of *A. polyphaga* trophozoites after 2 days of coinubation. Green fluorescence indicates live bacterial cells that have an intact membrane.

the remaining trophozoites, immature cysts, and extracellular bacteria, a treatment with hydrochloric acid (3%, vol/vol) was performed. Examination of thin sections by transmission electron microscopy revealed double-walled cysts containing one to nine mycobacteria within the amoeba outer cell wall (Fig. 5). *E. coli* was not found within cysts. Our observations that *L. pneumophila* was within the cysts but was not between the cell walls of the cysts were consistent with the description given in the previous study. After 7 days in PYG medium the *Acanthamoeba* cysts were able to excyst and replicate. Legionellae and mycobacteria (at lower numbers), but not *E. coli*, were culture positive on growth plates after the excystment of the amoebae.

DISCUSSION

The importance of protozoans in soil and water ecosystems has been recognized for several decades, and the relevance of predatory protozoans in the control of bacterial populations is widely acknowledged (11, 20, 30, 31). However, the potential role of protozoans as reservoirs for human pathogens has only recently received adequate attention (2, 4, 17, 25). The best-studied example of a protozoan-bacterial pathogen interaction is the intracellular parasitism of *L. pneumophila*. Adaptation of *L. pneumophila* to parasitism of free-living amoebae might have led to the ability of this environmental bacterium to infect human lung macrophages (4). This observation led to the concepts that selection for resistance to digestion by predatory

protozoans is a driving force in the evolution of pathogenic environmental bacteria and that protists are the "missing link between ecology and pathology" (4, 23, 25).

Since *M. avium* has been isolated from habitats where amoebae normally feed on bacteria (14, 26, 29, 32), we studied the interaction of *A. polyphaga* with *M. avium*. Our results show that *M. avium* does not behave in a manner analogous to *L. pneumophila*. Since *L. pneumophila* only grows intracellularly within amoebae, the parasitism of legionellae seems to be much more specific and more highly evolved than the simple growth enhancement of mycobacteria and *E. coli* by their association with amoebae. In pure buffer (parachamber experiments) *M. avium* and *E. coli* were able to grow as free-living saprophytes on products secreted by *A. polyphaga*. Although we demonstrated that the growth of *M. avium* in direct cocultures and the growth of *M. avium* in parachamber experiments are similar, we cannot eliminate the possibility that intracellular multiplication occurs. Consistent with results published recently (8), we found that *M. avium* is able to survive in the intracellular environment of amoebae, and consequently, these bacteria may receive nutrients and grow within the protozoans. Considering that *M. avium* is a slowly growing bacterium that lives in a habitat where amoebae feed on bacteria, survival in a hostile intracellular environment may be an important advantage.

Although there are important differences in the interactions of amoebae with legionellae and mycobacteria, intracellular




FIG. 5. Transmission electron micrograph of a mature *A. polyphaga* cyst containing *M. avium* serotype 4 bacilli (arrow) within the double cell wall (note that the outer cell wall is divided and surrounds the bacterial cells). Bar = 1 μ m.

survival provides a possible evolutionary explanation for how a saprozoic-saprophytic or parasitic environmental organism can acquire the ability to survive within human macrophages. *M. avium* and *L. pneumophila* are known to inhibit phagolysosomal vacuole fusion in amoebae and macrophages (5, 8, 13, 18, 21). Other defense mechanisms, such as bacterial toxin production, the presence of protective outer membrane structures, and the reduction of vacuole acidification by the bacteria, might also contribute to resistance in phagocytic cells. A comparison of the intracellular activities that occur in protozoans after ingestion of bacteria showed that these activities are very similar to the activities observed in macrophages (1, 3, 5, 7, 8, 25, 33).

It is generally accepted that engulfed bacteria may benefit from the protective coat conferred by protozoans (23). The presence of mycobacteria in domestic water supplies and the results of disinfection studies in which the authors tested various concentrations of sodium hypochlorite suggest that chlorination has little effect on mycobacteria (6, 9, 27). Previous studies have demonstrated that cysts of *A. polyphaga* can contain viable *L. pneumophila* cells which are protected from disinfection (22). Recovery of *M. avium* from HCl-treated cysts showed that the exploitation of this amoebal differentiation event is not restricted to *L. pneumophila*. Therefore, we suggest that the resistance of infected amoebal cysts to biocidal agents may additionally interfere with disinfection of domestic water supplies contaminated with mycobacteria. However, in contrast to *L. pneumophila*, *M. avium* is located within the

double walls of the cysts, which are known to be composed largely of polysaccharides (one-third is cellulose) (28). The numbers of bacterial cells in the cysts are much lower than the numbers of legionellae, and it is unclear how mycobacteria are distributed to this location.

To control and prevent the dissemination of nontuberculous mycobacterial infections, it is important to focus on the ecology of the bacterial environment. Although no data are available, protozoans like *Hartmannella* and *Naegleria* spp. very likely exhibit interactions with mycobacteria identical to those observed with acanthamoebae. The results of these and other studies in which pathogens were enhanced by associations with protozoans and perhaps biofilms indicate that there is a need for further research on the physiological ecology of these pathogens (16).

ACKNOWLEDGMENT

This work was supported by grant Ste 838/1-1 from the Deutsche Forschungsgemeinschaft.

REFERENCES

1. Allen, P. G., and E. A. Dawidowicz. 1990. Phagocytosis in *Acanthamoeba*. I. A mannose receptor is responsible for the binding and phagocytosis of yeast. *J. Cell. Physiol.* **145**:508-513.
2. Amman, R., N. Springer, W. Schonhuber, W. Ludwig, E. N. Schmid, K. D. Muller, and R. Michel. 1997. Obligate intracellular bacterial parasites of acanthamoebae related to *Chlamydia* spp. *Appl. Environ. Microbiol.* **63**:115-121.
3. Avery, S. V., J. L. Harwood, and D. Lloyd. 1995. Quantification and charac-

- terization of phagocytosis in the soil amoeba *Acanthamoeba castellanii* by flow cytometry. *Appl. Environ. Microbiol.* **61**:1124–1132.
4. **Barker, J., and M. R. W. Brown.** 1994. Trojan horses of the microbial world: protozoa and the survival of bacterial pathogens in the environment. *Microbiology* **140**:1253–1259.
 5. **Bozue, J. A., and W. Johnson.** 1996. Interaction of *Legionella pneumophila* with *Acanthamoeba castellanii*: uptake by coiling phagocytosis and inhibition of phagosome-lysosome fusion. *Infect. Immun.* **64**:668–673.
 6. **Carson, L. A., N. J. Petersen, M. S. Favero, and S. M. Agüero.** 1978. Growth characteristics of atypical mycobacteria in water and their comparative resistance to disinfectants. *Appl. Environ. Microbiol.* **36**:839–846.
 7. **Cianciotto, N. P., and B. S. Fields.** 1992. *Legionella pneumophila mip* gene potentiates intracellular infection of protozoa and human macrophages. *Proc. Natl. Acad. Sci. USA* **89**:5188–5191.
 8. **Cirillo, J. D., S. Falkow, L. S. Tompkins, and L. E. Bermudez.** 1997. Interaction of *Mycobacterium avium* with environmental amoebae enhances virulence. *Infect. Immun.* **65**:3759–3767.
 9. **Collins, C. H., J. M. Grange, and M. D. Yates.** 1984. A review: mycobacteria in water. *J. Appl. Bacteriol.* **57**:193–211.
 10. **Crowle, A. J., E. R. Ross, D. L. Cohn, J. Gilden, and M. H. May.** 1992. Comparison of the abilities of *Mycobacterium avium* and *Mycobacterium intracellulare* to infect and multiply in cultured human macrophages from normal and human immunodeficiency virus-infected subjects. *Infect. Immun.* **60**:3697–3703.
 11. **Curds, C. R., and G. J. Fey.** 1969. The effect of ciliated protozoa on the fate of *E. coli* in the activated-sludge process. *Water Res.* **3**:853–867.
 12. **De Chastellier, C., C. Frehel, C. Offredo, and C. Skamene.** 1993. Implication of phagosome-lysosome fusion in restriction of *Mycobacterium avium* growth in bone marrow macrophages from genetically resistant mice. *Infect. Immun.* **61**:3775–3784.
 13. **de Chastellier, C., T. Lang, and L. Thilo.** 1995. Phagocytic processing of the macrophage endoparasite *Mycobacterium avium* in comparison to phagosomes which contain *Bacillus* or latex beads. *Eur. J. Cell Biol.* **68**:167–182.
 14. **du Moulin, G. C., K. D. Stottmeier, P. A. Pelletier, A. Y. Tsang, and J. Hedley-Whyte.** 1988. Concentration of *Mycobacterium avium* by hospital hot water systems. *JAMA* **260**:1599–1601.
 15. **Escuyer, V., N. Haddad, C. Frehel, and P. Berche.** 1996. Molecular characterization of a surface-exposed superoxide dismutase of *Mycobacterium avium*. *Microb. Pathog.* **20**:41–55.
 16. **Falkinham, J. O.** 1996. Epidemiology of infection by nontuberculous mycobacteria. *Clin. Microbiol. Rev.* **9**:177–215.
 17. **Fields, B. S.** 1996. The molecular ecology of legionellae. *Trends Microbiol.* **4**:286–290.
 18. **Frehel, C., C. De Chastellier, T. Lang, and N. Rastogi.** 1986. Evidence for inhibition of fusion of lysosomal and prelysosomal compartments with phagosomes in macrophages infected with pathogenic *Mycobacterium avium*. *Infect. Immun.* **52**:252–262.
 19. **Gan, H., G. W. Newman, and H. G. Remold.** 1995. Human macrophages acquire a hyporesponsive state of tumor necrosis factor alpha production in response to successive *Mycobacterium avium* serovar 4 stimulation. *Infect. Immun.* **63**:1921–1926.
 20. **Habte, M., and M. Alexander.** 1977. Further evidence for the regulation of bacterial populations in soil by protozoa. *Arch. Microbiol.* **113**:181–183.
 21. **Horwitz, M. A.** 1983. The Legionnaires' disease bacterium (*Legionella pneumophila*) inhibits phagosome-lysosome fusion in human monocytes. *J. Exp. Med.* **158**:2108–2126.
 22. **Kilvington, S., and J. Price.** 1990. Survival of *L. pneumophila* within cysts of *Acanthamoeba polyphaga* following chlorine exposure. *J. Appl. Bacteriol.* **68**:519–525.
 23. **King, C. H., E. B. Shotts, R. E. Wooley, and K. G. Porter.** 1988. Survival of coliforms and bacterial pathogens within protozoa during chlorination. *Appl. Environ. Microbiol.* **54**:3023–3033.
 24. **Krishna-Prasad, B., and S. K. Gupta.** 1978. Preliminary report on engulfment and retention of mycobacteria by trophozoites of axenically grown *Acanthamoeba castellanii* Douglas. *Curr. Sci.* **47**:245–247.
 25. **Ly, T. M. C., and H. E. Müller.** 1990. Ingested *Listeria monocytogenes* survive and multiply in protozoa. *J. Med. Microbiol.* **33**:51–54.
 26. **Ma, P., G. S. Visvesvara, A. J. Martinez, F. H. Theodore, P. M. Daggett, and T. K. Sawyer.** 1990. Naegleria and Acanthamoeba infections: review. *Rev. Infect. Dis.* **12**:490–513.
 27. **Mollohan, C. S., and M. S. Romer.** 1961. Public health significance of swimming pool granuloma. *Am. J. Public Health* **51**:883–891.
 28. **Neff, R. J., S. A. Ray, W. F. Benton, and M. Wilborn.** 1964. Induction of synchronous encystment (differentiation) in *Acanthamoeba* sp., p. 55–83. *In* D. M. Prescott (ed.), *Methods in cell physiology*, vol. 1. Academic Press, London, United Kingdom.
 29. **Schulze-Robbecke, R.** 1993. Mykobakterien in der Umwelt. *Immun. Infekt.* **21**:126–131.
 30. **Singh, B. N.** 1941. Selectivity in bacterial food by soil amoebae in pure and mixed culture in sterilized soil. *Ann. Appl. Biol.* **28**:52–65.
 31. **Stout, J. D.** 1973. The relationship between protozoan populations and biological activity in soils. *Am. Zool.* **13**:191–201.
 32. **von Reyn, C. F., J. N. Maslow, T. W. Barber, J. O. Falkinham III, and R. D. Arbeit.** 1994. Persistent colonisation of potable water as a source of *Mycobacterium avium* infection in AIDS. *Lancet* **343**:1137–1141.
 33. **Wintermeyer, E., B. Ludwig, M. Steinert, B. Schmitt, G. Fischer, and J. Hacker.** 1995. Influence of site-specific altered Mip proteins on intracellular survival of *Legionella pneumophila* in eukaryotic cells. *Infect. Immun.* **63**:4576–4583.
 34. **Xu, S., A. Cooper, S. Sturgill-Koszycki, T. van Heyningen, D. Chatterjee, L. Orme, P. Allen, and D. G. Russel.** 1994. Intracellular trafficking in *Mycobacterium tuberculosis* and *Mycobacterium avium*-infected macrophages. *J. Immunol.* **153**:2568.