




Awareness and Willingness to use Condoms and Preexposure Prophylaxis among Gay, Bisexual, and Other Cisgendered Men who Have sex with men in Slum Communities in Ghana. BSGH-004

Journal of the International
Association of Providers of AIDS Care
Volume 22: 1-12
© The Author(s) 2023
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/23259582231209649
journals.sagepub.com/home/jia



Gamji Rabi Abu-Ba'are, PhD^{1,2,3,4}, Osman Wumpini Shamrock, PhD^{1,2} , Amos Apreku, MSc⁵ , George Rudolph Kofi Agbemedu, BA⁶, Edem Yaw Zigah, BA⁶, Oliver C. Ezechi, PhD⁷, LaRon E. Nelson, PhD^{3,8} , and Kwasi Torpey, PhD⁵

Abstract

Introduction: Research has begun to examine human immunodeficiency virus (HIV) prevention strategies within the Ghanaian context. Still, little is known about specific populations such as gay, bisexual, and other cisgender men who have sex with men (GBMSM) living in slum communities. We studied HIV prevention strategies such as condoms and pre-exposure prophylaxis (PrEP) in slum communities and the awareness and willingness to use these choices among GBMSM. This qualitative study examines HIV prevention strategies, specifically examining PrEP and condom use behaviors among GBMSM in Ghana. **Methods:** We conducted in-depth face-to-face interviews among 12 GBMSM from slums in Accra and Kumasi cities in Ghana. Data were analyzed through a summative content analysis with multiple reviewers to develop codes. Data were collected from participants in January 2022. **Results:** We found the fear and perceived risk of infection were motivators for consistent condom use, especially during anal sex. GBMSM living with HIV receiving antiretroviral therapy were more inclined to use condoms. We found motivations for using PrEP were influenced by the type of sexual activity and a history of negative HIV status. Also, the barriers to PrEP for GBMSM included limited access to healthcare facilities and the distance to these facilities. **Conclusions:** To improve condom and PrEP access and uptake, we recommend addressing structural barriers by increasing the number of health facilities and implementing targeted interventions to address the lack of information on HIV awareness and prevention. Involving peer educators may also effectively promote HIV prevention strategies, especially in communities with limited access to healthcare such as slums. Overcoming these access constraints could significantly enhance awareness and prevention of HIV, leading to improved health outcomes for GBMSM living in slum communities.

¹ School of Nursing, University of Rochester, Rochester, USA

² Behavioral, Sexual and Global Health Lab, School of Nursing, University of Rochester, Rochester, USA

³ Center for Interdisciplinary Research on AIDS, Yale School of Public Health, University of Ghana, Accra, Ghana

⁴ Department of Public Health Sciences, University of Rochester Medical Center, Rochester, New York, USA

⁵ Department of Population, Family and Reproductive Health, School of Public Health, University of Ghana, Accra, Ghana

⁶ Behavioral, Sexual and Global Health Lab, Accra, Ghana

⁷ Nigeria Institute of Medical Research, Lagos, Nigeria

⁸ School of Nursing, Yale University, New Haven, Connecticut, USA

Corresponding Author:

Osman Wumpini Shamrock, PhD, School of Nursing, University of Rochester, Rochester, USA.

Email: osmanwumpini_shamrock@urmc.rochester.edu



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access page (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Keywords

bisexual, condom, HIV, PrEP, gay

Date received: 17 August 2023; revised: 3 October 2023; accepted: 6 October 2023.

Introduction

Globally, strategies to prevent human immunodeficiency virus (HIV) have been a significant focus in many countries.^{1–4} According to the World Health Organization [WHO]⁵, the number of people living with HIV at the close of 2021 reached an estimated 38.4 million with 0.7% of this population aged 15–49 years. While this number continues to rise over the years, the WHO (2022) report shows the African continent has been severely affected with an estimated (3.4%) 1 in 35 adults living with HIV. These projections position the African continent to account for two-thirds of all cases worldwide. Within Africa, the Sub-Saharan region is known to have the highest affected rate of HIV infection, accounting for an estimated 70% of HIV globally.^{6,7} Given these statistics, the calls for strategic steps to curb the high epidemic incidence have increased, particularly within Africa's Sub-Saharan region.^{8–10}

Like many other African countries, Ghana has adopted some intervention strategies into its policy such as The U.S. President Emergency Plan For AIDS Relief (PEPFAR) and the (UNAIDS) 90–90–90 to reduce the epidemic spread of HIV.^{11–13} PEPFAR was a United States-funded initiative to support and expand Ghana's efforts to prevent the spread of HIV/AIDS, care for persons living with HIV, and provide support to improve coverage of the country's response to the epidemic.^{11,14–16} Ghana adopted the (UNAIDS) 90-90-90 achieve the United Nations Programme (90% of all people who are HIV-positive know their status, 90% of those diagnosed with HIV are on treatment, and 90% of those diagnosed with HIV are virally suppressed).^{12,17–19} These policy efforts among other efforts from the Ministry of Health, Ghana AIDS Commission and other community and nongovernmental organization have contributed to the low-level epidemic rate in the country.

Among many interventional strategies to prevent the spread of HIV, administering preexposure prophylaxis (PrEP) has been identified as a practical way to reduce incidence.^{20–22} PrEP can be presented as a daily medication to prevent a diagnosed HIV-negative person from getting infected.²³ PrEP is the use of antiretrovirals in individuals who do not have HIV to prevent infection. PrEP is taken in anticipation of exposure to HIV and is mainly recommended for individuals at increased risk of exposure.²⁴ PrEP comes in many forms and can be administered (oral or injectable) through various therapeutic options such as using tenofovir/emtricitabine in combination with maraviroc, using long-acting rilpivirine administered every 8 weeks, using cabotegravir/vocabria, which is an Integrase Strand Transfer Inhibitor, and using VRC-HIVMAB060-00-AB.²⁵ Effective administration of antiretrovirals can support individuals living with HIV to

achieve a state where the virus has been suppressed to undetectable levels, low enough to prevent transmission.^{26,27} Meaning the virus cannot be transmitted through sexual contact.^{26–29} This scientific evidence is based on the concept of U=U (undetectable equals untransmittable).^{26–30} Despite the evidence showing U=U significance in reducing the risk of transmission, many have stressed that it does not eliminate the risk of infection completely, emphasizing the need for regular testing and safer sexual practices such as condom use.^{31,32} U=U significance has also been questioned, as it does not prevent other sexually transmitted infections.^{31–33}

While the PrEP strategy has proven effective in controlling the spread of HIV, it comes with many challenges. Awareness and usage of PrEP have been reportedly low in certain contexts,^{24,34–36} particularly among low- and middle-income countries.^{37,38} The uptake of antiretrovirals as an effective means of preventing the spread of HIV is reported to be globally low.^{20,39–41} Many other barriers to PrEP implementation and antiretroviral usage are seen in the stigma surrounding its use among individuals at increased risk of HIV, the possible drug-to-drug interaction, the risk of sexually transmitted infection, the cost of antiretrovirals, and its reported side effects.^{41–43}

Additionally, the cost of antiretrovirals has been shown to significantly affect PrEP implementation among key Ghanaian population.⁴⁴ Barriers related to the cost of antiretrovirals among key Ghana populations have been associated with long-term financial implications and poor health insurance coverage.⁴⁴ Other cost barriers related to antiretrovirals were associated with those incurred by the patients such as laboratory fees, travel-related expenses, and testing for sexually transmitted diseases.^{44–46} Recommendations to improve antiretrovirals uptake in Ghana include the government subsidizing the cost of the medication, or including it in the National Health Insurance Scheme.⁴⁴

Past studies have also shown insufficient knowledge of PrEP among MSM as a primary reason for its low acceptability, even when the antiretrovirals are presented as a reliable means of preventing the spread of HIV.^{47–49} In Ghana, available research indicates that PrEP rejection is high among individuals at an increased risk of infection.²⁰ According to Dako-Gyeke et al⁵⁰ and Kumar et al⁵¹, MSM in Ghana form a significant population (17.5%) of people living with HIV in the country and have a higher chance of not getting tested or engaging in HIV-related in treatment. These studies highlight an increasing need for MSM to explore preventive strategies when engaged in activities that increase their risk of infection. While PrEP implementation is coupled with many challenges within Sub-Saharan Africa,⁴⁸ understanding perceptions, awareness and preferences

among MSM is crucial to PrEP acceptance as a reliable strategy of HIV prevention.^{52–54} Given Ghana's conscious efforts to present PrEP as an alternative strategy to preventing the spread of HIV, leaders in the implementation drive remain hesitant due to the lack of context-based evidence from data to support a scale-up of antiretrovirals in the country.^{55,56}

Similarly, condom use as an intervention technique against the spread of HIV has been emphasized in many countries, especially within sub-Saharan Africa, where HIV continues to be high among sexual minorities.^{48,49,57,58} Condom use has been identified as one of the most effective means of preventing the spread of HIV.^{59,60} According to the United States Agency for International Development⁶¹, condoms are effective when used correctly to reduce HIV transmission by 90%. Among key populations such as gay men, condoms effectively reduce the spread of HIV by 70% for individuals with HIV-positive partners who reported consistent use, 72% for those who indicated they were bottoming, and 63% for those who were topping.⁶² Hence, many health providers and promoters urged its use, especially among individuals at an increased risk of infection.^{63–65} Several interventions have been employed in Ghana to encourage condom use.^{48,49,63,66,67} One of the major intervention approaches in the country to increase condom use, especially among individuals who are at an increased risk of infection in Ghana is the Ghana Nation Condom and Water Based Lubricant Programming Strategy 2014–2019. The intervention was a conscious approach by the government in collaboration with multiple nongovernmental and community-based organizations to identify barriers within Ghana that impacted condom use and employ strategies that addressed the needs of different segments of the Ghanaian population.⁶⁶ The report captures key populations (23%) as the second largest group in HIV new infection in the country.⁶⁶

Despite employing strategies such as the Ghana Nation Condom and Water Based Lubricant Programming Strategy 2014–2019, the National HIV and AIDS Strategic Plan (2011–2015), and the Ghana MARP Strategic Framework 2011–2015, among others, HIV among key populations, particularly MSM in Ghana continue to record increasing rate of infection.⁶⁸ The USAID recommends HIV prevention services such as condom use and PrEP be combined to maximize the odds of viral transmission since no barrier method can guarantee 100% prevention.

Slums in Ghana are known to be under-resourced, evident in the lack of health facilities, leading to adverse health outcomes.⁶⁹ Slums in Ghana are characterized by poor access to drinking water, schools, transportation services, and inadequate housing settlements.^{70–74} The individual or combination of these slum characteristics can affect how gay, bisexual and other cisgender men who have sex with men (GBMSM) access HIV prevention services. Understanding how these implications manifest for highly stigmatized groups such as GBMSM is imperative when suggesting interventions to increase HIV prevention strategies, particularly concerning PrEP, and condom use. This research was conducted in selected slum communities due to lacking information on HIV

prevention strategies for GBMSM living in these areas. We attempt to use this study to understand the current state of HIV prevention strategies for GBMSM in these slum communities and how the unique characteristic of slums in Ghana affects the utilization of HIV prevention strategies among GBMSM.

Ghana has been described as a country where the safety of GBMSM cannot be guaranteed as the laws of the country do not provide protection for LGBTQ groups or individuals or groups advocating for their wellbeing.⁷⁵ Ghana's laws currently criminalize same-sex penetrative behaviors.^{76,77} The country's current legislation on sexuality is vague; hence, the policing of sexual activities, particularly among GBMSM is left to the discretion of law enforcement officers.^{76,78} Ghanaian criminal law is fraught with ambiguity, resulting in individuals and law enforcement officials resorting to personal interpretation to determine its application to people in the LGBTQ community.⁷⁶ The phrase “unnatural carnal knowledge,” found in the Ghanaian Criminal Code Amendment Act of 2003, does not explicitly reference LGBTQ individuals.⁷⁶ However, it has been invoked to justify criminal assaults and the perpetuation of stigmatization against members of these communities.^{76,79,80}

The current stigmatizing climate in Ghana towards LGBTQ or individuals in close relation to these groups translates into challenges in health services.^{81,82} Accessing preventive services such as PrEP can be discouraging for these groups of individuals due to the stigma they face at the health facilities.⁸¹ Though calls have been made for the Government of Ghana to subsidize these rates or include them fully into the National Health Insurance Policy, the fear associated with high stigma may prevent GBMSM from accessing these services.⁴⁴

Methods

Research Design

The research used qualitative interviews⁸³ to understand the lived experiences of GBMSM in Ghanaian slum communities. We collected first-hand accounts from participants about their day-to-day experiences of HIV prevention strategies related to condoms and PrEP use in the slum communities. We also investigated how they interpreted their experiences concerning their sexual orientation and the barriers or willingness to HIV prevention.

Sampling and Recruitment Procedure

To reach and recruit GBMSM in slum communities, we partnered with our community organizations in Accra and Kumasi, using the time location sampling method.⁸⁴ A sampling method that has been proven effective in researching similar populations in the past.^{85–87} The time location sampling method in this study was purposively used due to the stigmatizing nature of GBMSM in Ghana. This approach allowed researchers reach the sample at locations and times that provided them the maximum amount of safety. Aside from providing security, the time location sampling method ensured diversity as GBMSM were randomly chosen from these

secured locations, ensuring equal chances of being included in the study. Additionally, multiple visits from the researchers to recruit participants at times that were not chosen by the research team increased the likelihood of recruiting a diverse pool of respondents into the study. Community partners and research assistants came from PORSH (Priorities on Rights and Sexual Health) in Accra and YAHR (Youth Alliance on Health and Human Rights) in Kumasi. Research assistants working with our community partner organizations screened and invited GBMSM to participate in interview sessions during one of the organizations' activities when GBMSM visited the site at times and locations that had been previously determined as safe. We have a long history of partnering with these two organizations to recruit participants and conduct studies among GBMSM in Ghana.^{48,49,88} Initially, we had planned to recruit 19 participants, but after the eighth interview, we reached saturation in responses to fill the information gap. We continued to capture four more participants to ensure complete saturation, bringing the final number of transcripts to 12. We decided not to conduct further interviews with our samples when we realized the data were repetitive, and information received from GBMSM were recurring, similar to those from previous participants. Participants who participated in the study were given 80 Ghana Cedis each as compensation to cater for their transportation to and from the interview site.

Inclusion Criteria

Before enrollment, all participants were 18 years or older and resided in a slum community within Ghana's Greater Accra regional capital. Participants self-identified as a cisgender man and fell under the GBMSM category having had sexual relations with other cisgender men for reasons other than sexual orientation in the past 6 months. The HIV status of participants did not affect their eligibility to be included in this study.

Data Collection

Procedure. We conducted in-depth face-to-face interviews to collect participant data.⁸⁹ Following screening, research assistants gave participants consent forms to read. The research assistants read the consent forms aloud to ensure clarity and gave additional explanations. Before beginning the interviews, research assistants responded to participants' questions and obtained their signatures to confirm their consent to participate and enable audio recording. All conversations took place in private spaces owned by the community partners. Four interviews were recorded in a local Ghanaian language called *Twi* because some participants found it challenging to communicate in English; all other interviews in the study were conducted in English. Data were collected in January 2022.

Nature of Questions. The checklist created for the study served as the basis for the qualitative interview training given to the

research assistants. Consistent with our design, the checklist encouraged free and open discussion over the more conventional question-and-answer interview format. Participants were invited to share personal narratives about their background and family, sex and gender expectations, experiences of stigma within the family, openness about sexuality and sexual behavior, acceptance within the family, and how they deal with stigma or live as GBMSM.

Analytical Strategy. The audio interview recordings were directly transcribed verbatim by trained research assistants, who also deidentified the transcripts by removing details that could be used to identify the participants. After that, we conducted a summative content analysis with multiple reviewers on the transcripts.⁹⁰ We have effectively used this analytical procedure to understand essential elements in participant accounts.⁴⁹ We allocated at least two reviewers to each transcript. Each reviewer examined the interview checklist before reviewing the transcripts to find the most important points brought up by the participants, which they then independently reported in between 100 and 200 words. The first author went over each summary and compiled its key points into a data spreadsheet, which helped identify clusters in the qualitative data and highlighted the factors that frequently came up in transcripts and summaries.

Ethical Approval and Informed Consent. IRB approvals were obtained from the University of Rochester (IRES IRB #RNI00002010) and the Ghana Health Service (GHS-ERC 001/10/21) approved the study. Before any information was collected, the study's interviewers ensured that every participant had read and understood the informed consent form and given approval for information to be collected on HIV prevention strategies in condom use and PrEP.

Results

Description of Participants

Six GBMSM reported being Christian, two were Muslim, and the remaining four practiced a combination of both religions. Concerning formal education, only one had an education level below a Junior High School, six had completed Senior High School Education, and five had earned a university degree. Six participants reported being unemployed, while the other six worked part-time or full-time jobs.

Description of Categories and Subcategories

Two categories emerged from the study. The first category was condom use determinants. Under this category were subcategories: (a) familiarity, trust, and longevity, (b) fear and perceived risk of infection, and (c) preventing HIV transmission to others. The second category our data analyzed was PrEP use determinants. Subcategories under this team were (a) motivators to use PrEP and (b) access constraints to PrEP.

Condom Use Determinants

We categorized condom use drivers into three subcategories. The first was familiarity, trust, and longevity, the second was the fear and perceived risk of infection, and the third was preventing HIV transmission to others.

Familiarity, Trust, and Longevity. Our interviews with participants indicated the decision to use condoms was complex and influenced by various factors, including familiarity with partners, trust, and longevity with an intimate partner. Participants reported they prioritize safety and use condoms mostly. Still, they may be persuaded to have sex without condoms if they have been with the same partner for a while and the partner convinces them to trust them. Another participant shared that they sometimes chose not to use condoms when they were strongly attracted to someone and wanted a more intimate experience.

For safety measures, I use condoms mostly, but when I have sex continuously, and the person is trying to tell me to trust him because he's not flirting around, then I give in for raw sex (GBMSM participant I).

Sometimes but not always, though. It gets to when you are attracted to someone you really like and want to feel the person. That's when I decide not to use the condom. But not everyone I allow to go raw like that (GBMSM participant H).

Fear and Perceived Risk of Infection. Participants expressed a strong commitment to condom use, stating they use condoms every time they engage in anal sex. However, some participants reported using condoms only when necessary, as they had regular partners and did not engage in high-risk sexual behaviors. The risk of sexual infections also motivated participants to consistently use condoms during anal sex, citing previous experiences with gonorrhea as a motivator for consistent condom use.

I always use condoms and I use them every time. I engage in anal sex. But during sex, what I ensure that the sex is safe for me without any harm, is that I use condoms. I usually will use condoms. But let me say I will give it like 80% condom use. Because the world is a little scary, you don't know when you will contract something. (GBMSM participant A)

I engage in anal sex. I'm always on top. I actually use a condom. But I don't move around a lot. I have a regular partner (GBMSM participant F)

I always go for a condom during sex. I do anal sex. There was a time I ended up having gonorrhea and from that time no matter what you claim to be, whoever you are, you have to wear a condom. (GBMSM participant C)

Preventing HIV Transmission to Others. The results showed some participants who reported being HIV-positive still used condoms during intimate relationships. Participants who disclosed their HIV-positive status mentioned using

condoms consistently after learning about their situation and being advised by a healthcare provider on the risk of spreading HIV. Participants mentioned insisting on using a condom when their intimate partners wanted to engage in condomless sex. Results also showed participants preferred to move away from current intimate partners who were insisting on condomless sex to those who preferred condoms use for the sole purpose of preventing further infections.

I love sex. At first, not so much, but now yes, Because I am on ARTs, I use condoms. If I am with someone who doesn't want condoms, I will leave and do it with another person who is okay with condoms. As for condom use, 100%. (GBMSM participants G)

Because I am on ARTs, we use condoms. We use gold circles (condom brand). Previously, I had sex with no condoms depending on my partner. When I got to know my status, I was always advised by a health person to use a condom and I have been practicing that. (GBMSM participant B)

PrEP use Determinants

The results show PrEP presented a significant role in HIV prevention for GBMSM. Determinants of PrEP use were captured within two subcategories. The first was the motivators to use PrEP. The second was Access constraints to PrEP use.

Motivators to use PrEP. GBMSM were motivated to adhere to PrEP due to the nature of sex they engaged with their intimate partners. Other actors, such as peer educators in the lives of GBMSM were instrumental in how they decided to adhere to PrEP use by encouraging them to get tested for HIV to keep being eligible for PrEP. Other reasons cited by GBMSM were the nature of how they engaged in sexual activities and the associated risk of acquiring HIV.

I have engaged in anal sex, hence, take PrEP every day. I'm a top now even though I used to be at the bottom, you don't know when you will contract something. (GBMSM participant D)

I'm HIV-negative, hence, use PrEP. I have a peer educator in my community which I have been randomly doing tests with him. And I'm on the safe side. (GBMSM participant E)

Yes, I'd like to be on it because of how rough I can be at times during sex. I'm not on PrEP. But I know what PrEP is and would like to be on it. (GBMSM participant K)

Access Constraints to PrEP use. GBMSM indicated structural barriers such as the lack of health facilities and distance to a health facility limited their PrEP access. Responses from our participants show if these structural challenges were tackled adequately, it would go a long way to encourage access and adherence to PrEP. GBMSM also indicated that additional constraints connected to time prevented them from accessing anti-retrovirals from individuals who had the drug available. According to participants, the inability to find the time for medication pickups meant staying on other sexual protective mechanisms to prevent infection. Another significant finding from

GBMSM showed the challenge of not knowing where to go for PrEP presented as a constraint GBMSM living in slums mentioned as a reason for not taking the medication.

I used to be on PrEP but I have stopped because of the distance to the health facility. I will continue if there is a facility close by to pick up the drug. (GBMSM participant N)

I know PrEP. A friend told me about it when I was in school. I thought he could get some for me, but he said unless I come there. But I haven't had time to go, so I make sure I protect myself. I know my status. (GBMSM participant K)

I'm not on PrEP. But I know what PrEP is and would like to be a part of it. I have

heard about it, but I don't know how to get some. (GBMSM participant J)

Discussion

Living in slums are filled with multiple challenges.^{70,91,92} Empirical information in Ghana on PLHIV in slums is not extensive, making it difficult to understand the recent experiences of PLHIV and HIV prevention strategies among GBMSM. This study revealed GBMSM familiarity, trust, longevity with intimate partners, the fear and perceived risk of infection and preventing HIV transmission to others significantly affected condom use. The study also identified GBMSM engaging in anal sex and a history of testing negative for HIV as motivations for PrEP use. Structural barriers such as inadequate health facilities, the limited proximity to services, and time constraints associated with obtaining PrEP limited its use.^{45,46}

Previous studies in Africa have shown that strong relationships may affect condom use negotiations due to the desire to seek emotional intimacy with a partner.^{93,94} During our interviews, we found familiarity, trust, and longevity emerging as significant determinants when deciding whether or not to use a condom. This finding was similar to those by Peasant et al⁹⁵, who found that self-sacrificing was a strategy to earn a partner's trust, or nurture a romantic relationship that could result in less condom use among sexually active partners. Other studies have found proposing condom use, especially among gay men can be perceived as less romantic, with condomless sex seen as a way of showing trust and emotional connection.^{94,96} Though not a focus in this study, familiarity, and trust has emerged as a significant factor in HIV transmission, especially in the areas of U = U (undetectable = untransmittable) with findings showing 42.3% trusted it, 19.8% did not trust it, and 38.0% were not sure.^{30,97,98} We found condoms use in Ghanaian slum communities was also determined by the longevity with intimate partners, the sexual attractiveness of the partner, and the partner's characteristics. Participants who engaged in sexual activity consistently with a partner were likely to develop trust and stop using condoms. Our findings highlight the role of familiarity, trust, and longevity in

condom use decisions among GBMSM. Our results also suggest consistent condom use may be challenging for individuals engaged in long-term relationships or regular partners. Findings on familiarity, trust, and longevity to consistent condom use and long-term support the call by earlier researchers to investigate the attitudes of GBMSM when considering interventions in HIV/STI in the country.^{10,68,99,100}

Studies in Africa have shown that HIV-positive individuals appear to use condoms more consistently after knowing their status, with the main aim of not getting new infections or transmitting to their intimate partners.^{101,102} Previous research has also indicated that GBMSM were more likely to use condoms if they knew about their HIV-positive status, were directed by health professionals, or knew where to access condoms.^{103,104} Our participants on antiretroviral therapies (ARTs) appeared to be more consistent with condom use after learning about their HIV-positive status. They also assumed the responsibility of ensuring no further infections of HIV by either insisting on condom use with their partners or breaking off with individuals who did not want to engage in condomless sex. Consistent condom use among HIV-positive GBMSM was also driven by advice from healthcare professionals as a critical step in preventing the spread of HIV. When paying attention to the lack of infrastructure in health facilities and services (counseling, HIV testing, condom provision, and use education) and health personnel for GBMSM in HIV prevention, we find these results informative in tailoring interventions for HIV prevention among GBMSM living in Ghanaian slum communities.

Previous research has shown that individuals who are at an increased risk of infection, such as men who have sex with men stand a higher chance of HIV transmission through sexual intercourse.^{105–107} Studies have also indicated anal sexual intercourse has higher chances of transmitting HIV compared to vaginal sex.^{107,108} While highlighting the risk of anal sex in HIV transmission, some studies show other infections, such as gonorrhea, can present during sexual activities among sexual minorities.¹⁰⁹ In our study, the risk of HIV infection through anal sex for GBMSM came up as a vital determinant for condom use. Anal intercourse for GBMSM was perceived as a sexual activity that could put one at significant risk of HIV infection. The fear associated with infection and further spread of HIV through anal sexual intercourse among GBMSM reportedly showed consistent condom use among participants. Some participants also indicated that the high rates of HIV infections globally put enough fear in them to encourage consistent condom use. Closely related to this finding was GBMSM history of STI and constant use of condoms. Our finding suggests a historical experience of an STI such as gonorrhea can have profound implications on sexual practices, leading to a more cautious and protective sexual lifestyle.¹¹⁰ Our findings demonstrate that the lack of health facilities and personnel for GBMSM living in slum communities could increase the risk of HIV infection if interventions are not tailored, providing these amenities and making them accessible to these populations. Given the stigmatized context of slum communities,¹¹¹ sensitization campaigns surrounding condom

use in anal sex activities must be prioritized to sustain our current finding.^{48,49} We also urge bringing on board community stakeholders such as community religious leaders and political heads to reduce stigma and discrimination when these educational programs are organized around condom use and HIV prevention.^{112–114}

Safer sex practices in previous research among sexual minorities, particularly HIV-positive MSM, are primarily concerned with exposing other intimate partners to the risk of HIV infection.^{115,116} Our findings show having casual and/or multiple sex partners, the risk of contracting, and the fear of infecting a partner encouraged consistent condom use among GBMSM who engaged in sexual intimacy. According to Serovich et al¹¹⁷, personality traits among sexual minorities with HIV-positive status who had casual intimate partners played a significant role when insisting on or not using a condom for sexual intimacy. Our findings show GBMSM interviewed had developed a sense of personal responsibility to prevent their partner from contracting HIV and used condoms. These findings demonstrate to some extent how individual personality traits can play a role in condom use among GBMSM. The results also indicate higher condom use preference when the risk of infecting others or being infected is high among GBMSM. For GBMSM living in slum communities, the resource constraint of these areas, especially regarding health facilities and personnel could hinder services such as HIV/STI counseling in promoting favorable attitudes towards condom use in HIV prevention.

For GBMSM, PrEP uptake is a recommended preferred means of HIV prevention as PrEP effectively reduces the risk of HIV infection, especially among GBMSM.^{47,118} Whereas no studies were conducted in Ghana among GBMSM in slums on PrEP, the only study in the country on PrEP among the GBMSM showed a high willingness to take up PrEP.⁴⁷ This willingness has been further highlighted in research among MSM, who suggest antivirals be subsidized by the Government of Ghana or included fully in the National Health Insurance policy.⁴⁴ Consistently the GBMSM participants in our study were motivated to accept PrEP because of the HIV risk associated with same-sex sexual activities and the belief in the efficacy of antiretrovirals in preventing HIV infection.

Participants also described the role of peer educators as crucial in antiretroviral uptake and adherence through HIV testing to remain eligible for the drug. The role of peer educators has been stressed in multiple studies in PrEP follow-ups and uptake among at-risk individuals.^{119–121} Although no published work on PrEP and peer mentors in slums exist in Ghana, GBMSM peers have played a critical role in previous interventions that address HIV prevention and linkage to care GBMSM in Ghana.^{9,48,49,67,88} Despite the favorable attitude toward PrEP uptake among our participants, we urge the training of more peer educators to strengthen use and adherence. Due to the location of this study, we propose peer educators be trained from within the slum to reduce accessibility challenges, increase relatability and trust, and facilitate linkage to care for GBMSM in HIV prevention and care services.⁴⁸

Challenges associated with PrEP use are complex and range within the individual, community, and health systems.^{122,123} Our results about PrEP access and use were closely aligned with the challenges surrounding health systems.⁹ We found the proximity of PrEP access points limited access and use. Ahouada et al (2020) also arrived at these conclusions, indicating distance and sites for PrEP distribution significantly affected PrEP access and use in HIV prevention. Though we couldn't find previous literature on the case of GBMSM living in Ghanaian slums, we trust that properly addressing health system challenges could promote PrEP access and adherence in these communities.

Conclusion

Our findings show familiarity, trust, and the attractiveness and characteristics of GBMSM intimate partners influenced condom use in HIV prevention. Fear and perceived risk of infection came up as motivators for consistent condom use, particularly when the mode of sexual intimacy was primarily anal sex. Our findings also revealed GBMSM with HIV-positive status and receiving ARTs were inclined to condom use. The study shows motivations for PrEP use were influenced by the nature of sex GBMSM engaged in and a history of HIV negative status. PrEP use barriers included limited access to health-care facilities and the distance to these facilities.

Our findings suggest that efforts to improve PrEP access and uptake among GBMSM should prioritize addressing structural barriers, such as increasing the number of health facilities providing PrEP and developing targeted interventions to address time constraints and lack of information about antiretrovirals. Additionally, engaging peer educators may effectively promote PrEP use among GBMSM, especially in communities with limited access to health facilities. Addressing these access constraints could significantly enhance the uptake and adherence to PrEP to improve HIV prevention outcomes among GBMSM living in slum communities.


Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.


Funding

The authors disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by the Yale University FLAGS grant,

ORCID iDs

Osman Wumpini Shamrock  <https://orcid.org/0000-0003-3127-9596>

Amos Apreku  <https://orcid.org/0000-0001-8115-9125>

LaRon E. Nelson  <https://orcid.org/0000-0002-2630-602X>

References

1. Excler JL, Robb ML, Kim JH. Prospects for a globally effective HIV-1 vaccine. *Vaccine*. 2015;33. doi:10.1016/j.vaccine.2015.03.059

2. Gartner MJ, Roche M, Churchill MJ, Gorry PR, Flynn JK. Understanding the mechanisms driving the spread of subtype C HIV-1. *EBioMedicine*. 2020;53. doi:10.1016/j.ebiom.2020.102682
3. Mahamboro DB, Fauk NK, Ward PR, Merry MS, Siri TA, Mwanri L. HIV stigma and moral judgement: qualitative exploration of the experiences of HIV stigma and discrimination among married men living with HIV in Yogyakarta. *Int J Environ Res Public Health*. 2020;17(2): 636. doi:10.3390/ijerph17020636
4. Rotheram-Borus MJ, Davis E, Rezai R. Stopping the rise of HIV among adolescents globally. *Curr Opin Pediatr*. 2018;30(1): 131. doi:10.1097/MOP.0000000000000580
5. World Health Organization. Data on the size of the HIV epidemic.
6. Ba DM, Ssentongo P, Sznajder KK. Prevalence, behavioral and socioeconomic factors associated with human immunodeficiency virus in Ghana: a population-based cross-sectional study. *J Glob Health Rep*. 2019;3. doi:10.29392/joghr.3.e2019092.
7. Joint United Nations Programme on HIV/AIDS. *Global Report: UNAIDS Report on the Global AIDS Epidemic 2010*. https://unaids-test.unaids.org/sites/default/files/unaids/contentassets/documents/unaidspublication/2010/20101123_globalreport_en%5b1%5d.pdf
8. Abubakari GM, Dada D, Nur J, et al. Intersectional stigma and its impact on HIV prevention and care among MSM and WSW in sub-Saharan African countries: a protocol for a scoping review. *BMJ Open*. 2021;11(8):e047280. doi:10.1136/bmjopen-2020-047280
9. Nelson LRE, Nyblade L, Torpey K, et al. Multi-level intersectional stigma reduction intervention to increase HIV testing among men who have sex with men in Ghana: protocol for a cluster randomized controlled trial. *PLoS One*. 2021;16(11 November). doi:10.1371/journal.pone.0259324
10. Nelson LE, Ogunbajo A, Abu-Ba'are GR, et al. Using the implementation research logic model as a lens to view experiences of implementing HIV prevention and care interventions with adolescent sexual minority men—A global perspective. *AIDS Behav*. 2022;27:128-143.
11. Ali H, Amoyaw F, Baden D, et al. Ghana's HIV epidemic and PEPFAR's contribution towards epidemic control. *Ghana Med J*. 2019;53(1): 59-62. doi:10.4314/gmj.v53i1.9
12. UN Joint Programme on HIV/AIDS (UNAIDS). 90-90-90 An ambitious treatment target to help end the AIDS epidemic to help end the AIDS epidemic. *United Nations*. 2014. Published online 2014.
13. Owusu SA, Enimil A. Prevalence and determining factors of HIV infection among HIV exposed infants in a teaching hospital in Ghana. *Afr J Curr Med Res*. 2022;5(1). doi:10.31191/afrijcmr.v5i1.91
14. Cantelmo CB, Lee B, Dutta A. Smart cascades: using cost analysis to improve HIV care and treatment interventions to achieve global 95-95-95 goals. *Afr J AIDS Res*. 2019;18(4): 350-359. doi:10.2989/16085906.2019.1679201
15. Nkrumah B, van der Puije B, Bekoe V, et al. Building local human resources to implement SLMTA with limited donor funding: the Ghana experience. *Afr J Lab Med*. 2014;3(2): 1-7. doi:10.4102/ajlm.v3i2.214
16. Kaiser Family Foundation. The US President's Emergency Plan for AIDS Relief (PEPFAR) Fact Sheet, 2015. 2015;(December). <https://www.kff.org/global-health-policy/fact-sheet/the-u-s-presidents-emergency-plan-for-aids-relief-pepfar/>
17. UNAIDS. UNAIDS Goal 90-90-90: An ambitious treatment target to help end the AIDS epidemic. <https://www.unaids.org/en/resources/909090#:~:text=Powerful%20momentum%20is%20now%20building,will%20receive%20sustained%20antiretroviral%20therapy>. Published online 2014.
18. UNAIDS. To help end the AIDS epidemic. *United Nations*. 2014. Published online 2014.
19. Joint United Nations Program on HIV/AIDS. To help end the AIDS epidemic. *United Nations*. 2014. Published online 2014.
20. Guure C, Afagbedzi S, Torpey K, Chaurasia A. Willingness to take and ever use of preexposure prophylaxis among female sex workers in Ghana. *Medicine (United States)*. 2022;101(5). doi:10.1097/MD.00000000000028798
21. Eakle R, Venter F, Rees H. Preexposure prophylaxis (PrEP) in an era of stalled HIV prevention: can it change the game? *Retrovirology*. 2018;15(1): 1-10. doi:10.1186/s12977-018-0408-3
22. Mimiaga MJ, Case P, Johnson C, Safren SA, Mayer KH. Preexposure antiretroviral prophylaxis attitudes in high-risk Boston area men who report having sex with men: limited knowledge and experience but potential for increased utilization after education. *J Acquir Immune Defic Syndr (1988)*. 2009;50(1): 77-83. doi:10.1097/QAI.0b013e31818d5a27
23. Weitzman PF, Zhou Y, Kogelman L, Rodarte S, Vicente SR, Levkoff SE. MHealth for preexposure prophylaxis adherence by young adult men who have sex with men. *Mhealth*. 2021;7. doi:10.21037/mhealth-20-51.
24. Collins LC. Preexposure prophylaxis (PrEP) and 'risk' in the news. *J Risk Res*. 2022;25(3): 379-394. doi:10.1080/13669877.2021.1894470
25. Landovitz RJ. Preexposure prophylaxis for HIV prevention: what we know and what we still need to know for implementation. *Top Antivir Med*. 2015;23(2): 85.
26. Coyne R, Walsh JC, Noone C. Awareness, understanding and HIV stigma in response to undetectable = untransmittable messages: findings from a nationally representative sample in the United Kingdom. *AIDS Behav*. 2022;26(12): 85. doi:10.1007/s10461-022-03710-9
27. Huntingdon B, De Wit J, Duracinsky M, Juraskova I. Belief, covariates, and impact of the "Undetectable = Untransmittable" message among people living with HIV in Australia. *AIDS Patient Care STDS*. 2020;34(5): 205-212. doi:10.1089/apc.2019.0300
28. Sauermilch D, Siegel K, Hoppe T, Roth G, Meunier É. Attitudes toward HIV-positive status disclosure among U=U-aware sexual and gender minority individuals in the USA: a consensual qualitative research approach. *Sex Res Soc Policy*. 2023;20(2): 692-704. doi:10.1007/s13178-022-00710-1
29. Agaku I, Nkosi L, Gwar JN, Tsafa T. A cross-sectional analysis of U=U as a potential educative intervention to mitigate HIV

- stigma among youth living with HIV in South Africa. *Pan Afr Med J*. 2022;41. doi:10.11604/pamj.2022.41.248.33079
30. MacGibbon J, Bavinton BR, Broady TR, et al. Familiarity with, perceived accuracy of, and willingness to rely on undetectable=untransmittable (U=U) among gay and bisexual men in Australia: results of a national cross-sectional survey. *Sex Health*. 2023. Published online 2023;20(4): 373-373. doi:10.1071/sh23050
 31. Behrens GMN, Aebi-Popp K, Babiker A. Close to zero, but not zero: what is an acceptable HIV transmission risk through breastfeeding? *J Acquir Immune Defic Syndr (1988)*. 2022;89(4). doi:10.1097/QAI.0000000000002887
 32. Ngunjiri K, Ong'ola F, Dolla A, et al. I just believe there is a risk' understanding of undetectable equals untransmittable (U = U) among health providers and HIV-negative partners in serodiscordant relationships in Kenya. *J Int AIDS Soc*. 2020;23(3). doi:10.1002/jia2.25466
 33. Kowalska JD, Krankowska DC, Wróblewska A, Firląg-Burkacka E. Undetectable HIV equals untransmittable HIV (U = U) statement in women. Are we there yet? *Acta Scientific Women's Health*. 2022;4:44-48. Published online June 1, 2022. doi:10.31080/aswh.2022.04.0374
 34. Walsh-Buhi E, Houghton RF, Lange C, Hockensmith R, Ferrand J, Martinez L. Preexposure prophylaxis (PrEP) information on Instagram: content analysis. *JMIR Public Health Surveill*. 2021;7(7). doi:10.2196/23876
 35. Golub SA, Fikslin RA, Starbuck L, Klein A. High rates of PrEP eligibility but low rates of PrEP access among a national sample of transmasculine individuals. *J Acquir Immune Defic Syndr (1988)*. 2019;82(1). doi:10.1097/QAI.0000000000002116
 36. Parsons JT, Rendina HJ, Lassiter JM, Whitfield THF, Starks TJ, Grov C. Uptake of HIV preexposure prophylaxis (prep) in a national cohort of gay and bisexual men in the United States. *J Acquir Immune Defic Syndr (1988)*. 2017;74(3): 285-292. doi:10.1097/QAI.0000000000001251
 37. Yi S, Tuot S, Mwai GW, et al. Awareness and willingness to use HIV preexposure prophylaxis among men who have sex with men in low- and middle-income countries: a systematic review and meta-analysis. *J Int AIDS Soc*. 2017;20(1). doi:10.7448/IAS.20.1.21580
 38. Young I, McDaid L. How acceptable are antiretrovirals for the prevention of sexually transmitted HIV?: a review of research on the acceptability of oral preexposure prophylaxis and treatment as prevention. *AIDS Behav*. 2014;18(2):195-216. doi:10.1007/s10461-013-0560-7
 39. Ojikutu BO, Bogart LM, Mayer KH, Stopka TJ, Sullivan PS, Ransome Y. Spatial access and willingness to use preexposure prophylaxis among black/African American individuals in the United States: cross-sectional survey. *JMIR Public Health Surveill*. 2019;5(1). doi:10.2196/12405
 40. Poon AN, Han L, Li Z, et al. Acceptability and willingness of HIV preexposure prophylaxis amongst female sex workers in China. *AIDS Care - Psychol Socio-Med Aspects AIDS/HIV*. 2019;31(12). doi:10.1080/09540121.2019.1612001
 41. Zhao Z, Sun Y, Xue Q, et al. Acceptability of preexposure prophylaxis among female sex workers in Xinjiang. *Zhejiang Da Xue Xue Bao Yi Xue Ban*. 2011;40(3). doi:10.3785/j.issn.1008-9292.2011.03.009
 42. Jain JP, Strathdee SA, Patterson TL, et al. Perceived barriers to pre-exposure prophylaxis use and the role of syndemic factors among female sex workers in the Mexico-United States border region: a latent class analysis. *AIDS Care - Psychol Socio-Med Aspects AIDS/HIV*. 2020;32(5): 557-566. doi:10.1080/09540121.2019.1626338
 43. Emmanuel G, Folayan M, Undelikwe G, et al. Community perspectives on barriers and challenges to HIV preexposure prophylaxis access by men who have sex with men and female sex workers access in Nigeria. *BMC Public Health*. 2020;20(1): 1-10. doi:10.1186/s12889-020-8195-x
 44. Nagai H, Adiibokah E, Tagoe H, et al. Policymakers' and health-care providers' perspectives on the introduction of oral preexposure prophylaxis for key populations in Ghana. *BMC Public Health*. 2023;23(1): 1-9. doi:10.1186/s12889-023-15871-w
 45. Gomez GB, Borquez A, Case KK, Wheelock A, Vassall A, Hankins C. The cost and impact of scaling up preexposure prophylaxis for HIV prevention: a systematic review of cost-effectiveness modelling studies. *PLoS Med*. 2013;10(3). doi:10.1371/journal.pmed.1001401
 46. van de Vijver DAMC, Richter AK, Boucher CAB, et al. Cost-effectiveness and budget effect of preexposure prophylaxis for HIV-1 prevention in Germany from 2018 to 2058. *Eurosurveillance*. 2019;24(7). doi:10.2807/1560-7917.ES.2019.24.7.1800398
 47. Ogunbajo A, Leblanc NM, Kushwaha S, et al. Knowledge and acceptability of HIV preexposure prophylaxis (PrEP) among men who have sex with men (MSM) in Ghana. *AIDS Care - Psychol Socio-Med Aspects AIDS/HIV*. 2020;32(3): 330-336. doi:10.1080/09540121.2019.1675858
 48. Abubakari GMR, Nelson LRE, Ogunbajo A, et al. Implementation and evaluation of a culturally grounded group-based HIV prevention programme for men who have sex with men in Ghana. *Glob Public Health*. 2021;16(7): 1028-1045. doi:10.1080/17441692.2020.1832555
 49. Abubakari GM, Turner D, Nelson LE, et al. An application of the ADAPT-ITT model to an evidence-based behavioral HIV prevention intervention for men who have sex with men in Ghana. *Int Health Trends Perspect*. 2021;1(1): 1-16. doi:10.32920/ihtp.v1i1.1412
 50. Dako-Gyeke P, Snow R, Yawson AE. Who is utilizing antiretroviral therapy in Ghana: an analysis of ART service utilization. *Int J Equity Health*. 2012;11(1): 1-8. doi:10.1186/1475-9276-11-62
 51. Kumar N, Reece R, Norman B, Kwara A, Flanigan T, Rana A. Delayed entry to care by men with HIV infection in Kumasi, Ghana. *Pan Afr Med J*. 2015;22. doi:10.11604/pamj.2015.22.107.7010
 52. Sun Z, Gu Q, Dai Y, et al. Increasing awareness of HIV preexposure prophylaxis (PrEP) and willingness to use HIV PrEP among men who have sex with men: a systematic review and meta-analysis of global data. *J Int AIDS Soc*. 2022;25(3). doi:10.1002/jia2.25883
 53. Plotzker R, Seekaew P, Jantarapakde J, et al. Importance of risk perception: predictors of PrEP acceptance among Thai MSM and

- TG women at a community-based health service. *J Acquir Immune Defic Syndr (1988)*. 2017;76(5): 473–481. doi:10.1097/QAI.0000000000001536
54. Whitfield DL, Beauchamp G, Fields S, et al. Risk compensation in HIV PrEP adherence among Black men who have sex with men in HPTN 073 study. *AIDS Care - Psychol Socio-Med Aspects AIDS/HIV*. 2021;33(5): 633-638. doi:10.1080/09540121.2020.1810618
 55. Nagai H, Tagoe H, Tun W, et al. Perspectives of policymakers on the introduction and scale-up of HIV self-testing and implication for national HIV programming in Ghana. *Front Public Health*. 2021;9. doi:10.3389/fpubh.2021.694836
 56. Koduah A, Agyepong IA, van Dijk H. Towards an explanatory framework for national level maternal health policy agenda item evolution in Ghana: an embedded case study. *Health Res Policy Syst*. 2018;16(1): 1-16. doi:10.1186/s12961-018-0354-5
 57. Winskell K, Sabben G. Sexual stigma and symbolic violence experienced, enacted, and counteracted in young Africans' writing about same-sex attraction. *Soc Sci Med*. 2016;161: 143-150. doi:10.1016/j.socscimed.2016.06.004
 58. Beyrer C, Baral SD, van Griensven F, et al. Global epidemiology of HIV infection in men who have sex with men. *Lancet*. 2012;380(9839): 367-377. doi:10.1016/S0140-6736(12)60821-6
 59. Wiyeh AB, Mome RKB, Mahasha PW, Kongnyuy EJ, Wiysonge CS. Effectiveness of the female condom in preventing HIV and sexually transmitted infections: a systematic review and meta-analysis. *BMC Public Health*. 2020;20(1): 1-17. doi:10.1186/s12889-020-8384-7
 60. Giannou FK, Tsiara CG, Nikolopoulos GK, et al. Condom effectiveness in reducing heterosexual HIV transmission: a systematic review and meta-analysis of studies on HIV serodiscordant couples. *Expert Rev Pharmacoecon Outcomes Res*. 2016;16(4): 489-499. doi:10.1586/14737167.2016.1102635
 61. United States Agency for International Development. *Condom Fact Sheet*; 2015. <https://www.usaid.gov/sites/default/files/2022-05/condomfactsheet.pdf>
 62. Barcavage SH. How well do condoms protect gay men from HIV? [https://www.sfaf.org/collections/beta/how-well-do-condoms-protect-gay-men-from-hiv/#:~:text=They%20still%20have%20a%20role,i.e.%20breakage\)%20resulting%20in%20infections](https://www.sfaf.org/collections/beta/how-well-do-condoms-protect-gay-men-from-hiv/#:~:text=They%20still%20have%20a%20role,i.e.%20breakage)%20resulting%20in%20infections).
 63. Ahinkorah BO, Hagan JE, Seidu AA, et al. Understanding the association between exposure to family planning messages and consistent condom use among never married men in Ghana. *PLoS One*. 2021;16(8 August 2021). doi:10.1371/journal.pone.0255325
 64. Davis KR, Weller SC. The effectiveness of condoms in reducing heterosexual transmission of HIV. *Fam Plann Perspect*. 1999;31(6): 272-279. doi:10.2307/2991537
 65. Wald A, Langenberg AGM, Link K, et al. Effect of condoms on reducing the transmission of herpes simplex virus type 2 from men to women. *JAMA*. 2001;285(24): 3100-3106. doi:10.1001/jama.285.24.3100
 66. Ministry of Health. *Ghana National Condom and Water-Based Lubricant Programming Strategy-2014–2019*. <https://www.ccmghana.net/index.php/strategic-plans-reports?download%3D35:national-condom-and-lubricant-strategy-2014-2019&tbn=ilp&sa=X&ved=2ahUKEwjks8eDuf2BAxXaDFkFHbYBCukQv5AHegQIABAE>
 67. Abubakari GMR, Turner DA, Ni Z, et al. Community-based interventions as opportunities to increase HIV self-testing and linkage to care among men who have sex with men – Lessons from Ghana, West Africa. *Front Public Health*. 2021;9. doi:10.3389/fpubh.2021.660256
 68. Nelson LRE, Wilton L, Agyarko-Poku T, et al. Predictors of condom use among peer social networks of men who have sex with men in Ghana, West Africa. *PLoS One*. 2015;10(1). doi:10.1371/journal.pone.0115504
 69. Go VF, Srikrishnan AK, Sivaram S, et al. High HIV prevalence and risk behaviors in men who have sex with men in Chennai, India. *J Acquir Immune Defic Syndr (1988)*. 2004;35(3): 314-319. doi:10.1097/00126334-200403010-00014
 70. Shah N. *Characterizing slums and slum-dwellers: exploring household-level Indonesian data*. University of California; 2012, Published online 2012.
 71. Appiah-Kubi J. Challenges encountered in community development in urban slums: a study of Ashaiman, Ghana. *Int J Sci: Basic Appl Res* 2018;4531(October 2018): 81-93.
 72. Jankowska MM, Weeks JR, Engstrom R. Do the most vulnerable people live in the worst slums? A spatial analysis of Accra, Ghana. *Ann GIS*. 2011;17(4): 221-235. doi:10.1080/19475683.2011.625976
 73. Owusu G, Agyei-Mensah S, Lund R. Slums of hope and slums of despair: mobility and livelihoods in Nima, Accra. *Norsk Geografisk Tidsskrift*. 2008;62(3): 180-190. doi:10.1080/00291950802335798
 74. Ezeh A, Oyebode O, Satterthwaite D, et al. The history, geography, and sociology of slums and the health problems of people who live in slums. *Lancet*. 2017;389(10068): 547-558. doi:10.1016/S0140-6736(16)31650-6
 75. Zigah EY, Abu-Ba'are GR, Shamrock OW, et al. For my safety and wellbeing, I always travel to seek health care in a distant facility—the role of place and stigma in HIV testing decisions among GBMSM – BSGH 002. *Health Place*. 2023;83:103076. doi:10.1016/j.healthplace.2023.103076
 76. Atuguba RA. Homosexuality in Ghana: morality, law, human rights. *J Pol & L*. 2019;12:113:103076.
 77. Opoku JK, Manu E, Appiah DM. Same-sex marriage in Ghana: scripture, health law and bioethics. *Developing Country Studies*. 2021;11(4).
 78. Nartey M. Marginality and otherness: the discursive construction of LGBT issues/people in the Ghanaian news media. *Media Cult Soc*. 2022;44(4): 785-801. doi:10.1177/01634437211045552
 79. Oppong S. Examining attitudes and the law on homosexuality in non-Western Societies: the example of Ghana in West Africa. *Polish Psychol Bullet*. 2018;49(4): 416-423. doi:10.24425/119510
 80. Dai-Kosi AD, Asamani L, Adomako B. Ghanaian perspectives on the present day dynamics of homosexuality. *Afr Res Rev*. 2016;10(5): 1-12. doi:10.4314/afrev.v10i5.1
 81. Gyamerah AO, Kinzer E, Aidoo-Frimpong G, et al. PrEP knowledge, acceptability, and implementation in Ghana: perspectives of HIV service providers and MSM, trans women, and gender

- diverse individuals living with HIV. *PLoS Global Public Health*. 2023;3(6). doi:10.1371/journal.pgph.0001956
82. Laar A, Debruin D. Key populations and human rights in the context of HIV services rendition in Ghana. *BMC Int Health Hum Rights*. 2017;17(1): 1-10. doi:10.1186/s12914-017-0129-z
 83. Colorafi KJ, Evans B. Qualitative descriptive methods in health science research. *Health Environ Res Design J*. 2016;9(4):16–25. doi:10.1177/1937586715614171
 84. Leon L, Jauffret-Roustide M, Le Strat Y. Design-based inference in time-location sampling. *Biostatistics*. 2014;16(3): 565-579. doi:10.1093/biostatistics/kxu061
 85. Trouiller P, Velter A, Saboni L, et al. Injecting drug use during sex (known as “slamming”) among men who have sex with men: results from a time-location sampling survey conducted in five cities, France. *Int J Drug Policy*. 2020;79. doi:10.1016/j.drugpo.2020.102703
 86. Zhao J, Cai R, Chen L, et al. A comparison between respondent-driven sampling and time-location sampling among men who have sex with men in Shenzhen, China. *Arch Sex Behav*. 2015;44(7): 2055-2065. doi:10.1007/s10508-014-0350-y
 87. Sommen C, Saboni L, Sauvage C, et al. Time location sampling in men who have sex with men in the HIV context: the importance of taking into account sampling weights and frequency of venue attendance. *Epidemiol Infect*. 2018;146(7): 913-919. doi:10.1017/S0950268818000675
 88. Abubakari GM, Owusu-Dampare F, Ogunbajo A, et al. HIV Education, empathy, and empowerment (HIVE3): a peer support intervention for reducing intersectional stigma as a barrier to HIV testing among men who have sex with men in Ghana. *Int J Environ Res Public Health*. 2021;18(24):13103. doi:10.3390/ijerph182413103
 89. Cypress B. Qualitative research methods: a phenomenological focus. *Dimens Crit Care Nurs*. 2018;37(6): 302–309. doi:10.1097/DCC.0000000000000322
 90. Holloway I, Freshwater H. *Narrative Research in Nursing*; 2009. doi:10.1002/9781444316513.
 91. Unger A, Riley LW. Slum health: from understanding to action. *PLoS Med*. 2007;4(10):e295. doi:10.1371/journal.pmed.0040295
 92. Kshetrimayum B, Bardhan R, Kubota T. Factors affecting residential satisfaction in slum rehabilitation housing in Mumbai. *Sustainability (Switzerland)*. 2020;12(6): 2344. doi:10.3390/su12062344
 93. Bell DC, Atkinson JS, Mosier V, Riley M, Brown VL. The HIV transmission gradient: relationship patterns of protection. *AIDS Behav*. 2007;11(6):789–811. doi:10.1007/s10461-006-9192-5
 94. Skakoon-Sparling S, Cramer KM. Are we blinded by desire? Relationship motivation and sexual risk-taking intentions during condom negotiation. *J Sex Res*. 2020;57(5): 545-558. doi:10.1080/00224499.2019.1579888
 95. Peasant C, Parra GR, Okwumabua TM. Condom negotiation: findings and future directions. *J Sex Res*. 2015;52(4): 470-483. doi:10.1080/00224499.2013.868861
 96. Starks TJ, Pawson M, Stephenson R, Sullivan P, Parsons JT. Dyadic qualitative analysis of condom use scripts among emerging adult gay male couples. *J Sex Marital Ther*. 2018;44(3): 470–483. doi:10.1080/0092623X.2017.1359713
 97. Smith P, Bottenheim A, Schmucker L, Bekker LG, Thirumurthy H, Davey DLJ. Undetectable=untransmittable (U=U) messaging increases uptake of HIV testing among men: results from a pilot cluster randomized trial. *AIDS Behav*. 2021;25(10):3128–3136. doi:10.1007/s10461-021-03284-y
 98. Carneiro PB, Westmoreland DA, Patel VV, Grov C. Awareness and acceptability of undetectable=untransmittable among a U.S. National sample of HIV-negative sexual and gender minorities. *AIDS Behav*. 2021;25(2):634–644. doi:10.1007/s10461-020-02990-3
 99. Saalim K, Amu-Adu P, Amoh-Otu RP, et al. Multi-level manifestations of sexual stigma among men with same-gender sexual experience in Ghana. *BMC Public Health*. 2023;23(1): 166. doi:10.1186/s12889-023-15087-y
 100. Nyblade L, Stockton MA, Saalim K, et al. Using a mixed-methods approach to adapt an HIV stigma reduction intervention to address intersectional stigma faced by men who have sex with men in Ghana. *J Int AIDS Soc*. 2022;25(S1). doi:10.1002/jia2.25908
 101. Ali BK. *Prevalence and predictors of virological non-suppression among key populations on HIV treatment in Zanzibar*. Muhimbili University of Health and Allied Sciences; 2019.
 102. Raiford JL, Wingood GM, DiClemente RJ. Correlates of consistent condom use among HIV positive African American women. *Women Health*. 2007;46(2-3): 41-58. doi:10.1300/J013v46n02_04
 103. Smith DK, Herbst JH, Zhang X, Rose CE. Condom effectiveness for HIV prevention by consistency of use among men who have sex with men in the United States. *J Acquir Immune Defic Syndr (1988)*. 2015;68(3): 337-344. doi:10.1097/QAI.0000000000000461
 104. Manguro GO, Musau AM, Were DK, et al. Increased condom use among key populations using oral PrEP in Kenya: results from large scale programmatic surveillance. *BMC Public Health*. 2022;22(1): 328-340. doi:10.1186/s12889-022-12639-6
 105. Dudley MG, Rostosky SS, Korfhage BA, Zimmerman RS. Correlates of high-risk sexual behavior among young men who have sex with men. *AIDS Educ Prev*. 2004;16(4):328-340. doi:10.1521/aeap.16.4.328.40397
 106. Baral S, Sifakis F, Cleghorn F, Beyrer C. Elevated risk for HIV infection among men who have sex with men in low- and middle-income countries 2000-2006: a systematic review. *PLoS Med*. 2007;4(12). doi:10.1371/journal.pmed.0040339
 107. Musinguzi G, Bastiaens H, Matovu JKB, et al. Barriers to condom use among high risk men who have sex with men in Uganda: a qualitative study. *PLoS One*. 2015;10(7). doi:10.1371/journal.pone.0132297
 108. Baggaley RF, White RG, Boily MC. HIV transmission risk through anal intercourse: systematic review, meta-analysis and implications for HIV prevention. *Int J Epidemiol*. 2010;39(4):1048-1063. doi:10.1093/ije/dyq057
 109. Hui B, Fairley CK, Chen M, et al. Oral and anal sex are key to sustaining gonorrhoea at endemic levels in MSM populations:

- a mathematical model. *Sex Transm Infect.* 2015;91(5): 365-369. doi:10.1136/sextrans-2014-051760
110. Patel VL, Gutnik LA, Yoskowitz NA, O'Sullivan LF, Kaufman DR. Patterns of reasoning and decision making about condom use by urban college students. *AIDS Care - Psychol Socio-Med Aspects AIDS/HIV.* 2006;18(8): 918-930. doi:10.1080/09540120500333509
 111. Kibaso JN. *Levels of HIV/AIDS-Related Stigma and Its Association with Individual HIV Status in Nairobi City Slums.* Order No. 10193765 ed. Loma Linda University; 2016. <https://ezp.lib.rochester.edu/login?url=https://www.proquest.com/dissertations-theses/levels-hiv-aids-related-stigma-association-with/docview/2135749943/se-2>
 112. Nyblade L, Carr D. *Towards a stronger response to HIV and AIDS: challenging stigma.* International Center for Research on Women; 2004, Published online 2004.
 113. Hartwig KA, Kissioki S, Hartwig CD. Church leaders confront HIV/AIDS and stigma: a case study from Tanzania. *J Community Appl Soc Psychol.* 2006;16(6): 492-497. doi:10.1002/casp.897
 114. Surur F, Kaba M. The role of religious leaders in HIV/AIDS prevention, control, and patient care and support: a pilot project in Jimma Zone. *Northeast Afr Stud.* 2000;7(2): 59-79. doi:10.1353/nas.2004.0021
 115. van Kesteren NMC, Hospers HJ, Kok G, van Empelen P. Sexuality and sexual risk behavior in HIV-positive men who have sex with men. *Qual Health Res.* 2005;15(2):145-168. doi:10.1177/1049732304270817
 116. Van Kesteren NMC, Hospers HJ, Van Empelen P, Van Breukelen G, Kok G. Sexual decision-making in HIV-positive men who have sex with men: how moral concerns and sexual motives guide intended condom use with steady and casual sex partners. *Arch Sex Behav.* 2007;36(3): 437-449. doi:10.1007/s10508-006-9125-4
 117. Serovich JM, Oliver DG, Smith SA, Mason TL. Methods of HIV disclosure by men who have sex with men to casual sexual partners. *AIDS Patient Care STDS.* 2005;19(12):823-832. doi:10.1089/apc.2005.19.823.
 118. Hillis A, Germain J, Hope V, McVeigh J, Van Hout MC. Preexposure prophylaxis (PrEP) for HIV prevention among men who have sex with men (MSM): a scoping review on PrEP service delivery and programming. *AIDS Behav.* 2020; 24(11):3056-3070. doi:10.1007/s10461-020-02855-9
 119. Sun Y, Lu H, Ye J, Li D, Li G. Awareness and use of HIV pre-exposure prophylaxis and factors associated with awareness among MSM in Beijing, China. *Sci Rep.* 2023;13(1):554. doi:10.1038/s41598-023-27485-8
 120. Eubanks A, Coulibaly B, Keita BD, et al. Loss to follow-up from HIV preexposure prophylaxis care in men who have sex with men in West Africa. *Viruses.* 2022;14(11): 2380. doi:10.3390/v14112380
 121. Atkins K, Musau A, Mugambi M, et al. Health system opportunities and challenges for PrEP implementation in Kenya: a qualitative framework analysis. *PLoS One.* 2022;17(10 October). doi:10.1371/journal.pone.0259738
 122. Ahouada C, Diabaté S, Mondor M, et al. Acceptability of pre-exposure prophylaxis for HIV prevention: facilitators, barriers and impact on sexual risk behaviors among men who have sex with men in Benin. *BMC Public Health.* 2020;20(1):1-17. doi:10.1186/s12889-020-09363-4
 123. Hannaford A, Lipshie-Williams M, Starrels JL, et al. The use of online posts to identify barriers to and facilitators of HIV preexposure prophylaxis (PrEP) among men who have sex with men: a comparison to a systematic review of the peer-reviewed literature. *AIDS Behav.* 2018;22(4):1080-1095. doi:10.1007/s10461-017-2011-3