

Research article

Open Access

Histopathological features of breast tumours in *BRCA1*, *BRCA2* and mutation-negative breast cancer familiesHannaleena Eerola^{1,3}, Päivi Heikkilä², Anitta Tamminen³, Kristiina Aittomäki⁴, Carl Blomqvist¹ and Heli Nevanlinna³¹Department of Oncology, Helsinki University Central Hospital, Helsinki, Finland²Department of Pathology, Helsinki University Central Hospital, Helsinki, Finland³Department of Obstetrics and Gynaecology, Helsinki University Central Hospital, Helsinki, Finland⁴Department of Clinical Genetics, Helsinki University Central Hospital, Helsinki, FinlandCorresponding author: Hannaleena Eerola, hannaleena.eerola@hus.fi

Received: 15 Jun 2004 Revisions requested: 4 Aug 2004 Revisions received: 31 Aug 2004 Accepted: 30 Sep 2004 Published: 19 Nov 2004

Breast Cancer Res 2005, **7**:R93-R100 (DOI 10.1186/bcr953)© 2004 Eerola *et al.*; licensee BioMed Central Ltd.This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.**Abstract**

Introduction Histopathological features of *BRCA1* and *BRCA2* tumours have previously been characterised and compared with unselected breast tumours; however, familial non-*BRCA1/2* tumours are less well known. The aim of this study was to characterise familial non-*BRCA1/2* tumours and to evaluate routine immunohistochemical and pathological markers that could help us to further distinguish families carrying *BRCA1/2* mutations from other breast cancer families.

Methods Breast cancer tissue specimens ($n = 262$) from 25 *BRCA1*, 20 *BRCA2* and 74 non-*BRCA1/2* families were studied on a tumour tissue microarray. Immunohistochemical staining of oestrogen receptor (ER), progesterone receptor (PgR) and p53 as well as the histology and grade of these three groups were compared with each other and with the respective information on 862 unselected control patients from the archives of the Pathology Department of Helsinki University

Central Hospital. Immunohistochemical staining of erbB2 was also performed among familial cases.

Results *BRCA1*-associated cancers were diagnosed younger and were more ER-negative and PgR-negative, p53-positive and of higher grade than the other tumours. However, in multivariate analysis the independent factors compared with non-*BRCA1/2* tumours were age, grade and PgR negativity. *BRCA2* cases did not have such distinctive features compared with non-*BRCA1/2* tumours or with unselected control tumours. Familial cases without *BRCA1/2* mutations had tumours of lower grade than the other groups.

Conclusions *BRCA1* families differed from mutation-negative families by age, grade and PgR status, whereas ER status was not an independent marker.

Keywords: breast cancer, oestrogen receptor, hereditary, p53, progesterone receptor**Introduction**

Women predisposed to hereditary or familial breast cancer form a heterogeneous group. It would be useful if we could identify carriers of the high-risk *BRCA1* and *BRCA2* genes and target the expensive and time-consuming genetic testing to individuals who most probably carry those mutations. Besides family history, histopathological markers could also be useful in distinguishing patients and families likely to carry a *BRCA1/2* germline mutation from mutation-negative families and breast cancer patients in general.

Several studies have compared the characteristics of breast cancers in *BRCA1* carriers and in sporadic controls. Distinct features between *BRCA1*-associated tumours have been found, such as high tumour grade, oestrogen receptor (ER) negativity, and overexpression of p53 [1-3]. In addition, negativity for progesterone receptor (PgR) [3,4], a higher proportion of medullary and atypical medullary carcinomas [5,6], and tumours with a low expression of c-erbB-2 [1,4,6] have been detected. Besides the higher proportion of medullary histology, a higher frequency of ductal carcinoma has also been reported [7,8]. Recently, cDNA expression analyses have suggested a basal

epithelial phenotype for *BRCA1* tumours [9] and expression of cytokeratins 5/6 has been associated with *BRCA1* tumours [10].

Among *BRCA2*-associated tumours, a slight increase has been observed in the incidence of lobular or tubulolobular carcinomas [11,12]. However, results are inconsistent, and in most cases no significant difference has been found between *BRCA2*-associated tumours and sporadic cancers [1,4,6,13]. Complementary DNA (cDNA) expression analysis has suggested distinct expression profiles for *BRCA2* tumours as well [14].

It has been clear for some time that in many families hereditary susceptibility is not due to the *BRCA1* or *BRCA2* genes [15,16]. Only a few studies have evaluated the features of this large group of families. It would be crucial for genetic counselling and for our understanding of tumour development to learn more about these patients. Lakhani and colleagues [17] studied the pathology of 82 familial breast cancers not attributable to *BRCA1* or *BRCA2* mutations, and they found these non-*BRCA1/2* cancers to be of lower grade, to show less pleomorphism and to have a lower mitotic count than sporadic cancers or *BRCA* mutation-positive cancers. No other features differed significantly in their study. However, they did not examine immunohistochemical characteristics. There has been only one study so far that has characterised the immunohistochemical features of familial non-*BRCA1/2* cancers: Palacios and colleagues [18] studied immunohistochemical staining and histopathology and compared 37 non-*BRCA1/2* cancers with 20 *BRCA1*-associated and 18 *BRCA2*-associated cancers, and also with unselected control cancers. They similarly found those to be of lower grade than in all the other three groups. In comparison with sporadic cancers they were also more frequently p53-negative and erbB2-negative, and expressed reduced E-cadherin and β -catenin. However, the number of patients in this study was small and it was restricted to only a univariate analysis of the studied parameters.

Most of the previous studies on *BRCA1*-associated and *BRCA2*-associated cancers have studied highly selected patient groups, but in the present study families were collected with a simple criterion of at least three first-degree or second-degree relatives with breast or ovarian cancer with no restriction on age. We studied an extensive material of 152 non-*BRCA1/2* tumours, and also 110 tumours from *BRCA1/2* families for histopathological features as well as for the immunohistochemical expression of ER, PgR, p53 and erbB2. We describe here the histopathological profile of the tumours originating from non-*BRCA1/2* breast cancer families and also present a multivariate analysis to find the independent markers that can further help in distin-

guishing especially *BRCA1* mutation-positive families from other familial cases.

Patients and methods

Familial breast cancer patients were identified and collected by a systematic screening for family history at the Department of Oncology, Helsinki University Central Hospital, as described previously [19]. We defined breast cancer families by the selection criterion of at least three first-degree or second-degree relatives with breast or ovarian cancer (including the proband). We confirmed the genealogy of the families through population registries, and cancer diagnoses through the Finnish Cancer Registry. In this study we included 25 *BRCA1* families, 20 *BRCA2* families and 74 families not associated with either of these genes (non-*BRCA1/2* families) (Table 1). All families had previously been tested for *BRCA1* and *BRCA2* mutations by mutation analysis of the whole coding sequences and exon/intron boundaries of the genes as described [16,20] or tested for all previously reported 18 Finnish *BRCA1* and *BRCA2* mutations [16,20-22].

We collected all the paraffin blocks of all the primary breast cancers that were available ($n = 262$) from these families. However, cases tested to be non-carriers in the mutation-positive families were excluded. In total, 51 cancers from the 25 *BRCA1* families, 59 cancers from the 20 *BRCA2* families and 152 cancers from the 74 non-*BRCA1/2* families were included in this study. We studied the haematoxylin and eosin sections of the original blocks to achieve histological diagnosis and grading. Grading was performed according to Scarff-Bloom-Richardson modified by Elston and Ellis [23]. The most representative area of the tumour was punched to produce a hereditary breast cancer tissue microarray including two cores (diameter 0.6 mm) from all of the original blocks. The array block of non-*BRCA1/2* cases was described previously by Vahteristo and colleagues [24]. All of the microarray slides were stained with routine methods by antibodies against ER, PgR, p53 and erbB2 in the same pathology laboratory as our controls. In brief, 5 μ m sections were cut from paraffin-embedded blocks, deparaffinated in xylene, and dehydrated in a series of graded alcohols. The sections were pretreated in a microwave oven and incubated overnight with antibody. Antibodies for ER (dilution 1:50) and c-erbB-2 (NCL-CB11; dilution 1:400) were purchased from Novocastra (Newcastle upon Tyne, UK), and those for PgR (dilution 1:250) and p53 (dilution 1:100) were from Dako (Copenhagen, Denmark). The evaluation of the staining results was similar to that used in routine diagnostics and samples were considered positive when 10%, 10% and 20% of the cells were stained for ER, PgR and p53, respectively. Samples having a moderate or intense staining of the entire membrane in more than 10% of the tumour cells (2+ and

3+) were considered to be c-erbB-2-positive. Other staining patterns were considered to be negative (0 and 1+).

As a control group we drew from the archives of the Pathology Department at Helsinki University 862 unselected breast cancer tumours from the years 1997–2001 that were scored by the same pathologist (PH) as the tumours of the hereditary breast cancer patients.

Statistical analysis was conducted with SPSS version 8.0 for Windows. We tested the differences in continuous variables by the Mann–Whitney test and in dichotomous variables by the χ^2 test or Fisher's exact test. In multivariate analysis, we used logistic regression analysis (stepwise backwards logistic regression, 99%). All *P* values are two-sided.

Permissions for this study were obtained from the ethics committees of the Department of Oncology and the Department of Obstetrics and Gynaecology, Helsinki University Central Hospital, and of the Ministry of Social Affairs and Health in Finland. Blood and tumour samples were used in this study with written informed consent from probands and family members.

Results

BRCA1-associated cancers were diagnosed at a younger age than unselected breast cancers (median ages 44 and 56 years, respectively; $P \leq 0.0005$) and they were more often ER-negative and PgR-negative, p53-positive and of higher grade (Table 2). The frequency of medullary histology was also higher. In logistic regression analysis, taking into account all of these factors, the independent factors were age ($P \leq 0.0005$), ER status ($P = 0.0597$), PgR status ($P = 0.0170$) and medullary histology ($P = 0.0636$).

In comparison with familial non-*BRCA1/2* cancers (median age 55 years), a univariate analysis of *BRCA1* tumours showed similar differences from unselected cancers (Table 2). However, in multivariate analysis, taking into account the same factors, the independent factors were age ($P = 0.0012$), grade ($P = 0.0014$) and PgR negativity ($P = 0.0196$) (Table 3). We did not find any significant differences between groups of tumours from carriers with different mutations or when comparing tumours from the carriers of the mutations at the 5' end or the 3' end of the gene.

BRCA2-associated cancers were diagnosed at younger age (median age 47 years) than unselected breast cancers (median age 56 years, $P \leq 0.0005$). They were also more often ER-negative and PgR-negative (Table 2). In multivariate analysis, the independent factors were age ($P \leq 0.0005$), PgR status ($P = 0.0365$) and p53 status ($P = 0.0318$).

When compared with familial non-*BRCA1/2* cancers, the *BRCA2*-associated cancers were diagnosed at a younger age (median age 47 years; among non-*BRCA1/2* patients the median age was 55 years; $P \leq 0.0005$). No other variable differed significantly from non-*BRCA1/2* cancers (Table 2). We did not find any significant differences between tumours originating from different mutation carriers or when comparing tumours from the carriers of mutations from the OCCR (Ovarian Cancer Cluster region) ($n = 7$) and the end of the gene. In a logistic regression analysis comparing *BRCA2*-associated cancers with non-*BRCA1/2* cancers and taking into account age, grade and all tested histological and immunohistochemical factors, the only independently significant factor was age ($P = 0.0001$) (Table 2).

Among familial non-*BRCA1/2* patients, the median age of onset was marginally younger (55.0 years) than among the unselected controls (56.0 years, $P = 0.060$). The frequency of grade I and II tumours was much higher among the non-*BRCA1/2* group than in the unselected control group (odds ratio 1.8, $P = 0.009$) (Table 2) or in the *BRCA1* and *BRCA2* groups. In a multiple regression analysis comparing non-*BRCA1/2* tumours with unselected controls, grade (odds ratio 0.54, $P \leq 0.00005$) and ER status (odds ratio 2.4 for negative ER status, $P = 0.0006$) were the independent significant factors.

The erbB2 results were very similar in the three groups of familial cases, with 18.6%, 15.1% and 17.4% of the *BRCA1*, *BRCA2* and non-*BRCA1/2* tumours, respectively, expressing the erbB2 antigen (Table 2).

Discussion

Although genetic testing for *BRCA1* and *BRCA2* mutations is available, it is expensive, time-consuming and stressful to patients. Several models have therefore been developed for evaluating the probability of carrying a *BRCA1* or *BRCA2* mutation [25–30]. In our previous study of Finnish breast cancer families [15,29], multivariate analysis suggested simple family history criteria for breast cancer onset under the age of 40 years and the presence of ovarian cancer to be most strongly associated with *BRCA1/2* mutation status [29]. However, in addition to family history, it is important to find other markers that could help to identify mutation carriers. In many countries, markers that are already routinely used, such as ER, PgR, p53 and the grade of the tumour, could serve as an excellent tool to aid distinguishing families because this information is easily available. In this study, we found that *BRCA1*-associated cancers have a different histological profile and can be distinguished from other familial cancers. Specifically, multivariate analysis revealed age, grade and PgR negativity as the independent factors distinguishing *BRCA1* tumours from familial non-*BRCA1/2* tumours.

Table 1**Mutation spectrum and numbers of families and patients by mutation**

| Gene | Mutation | Number of families | Number of patients |
|---------------------|-----------------------|--------------------|--------------------|
| <i>BRCA1</i> | ex 10 782 del AA | 1 | 1 |
| | ex 11 1047 C→T | 1 | 1 |
| | ex 11 1731 C→T | 1 | 2 |
| | ex 11 1806 C→T | 1 | 1 |
| | ex 11 1924 del A | 1 | 1 |
| | ex 11 2592 ins A | 1 | 2 |
| | ex 11 2803 del AA | 1 | 3 |
| | ex 11 3604 del A | 3 | 3 |
| | ex 11 3744 del T | 2 | 2 |
| | ex 11 3904 C→A | 1 | 1 |
| | ex 11 4153 del A | 1 | 1 |
| | int 11 4216 (- 2) A→G | 4 | 9 |
| | ex 13 4446 C→T | 2 | 14 |
| | ex 14 4599 G>T | 1 | 3 |
| | ex 17 5145 del 11 | 1 | 4 |
| | ex 20 5370 C→T | 1 | 1 |
| | ex 20 5382 ins C | 2 | 2 |
| | Any | 25 | 51 |
| <i>BRCA2</i> | ex 9 999 del TCAAA | 5 | 22 |
| | ex 10 1822 G→T | 1 | 1 |
| | ex 11 4075 del GT | 1 | 1 |
| | ex 11 4081 ins A | 1 | 2 |
| | ex 11 5797 G→T | 1 | 1 |
| | ex 11 6496 del CA | 1 | 2 |
| | ex 11 6503 del TT | 1 | 1 |
| | ex 15 7708 C→T | 3 | 12 |
| | ex 18 8555 T→G | 1 | 6 |
| | int 23 9346 (- 2) A→G | 5 | 11 |
| | Any | 20 | 59 |
| Non- <i>BRCA1/2</i> | 74 | 152 | |

BRCA2-associated cancers did not differ greatly, but those were also significantly younger and there was a trend towards higher grade than among familial non-*BRCA1/2* cancers.

ER negativity has previously been highlighted to be linked to *BRCA1* tumours. It was obvious in this study too. Our

results are consistent with, for example, the study of Lakhani and colleagues [4], in which *BRCA1* cancers were clearly more ER-negative (90%) than unselected control cancers (35%). However, in our study the overall percentages of ER-negative cancers were much lower for both groups (67% among *BRCA1* and 19% among unselected controls). This might be due to a different age distribution

Table 2**Analysis of features among BRCA1-, BRCA2- and non-BRCA1/2-associated cancers and in unselected cancers**

| Feature | N (frequency, %) | | | | P |
|-----------------------------|------------------|-----------|-------------|------------|---|
| | BRCA1 | BRCA2 | Non-BRCA1/2 | Unselected | |
| Histology | | | | | |
| Ca ductale | 37 (72.5) | 37 (62.7) | 102 (67.1) | 561 (65.1) | |
| Ca lobulare | 8 (15.7) | 17 (28.8) | 30 (19.7) | 219 (25.4) | |
| Ca medullare | 5 (9.8) | - | 3 (2.0) | 20 (2.3) | 0.013*, 0.001** |
| Others | 1 (2.0) | 5 (8.5) | 17 (11.2) | 62 (7.2) | |
| Grade | | | | | |
| I | 3 (6.1) | 12 (23.1) | 46 (32.4) | 152 (23.3) | |
| II | 11 (22.4) | 26 (50.1) | 66 (46.5) | 289 (44.4) | |
| III | 35 (71.4) | 14 (26.9) | 30 (21.1) | 210 (32.3) | |
| I-II | 14 (28.6) | 38 (73.1) | 112 (78.9) | 441 (67.7) | ≤ 0.0005*, <0.0005**, 0.009**** |
| Immunohistochemistry | | | | | |
| ER ⁻ | 28 (66.7) | 17 (32.1) | 40 (27.0) | 167 (19.4) | ≤ 0.0005*, <0.0005**, 0.026***, 0.035**** |
| ER ⁺ | 14 (33.3) | 36 (67.9) | 108 (73.0) | 692 (80.6) | |
| PgR ⁻ | 37 (84.1) | 28 (51.9) | 67 (45.3) | 312 (36.2) | ≤ 0.0005*, <0.0005**, 0.021***, 0.036**** |
| PgR ⁺ | 7 (15.9) | 26 (48.1) | 81 (54.7) | 549 (63.8) | |
| p53 ⁻ | 27 (62.8) | 42 (82.4) | 118 (78.7) | 628 (74.0) | 0.034*, <0.0005** |
| p53 ⁺ | 16 (37.2) | 9 (17.6) | 32 (21.3) | 221 (26.0) | |
| erbB2 ⁻ | 35 (81.4) | 45 (84.9) | 109 (82.6) | NA | |
| erbB2 ⁺ | 8 (18.6) | 8 (15.1) | 23 (17.4) | NA | |

*BRCA1 versus non-BRCA1/2 tumours; **BRCA1 versus unselected breast tumours; ***BRCA2 versus unselected breast tumours; ****non-BRCA1/2 tumours versus unselected breast tumours.

Ca, carcinoma; ER, oestrogen receptor; NA, not available; PgR, progesterone receptor.

because the prevalence of ER-negative cases is higher among young breast cancer patients [31,32] and our familial cases were not selected on the basis of young age of diagnosis. Furthermore, there is a very high coverage mammography screening (in the general age group 50–60 years) in Finland, aiding in finding asymptomatic cancers, which are more often ER-positive as well as being of lower grade and PgR-positive [33].

In this study, we specifically sought to compare BRCA1 cancers with familial non-BRCA1/2 cancers as well, which is the relevant question in a clinical and genetic counselling setting. Surprisingly, in this analysis the ER status was not an independent marker in multiple regression. More important markers were age, PgR status and grade. ER status was dependent on age and grade. Palacios and colleagues [18] also compared BRCA1 cases with non-BRCA1/2 cases (cases from families with at least three affected with breast cancer, one of them diagnosed under 50 years of

age). In a univariate analysis, the results showing ER and PgR negativity, p53 overexpression and high grade were very similar to ours. However, multivariate analysis was not used for evaluating independent markers. Recently, cDNA expression analyses have suggested a basal epithelial phenotype for BRCA1 tumours [9]. Epithelial phenotype is associated with breast cancers that express neither ER nor erbB2, a feature that also occurs in BRCA1-mutation carriers [10]. A large majority (20 of 22 tumours) of BRCA1-associated tumours were also ER-negative and erbB2-negative in our study, in accordance with the high frequency of ER-negative tumours among BRCA1 carriers. The frequency of erbB2 expression was similar in all three groups of familial tumours.

BRCA2-associated tumours did not differ significantly from familial non-BRCA1/2 tumours, although they were diagnosed at an earlier age. Our results on BRCA2 cancers were quite similar to those of Palacios and colleagues [18]

Table 3

Logistic regression analysis of features of breast cancers among BRCA1- and BRCA2-associated cases in comparison with non-BRCA1/2 cases

| Feature | Odds ratio (95% confidence interval, <i>P</i>) | |
|---------------------------------------|---|--------------------------|
| | <i>BRCA1</i> | <i>BRCA2</i> |
| First step: | | |
| Age (continuous) | 0.94 (0.91–0.98, 0.0009) | 0.94 (0.91–0.97, 0.0001) |
| Grade (continuous) | 3.78 (1.70–8.39, 0.0011) | 1.27 (0.71–2.26, 0.42) |
| ER ⁺ | 1.0 (referent) | 1.0 (referent) |
| ER ⁻ | 0.91 (0.28–2.90, 0.88) | 0.85 (0.32–2.29, 0.75) |
| PgR ⁺ | 1.0 (referent) | 1.0 (referent) |
| PgR ⁻ | 4.18 (1.26–13.87, 0.019) | 1.82 (0.75–4.38, 0.18) |
| p53 ⁺ | 1.0 (referent) | 1.0 (referent) |
| p53 ⁻ | 1.96 (0.74–5.18, 0.18) | 1.61 (0.63–4.12, 0.34) |
| Final step after stepwise regression: | | |
| Age (continuous) | 0.94 (0.92–0.98, 0.0012) | 0.94 (0.92–0.97, 0.0001) |
| Grade (continuous) | 3.25 (1.58–6.69, 0.0014) | |
| PgR ⁺ | 1.0 (referent) | |
| PgR ⁻ | 3.45 (1.22–9.80, 0.02) | |

ER, oestrogen receptor; PgR, progesterone receptor.

and Lakhani and colleagues [17], although in the former study the *BRCA2* cancers were more ER-positive and PgR-positive (on the basis of smaller sample set of 14 cases).

BRCA1 and *BRCA2* mutations have been detected in quite a low proportion of breast cancer families. In Finland, among families with three affected first-degree or second-degree relatives without age restrictions, the proportions were 10% and 11% for *BRCA1* and *BRCA2*, respectively [16,20]. Thus an important and large group of families is not due to mutations in the *BRCA1* and *BRCA2* genes. In this study, the familial non-*BRCA1/2* cancers were diagnosed at a marginally younger age than those among unselected cases, and were more often of lower grade than the control cancers or *BRCA1* and *BRCA2* cancers. These factors might be influenced by recall bias because patients having affected relatives undergo diagnostic procedures earlier than women with no family history of breast cancer. However, no such influence is seen for *BRCA1* or *BRCA2* cases, which often have a much stronger family background of cancer. Furthermore, a lower grade among familial non-*BRCA1/2* cases has been detected elsewhere as well [17,18].

In comparison with *BRCA1* or *BRCA2* cancers, familial non-*BRCA1/2* cancers represented a much greater

number of ER-positive and PgR-positive cases; however, in comparison with the unselected control group they were more receptor-negative. The frequency of ER-positive cancers among our control cancers is quite high, which might be due to a later year of diagnosis than for non-*BRCA1/2* cancers and in general because of age distribution [34], mammography screening [33] or ethnic differences [35] in different populations.

Conclusions

In this report we have characterised familial non-*BRCA1/2* tumours and evaluated routine immunohistochemical and pathological markers that could help us to further distinguish families carrying *BRCA1/2* mutations from other breast cancer families. It is noteworthy here that, although ER negativity has been considered a hallmark of *BRCA1* tumours, logistic regression analysis indicated that this was not an independent marker but was dependent on the age of diagnosis and tumour grade. When considering the possibility of mutation testing in the context of genetic counselling, for instance, it would be important to consider the tumour characteristics specifically in comparison with those from other breast cancer families. It also seems crucial to consider the histopathological features with regard to the age of the patients. In this study, the independent markers that distinguished *BRCA1* carrier tumours from familial non-*BRCA1/2* tumours were earlier age of diagno-

sis, negative PgR status and higher grade. These immunohistochemical and pathological characteristics of the tumours, which are available in routine pathological diagnostics, should be of value in evaluating the possibility of mutation and in targeting mutation screening in such families, especially when considering the characteristics of several tumours in the family and combined with the family history of cancer.

Competing interests

The author(s) declare that they have no competing interests.

Acknowledgements

We wish to thank families who participated in this study, Minna Merikivi for her help in sample collection, and the Finnish Cancer Registry for cancer diagnosis and identification numbers for archival material from the pathology laboratories. Financial support for this study was provided by grants from the Academy of Finland, from the Clinical Research Fund of Helsinki University Central Hospital, from the Finnish Breast Cancer Group, from the Finnish Cancer Society, from the Finnish-Norwegian Medical Foundation and from the Sigrid Juselius Foundation.

References

- Noguchi S, Kasugai T, Miki Y, Fukutomi T, Emi M, Nomizu T: **Clinicopathologic analysis of BRCA1- or BRCA2 associated hereditary breast carcinoma in Japanese women.** *Cancer* 1999, **85**:2200-2205.
- Foulkes WD, Chappuis PO, Wong N, Brunet JS, Vesprini D, Rozen F, Yuan ZQ, Pollak MN, Kuperstein G, Narod S, *et al.*: **Primary node negative breast cancer in BRCA1 mutation carriers has a poor outcome.** *Ann Oncol* 2000, **11**:307-313.
- Loman N, Johannsson O, Bendahl PO, Borg A, Ferno M, Olsson H: **Steroid receptors in hereditary breast carcinomas associated with BRCA1 or BRCA2 mutations or unknown susceptibility genes.** *Cancer* 1998, **83**:310-319.
- Lakhani SR, Van de Vijver MJ, Jacquemier J, Anderson TJ, Osin PP, McGuffog L, Easton D: **The pathology of familial breast cancer: predictive value of immunohistochemical markers estrogen receptor, progesterone receptor, HER-2, and p53 in patients with mutations in BRCA1 and BRCA2.** *J Clin Oncol* 2002, **20**:2310-2318.
- Eisinger F, Nogues C, Birnbaum D, Jacquemier J, Sobol H: **BRCA1 and medullary breast cancer.** *JAMA* 1998, **280**:1227.
- Phillips K, Andrulis IL, Goodwin PJ: **Breast carcinomas arising in carriers of mutations in BRCA1 or BRCA2: are they prognostically different?** *J Clin Oncol* 1999, **17**:3653-3663.
- Eisinger F, Stoppa-Lyonnet D, Longy M, Kerangueven F, Noguchi T, Bailly C, Vincent-Salomon A, Jacquemier J, Birnbaum D, Sobol H: **Germ line mutation at BRCA1 affect the histoprognostic grade in hereditary breast cancer.** *Cancer Res* 1996, **56**:471-474.
- Johannsson OT, Idvall I, Anderson C, Borg A, Barkardottir RB, Egilsson V, Olsson H: **Tumour biological features of BRCA1-induced breast and ovarian cancer.** *Eur J Cancer* 1997, **33**:362-371.
- Sorlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, Deng S, Johnsen H, Pesich R, Geisler S, *et al.*: **Repeated observation of breast tumour subtypes in independent gene expression data sets.** *Proc Natl Acad Sci USA* 2003, **100**:8418-8423.
- Foulkes W, Stefansson IM, Chappuis PO, Begin LR, Goffin JR, Wong N, Trudel M, Akslen LA: **Germline BRCA1 mutations and basal epithelial phenotype in breast cancer.** *J Natl Cancer Inst* 2003, **95**:1482-1485.
- Marcus JN, Watson P, Page DL, Narod S, Lenoir G, Tonin P, Linder-Stephenson L, Salerno G, Conway TA, Lynch H: **Hereditary breast cancer: pathobiology, prognosis, and BRCA1 and BRCA2 gene linkage.** *Cancer* 1996, **77**:697-709.
- Armes JE, Egan AJ, Southey MC, Dite GS, McCredie MR, Giles GG, Hopper JL, Venter DJ: **The histologic phenotypes of breast carcinoma occurring before age 40 years in women with and without BRCA1 or BRCA2 germline mutations: a population-based study.** *Cancer* 1998, **83**:2335-2345.
- Syrjäkoski K, Vahteristo P, Eerola H, Tamminen A, Kivinummi K, Sarantaus L, Holli K, Blomquist C, Kallioniemi OP, Kainu T, Nevanlinna H: **Population-based study of BRCA1 and BRCA2 mutations in 1035 unselected Finnish breast cancer patients.** *J Natl Cancer Inst* 2001, **93**:152-154.
- Hedenfalk I, Duggan D, Chen Y, Radmacher M, Bittner M, Simon R, Meltzer P, Gusterson B, Esteller M, Kallioniemi O, *et al.*: **Gene-expression profiles in hereditary breast cancer.** *N Engl J Med* 2001, **344**:539-548.
- Ford D, Easton DF, Stratton M, Narod S, Goldgar D, Devilee P, Bishop DT, Weber B, Lenoir G, Chang-Claude J, *et al.*: **Genetic heterogeneity and penetrance analysis of the BRCA1 and BRCA2 genes in breast cancer families.** *Am J Hum Genet* 1998, **62**:676-689.
- Vehmanen P, Friedman LS, Eerola H, McClure M, Ward B, Sarantaus L, Kainu T, Syrjäkoski K, Pyrhönen S, Kallioniemi O, *et al.*: **Low proportion of BRCA1 and BRCA2 mutations in Finnish breast cancer families: evidence for additional susceptibility genes.** *Hum Mol Genet* 1997, **6**:2309-2315.
- Lakhani SR, Gusterson BA, Jacquemier J, Sloane JP, Anderson TJ, van de Vijver MJ, Venter DJ, Freeman A, Antoniou AC, McGuffog L, *et al.*: **The pathology of familial breast cancer: histological features of cancers in families not attributable to mutations in BRCA1 or BRCA2.** *Clin Cancer Res* 2000, **6**:782-789.
- Palacios J, Honrado E, Osorio A, Cazorla A, Sarrío D, Barroso A, Rodrigues S, Cigudosa JC, Diez O, Alonso C, *et al.*: **Immunohistochemical characteristics defined by tissue microarray of hereditary breast cancer not attributable to BRCA1 or BRCA2 mutations: differences from breast carcinomas arising in BRCA1 and BRCA2 mutation carriers.** *Clin Cancer Res* 2003, **9**:3606-3614.
- Eerola H, Blomqvist C, Pukkala E, Pyrhönen S, Nevanlinna H: **Familial breast cancer in southern Finland: how prevalent are breast cancer families and can we trust the family history reported by patients?** *Eur J Cancer* 2000, **36**:1143-1148.
- Vehmanen P, Friedman LS, Eerola H, Sarantaus L, Pyrhönen S, Ponder BAJ, Muhonen T, Nevanlinna H: **A low proportion of BRCA2 mutations in Finnish breast cancer families.** *Am J Hum Genet* 1997, **60**:1050-1058.
- Huuskio P, Pääkkönen K, Launonen V, Pöyhönen M, Blanco G, Kauppila A, Puistola U, Kiviniemi H, Kujala M, Leisti J, *et al.*: **Evidence of founder mutations in Finnish BRCA1 and BRCA2 families.** *Am J Hum Genet* 1998, **62**:1544-1548.
- Sarantaus L, Huuskio P, Eerola H, Launonen V, Vehmanen P, Rappakko K, Gillanders E, Syrjäkoski K, Kainu T, Vahteristo P, *et al.*: **Multiple founder effects and geographical clustering of BRCA1 and BRCA2 families in Finland.** *Eur J Hum Genet* 2000, **8**:757-763.
- Elston C, Ellis I: **Pathological prognostic factors in breast cancer. I. The value of histological grade in breast cancer: experience from a large study with long-term follow-up.** *Histopathology* 1991, **19**:403-410.
- Vahteristo P, Bartkova J, Eerola H, Syrjäkoski K, Ojala S, Kilpivaara O, Tamminen A, Kononen J, Aittomäki K, Heikkilä P, *et al.*: **A CHK2 genetic variant contributing to a substantial fraction of familial breast cancer.** *Am J Hum Genet* 2002, **71**:432-438.
- Shattuck-Eidens D, Oliphant A, McClure M, McBride C, Gupte J, Rubano T, Pruss D, Tavtigian SV, Teng DH, Adey N, *et al.*: **BRCA1 sequence analysis in women at high risk for susceptibility mutations. Risk factor analysis and implications for genetic testing.** *JAMA* 1997, **278**:1242-1250.
- Couch FJ, DeShano ML, Blackwood MA, Calzone K, Stopfer J, Campeau L, Ganguly A, Rebbeck T, Weber BL: **BRCA1 mutations in women attending clinics that evaluate the risk of breast cancer.** *N Engl J Med* 1997, **336**:1409-1415.
- Chan-Claude J, Dong J, Schmidt S, Shayeghi M, Komitowski D, Becher H, Stratton MR, Royer-Pokora B: **Using gene carrier probability to select high risk families for identifying germline mutations in breast cancer susceptibility genes.** *J Med Genet* 1998, **35**:116-121.

28. Parmigiani G, Berry D, Aguilar O: **Determining carrier probabilities for breast cancer-susceptibility genes BRCA1 and BRCA2.** *Am J Hum Genet* 1998, **62**:145-158.
29. Vahteristo P, Eerola H, Tamminen A, Blomqvist C, Nevanlinna H: **A probability model for predicting BRCA1 and BRCA2 mutations in breast and breast-ovarian cancer families.** *Br J Cancer* 2001, **84**:704-708.
30. Gilpin CA, Carson N, Hunter AGW: **A preliminary validation of a family history assessment form to select women at risk for breast or ovarian cancer for referral to a genetic center.** *Clin Genet* 2000, **58**:299-308.
31. Nixon AJ, Neuberger D, Hayes DF, Gelman R, Connolly JL, Schnitt S, Abner A, Recht A, Vicini F, Harris JR: **Relationship of patient age to pathologic features of the tumor and prognosis for patients with stage I or II breast cancer.** *J Clin Oncol* 1994, **12**:888-894.
32. Ferno M, Borg A, Johansson U, Norgren A, Olsson H, Ryden S, Sellberg G: **Estrogen and progesterone receptor analyses in more than 4,000 human breast cancer samples. A study with special reference to age at diagnosis and stability of analyses.** *Acta Oncol* 1990, **29**:129-135.
33. Molino A, Pavarana M, Micciolo R, Nortilli R, Pedersini R, Manno P, Bozzo P, Bonetti F, Piubello Q, Cetto GL: **Comparative study of clinical, pathological, and biological characteristics of symptomatic versus asymptomatic breast cancers.** *Ann Oncol* 2000, **11**:581-586.
34. Talley LI, Grizzle WE, Waterbor JW, Brown D, Weiss H, Frost AR: **Hormone receptors and proliferation in breast carcinomas of equivalent histologic grades in pre- and postmenopausal women.** *Int J Cancer* 2002, **98**:118-127.
35. Chu KC, Anderson WF, Fritz A, Ries LAG, Brawley OW: **Frequency distributions of breast cancer characteristics classified by estrogen receptor and progesterone receptor status for eight racial/ethnic groups.** *Cancer* 2001, **92**:37-45.