



Review

Nutritional and activity recommendations for the child with normal weight, overweight, and obesity with consideration of food insecurity: An Obesity Medical Association (OMA) Clinical Practice Statement 2022



Nancy T. Browne^{a,*}, Suzanne E. Cuda^b

^a 25 Andrews Avenue, Falmouth, ME, 04105, USA

^b Alamo City Healthy Kids and Families, 1919 Oakwell Farms Parkway, Ste 145, San Antonio, TX, 78218, USA

ARTICLE INFO

Keywords:

Activity recommendation
Assessment
Food insecurity
Nutritional evaluation
Obesity
Pediatric

ABSTRACT

Background: This Obesity Medicine Association (OMA) Clinical Practice Statement (CPS) details nutritional and activity recommendations for the child with normal weight, overweight, and obesity (Appendix A) with consideration of food insecurity. This CPS is intended to provide clinicians with an overview of clinical practices applicable to children and adolescents with body mass indices in the normal range and body mass indices greater than or equal to the 85th percentile for their ages, particularly those with adverse consequences resulting from increased body mass. The information in this CPS is based on scientific evidence, supported by the medical literature, and derived from the clinical experiences of members of the OMA.

Methods: The scientific information and clinical guidance in this CPS is based upon referenced evidence and derived from the clinical perspectives of the authors.

Results: This OMA Clinical Practice Statement details nutritional and activity recommendations for the child with normal weight, overweight, and obesity with consideration of food insecurity. In addition, this CPS addresses nutritional recommendations for complications related to the disease of obesity as well as providing guidance on food insecurity as it impacts children with obesity and their families.

Conclusions: This OMA Clinical Practice Statement on nutritional and activity recommendations for the child with normal weight, overweight, and obesity with consideration of food insecurity is an overview of current recommendations. These recommendations provide a roadmap to the improvement of the health of children and adolescents with obesity, especially those with metabolic, physiological, and psychological complications.

1. Introduction

The purpose of this CPS regarding nutritional and activity recommendations for the child with normal weight, overweight, and obesity with consideration of food insecurity is to provide clinicians with tools to clinically assess and manage children with obesity. The OMA is an organization of providers in the field of obesity medicine dedicated to the comprehensive care of patients with obesity. OMA members are physicians, nurse practitioners, physician assistants, and other healthcare providers who take a comprehensive, evidence-based approach to treating obesity. This approach is comprised of the four pillars of nutrition, physical activity, behavior, and medication. While it is hoped many clinicians may find the recommendations in this CPS helpful, the final decision regarding the optimal care of the patient with overweight or

obesity is dependent upon the individual clinical presentation and the judgment of the clinician who is tasked with directing a treatment plan that is in the best interest of the patient.

2. Nutritional recommendations for children of normal weight

General nutritional recommendations for children and adolescents fall into age categories that are related to overall growth and development stages. For infants from 0 to 12 months of age, guidelines are further divided into birth to 4 months, 4–6 months, 6–8 months, 8–10 months, and 10–12 months due to the rapid physiologic and metabolic changes during the child's first year of life [1]. Recommendations from 0 to 12 months emphasize breastfeeding whenever possible. Fortified formula may be used when breastmilk is not available. Cereals, breads,

* Corresponding author. Andrews Avenue, Falmouth, ME, 04105, USA.

E-mail addresses: nancytkacz@sbcglobal.net (N.T. Browne), suzanne@alamocityhealthykid.com (S.E. Cuda).

<https://doi.org/10.1016/j.obpill.2022.100012>

Received 28 February 2022; Accepted 4 March 2022

2667-3681/© 2022 The Authors. Published by Elsevier Inc. on behalf of Obesity Medicine Association. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

starches, fruits, vegetables, and proteins (meat or legumes) are gradually introduced from 6 to 12 months of age [2].

From 1 to 4 years of age, children gradually transition to milk/milk products from breastmilk or formula. Intake of breads/starches, fruits, vegetables, and proteins gradually increases with age. Of note, low-fat products are not recommended in the first two years of life, with 3–4 teaspoons per day of fats and oil recommended from ages 3–5 years. Desserts, sweets, soft drinks, candy, jams, jelly and added sugar to foods are not advised during the first 5 years of life [1,2].

During ages 5–18 years, volume in all food categories increases as the child/adolescent grows and develops. Vegetables continue to be encouraged, particularly non-starchy vegetables such as broccoli, spinach, green beans, and tomatoes. Sweets and added sugars (including in all fluids) are not recommended [1,2]. Tables 1–3 offer more specific guidance, and Table 4 shows the top takeaways from the OMA regarding nutritional principles for children of normal weight.

3. Activity recommendations for children of normal weight, overweight, and obesity

Activity recommendations for children and adolescents of normal weight and for those with overweight/obesity also fall into age categories that correspond to growth and development stages. Caregivers should follow recommendations from pediatric providers skilled in growth and development principles.

For infants from 0 to 12 months of age, movement and active play with caregivers is encouraged for short periods several times a day. The setting should encourage and stimulate movement experiences, and the environment should meet or exceed recommended safety standards for performing large-muscle activities. These activities should promote skill development appropriate to the specific developmental skill of the infant. The key focus is exploring movement through the infant's environment through structured and unstructured physical activity [3,4].

For the child 12–36 months of age, activity should include a minimum of 30 minutes of structured and 60 minutes of unstructured physical activity per day. During this growth phase, activities should maximize any opportunity to develop movement skills that will serve as the building blocks for future motor skillfulness and physical activity. Children benefit from access to indoor and outdoor areas for performing large-muscle activities. Caretakers should promote movement skills by providing activity and physical movement experiences [3,4].

Recommendations for the child ages 3–5 years are similar, with structured physical activity increasing to 60 minutes daily. For children ages 5 through adolescence, a minimum of 60 minutes, with a preference of several hours, is recommended for age-appropriate physical activity and play. These activities should include several bouts of physical activity lasting 15 minutes or more each day designed to achieve optimal health,

Table 1

General intake guidelines (normal weight): 0 to 12 months. Infants from 0 to 12 years of age have unique nutritional considerations; nutritional recommendations are divided into categories for 0–4 months, 4–6 months, 6–8 months, 8–10 months, and 10–12 months [1,2].

	Birth–4 months	4–6 months	6–8 months	8–10 months	10–12 months
Breast milk and/or fortified infant formula	8 to 12 feedings *2 to 6 oz per feeding (18–32 oz per day)	4 to 6 feedings *4 to 6 oz per feeding (27–45 oz per day)	3 to 5 feedings *6 to 8 oz per feeding (24–32 oz per day)	3 to 4 feedings *7 to 8 oz per feeding (24–32 oz per day)	3 to 4 feedings *24–32 oz per day
Cereal, breads, starches	None	None	2–3 servings of iron-fortified baby cereal (serving = 1–2 tbsp)	2–3 servings of iron-fortified baby cereal (serving = 1–2 tbsp)	4 servings of iron-fortified bread or other soft starches or baby cereal (serving = 1–2 tbsp)
Fruits and vegetables	None	None	Offer plain, cooked, mashed, or strained baby foods vegetables and fruits. Avoid combination foods. No juice.	2–3 servings (1–2 tbsp) of soft, cut-up, and mashed vegetables and fruits daily. No juice.	4 servings (2–3 tbsp) daily of fruits and vegetables. No juice.
Meats and other protein sources	None	None	Begin to offer plain-cooked meats. Avoid combination dinners.	Begin to offer well-cooked, soft, finely chopped meats.	1–2 oz daily of soft, finely cut or chopped meats or other protein foods

While there is no comprehensive research indication of which complimentary foods are best to introduce first, the focus should be on first foods that are higher in iron and zinc, such as pureed meats and fortified-iron rich foods.

Table 2

General intake guidelines (normal weight): 1 to 4 years. Children from 1 to 4 years of age gradually transition to milk and milk products and begin to incorporate more proteins, breads, grains, starches, fruits, and vegetables [1,2].

	12–23 months	2–3 years	3–4 years
Milk and Milk Products	2 cups/day (whole milk or milk products)	2–2.5 cups/day	2.5–3 cups/day
Serving: 1 cup of milk or cheese	1½ oz of natural cheese	1/3 cup shredded cheese	
Meat and Other Protein Foods	1½ oz/day	2 oz/day	2–3 oz/day
Serving: (1 oz equivalent) = 1 oz beef, poultry, fish, ¼ cup cooked beans, 1 egg, 1 tbsp peanut butter*, ½ oz of nuts*			
*peanut butter and nuts may be a choking hazard under the age of three			
Breads, Cereal, and Starches	2 oz/day	2 oz/day	2–3 oz/day
1 oz = 1 slice whole grain bread, ½ cup cooked cereal, rice, pasta or 1 cup dry cereal			
Fruits	1 cup/day	1 cup/day	1–1½ cups/day
Serving: 1 cup of fruit or ½ cup dried fruit; NO JUICE			
Vegetables (non-starchy vegetables to include sources of vitamin C and A)	3/4 cup/day	1 cup/day	1–1½ cups/day
Serving: (1 cup equivalent) = 1 cup of raw or cooked vegetables; 2 cups of raw leafy green greens			
Fats and Oil	Do not limit* *Low-fat products are not recommended under the age of 2	3 teaspoons	3–4 teaspoons/day
Miscellaneous desserts, sweets, soft drinks, candy, jams, jelly	none	none	None

wellness, fitness, and performance benefits. Extended periods of inactivity are discouraged, especially during the daytime hours [3,4]. Table 5 shows the top takeaways from the OMA regarding activity recommendations for children for normal weight.

4. Nutrition management options for children with overweight and obesity

Evidence suggests that weight loss in children with obesity is possible regardless of the macronutrient distribution in an energy-reduced diet [6]. Similar results have been found in adult studies [7]. Tailoring the macronutrient content to address specific obesity-related complications may be helpful, but more research is needed. The pediatric provider uses the principles of precision medicine to consider which nutritional option best addresses a particular child and family's needs. Examples of macronutrient options and comparisons of common recommendations follow in Figs. 1 and 2. Table 6 shows the top takeaways from the OMA regarding nutritional interventions for the child with obesity. All

Table 3

General intake guidelines (normal weight): 5 to 18 years. From 5 to 18 years of age, the recommendations for children in all food categories increase in volume as the children grow and develop. There is a continued emphasis on the inclusion of non-starchy vegetables; desserts, sweets, and other added-sugar foods are best avoided [1,2].

	12–23 months	2–3 years	3–4 years
Milk and Milk Products	2 cups/day (whole milk or milk products)	2–2.5 cups/day	2.5–3 cups/day
Serving: 1 cup of milk or cheese, 1½ oz of natural cheese, 1/3 cup shredded cheese			
Meat and Other Protein Foods	1½ oz/day	2 oz/day	2–3 oz/day
Serving: (1 oz equivalent) = 1 oz beef, poultry, fish, ¼ cup cooked beans, 1 egg, 1 tbsp peanut butter*, ½ oz of nuts*			
*peanut butter and nuts may be a choking hazard under the age of three			
Breads, Cereal, and Starches	2 oz/day	2 oz/day	2–3 oz/day
Serving: 1 oz = 1 slice whole grain bread, ½ cup cooked cereal, rice, pasta or 1 cup dry cereal			
Fruits	1 cup/day	1 cup/day	1–1½ cups/day
Serving: 1 cup of fruit or ½ cup dried fruit; NO JUICE			
Vegetables (non-starchy vegetables to include sources of vitamin C and A)	¾ cup/day	1 cup/day	1–1½ cups/day
Serving: (1 cup equivalent) = 1 cup of raw or cooked vegetables; 2 cups of raw leafy green greens			
Fats and Oil	Do not limit* *Low-fat products are not recommended under the age of 2	3 teaspoons	3–4 teaspoons/day
Miscellaneous desserts, sweets, soft drinks, candy, jams, jelly	None	none	None

Table 4

Top takeaways: nutrition recommendations for children of normal weight. Shown are the top takeaways from the OMA regarding nutrition for children of normal weight, including increases in food quantities, encouragement of vegetables, and avoidance of sweets and added sugars.

1. Dietary recommendations for infants and children reflect their stages of growth and development.
2. Breastfeeding is encouraged during the first 12 months of life, with the gradual introduction of solids after 6 months of age.
3. Low-fat products are not recommended under the age of 2 years. Fats and oils are gradually increased from 3 to 6 teaspoons per day from ages 2–18 years.
4. Desserts, sweets, and added sugar, especially in fluids, are not recommended at any age.
5. Vegetables, especially non-starchy vegetables, are encouraged in various forms starting at 6 months of age and continuing through adulthood, but juices should be avoided.

Table 5

Top takeaways: activity recommendations for children of normal weight, overweight, and obesity. Shown are the OMA's top takeaway messages regarding activity recommendations for children of normal weight [5].

1. Activity recommendations exist for children and adolescents of normal weight and overweight/obesity at each developmental stage.
2. Structured activity and unstructured play are equally key components of an optimal activity plan.
3. Extended periods of inactivity are discouraged at any age.

nutritional interventions are more effective in combination with activity and behavioral counseling, which includes addressing conditions detrimental to the child and family's social determinants of health.

5. Nutritional recommendations for complications associated with obesity in children

Precision medicine uses traditional principles of dietary management

of obesity-related complications, tailoring them to specific children, complications, and circumstances. General dietary management for children with obesity-related complications (type 2 diabetes, hypertension, metabolic syndrome, familial hypercholesterolemia, non-alcoholic fatty liver disease, and attention deficit hyperactivity disorder) follow; these can be adapted for the specific developmental age, disease progression, and individual response of the patient.

5.1. Type 2 diabetes

Conventional management of type 2 diabetes (T2D) suggests a nutrient-dense diet consisting of high fiber and low sugar, with approximately 30 grams of carbohydrate (CHO) per meal. If the baseline fasting insulin level is elevated, an alternative meal plan might include low CHO content with a well-balanced diet and nonrestricted caloric intake. The source of CHO is important and varies with individuals in maintaining glucose control. Frequent monitoring is recommended. Guidance from nutritional professionals is critical for long-term success [23,24].

5.2. Hypertension

The Dietary Approaches to Stop Hypertension (DASH) diet is a nutritional plan based on an intake of 2300 mg of sodium per day or less (1500 mg/day or less for individuals identifying as non-Hispanic Black). Adherence to the plan is associated with a lowering of blood pressure. The plan concentrates on vegetables, fruits, whole grains, low-fat meats/proteins, fish, and low-fat dairy with limitations on sugar, sweets, and sodium [25–27].

5.3. Metabolic syndrome

The Framingham Study is a long-term, population-based study that began in 1948 to investigate cardiovascular disease; in this study, whole grain intake was inversely associated with metabolic disease prevalence while glycemic index was positively associated with prevalence [28].

The source and quality of dietary carbohydrates may differentially optimize insulin action. Fasting insulin concentrations are lower among individuals reporting higher dietary fiber or whole grain intake [29].

5.4. Familial hypercholesterolemia (FH)

The National Heart, Lung, and Blood Institute (NHLBI) Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents has created recommendations to help clinicians promote cardiovascular (CV) health and manage CV risk factors in children and adolescents [30]. Once the diagnosis of FH is made, these recommendations suggest initiating the NHLBI Child 2 diet plan based on less than 7% saturated fat and less than 200 mg/day of cholesterol per day. Since most children with obesity have LDL <140 mg/dl, if higher levels are found, the clinician should suspect heterozygous FH [31]. It is important for both children and parents to know that while reducing dietary saturated fat and dietary cholesterol may have lifelong health benefits, the cholesterol blood levels of children and adults is mainly dependent upon the clinical impact of the genetic abnormality causing FH. No child with FH or parent of a child with FH should have expectations that cholesterol blood levels will substantially improve with diet alone [32]. Failure to convey this message may lead to unwarranted blame and guilt, as well as unnecessary delay in starting pharmacological therapy (e.g., statins) [33,34].

5.5. Non-alcoholic fatty liver disease (NAFLD)

The spectrum of liver disease related to obesity varies considerably in the degree of pathology and genetic predisposition. The dietary intervention for known or presumed NAFLD is a nutrient-dense diet rich in fiber (legumes, whole grains), protein, and unsaturated fats. A significant

Portion Control or Balanced	CHO Restricted or Reduced	Low Glycemic Index	Low Fat	Elimination
<ul style="list-style-type: none"> • Tolerance is high • Useful in toddlers and young children • Small amount of weight loss 	<ul style="list-style-type: none"> • Adherence to diet is approximately 50% • Weight loss is moderate to good • Lowers fasting insulin and triglyceride levels • Amount of protein is not associated with effect on muscle sparing 	<ul style="list-style-type: none"> • Tolerance is high • Small amount of weight loss • Decreases postprandial glucose response, slows insulin secretion, and delays gastric emptying leading to longer satiety • Favorable for high fasting insulin level 	<ul style="list-style-type: none"> • Tolerance is high • Limited trials: in 3 RCTs, small reduction in BMI, total and LDL Cholesterol • No weight loss or minimal • No difference between low fat & low glycemic in reduction of steatosis 	<ul style="list-style-type: none"> • Used mostly in fast food or cafeteria type settings • No clear evidence for effectiveness • Weight loss is small to moderate • Tolerance is high • Not same as elimination diets for food allergies/intolerance

Fig. 1. Nutritional Therapy: Comparison of Common Recommendations I. Nutritional management for pediatric patients can take several forms. These include portion control, CHO-restricted or reduced diets, low glycemic index foods, low-fat diets, and elimination diets; information, pros, and cons for each are shown [6, 8–19].

CHO: carbohydrate; RCT: randomized controlled trial; P: protein; KD: Ketogenic diet; BMI: body mass index; LDL: low-density lipoprotein.

Ketogenic Diet (KD)	Ketogenic Diet (KD)	Intermittent Fasting or Time-Restricted Feeding	Plant-Based
<ul style="list-style-type: none"> • Used in children for epilepsy; few RCTs in obesity • Ketosis ratio: $(0.9F+0.46P)/(1.0C+0.58P+0.1F)$, where F,P,C are grams of fat, protein, carbohydrates (CHO) • Current popular version is 20–50 grams of CHO/day • Fat content 70–80% • Protein 1.2–1.5 g/kg/d to maintain ketosis • Levels of 20 g/d CHO require eliminating fiber-rich starchy vegetables, most fruit, legumes and whole grains 	<ul style="list-style-type: none"> • Children may gain weight if they increase their meat consumption significantly, thereby increasing saturated fat • KD may impair glucose tolerance compared to diets containing >20 g of CHOs • KD may achieve improved metabolic control quickly but may be more difficult for a child to stay on long term 	<ul style="list-style-type: none"> • Types: 5 days on, 2 days off; alternate days fasting and ad lib eating; eating for 6 hours and fasting for 18 hours • Some studies show consumption is greater on fasting days than planned and less on non fasting days • Most studies show poor adherence • Not enough research in children to recommend • May lead to disordered eating • Not a great option for those with binge eating disorder or depression 	<ul style="list-style-type: none"> • Seen as a population-wide risk factor strategy to reduce CV disease • Excludes products of animal origin, e.g., meat, eggs, and fish • Promoted as more sustainable for the planet • Lower energy density than an omnivore diet if high in vegetables, fruit, legumes, and nuts • No significant difference in weight status in children in a small 4-week study • More studies needed

Fig. 2. Nutritional Therapy: Comparison of Common Recommendations II. Nutritional management for pediatric patients can take several forms, including diets that are popular with adults. They may include the ketogenic diet, intermittent fasting/time-restricted feeding, and plant-based diets; information, pros, and cons are shown [20–22].

CHO: carbohydrate; RCT: randomized controlled trial.

Table 6
Top takeaways: nutritional interventions for the child with overweight and obesity. Shown are the top takeaways from the OMA regarding nutritional interventions for the child with obesity.

1. Improvement in weight status in children with overweight and obesity is possible regardless of the macronutrient distribution in an energy-reduced diet.
2. The choice of which macronutrient option to implement is based on a thorough assessment with the family along with shared decision making.
3. Frequent evaluation of the macronutrient choice allows for adjustment or change to a better option, reducing wasted time and frustration.
4. All nutritional interventions are more effective in combination with activity and behavioral counseling, which includes addressing conditions detrimental to the child/family's social determinants of health.

reduction in foods and beverages with added sugar, salt, saturated fat, and refined carbohydrates (high glycemic index) is encouraged [35].

5.6. Attention deficit hyperactivity disorder (ADHD)

Overall, children with ADHD should follow a healthy diet with well-balanced nutrients that is low in added sugar, low in sodium, and high in fiber. While no specific diet has been shown to improve symptoms of ADHD, some research suggests that ADHD is associated with disordered eating that may contribute to weight gain [36]. Some children with ADHD demonstrate difficulty with mundane and repetitive tasks such as following a dietary plan. Strategies such as encouraging choices in diet

Table 7

Top Takeaways: Nutritional Recommendations for Complications Associated with Obesity in Children. Shown are the top takeaways from the OMA regarding nutritional recommendations for complications associated with obesity in children.

1. Precision medicine uses traditional principles of dietary management of obesity-related complications, tailoring them to specific children, complications, and circumstances.
2. The source of CHO is important in managing T2D and varies with individuals in maintaining glucose control.
3. The DASH diet is a nutritional plan based on a 2300 mg of sodium/day (1500 mg/day for individuals identifying as non-Hispanic Black).
4. The NHBLI Expert Panel recommends the Child 2 Diet Plan to help clinicians promote cardiovascular (CV) health and manage children and adolescents at risk for CV disease.
5. Dietary intervention for known or presumed NAFLD is a nutrient-dense diet rich in fiber (legumes, whole grains), protein, and unsaturated fats.
6. Children with ADHD should follow a healthy diet with well-balanced nutrients that is low in added sugar, low in sodium, and high in fiber, as no specific diet has been shown to improve symptoms of ADHD.

T2D: type 2 diabetes; CHO: carbohydrates; NHBLI: National Heart, Lung, and Blood Institute; NAFLD: non-alcoholic fatty liver disease; ADHD: attention deficit hyperactivity disorder.

selection offers a needed sense of control for the child [37]. Table 7 shows the top takeaways from the OMA regarding nutritional recommendations for complications associated with obesity in children.

6. Activity management for the pediatric patient with overweight and obesity

There is ongoing evidence that youth with obesity may experience a range of neuromuscular injuries and physical complications including increased pain, reduced muscle strength, postural malalignment, increased fatigue, reduced flexibility, and impaired balance, motor skills, and gait [38]. A review of 21 systematic reviews found a substantial impact on the physical health of children with overweight and obesity, predominantly represented by impairments in body structures and function [39]. Children with overweight and obesity are often compromised during weight-bearing aerobic exercise; specifically, they are metabolically compromised due to impaired fat oxidation and insulin sensitivity, biomechanically disadvantaged during walking and running, and emotionally compromised due to teasing and bullying [40,41]. Neuromuscular impairments may limit engagement in effective obesity treatment and require a redesign of obesity interventions specific to the needs and capabilities of the child and family.

The impact of physical activity on metabolism is dependent on the type, intensity and volume of activity pursued. Impact is also dependent on the genetic potential, gender and age, health, and current training status of the individual child. The benefits of long-term exercise training, particularly for youth with obesity, include reduction in fat mass, increases in fat free mass and fat oxidation, improvement in metabolic function and insulin sensitivity, and control of low-grade systemic inflammation [42].

Tools are available to support health professionals in assessing, adapting, and evaluating their treatment plan related to neuromusculoskeletal health in children with overweight and obesity [38,42]. Collaborative goal setting and use of the FITT-VP principles (frequency, intensity, time, type, volume, and progression) is recommended [38,43].

Strategies for youth with any degree of physical inactivity include using an incremental approach to reach the recommended 60 minutes per day by increasing activity by 10% per week [42,44]. Progressing too quickly is counterproductive and may lead to injury. Realistic and obtainable physical activity goals comparable to the child's own baseline abilities, not normal weight peers, allow for individualized management [44]. Regular assessment of the child's physical activity level allows the health care professional to provide positive feedback and adjust recommendations as needed. Other characteristics of successful activities include:

- Fun and entertaining
- Developmentally appropriate
- Promote aerobic fitness, muscular strength and endurance, and flexibility
- Consider the physical and emotional limits of the child

Table 8 shows the top takeaways from the OMA regarding activity management options for the pediatric patient with obesity.

7. Summary of recommendations for the child above the 85th percentile (overweight & obesity)

The short-term goals in treating children with overweight and obesity include interrupting the trajectory of abnormal weight gain and assessing and managing any obesity-related complications. The long-term goal in treating children with overweight and obesity is slow, steady reduction of BMI percentile until obesity-related complications are resolved followed by weight maintenance [45]. Multidisciplinary interventions include nutritional strategies, activity modifications, behavioral strategies, and improved sleep hygiene. Current evidence suggests that improved weight status can be achieved in children and adolescents with overweight or obesity irrespective of the macronutrient distribution of a reduced-energy diet. Tailoring the macronutrient content to target specific cardio-metabolic risk factors, such as a low-carbohydrate diet to treat insulin resistance, may be possible [6]. Figs. 3–7 show management strategies for children at different stages of development, and Table 9 shows a summary of OMA recommendations for children above the 85th percentile (overweight & obesity).

8. Food insecurity: impact and guidance

Food insecurity is defined as being uncertain of having, or unable to acquire, enough food to meet the needs of all family members because of insufficient money or other resources for food [48]. As of 2020, the US Department of Agriculture reports that 10.5% of Americans are food insecure, with 6.6% reporting low food security and 3.9% reporting very low food security; in households with children, 14.8% reported food insecurity, which represents 6.1 million children [49]. A 2017 study showed that the odds of being obese were 5 times higher for children from food insecure households when compared to children from food-secure households [50]. Those with greater risk of food insecurity include children in immigrant families, families headed by single women, increased members in household, and parental separation/divorce [51]. While food insecurity is prevalent in these groups, families with greater resources are also periodically at risk. 16% of low-income families do not receive federal support to supplement household food [52]. Participation in a Food Assistance Program (such as NSLP/WIC/SNAP) shows a reduction in food insecurity [53]. Intermittent access to adequate food can result in unhealthy eating patterns and increased stress, anxiety, and depression [54–59]. Other sequelae from the chronic stress of FI include dysregulated behavior, emotional distress, and suicidal ideation

Table 8

Top Takeaways for Activity Management Options for the Pediatric Patient with Overweight and Obesity. Shown are the top takeaways from the OMA regarding activity management options for the pediatric patient with overweight and obesity.

1. Appropriate and reliable physical activity assessment tools are available for pediatric patients with overweight and obesity.
2. Physiologic, metabolic, and psychologic factors may limit physical activity in pediatric patients with overweight and obesity.
3. Evidence-based recommendations and/or protocols exist for physical activity in the treatment of the pediatric patient with overweight and obesity.
4. Physical activity health professionals are available for appropriate referrals for the child with overweight and obesity.




Fig. 3. Management of the Infant with Overweight and Obesity: 0 to 24 Months. Recommendations for the infant with overweight and obesity (0–24 months) address optimal behavior and sleep patterns, food intake, and activity levels [1,2,46,47].



Fig. 4. Management of the Toddler with Overweight and Obesity: 2 to 4 Years. Recommendations for the toddler with overweight and obesity (2–4 years) address optimal behavior and sleep patterns, food intake, and activity levels [1,2,46,47].




Fig. 5. Management of the Young Child with Overweight and Obesity: 5 to 9 Years. Recommendations for the young child with overweight and obesity (5–9 years) address optimal behavior and sleep patterns, pharmacology, food intake, and activity levels [1,2,46,47]. ACE: Angiotensin-converting enzyme.

(particularly in adolescents) [52,60,61].

The American Academy of Pediatrics (AAP) recommends that all

children seen by their health professional are screened for food insecurity using a validated tool [51]. A positive screen for FI includes a positive




Fig. 6. Management of the Pubertal Child with Overweight and Obesity: 10 to 14 Years. Recommendations for the pubertal child with obesity (10–14 years) address optimal behavior and sleep patterns, pharmacology, food intake, and activity levels [1,2,46,47]. ACE: Angiotensin-converting enzyme; SGA: second-generation antipsychotics; T2D: type 2 diabetes.




Fig. 7. Management of the Adolescent Child with Overweight and Obesity: 15 to 18 Years. Recommendations for the adolescent child with overweight and obesity (15–18 years) address optimal behavior and sleep patterns, pharmacology, food intake, and activity levels [1,2,46,47]. ACE: Angiotensin-converting enzyme; SGA: second-generation antipsychotics; T2D: type 2 diabetes.

response to either or both of the following 2 questions [62]:

1. Within the past 12 months, we worried whether our food would run out before we got money to buy more. (Yes or No)
2. Within the past 12 months, the food we bought just didn't last and we didn't have money to get more. (Yes or No)

The AAP recommendations for a positive screening include being familiar with community, state, and federal resources; having an

awareness of nutritional content of these resources; and having an awareness of factors that increase vulnerability for FI. The AAP also encourages pediatric providers to advocate for their patients with FI at the local, state, and national levels [51]. Table 10 shows the top takeaways from the OMA regarding food insecurity.

9. Conclusions

This Clinical Practice Statement on nutritional and activity

Table 9

Top Takeaways: Summary Recommendations for the Child Above the 85th Percentile (Overweight and Obesity). Shown are the top takeaways from the OMA regarding recommendations for the child above the 85th percentile, including goals and interventions.

1. The short-term goals in treating children with overweight and obesity include:
 - a Interrupting the trajectory of abnormal weight gain
 - b Assessing and managing any obesity-related complications
2. The long-term goal in treating children with obesity is slow, steady weight loss until obesity-related complications are resolved and then weight maintenance.
3. Interventions include nutritional strategies, activity modifications, behavioral strategies, and improved sleep hygiene.

Table 10

Top Takeaways: Food Insecurity: Impact and Guidance. Shown are the top takeaways from the OMA regarding food insecurity as it relates to pediatric patients with overweight and obesity.

1. Food insecurity is defined as being uncertain of having, or unable to acquire, enough food to meet the needs of all family members because of insufficient money or other resources for food [48].
2. As of 2020, 10.5% of Americans are food insecure, 6.6% report low food security, 3.9% report very low food security [49].
3. In households with children, 14.8% reported food insecurity representing 6.1 million children [49].
4. The odds of children having obesity are 5 times higher in food insecure households compared to food-secure households [50].
5. Children from immigrant families, families headed by single women, families with increased members in household, and parental separation/divorce have greater risk of FI [51].
6. Families with greater resources are also periodically at risk for FI [52].
7. Participation in a food assistance program (such as NSLP/WIC/SNAP) shows a reduction in FI [53].
8. Intermittent access to adequate food can result in unhealthy eating patterns and increased stress, anxiety, and depression [54–59].
9. The American Academy of Pediatrics (AAP) recommends that all children seen by their health professional are screened for food insecurity using a validated tool [51].
10. A positive screen for FI includes a positive response to either or both of the following 2 questions [62]:
 1. Within the past 12 months, we worried whether our food would run out before we got money to buy more. (Yes or No)
 2. Within the past 12 months, the food we bought just didn't last and we didn't have money to get more. (Yes or No)

recommendations for the child with normal weight, overweight, and obesity with consideration of food insecurity provides clinicians with recommendations regarding their pediatric patients. The nutritional and activity interventions presented may lead to improvements in the health and wellbeing of children and adolescents with overweight and obesity, especially those with metabolic, physiological, and psychological complications.

Transparency [63]

This manuscript was largely derived and edited from the 2020–2022 Obesity Medicine Association (OMA) Pediatric Obesity Algorithm. Beginning in 2016, the OMA created and maintained an online Pediatric “Obesity Algorithm” (i.e., educational slides and eBook) that underwent updates approximately every two years by OMA authors and was reviewed and approved annually by the OMA Board of Trustees. Authors of prior years’ versions are included in Supplement #1. This manuscript is the first published version of the applicable chapter/s of the 2020–2022 OMA Pediatric Obesity Algorithm.

Group composition

Over the years, the authors of the OMA Pediatric Obesity Algorithm have represented a diverse range of clinicians, allied health professionals, clinical researchers, and academicians. (Supplement #1) The authors

reflect a multidisciplinary and balanced group of experts in obesity science, patient evaluation, and clinical treatment.

Author contributions

NTB transcribed the first draft of this CPS from the 2021 OMA Pediatric Obesity Algorithm. SEC then reviewed, edited, and approved the document for pre-peer review submission and post-peer review publication.

Disclosures (declaration of potential competing interest)

Potential dualities or conflicts of interest of the authors are listed in the Individual Disclosure section. Assistance of a medical writer paid by the Obesity Medicine Association is noted in the Acknowledgements section. Neither the prior OMAOMA Pediatric Algorithms nor the publishing of this Clinical Practice Statement received outside funding. The authors of prior OMA Pediatric Obesity Algorithms never received payment for their writing, editing, and publishing work. Authors of this Clinical Practice Statement likewise received no payment for their writing, editing, and publishing work. While listed journal Editors received payment for their roles as Editors, they did not receive payment for their participation as authors.

Individual Disclosures

SEC declares a relationship with Novo Nordisk as a member of an Advisory Board and a relationship with Rhythm Pharmaceuticals as a member of their Gold Panel. NTB reports no disclosures pertaining to this project.

Evidence

The content of the OMA Pediatric Obesity Algorithm and this manuscript is supported by citations, which are listed in the References section.

Ethics review

After approval by the authors, a draft manuscript was peer-reviewed and approved by the OMA Board of Trustees prior to publication. This submission did not involve human test subjects or volunteers.

Conclusions and recommendations

This Clinical Practice Statement is intended to be an educational tool that incorporates the current medical science and the clinical experiences of obesity specialists. The intent is to better facilitate and improve the clinical care and management of patients with pre-obesity and obesity. This Clinical Practice Statement should not be interpreted as “rules” and/or directives regarding the medical care of an individual patient. The decision regarding the optimal care of the patient with overweight and obesity is best reliant upon a patient-centered approach, managed by the clinician tasked with directing an individual treatment plan that is in the best interest of the individual patient.

Updating

It is anticipated that sections of this Clinical Practice Statement may require future updates. The timing of such an update will depend on decisions made by Obesity Pillars Editorial team, with input from the OMA members and OMA Board of Trustees.

Disclaimer and limitations

Both the OMA Obesity Algorithms and this Clinical Practice

Statement were developed to assist health care professionals in providing care for patients with pre-obesity and obesity based upon the best available evidence. In areas regarding inconclusive or insufficient scientific evidence, the authors used their professional judgment. This Clinical Practice Statement is intended to represent the state of obesity medicine at the time of publication. Thus, this Clinical Practice Statement is not a substitute for maintaining awareness of emerging new science. Finally, decisions by practitioners to apply the principles in this Clinical Practice Statement are best made by considering local resources, individual patient circumstances, patient agreement, and knowledge of federal, state, and local laws and guidance.

Declaration of competing interest

The authors declare the following financial interests/personal

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.obpill.2022.100012>.

Appendix B

Weight Category (2–19 years)	BMI Percentile Range for Age & Gender (2–19 years)
Underweight	Less than 5 th percentile
Healthy Weight	5 th to ≤ 85 th percentile
Overweight	85 th to ≤ 95 th percentile
Obesity	95 th to ≤ 99 th percentile
Severe Obesity	Above 99 th percentile

Glossary: Definition of terms in pediatric obesity.

References

- [1] Davis MM, Gance-Cleveland B, Hassink S, Johnson R, Paradis G, Resnicow K. Recommendations for prevention of childhood obesity. *Pediatrics* 2007;120(Suppl 4):S229–53.
- [2] Krebs NF, Himes JH, Jacobson D, Nicklas TA, Guilday P, Styne D. Assessment of child and adolescent overweight and obesity. *Pediatrics* 2007;120(Suppl 4):S193–228.
- [3] SHAPE America. Active start: a statement of physical activity guidelines for children from birth to age 5 nEC. IL: Human Kinetics Publishing; 2009.
- [4] Pediatrics AAo. Active healthy living: prevention of childhood obesity through increased physical activity. *Pediatrics* 2006;117:1834–42.
- [5] Yogman ea. Physical activity guidelines for Americans, 2nd edition. Committee on psychosocial aspects of child and family health and council on communications and media. *Pediatrics* 2018;142(3):e20182058.
- [6] Gow ML, Ho M, Burrows TL, Baur LA, Stewart L, Hutchesson MJ, et al. Impact of dietary macronutrient distribution on BMI and cardiometabolic outcomes in overweight and obese children and adolescents: a systematic review. *Nutr Rev* 2014;72:453–70.
- [7] Ge L, Sadeghirad B, Ball GDC, da Costa BR, Hitchcock CL, Svendrovski A, et al. Comparison of dietary macronutrient patterns of 14 popular named dietary programmes for weight and cardiovascular risk factor reduction in adults: systematic review and network meta-analysis of randomised trials. *BMJ* 2020;369:m696.
- [8] Gundersen CZJ. Childhood food insecurity in the U.S.: trends, causes, and policy options. Retrieved from, <http://www.futureofchildren.org/publications/docs/ResearchReport-July.2018.pdf>; 2014.
- [9] Ramon-Krauel M, Salsberg SL, Ebbeling CB, Voss SD, Mulkern RV, Apura MM, et al. A low-glycemic-load versus low-fat diet in the treatment of fatty liver in obese children. *Child Obes* 2013;9:252–60.
- [10] Epstein LH, Paluch RA, Beecher MD, Roemmich JN. Increasing healthy eating vs. reducing high energy-dense foods to treat pediatric obesity. *Obesity* 2008;16:318–26.
- [11] Krebs NF, Gao D, Gralla J, Collins JS, Johnson SL. Efficacy and safety of a high protein, low carbohydrate diet for weight loss in severely obese adolescents. *J Pediatr* 2010;157:252–8.
- [12] Fisher JO, Goran MI, Rowe S, Hetherington MM. Forefronts in portion size. An overview and synthesis of a roundtable discussion. *Appetite* 2015;88:1–4.
- [13] Kirk S, Woo JG, Brehm B, Daniels SR, Saelens BE. Changes in eating behaviors of children with obesity in response to carbohydrate-modified and portion-controlled diets. *Child Obes* 2017;13:377–83.
- [14] Kral TV, Hetherington MM. Variability in children's eating response to portion size. A biobehavioral perspective. *Appetite* 2015;88:5–10.
- [15] Watowicz RPTR, Hummel JC. The protein-sparing modified fast for adolescents with severe obesity. *Infant Child Adolesc Nutr* 2015;7:233–41. 2015.
- [16] Visuthranukul C, Sirimongkol P, Prachansuwan A, Pruksananonda C, Chomtho S. Low-glycemic index diet may improve insulin sensitivity in obese children. *Pediatr Res* 2015;78:567–73.
- [17] Schwingshackl L, Hoffmann G. Long-term effects of low-fat diets either low or high in protein on cardiovascular and metabolic risk factors: a systematic review and meta-analysis. *Nutr J* 2013;12:48.
- [18] Jenkins DJ, Kendall CW, Augustin LS, Franceschi S, Hamidi M, Marchie A, et al. Glycemic index: overview of implications in health and disease. *Am J Clin Nutr* 2002;76:266s–73s.
- [19] Dodds P, Wolfenden L, Chapman K, Wellard L, Hughes C, Wiggers J. The effect of energy and traffic light labelling on parent and child fast food selection: a randomised controlled trial. *Appetite* 2014;73:23–30.
- [20] Grandl G, Straub L, Rudigier C, Arnold M, Wueest S, Konrad D, et al. Short-term feeding of a ketogenic diet induces more severe hepatic insulin resistance than an obesogenic high-fat diet. *J Physiol* 2018;596:4597–609.
- [21] Stockman MC, Thomas D, Burke J, Apovian CM. Intermittent fasting: is the wait worth the weight? *Curr Obes Rep* 2018;7:172–85.
- [22] Desmond MA, Sobiecki J, Fewtrell M, Wells JCK. Plant-based diets for children as a means of improving adult cardiometabolic health. *Nutr Rev* 2018;76:260–73.
- [23] Gardner CD, Trepanowski JF, Del Gobbo LC, Hauser ME, Rigdon J, Ioannidis JPA, et al. Effect of low-fat vs low-carbohydrate diet on 12-month weight loss in overweight Adults and the association with genotype pattern or insulin secretion: the DIETFITS randomized clinical trial. *JAMA* 2018;319:667–79.
- [24] Ebbeling CB, Feldman HA, Klein GL, Wong JMW, Bielak L, Steltz SK, et al. Effects of a low carbohydrate diet on energy expenditure during weight loss maintenance: randomized trial. *BMJ* 2018;363:k4583.
- [25] Cohen JFW, Lehnert ME, Houser RF, Rimm EB. Dietary Approaches to Stop hypertension diet, weight status, and blood pressure among children and adolescents: national health and nutrition examination surveys 2003–2012. *J Acad Nutr Diet* 2017;117:1437–44.e2.
- [26] Carey RM, Calhoun DA, Bakris GL, Brook RD, Daugherty SL, Dennison-Himmelfarb CR, et al. Resistant hypertension: detection, evaluation, and management: a scientific statement from the American Heart association. *Hypertension* 2018;72:e53–90.
- [27] Alexander L, Christensen SM, Richardson L, Ingersoll AB, Burrige K, Golden A, et al. Nutrition and physical activity: an obesity medicine association (OMA) clinical practice statement 2022. *Obesity Pillars* 2022:1.
- [28] McKeown NM, Meigs JB, Liu S, Saltzman E, Wilson PW, Jacques PF. Carbohydrate nutrition, insulin resistance, and the prevalence of the metabolic syndrome in the Framingham Offspring Cohort. *Diabetes Care* 2004;27:538–46.

- [29] Pereira MA, Jacobs Jr DR, Pins JJ, Raatz SK, Gross MD, Slavin JL, et al. Effect of whole grains on insulin sensitivity in overweight hyperinsulinemic adults. *Am J Clin Nutr* 2002;75:848–55.
- [30] Expert Panel on Integrated Guidelines for Cardiovascular H, Risk Reduction in C, Adolescents, National Heart L, Blood I. Expert panel on integrated guidelines for cardiovascular health and risk reduction in children and adolescents: summary report. *Pediatrics* 2011;128(Suppl 5):S213–56.
- [31] Harada-Shiba M, Ohta T, Ohtake A, Ogura M, Dobashi K, Nohara A, et al. Guidance for pediatric familial hypercholesterolemia 2017. *J Atheroscler Thromb* 2018;25:539–53.
- [32] Barkas F, Nomikos T, Liberopoulos E, Panagiotakos D. Diet and cardiovascular disease risk among individuals with familial hypercholesterolemia: systematic review and meta-analysis. *Nutrients* 2020;12.
- [33] Vuorio A, Kuoppala J, Kovanen PT, Humphries SE, Tonstad S, Wiegman A, et al. Statins for children with familial hypercholesterolemia. *Cochrane Database Syst Rev* 2014;CD006401.
- [34] Maliachova O, Stabouli S. Familial hypercholesterolemia in children and adolescents: diagnosis and treatment. *Curr Pharm Des* 2018;24:3672–7.
- [35] Ullah R, Rauf N, Nabi G, Ullah H, Shen Y, Zhou YD, et al. Role of nutrition in the pathogenesis and prevention of non-alcoholic fatty liver disease: recent updates. *Int J Biol Sci* 2019;15:265–76.
- [36] Levy LD, Fleming JP, Klar D. Treatment of refractory obesity in severely obese adults following management of newly diagnosed attention deficit hyperactivity disorder. *Int J Obes* 2009;33:326–34.
- [37] Bowling A, Davison K, Haneuse S, Beardslee W, Miller DP. ADHD medication, dietary patterns, physical activity, and BMI in children: a longitudinal analysis of the ECLS-K study. *Obesity* 2017;25:1802–8.
- [38] O'Malley GC, Shultz SP, Thivel D, Tsiros MD. Neuromusculoskeletal health in pediatric obesity: incorporating evidence into clinical examination. *Curr Obes Rep* 2021;10:467–77.
- [39] Tsiros MD, Tian EJ, Shultz SP, Olds T, Hills AP, Duff J, et al. Obesity, the new childhood disability? An umbrella review on the association between adiposity and physical function. *Obes Rev* 2020;21:e13121.
- [40] Chung ST, Onuzuruike AU, Magge SN. Cardiometabolic risk in obese children. *Ann N Y Acad Sci* 2018;1411:166–83.
- [41] Puhl RM, Lessard LM. Weight stigma in youth: prevalence, consequences, and considerations for clinical practice. *Curr Obes Rep* 2020;9:402–11.
- [42] Sothorn M. Profile of the overweight child. In: *Safe and effective exercise for overweight youth*. CRC Press; 2014.
- [43] Bushman B. Developing the P (for progression) in a FITTVP exercise prescription. *ACSM's Health & Fit J* 2018;22(3):6–9. <https://doi.org/10.1249/fit.00000.00000.000378.2018>.
- [44] Han A, Fu A, Cogley S, Sanders RH. Effectiveness of exercise intervention on improving fundamental movement skills and motor coordination in overweight/obese children and adolescents: a systematic review. *J Sci Med Sport* 2018;21:89–102.
- [45] Institute for Healthy Childhood Weight. Algorithm for the assessment and management of childhood obesity in patients 2 years and older. Available at: https://www.paaap.org/uploads/1/2/4/3/124369935/551b74_5a52cf9033cb48b09aba3c0280a15402.pdf. [Accessed December 2021].
- [46] Barlow SE. Expert committee recommendations regarding the prevention, assessment, and treatment of child and adolescent overweight and obesity: summary report. *Pediatrics* 2007;120(Suppl 4):S164–92.
- [47] Spear BA, Barlow SE, Ervin C, Ludwig DS, Saelens BE, Schetzina KE, et al. Recommendations for treatment of child and adolescent overweight and obesity. *Pediatrics* 2007;120(Suppl 4):S254–88.
- [48] Browne NT. Food insecurity: assessment and intervention. *J Pediatr Surg Nurs* 2017;6:7–10.
- [49] Coleman-Jensen ARM, Gregory CA, Singh A. Household food security in the United States in 2020, ERR-298. U.S. Department of Agriculture, Economic Research Service; 2021.
- [50] Kral TVE, Chittams J, Moore RH. Relationship between food insecurity, child weight status, and parent-reported child eating and snacking behaviors. *J Spec Pediatr Nurs* 2017;22.
- [51] Barnidge E, LaBarge G, Krupsky K, Arthur J. Screening for food insecurity in pediatric clinical settings: opportunities and barriers. *J Community Health* 2017;42:51–7.
- [52] American Academy of Pediatrics. Promoting food security for all children. *Pediatrics* 2015;136:e1431–8.
- [53] Food assistance programs (NSLP/WIC/SNAP) participation shows reduction in FI. **Healthy people**. <https://www.healthypeople.gov/2020/topics-objectives/topic/social-determinants-health/interventions-resources/food-insecurity>. [Accessed 6 June 2020].
- [54] Cook JT, Black M, Chilton M, Cutts D, Ettinger de Cuba S, Heeren TC, et al. Are food insecurity's health impacts underestimated in the U.S. population? Marginal food security also predicts adverse health outcomes in young U.S. children and mothers. *Adv Nutr* 2013;4:51–61.
- [55] Eicher-Miller HA, Mason AC, Weaver CM, McCabe GP, Boushey CJ. Food insecurity is associated with iron deficiency anemia in US adolescents. *Am J Clin Nutr* 2009;90:1358–71.
- [56] Eisenmann JC, Gundersen C, Lohman BJ, Garasky S, Stewart SD. Is food insecurity related to overweight and obesity in children and adolescents? A summary of studies, 1995–2009. *Obes Rev* 2011;12:e73–83.
- [57] Fox CK, Cairns N, Sunni M, Turnberg GL, Gross AC. Addressing food insecurity in a pediatric weight management clinic: a pilot intervention. *J Pediatr Health Care* 2016;30:e11–5.
- [58] Gundersen C, Kreider B. Bounding the effects of food insecurity on children's health outcomes. *J Health Econ* 2009;28:971–83.
- [59] Laraia BA. Food insecurity and chronic disease. *Adv Nutr* 2013;4:203–12.
- [60] Howard LL. Transitions between food insecurity and food security predict children's social skill development during elementary school. *Br J Nutr* 2011;105:1852–60.
- [61] Gross RS, Mendelsohn AL. Food insecurity during early childhood: marker for disparities in healthy growth and development. *Pediatrics* 2019;144.
- [62] Hager ER, Quigg AM, Black MM, Coleman SM, Heeren T, Rose-Jacobs R, et al. Development and validity of a 2-item screen to identify families at risk for food insecurity. *Pediatrics* 2010;126:e26–32.
- [63] Institute of Medicine (U.S. Committee on standards for developing trustworthy clinical practice guidelines., graham R. *Clinical practice guidelines we can trust*. Washington, DC: National Academies Press; 2011.