



Published in final edited form as:

*Breast Cancer Res Treat.* 2023 August ; 201(1): 139–145. doi:10.1007/s10549-023-06968-2.

## Validating the PROMIS Cognitive Function Short Form in Cancer Survivors

Ashley M. Henneghan, PhD, RN, FAAN<sup>1</sup>, Kathleen Van Dyk, PhD<sup>2,3</sup>, Xingtao Zhou, MS<sup>4</sup>, Raeanne C. Moore, PhD<sup>5</sup>, James C. Root, PhD<sup>6</sup>, Tim A. Ahles, PhD<sup>6</sup>, Zev M. Nakamura, MD<sup>7</sup>, Jeanne Mandelblatt, MD, MPH<sup>4</sup>, Patricia A. Ganz, MD<sup>3,8</sup>

<sup>1</sup>University of Texas at Austin, School of Nursing, Austin, TX, USA

<sup>2</sup>Semel Institute of Neuroscience and Human Behavior, University of California, Los Angeles, Los Angeles, CA, USA

<sup>3</sup>Jonsson Comprehensive Cancer Center, University of California Los Angeles, Los Angeles, CA, USA

<sup>4</sup>Georgetown Lombardi Comprehensive Cancer Center Georgetown University, Washington, DC

<sup>5</sup>Department of Psychiatry, University of California San Diego, CA

<sup>6</sup>Memorial Sloan Kettering Cancer Center, New York, NY

<sup>7</sup>Department of Psychiatry, University of North Carolina–Chapel Hill, Chapel Hill, NC

<sup>8</sup>Department of Health Policy and Management, UCLA Fielding School of Public Health, University of California at Los Angeles, Los Angeles, CA

### Abstract

**Purpose:** The Patient-Reported Outcome Measurement Information System Cognitive Function Short Form 8A (PROMIS Cog) could provide a shorter, useful alternative to the often used Functional Assessment of Cancer Therapy – Cognition (FACT-Cog) in research. This study aimed to determine the convergent validity and internal reliability of the PROMIS Cog in 3 separate samples of breast cancer survivors and to explore clinical cut points.

**Methods:** Data from three samples of breast cancer survivors (BCS) were used for this secondary analysis. Convergent validity was determined by evaluating correlation strength among the derived PROMIS Cog and measures of depression, anxiety, stress, fatigue, sleep, loneliness, the FACT

---

Correspondence to KVanDyk@mednet.ucla.edu.

**Author contributions:** AMH and KVD conceived and planned the secondary analysis design, conducted the analyses, drafted and revised the manuscript and coordinated review and feedback from all co-authors; JM, PAG, and AMH were the PIs on the 3 funded studies that provided the data for this study; AMH, KVD, and XZ conducted the psychometric analyses; KVD conducted the receiver operating characteristic curve analyses; AMH, KVD, XZ, RCM, JCY, TAA, J, ZMN, JM, and PAG all reviewed and provided critical feedback the drafts and final version of the manuscript.

**Ethics approval:** All study procedures and protocols were in accordance with the Declaration of Helsinki and were approved by local Institutional Review Boards (Georgetown University IRB # 2008-363; UCLA IRB #11-001178; UT Austin IRB #2015-10-0039).

**Consent to participate:** all participants in the studies provided written informed consent to participate and publish the findings.

**Competing interests:** R.C.M. is a co-founder of KeyWise AI, Inc. and a consultant for NeuroUX. The terms of this arrangement have been reviewed and approved by UC San Diego in accordance with its conflict of interest policies. The remaining authors declare that they have no competing interests.

Cognitive Function. Clinical cut-points for the PROMIS Cog were determined by plotting the receiver operating characteristic curves.

**Results:** 3 samples of breast cancer survivors (N=471, N=132, N=90) were included. Absolute values of correlations demonstrating convergent validity ranged from 0.21 to 0.82,  $p$ 's <.001, and were comparable to correlations with the full FACT Cognitive Function 18 item perceived cognitive impairments scale. ROC curve plots indicated a clinical cut off <34 for the combined sample.

**Conclusion:** The 8-item PROMIS Cog demonstrated good convergent validity and internal reliability in breast cancer survivors, comparable to the 18-item FACT Cog PCI. The PROMIS Cog 8a is a brief self-report measure that can be easily incorporated into CRCI research designs or used in clinical settings.

### Keywords

cancer-related cognitive impairment; self-report; PROMIS cognition; convergent validity; internal reliability

---

### Introduction

Cancer-related cognitive impairment (CRCI) in memory, attention, and other abilities is distressing and burdensome, and can negatively impact cancer survivors' quality of life, social functioning and occupational functioning [1, 2]. It is generally accepted that approximately 30% of newly diagnosed cancer patients, 75% of patients undergoing adjuvant therapies, and 35% of cancer survivors experience CRCI [3].

Patient reported outcomes (PROs) assessing cognition are being recognized for their contribution in evaluation of CRCI [4], without a consensus on choice of PRO instrument. Our working group, The Cancer Neuroscience Initiative, recently suggested the use of the Patient-Reported Outcome Measurement Information System (PROMIS) Cognitive Function Short Form 8A (PROMIS Cog) as a minimum measure in all CRCI research including cognitive PROs [5], in order to facilitate cross study comparisons and meta analyses.

Presently, many researchers use the 18-item Functional Assessment of Cancer Treatment-Cognitive Function Perceived Cognitive Impairment (PCI) scale. The brief 8-item PROMIS Cog includes most of the same items as the longer PCI, since they were developed from similar item banks and common developers[6]. The PROMIS Cog may therefore function as a useful, briefer assessment tool for CRCI, but there is currently limited psychometric evidence for this instrument in oncology populations. In this study, we determined the convergent validity and internal reliability of the PROMIS Cognitive Function Short Form 8a in 3 samples of breast cancer survivors and explored clinical cut points.

### Methods

Data from three samples of breast cancer survivors were used for this secondary post-hoc analysis: 1) the Thinking and Living with Cancer ("TLC") study [7], 2) the Mind Body Study ("MBS") [8]; and 3) the Cognition After Breast Cancer ("CABC") study

[9]. All study procedures and protocols were in accordance with the Declaration of Helsinki and were approved by local Institutional Review Boards. All participants provided written informed consent prior to data collection. Data for perceived cognitive functioning (measured with the FACT Cog) and other measures were included in these analyses. Since subjective and objective measures of CRCI are commonly unrelated [4], we focused on other cognitive PRO's and measures for mood, distress, and sleep to measure convergent validity as done in other studies since they are contemporaneously conceptualized as symptom clusters [10].

### **Thinking and Living with Cancer (“TLC”) Study**

The multi-site prospective TLC study cohort has been previously described in detail [7]. In brief, study criteria included females with a new diagnosis of stage 0-III breast cancer age 60 or older. We selected the 12-month assessment timepoint for analyses (following adjuvant treatment and/or initiation of hormone therapy). Demographic and clinical information were obtained via medical record. Other measures included the Pittsburgh Sleep Quality Index (PSQI), Functional Assessment of Chronic Illness Therapy – Fatigue (FACT-F), State Trait Anxiety Inventory – State (STAI-S), and the Center for Epidemiologic Studies Depression Scale (CESD).

### **Mind Body Study (“MBS”)**

The MBS study was a prospective longitudinal study examining the effects of endocrine therapy for treating breast cancer in females, and has been extensively described elsewhere [8]. Briefly, the study included participants aged 21-65 years, who completed primary cancer treatment in the last three months for breast cancer (stage 0-III). Data from the final follow-up assessment, 3-6 years after study enrollment, was used in this study. Demographic and clinical information was obtained via self-report, and self-report measures included the FACT-Cog, Beck Depression Inventory-2<sup>nd</sup> edition (BDI-2), Multidimensional Fatigue Symptom Inventory (MFSI), the PSQI, and STAI-S.

### **Cognition After Breast Cancer (“CABC”) Study**

All CABC study procedures have also been previously described [9]. Females between the aged 21 to 65 years with a history of breast cancer (stage 0-III) treated with chemotherapy within the past 10 years were enrolled. Data included demographic and clinical characteristics, the FACT-Cog, the PROMIS version 1.0 Emotional Distress-Anxiety short form 8a (PROMIS Anxiety) and Depression- short form 8a (PROMIS Depressive), Fatigue-short form 8a (PROMIS Fatigue), the Perceived Stress Scale (PSS), the PSQI, and the UCLA Loneliness Scale revised version 3 (UCLA Loneliness).

### **PROMIS Cog Scores**

For this analysis, the PROMIS Cog scores were derived from the FACT-Cog, which includes seven of the same items as the PROMIS Cog (CogA1; CogA3, CogC7, CogF23, CogC31, CogC32, CogMT2). See Table 1. We recoded the seven items to be congruent with the PROMIS Cog scale and have meaningful interpretation of the derived clinical cut points. Since 1 item of the PROMIS Cog is not included in the FACT-Cog, we created one

additional item using an average score across the 7 FACT Cog items to re-create the 8-item total score for the PROMIS Cog. The 8 items were summed, and total raw scores used in the analyses. This method of mean substitution was used based on the instructions for scoring FACT instruments[11] and is common practice to handle missing data in clinical measures[12]. If participants were missing more than two of the seven items, they were excluded from analyses.

## Data Analyses

Frequencies, means, and standard deviations were used to describe basic characteristics for the three samples. Comparative convergent validity was determined by evaluating correlations size (using a reference of 0.1 to 0.3 as small/weak correlation, 0.4 to 0.6 as a moderate correlation, and .7 or higher as a strong correlation)[13] between the derived PROMIS Cog and FACT-Cog PCI with measures others have used to demonstrate the validity of the FACT-Cog PCI [10] including depression, anxiety, stress, fatigue, sleep quality, loneliness and the FACT-Cog Cognitive Abilities, Quality of Life, and Cognitive Comments from Others subscales. Internal reliability was determined using the Cronbach alpha for total scales and with each item deleted. A threshold of 0.8 or greater was used for Cronbach alpha [14]. In order to identify comparable clinical cut-points for the PROMIS Cog, we plotted the receiver operating characteristic (ROC) curves of the PROMIS Cog using the published [15] and recently validated [16] FACT-Cog PCI 18 cut-point score of <54.

## Results

The TLC sample included 471 breast cancer survivors (average 1.3 years from diagnosis), the MBS sample included 132 breast cancer survivors (average 5 years from diagnosis), and the CABC study included 90 breast cancer survivors (on average 3.5 years since diagnosis) See Table 2.

The Cronbach's alphas for the PROMIS Cog ranged from 0.89 (TLC) to 0.91 (MBS) to 0.97 (CABC). Internal consistencies were also calculated for the PROMIS Cog with each item removed and ranged from 0.963 to 0.972 for the CABC study, 0.939 to .849 for the MBS study, and 0.913 to 0.889 for the TLC study (see Supplementary Materials Table 1). The Absolute values for the correlations among PROMIS Cog and convergent validity measures ranged from 0.21 to 0.82 ( $p'f < .001$ , see Table 3) across studies. ROC curve plots indicated a clinical cut-off <34 for the PROMIS Cog and area under the curves (AUC's) ranged from 0.96—0.98, indicating outstanding discrimination[17]. Sensitivity ranged from 0.89—0.98, and specificity ranged from 0.79—0.89, see Table 4. ROC curve plots were also calculated for the FACT Cog PCI 20, as some researchers use the 20 item subscale rather than the 18 item subscale (see Supplementary Materials Table 2).

## Discussion

This study provides evidence for strong internal consistency of the 8-item PROMIS Cog, similar comparative convergent validity as the FACT-Cog PCI with measures of depression, anxiety, stress, fatigue, sleep, loneliness, and a preliminary clinical cut point (<34).

Correlations between PROMIS Cog and other cognitive PRO's as well as measures of depression, anxiety, fatigue, stress, and sleep quality were consistent with previous reports in cancer survivors [4] and consistent with recent validity analyses for the FACT-Cog [10]. We found stronger correlations among these variables in the CABC and MBS samples than the TLC study which may be attributed differences in age (CABC and MBS were younger), differences in employment status (more were employed in the CABC and MBS samples), and/or differences in treatment histories—more than half of the samples in CABC and MBS were treated with chemotherapy compared to a quarter of the TLC study. Chemotherapy history, and employment status have been linked to poorer self-reported CRCI [4]. Younger age and employment status are also risk factors for greater psychosocial distress, and fatigue in BCS [18-20].

AUCs, sensitivity and specificity scores indicated using a cut-off of 34 for the PROMIS Cog. We used another cognitive PRO, the FACT-Cog PCI, to determine the clinical cut-off for the PROMIS Cog, which is common practice[15, 21, 22], however cut-points may be different if standardized cognitive tests or other functional measures are used, and should be further studied for clinical utility.

Study limitations should be considered. The PROMIS Cog scores were derived from seven overlapping items of the FACT-Cog PCI subscale and 1 mean substitution item, not the PROMIS Cog short form 8a itself, resulting in raw summed scores being used in the analyses. Results should be replicated with the full instrument and converted to T scores with standard errors (as recommended by the PROMIS scoring manual). The three samples used in these analyses only included breast cancer survivors after treatment, who were highly educated, limiting generalizability. Replication and validation of the published PROMIS Cog 8a are needed in other socio-demographically diverse cancer populations, including earlier and/or later in the cancer treatment trajectory age groups, and primary cancer diagnoses.

## Conclusion

These results provide preliminary psychometric evidence for using the PROMIS Cog and support recommendations to use the PROMIS Cog in CRCI research [5]. The PROMIS Cog 8a is a brief self-report measure that can be easily incorporated into CRCI research and clinical settings. These findings also offer a starting point for additional psychometric evaluation (e.g., sensitivity to change over time, test-retest reliability) and for determining classification methods for cognitive PROs in CRCI research. Using the short PROMIS Cog should enhance the ease of routine collection of cognitive data in clinical trials, geriatric assessments, and observational studies of broad populations of cancer survivors.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Funding:

Research reported in this publication was supported by the National Institute of Nursing Research of the National Institutes of Health under Award Numbers K01NR018970; F31NR015707 (AMH), R56AG068086 (JC, JM), R35CA197289 (JM), R01AG068193 (JM), the National Cancer Institute awards R01 CA 109650 (PAG), K08CA241337 (KVD), R01CA129769 (JM), the Loan Repayment Award (KVD), the American Cancer Society Award Number DSCN-15-072 (AMH), and the Breast Cancer Research Foundation (to PAG)The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the American Cancer Society.

## Data Availability:

The datasets analyzed in the current study are available from the corresponding author on reasonable request.

## List of Abbreviations

<b>AUC</b>	area under the curves
<b>BDI-2</b>	Beck Depression Inventory-2 <sup>nd</sup> edition
<b>CABC</b>	Cognition After Breast Cancer
<b>CESD</b>	Center for Epidemiologic Studies Depression Scale
<b>CRCI</b>	Cancer-related cognitive impairment
<b>FACT-Cog</b>	Functional Assessment of Cancer Therapy – Cognition
<b>FACT-F</b>	Functional Assessment of Chronic Illness Therapy – Fatigue
<b>MFSI</b>	Multidimensional Fatigue Symptom Inventory
<b>PCI</b>	Perceived Cognitive Impairment
<b>PSQI</b>	Pittsburgh Sleep Quality Index
<b>PROs</b>	Patient reported outcomes
<b>PROMIS Anxiety</b>	PROMIS version 1.0 Emotional Distress- Anxiety short form 8a
<b>PROMIS Cog</b>	Patient-Reported Outcome Measurement Information System Cognitive Function Short Form 8a
<b>PROMIS Depressive</b>	PROMIS version 1.0 Emotional Distress- Depressive short form 8a
<b>PROMIS Fatigue</b>	PROMIS version 1.0 Fatigue short form 8a
<b>ROC</b>	receiver operating characteristic
<b>STAI-S</b>	State Trait Anxiety Inventory – State

TLC Thinking and Living with Cancer

## References

1. Mayo SJ, Lustberg M, H MD, Nakamura ZM, Allen DH, Von Ah D, et al. Cancer-related cognitive impairment in patients with non-central nervous system malignancies: an overview for oncology providers from the MASCC Neurological Complications Study Group. *Support Care Cancer*. 2020. doi: 10.1007/s00520-020-05860-9.
2. Lange M, Joly F, Vardy J, Ahles T, Dubois M, Tron L, et al. Cancer-related cognitive impairment: an update on state of the art, detection, and management strategies in cancer survivors. *Ann Oncol*. 2019;30(12):1925–40. doi: 10.1093/annonc/mdz410. [PubMed: 31617564]
3. Janelsins MC, Kesler SR, Ahles TA, Morrow GR. Prevalence, mechanisms, and management of cancer-related cognitive impairment. *International review of psychiatry*. 2014;26(1):102–13. doi: 10.3109/09540261.2013.864260. [PubMed: 24716504]
4. Bray VJ, Dhillon HM, Vardy JL. Systematic review of self-reported cognitive function in cancer patients following chemotherapy treatment. *J Cancer Surviv*. 2018. doi: 10.1007/s11764-018-0692-x.
5. Henneghan AM, Van Dyk K, Kaufmann T, Harrison R, Gibbons C, Heijnen C, et al. Measuring Self-Reported Cancer-Related Cognitive Impairment: Recommendations from the Cancer Neuroscience Initiative Working Group. *JNCI: Journal of the National Cancer Institute*. 2021. doi: 10.1093/jnci/djab027.
6. Lai JS, Wagner LI, Jacobsen PB, Cella D. Self-reported cognitive concerns and abilities: two sides of one coin? *Psychooncology*. 2014;23(10):1133–41. doi: 10.1002/pon.3522. [PubMed: 24700645]
7. Mandelblatt JS, Small BJ, Luta G, Hurria A, Jim H, McDonald BC, et al. Cancer-Related Cognitive Outcomes Among Older Breast Cancer Survivors in the Thinking and Living With Cancer Study. *J Clin Oncol*. 2018;36(32):Jco1800140. doi: 10.1200/jco.18.00140. [PubMed: 30281396]
8. Van Dyk K, Crespi CM, Bower JE, Castellon SA, Petersen L, Ganz PA. The cognitive effects of endocrine therapy in survivors of breast cancer: A prospective longitudinal study up to 6 years after treatment. *Cancer*. 2019;125(5):681–9. doi: 10.1002/cncr.31858. [PubMed: 30485399]
9. Henneghan A, Stuifbergen A, Becker H, Kesler S, King E. Modifiable correlates of perceived cognitive function in breast cancer survivors up to 10 years after chemotherapy completion. *J Cancer Surviv*. 2018;12(2):224–33. doi: 10.1007/s11764-017-0661-9. [PubMed: 29116555]
10. Koch V, Wagner LI, Green HJ. Assessing neurocognitive symptoms in cancer patients and controls: psychometric properties of the FACT-Cog3. *Current Psychology*. 2021. doi: 10.1007/s12144-021-02088-6.
11. FACT-Cog Functional Assessment of Cancer Therapy - Cognitive Function. <https://www.facit.org/measures/FACT-Cog> Accessed 2022.
12. Downey RG, King C. Missing data in Likert ratings: A comparison of replacement methods. *J Gen Psychol*. 1998;125(2):175–91. doi: 10.1080/00221309809595542. [PubMed: 9935342]
13. Cohen J. *Statistical power analysis for the behavioral sciences* / Jacob Cohen. 2nd ed. ed. Hillsdale, N.J.: L. Erlbaum Associates; 1988.
14. Tavakol M, Dennick R. Making sense of Cronbach's alpha. *Int J Med Educ*. 2011;2:53–5. doi: 10.5116/ijme.4dfb.8dfd. [PubMed: 28029643]
15. Dyk KV, Crespi CM, Petersen L, Ganz PA. Identifying Cancer-Related Cognitive Impairment Using the FACT-Cog Perceived Cognitive Impairment. *JNCI Cancer Spectr*. 2020;4(1):pkz099. doi: 10.1093/jncics/pkz099. [PubMed: 32064458]
16. Fardell JE, Bray V, Bell ML, Rabe B, Dhillon H, Vardy JL. Screening for cognitive symptoms among cancer patients during chemotherapy: Sensitivity and specificity of a single item self-report cognitive change score. *Psychooncology*. 2022. doi: 10.1002/pon.5928.
17. Mandrekar JN. Receiver operating characteristic curve in diagnostic test assessment. *J Thorac Oncol*. 2010;5(9):1315–6. doi: 10.1097/JTO.0b013e3181ec173d. [PubMed: 20736804]
18. Syrowatka A, Motulsky A, Kurteva S, Hanley JA, Dixon WG, Meguerditchian AN, et al. Predictors of distress in female breast cancer survivors: a systematic review. *Breast Cancer Res Treat*. 2017;165(2):229–45. doi: 10.1007/s10549-017-4290-9. [PubMed: 28553684]

19. Joly F, Lange M, Dos Santos M, Vaz-Luis I, Di Meglio A. Long-Term Fatigue and Cognitive Disorders in Breast Cancer Survivors. *Cancers (Basel)*. 2019;11(12). doi: 10.3390/cancers11121896.
20. Islam T, Dahlui M, Majid HA, Nahar AM, Mohd Taib NA, Su TT, et al. Factors associated with return to work of breast cancer survivors: a systematic review. *BMC Public Health*. 2014;14 Suppl 3:S8. doi: 10.1186/1471-2458-14-S3-S8.
21. Cheung YT, Foo YL, Shwe M, Tan YP, Fan G, Yong WS, et al. Minimal clinically important difference (MCID) for the functional assessment of cancer therapy: cognitive function (FACT-Cog) in breast cancer patients. *Journal of clinical epidemiology*. 2014;67(7):811–20. doi: 10.1016/j.jclinepi.2013.12.011. [PubMed: 24656406]
22. Bell ML, Dhillon HM, Bray VJ, Vardy JL. Important differences and meaningful changes for the Functional Assessment of Cancer Therapy-Cognitive Function (FACT-Cog). *Journal of Patient-Reported Outcomes*. 2018;2(1):48. doi: 10.1186/s41687-018-0071-4.



**Table 1.**

Comparison of PROMIS Short Form 8a to FACT-Cog

<b>PROMIS Short Form 8a Items</b>	<b>Approximate FACT-Cog Items</b>
1. I have had trouble forming thoughts.	CogA1
2. My thinking has been slow.	CogA3
3. I have had trouble concentrating.	CogC7
4. I have had to work really hard to pay attention or I would make a mistake.	CogF23
5. I have had to work harder than usual to keep track of what I was doing.	CogC31
6. It has seemed like my brain was not working as well as usual (maps to "My thinking has been slower than usual." From the FACT-Cog PCI, however not exact match).	CogC32
7. I have had trouble shifting back and forth between different activities that require thinking.	CogMT2
8. I have had trouble adding or subtracting numbers in my head.	No PCI item, mean substitution used to create item

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 2.**

Demographic and Clinical Characteristics of the Samples (All Female Participants)

	<b>Thinking and Living with Cancer Study (N=471)</b>	<b>Mind Body Study (N=132)</b>	<b>Cognition After Breast Cancer Study (N=90)</b>
Age in years Mean (SD) Range	69.6 (5.7) 60-92	56.4 (8.0) 36-69	48.7 (9.0) 24-65
Years of Education	<16: 199 (42%) <sup>I</sup> 16: 112 (24%) >16: 159 (34%)	<16 26 (20%) 16 42 (32%) >16 63 (48%)	<16 18 (20%) 16 34 (38%) >16 35 (39%)
Identified as Minority/non-white	83 (18%)	27 (21%)	15 (17%)
Employed At Time of Study	143 (30%) <sup>I</sup>	85 (64%)	78 (87%)
Months Since Diagnosis Mean (SD) Range	15.7 (2.67) 6.0-33.0	58.0 (8.1) 41.8-82.6	42.0 (32.4) 6.8 to 120
Breast Cancer Stage			
0	79 (17%)	19 (15%)	
I	277 (60%)	58 (44%)	19 (21%)
II	88 (19%)	42 (32%)	51 (57%)
III	21 (5%)	12 (9%)	19 (21%)
Frequency of Chemotherapy	117 (25%)	71 (54%)	90 (100%)
Frequency of Radiation	287 (61%)	98 (75%)	59 (66%)
Frequency of Hormone Therapy	371 (80%)	96 (73%)	71 (79%)

<sup>I</sup> data available for n=470

Abbreviations: SD= standard deviation

**Table 3.** Convergent validity analyses comparing correlations between FACT-Cog Perceived Cognitive Impairments 18 item subscale and PROMIS Cog 8a

Construct	Thinking and Living with Cancer Study (N=471)		Mind Body Study (N=132)		Cognition After Breast Cancer Study (N=90)	
	FACT-Cog PCI-18	PROMIS Cog 8a	FACT-Cog PCI-18	PROMIS Cog 8a	FACT-Cog PCI-18	PROMIS Cog 8a
Depression <i>r</i>	-0.338**	-0.378**	-.646**	-.643**	-.55**	-.56**
(specific measure)	(CES-D)		(BDI-II)		(PROMIS-Emotional Distress-Depressive Short Form 8a)	
Anxiety <i>r</i>	-0.260**	-0.290**	-.455**	-.467**	-.52**	-.51**
(specific measure)	(STAI State-Anxiety subscale)		(STAI State-Anxiety subscale)		(PROMIS-Emotional Distress-Anxiety Short Form 8a)	
Stress <i>r</i>	-	-	-.534**	-.516**	-.66**	-.67**
(specific measure)	-		(PSS)		(PSS)	
Fatigue <i>r</i>	0.342**	0.366**	.542**	.559**	-.62**	-.62**
(specific measure)	(FACT-F)		(MFSI-Vigor subscale)		(PROMIS-Fatigue Short Form 8a)	
Sleep <i>r</i>	-0.205**	-0.215**	-.322**	-.315**	-.51**	-.48**
(specific measure)	(PSQI)		(PSQI)		(PSQI)	
Loneliness	-	-	-	-	-.51**	-.51**
(specific measure)	-		-		(UCLA Loneliness v 3)	
Cognitive Abilities	0.619**	0.596**	.746**	.690**	.82**	.82**
(specific measure)	(FACT-Cog PCA subscale)		(FACT-Cog PCA subscale)		(FACT-Cog PCA subscale)	
Quality of Life	0.554**	0.569**	.731**	.749**	.78**	.77**
(specific measure)	(FACT-Cog QOL subscale)		(FACT-Cog QOL subscale)		(FACT-Cog QOL subscale)	
Comments from Others about Cognition	0.340**	0.362**	.617**	.649**	.62**	.63**
(specific measure)	(FACT-Cog CO subscale)		(FACT-Cog CO subscale)		(FACT-Cog CO subscale)	
Cognitive Complaints: Squire Memory Questionnaire			.659**	.562**		
Cognitive Complaints: Patient's Assessment of Own Functioning Inventory Total			.710**	.665**		

Abbreviations: **BDI-II**: Beck Depression Inventory-II ; **CES-D**: Center for Epidemiologic Studies Depression Scale; **CO**: Comments from Others; **FACT-F**: Functional Assessment of Chronic Illness Therapy Fatigue; **FACT Cog**: Functional Assessment of Cancer Therapy - Cognitive Function; **MFSI**: Multidimensional Fatigue Symptom Inventory; **PCA**: Perceived Cognitive Abilities; **PCI**: Perceived

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Cognitive Impairments; **PROMIS**: Patient-Reported Outcomes Measurement Information System; **PSS**: Perceived Stress Scale; **PSQI**: Pittsburgh Sleep Quality Index; **STAI**: State-Trait Anxiety Inventory;  
QOL: Quality of Life

\*\*\*  
p < .001

**Table 4.**

Descriptive statistics, area under the curve for the receiver operating characteristics for the PROMIS Cog based on the FACT Cog PCI 18 (<54) clinical cut offs for 3 samples

	Thinking and Living with Cancer Study (N=471)	Mind Body Study (N=132)	Cognition After Breast Cancer Study (N=90)	Combined Samples
Raw total mean (SD), range	34.94 (5.77) 13.33-40	34.03 (6.13) 16-40	26.85 (9.96) 8-40	33.72 (7.04) 8-40
T-score mean (SD), range <sup>^</sup>	52.85 (8.36) 31.18-63.48	51.14 (8.03) 33.92-63.48	43.82 (11.10) 22.41-63.48	51.35 (9.18) 22.41-63.48
PCI-18 Cut Off	<34	<34	<34	<34
PCI-18 AUC	0.977	0.966	0.972	0.963
PCI-18 Sensitivity	0.928	0.921	0.982	.925
PCI-18 Specificity	0.892	0.844	0.794	.851

Abbreviations: **AUC**: area under the curve; **CABC**: Cognition After Breast Cancer; **MBS**: Mind Body Study; **PCI**: Perceived Cognitive Impairments; **ROC**: receiver operating characteristic; **TLC**: Thinking and Living with Cancer

<sup>^</sup> Truncated raw scores converted to T scores using PROMIS conversion table for the PROMIS Cog 8a